

Antimicrobial activity of iron-depriving pyoverdines against human opportunistic pathogens

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

[About eLife's process](#)

Reviewed preprint version 1

January 8, 2024 (this version)

Posted to preprint server

September 21, 2023

Sent for peer review

September 20, 2023

Vera Vollenweider, Karoline Rehm, Clara Chepkirui, Manuela Pérez-Berlanga, Magdalini Polymenidou, Jörn Piel, Laurent Bigler, Rolf Kümmerli 

Department of Quantitative Biomedicine, University of Zurich, Zurich, Switzerland • Department of Chemistry, University of Zurich, Zurich, Switzerland • Institute of Microbiology, Eidgenössische Technische Hochschule (ETH) Zurich, Zurich, Switzerland

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

The global rise of antibiotic resistance calls for new drugs against bacterial pathogens. A common approach is to search for natural compounds deployed by microbes to inhibit competitors. Here we show that the iron chelating pyoverdines, siderophores produced by environmental *Pseudomonas* spp., have strong antibacterial properties by inducing iron starvation and growth arrest in pathogens. A screen of 320 natural *Pseudomonas* isolates used against 12 human pathogens uncovered several pyoverdines with particularly high antibacterial properties and distinct chemical characteristics. The most potent pyoverdine effectively reduced growth of the pathogens *Acinetobacter baumannii*, *Klebsiella pneumoniae* and *Staphylococcus aureus* in a concentration- and iron-dependent manner. Pyoverdine increased survival of infected *Galleria mellonella* host larvae, and showed low toxicity for the host, mammalian cell lines, and erythrocytes. Furthermore, experimental evolution combined with whole-genome sequencing revealed reduced potentials for resistance evolution compared to an antibiotic. Thus, pyoverdines from environmental strains could become new sustainable antibacterials against human pathogens.

eLife assessment

This **important** study furthers our understanding of the antimicrobial properties of siderophores, and their potential use to battle opportunistic pathogens. The evidence supporting the conclusion is **solid**, based on rigorous biochemical, growth, and virulence assays. The work would benefit from a more in-depth discussion of the consequences and efficacy of 'siderophore therapy' in more complex communities/environments. The work will be of broad interest to colleagues in the fields of evolutionary ecology, microbiology, and medical sciences.

Introduction

There are approximately 4.95 million fatalities associated with antibiotic-resistant bacteria worldwide each year¹. This high mortality rate contrasts with the low development rate of new antibacterial agents for clinical applications^{2,3}. Thus, the development of alternative therapies with higher sustainability to counteract the rapidly dwindling treatment options due to resistance evolution is imperative^{4–6}. Numerous alternative approaches have been proposed including phage therapy, antimicrobial peptides and nanobodies, anti-virulence treatments to disarm pathogens, treatments that capitalize on ecological competition between susceptible and resistant bacteria, and the search for new antibacterial compounds produced by environmental microbes^{7–12}.

Mining natural microbial communities has become a promising endeavour in the hunt for novel antimicrobials because most microbes deploy bioactive compounds to contend with their competitors in the diverse assemblies they live in¹³. Competition often involves the secretion of secondary metabolites with specific antimicrobial properties against other microbes¹⁴. The isolation, characterization and synthesis of such secondary metabolites can result in novel prospective antibiotics for clinical applications^{15,16}. Secondary metabolites that act as toxins to kill competitors are often considered promising candidates. However, microbes compete through a variety of mechanisms other than toxins. For example, there is increasing evidence that competition for iron is a main determinant of species interactions in soil and freshwater communities^{17,18}. Competition for iron involves the secretion of siderophores, a class of secondary metabolites that scavenges environmental iron with high affinity¹⁹. Because siderophores are often species specific, they have two opposing effects: they make iron available for related community members possessing the matching receptor but withhold iron from competitors with non-matching receptors. Thus, siderophores can be competitive agents in inter-species interactions^{20–23}.

Here, we apply the concept of siderophore-mediated iron competition observed in natural communities to human opportunistic pathogens. Specifically, we investigate whether siderophores from non-pathogenic environmental bacteria can induce iron starvation and growth arrest in human pathogens. We focus on pyoverdines, a class of siderophores with high iron affinity that are produced and secreted by fluorescent *Pseudomonas* spp. under iron-limited conditions²⁴. Upon complexation with ferric iron, the iron-loaded pyoverdine is imported into the cell by receptors with high specificity and subsequently reduced to the bio-available ferrous iron^{25,26}. Pyoverdines show an extraordinary structural diversity with more than 70 described variants that differ in their peptide backbone^{27,28}, and have an extremely high affinity for iron ($K_a = 10^{32} \text{ M}^{-1}$, pyoverdine produced by *Pseudomonas fluorescens* biotype B)²⁹. For these reasons, we propose that pyoverdines from non-pathogenic *Pseudomonas* spp. could be potent agents to reduce the growth of opportunistic human pathogens by intensifying iron competition.

To test our hypothesis, we screened pyoverdines from a library of 320 environmental *Pseudomonas* strains, isolated from soil and freshwater habitats, for their activity against 12 strains of human opportunistic bacteria. For the most promising pyoverdine candidates, we elucidated the chemical structure to assess diversity and to understand the chemical properties important for iron competition. Subsequently, we assessed the efficacy of the three top pyoverdine candidates against four human opportunistic pathogens (*Acinetobacter baumannii*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*) *in vitro* and/or *in vivo* via infections of the Greater wax moth larvae (*Galleria mellonella*). We further assessed their toxicity in the host, towards two mammalian cell lines and red blood cells. Finally, we used experimental evolution combined with whole-genome sequencing to study the potential of pathogens evolving resistance against pyoverdine treatment.

Results

Pyoverdines can inhibit the growth of human opportunistic pathogens

To assess the antibacterial properties of pyoverdines, we performed a growth-inhibition screen using a well-characterized collection of 320 natural *Pseudomonas* isolates^{17,30–32}. The isolates stem from freshwater and soil communities and belong to the group of fluorescent pseudomonads (e.g. *P. fluorescens*, *P. putida*, *P. syringae*, *P. chlororaphis*) that are non-pathogenic to humans and can produce and secrete pyoverdines, a group of high iron-affinity siderophores³³.

In a first screen, we evaluated how the sterile pyoverdine-containing supernatants of the 320 isolates affect the growth of 12 human opportunistic pathogens (**Table 1**). We found high variation in the supernatant effect, ranging from complete pathogen growth inhibition to growth promotion compared to the mock control (**Figure 1A** and Figure S1A). The mean supernatant effect across all pathogens showed a bimodal distribution with two peaks at 0.47 – 0.58 (approximately 50% growth inhibition) and 1.00 – 1.10 (no growth inhibition) (Figure S1B). Our observations are consistent with previous studies, in which supernatants containing siderophores can have both growth inhibitory and stimulatory effects^{21,30}. For each supernatant, we quantified the number of pathogens that experienced a growth reduction of at least 20% compared to the mock control and found that 25 supernatants inhibited all 12 pathogens (**Figure 1A** and S1C).

To confirm that the growth-inhibiting properties of the 25 most potent supernatants are indeed caused by pyoverdines and not by other agents in the supernatant, we repeated the above assay, but this time supplemented the supernatant with 40 μM FeCl_3 . Under such iron-replete conditions, pyoverdines should not be able to substantially sequester environmental iron, and pathogen growth should therefore be restored. We identified seven top candidate supernatants for which a relative growth recovery of at least 0.2 occurred for all 12 pathogens (**Figure 1B**). Among the remaining 18 supernatants, the overall rate of growth recovery was also high, indicating that pyoverdines are responsible for pathogen growth inhibition under iron-limited conditions in most cases.

Growth-inhibiting pyoverdines show distinct chemical features

In previous studies, we developed a new method for pyoverdine structure elucidation from low-volume crude extracts using ultra-high-performance liquid chromatography high-resolution tandem mass spectrometry (UHPLC-HR-MS/MS)^{28,34}. These studies included the pyoverdines extracted from the seven top candidate supernatants and the analysis therein yielded five structurally different pyoverdines (**Figure 2A**). We found one novel pyoverdine type (1, from isolate 3A06) and four pyoverdine types already known from the literature (2–5). The pyoverdine from isolate 3G07 can occur in either a linear or cyclic conformation (**2a** and **2b**). Three isolates (s3b09, s3b10, s3b12), all originating from the same soil sample, had an identical pyoverdine type (**3**).

Here, we investigated whether the chemical properties of growth-inhibitory pyoverdines differ from pyoverdines that do not inhibit pathogen growth. To this end, we further elucidated the chemical structure of a set of six pyoverdines extracted from supernatants that inhibited none of the 12 pathogens, and one additional pyoverdine inhibiting the growth of 10 pathogens²⁸. We found that inhibitory pyoverdines had significantly larger collision-cross sections (CCS in $\text{\AA}^2$) than non-inhibitory ones ($F_{1,22} = 10.63$, $p = 0.0036$, measured by ion mobility spectrometry (IMS)) (**Figure 2B**). As CCS values correlate with the volume of a molecule, our results suggest that inhibitory pyoverdines are larger and have more complex molecule structures than non-inhibitory pyoverdines. In contrast, there was no significant difference between the two classes of




Figure 1.

Effect of supernatants from environmental *Pseudomonas* isolates on the growth of 12 human opportunistic pathogens.

(A) Screen to assess the extent to which pyoverdine-containing supernatants from 320 natural *Pseudomonas* isolates inhibit the growth of 12 human opportunistic pathogens. The heatmap depicts relative pathogen growth in the supernatant treatment relative to the mock control, with values ranging from stimulation (yellow) to inhibition (blue) based on four independent replicates. Grey bars above the heatmap show the number of pathogens that were at least 20% inhibited in their growth by a given supernatant. The screen returned 25 supernatant candidates that inhibited the growth of all pathogens. (B) Control screen with the 25 supernatant candidates to check whether pyoverdine causes the observed growth inhibition. The heatmap depicts the level of growth recovery in the 12 pathogens when iron was added to the supernatant. Growth recovery in iron-rich medium indicates that pyoverdines are involved in growth inhibition in iron-limited medium. Grey bars above the heatmap show the number of pathogens that experienced a relative growth recovery of at least 0.2. The screen returned 7 supernatant top candidates for which growth recovery occurred for all pathogens under iron-rich conditions.

Species and strain name	Class	Gram-staining	Frequency of inhibiting supernatants
<i>Acinetobacter johnsonii</i>	Gammaproteobacteria	negative	75.0
<i>Acinetobacter junii</i>	Gammaproteobacteria	negative	73.1
<i>Burkholderia cenocepacia</i> H111	Betaproteobacteria	negative	52.2
<i>Burkholderia cenocepacia</i> K56-2	Betaproteobacteria	negative	54.7
<i>Cronobacter sakazakii</i>	Gammaproteobacteria	negative	48.8
<i>Escherichia coli</i> K12	Gammaproteobacteria	negative	31.6
<i>Klebsiella michiganensis</i>	Gammaproteobacteria	negative	30.9
<i>Pseudomonas aeruginosa</i> PA14	Gammaproteobacteria	negative	9.7
<i>Pseudomonas aeruginosa</i> PAO1	Gammaproteobacteria	negative	20.9
<i>Shigella</i> sp.	Gammaproteobacteria	negative	57.8
<i>Staphylococcus aureus</i> Cowan	Bacilli	positive	47.8
<i>Staphylococcus aureus</i> JE2	Bacilli	positive	41.6

Table 1

Species and strains of human opportunistic pathogens used for the pyoverdine growth inhibition assay.




Figure 2.

Chemical structure of growth-inhibitory pyoverdines and their properties compared to non-inhibitory pyoverdines.

(A) The chemical structures of the seven top candidate pyoverdines were elucidated using UHPLC-HR-MS/MS and revealed 5 unique pyoverdine structures (1-5) differing in their peptide backbone (see²⁸ for an in-depth chemical analysis). Pyoverdine 1 is a novel structure (from isolate 3A06). Pyoverdine 2 can occur in either a linear 2a or a cyclic 2b form (from isolate 3G07). Pyoverdine 3 was found in three different isolates, originating from the same soil sample (from isolates s3b09, s3b10 and s3b12). Pyoverdine 4 and 5 are from isolate s3c13 and s3e20, respectively. (B) The CCS values of inhibitory pyoverdines are higher than for non-inhibitory pyoverdines. (C) Molecule polarity, measured as the chromatographic retention time, is higher for ferri-pyoverdines (iron loaded) than for apo-pyoverdines (iron free). (D) Iron complex stability, assessed by the collision induced unfolding of ferri-pyoverdines, was significantly higher for inhibitory than non-inhibitory pyoverdines. (E) The normalised collision energy (CE50) necessary to fragment 50% of ferri-pyoverdines was not different between inhibitory and non-inhibitory pyoverdines. Box plots show the median and the first and third quartiles across the 7 inhibitory and 6 non-inhibitory pyoverdines. Whiskers represent the 1.5x interquartile range.

pyoverdines regarding molecule polarity ($F_{1,22} = 2.00$, $p = 0.1711$, assessed by the chromatographic retention time on a reverse phase column) (Figure 2C), but molecule polarity was significantly higher for ferri- than for apo-pyoverdines ($F_{1,22} = 12.78$, $p = 0.0017$). Finally, we assessed the stability of ferri-pyoverdines to derive a proxy for iron affinity. We first measured the propensity of ferri-pyoverdines to undergo collision-induced unfolding (CIU) by IMS measurements at different $\Delta 6$ voltages and found that inhibitory pyoverdines are significantly more stable than non-inhibitory pyoverdines ($F_{1,11} = 7.55$, $p = 0.019$) (Figure 2D). The same trend was observed for our second measure of stability (CE50, normalised collision energy necessary to fragment 50% of ferri-pyoverdines), but in this case, the difference between the two pyoverdine classes was not significant ($F_{1,11} = 1.87$, $p = 0.198$) (Figure 2E). Taken together, the chemical analyses suggest that inhibitory pyoverdines are larger and have a higher affinity for iron than non-inhibitory pyoverdines.

Pyoverdines inhibit human pathogens in a concentration-dependent manner

We purified three out of the five most potent pyoverdines and tested their efficacy against the four human pathogens *A. baumannii*, *K. pneumoniae*, *P. aeruginosa* and *S. aureus*. For these experiments, we decided to include the most common (s3b09), the novel (3A06), and the cyclical-linear (3G07) pyoverdines. We crude-purified all three pyoverdines using established protocols^{17,35} and further purified the most common pyoverdine (s3b09) using preparative and analytical HPLC. During HPLC purification, we observed that ferribactin (the final pyoverdine precursor) was also highly abundant in the supernatant^{36,37}. Due to its compromised iron-binding capacity, we used it as a negative control for which no activity against pathogens is expected³⁷. As positive treatment control, we used the antibiotic ciprofloxacin. We subjected the four pathogens to the three pyoverdine variants, ferribactin, and ciprofloxacin across a concentration gradient and measured their effects on pathogen growth (Figure 3, Figure S2).

For *A. baumannii* and *S. aureus*, we observed classical dose-response curves in all pyoverdine assays, characterized by a decrease in bacterial growth at low and intermediate concentrations, followed by a complete halt of bacterial growth at higher pyoverdine concentrations. There was a good qualitative match between the dose-response curves obtained with the HPLC vs. the crude-purified pyoverdines. Note that pyoverdine concentration is unknown for the crude extracts and is thus expressed as relative concentrations. In contrast, the pyoverdine concentration is absolute for the HPLC-purified version, allowing the determination of an IC50 value for pyoverdine s3b09, which is 5.037 $\mu\text{g/mL}$ and 12.163 $\mu\text{g/mL}$ for *A. baumannii* and *S. aureus*, respectively.

For *K. pneumoniae*, the pyoverdines were less effective in reducing pathogen growth. The dose-response curves for the pyoverdines s3b09 (HPLC and crude) and 3A06 showed similar trajectories and the model fits showed a growth reduction between 9-12%. The strongest inhibitory effects were found for pyoverdine 3G07 with an estimated 54% growth reduction. One reason why pyoverdines might be less effective against *K. pneumoniae* is that this species produces enterobactin, a siderophore that has higher iron affinity than pyoverdine³⁸.

For *P. aeruginosa*, we observed that all pyoverdines were growth inhibitory at intermediate concentrations. The initial inhibition converted into growth promotions at high pyoverdine concentrations in three out of four cases. The only exception was pyoverdine 3G07, which yielded consistent growth inhibitions of about 66%. One reason for these non-standard dose-response curves could be that *P. aeruginosa* possesses two receptors (FpvA and FpvB) for pyoverdine uptake³⁹. It might thus capitalize on the supplemented pyoverdines, especially at high concentrations via the up-regulation of the heterologous promiscuous FpvB receptor^{40,41}. Our findings for *P. aeruginosa* are not unexpected and show that pyoverdine treatment should not be applied against this pathogen.




Figure 3.

Pyoverdine dose-response curves for *A. baumannii*, *K. pneumoniae*, *P. aeruginosa* and *S. aureus*.

We exposed the four human opportunistic pathogens to three pyoverdines (3A06, 3G07, s3b09) that were among the most potent ones. We used crude-purified extracts of all three pyoverdines and a HPLC-purified variant for pyoverdine s3b09. The absolute concentrations of the crude-purified extracts are unknown and therefore expressed relative to the weighed amount of 6 mg. The absolute concentration of the HPLC-purified variant is given in $\mu\text{g/mL}$. Growth values are scaled relative to the untreated control in CAA medium. Dots and error bars show mean values and standard errors, respectively, across a minimum of three replicates per concentration. Dose-response curves were fitted using 4-or 5-parameter logistic regressions.

Taken together, we found that all pyoverdines are highly potent against *A. baumannii* and *S. aureus*, and one pyoverdine (3G07) is potent against *K. pneumoniae*. Especially in the case of *A. baumannii* and *S. aureus*, the pyoverdine dose-response curves follow the standard trajectory observed for antibiotics (Figure S2 for ciprofloxacin). In support of our hypothesis that the inhibitory effects of pyoverdines operate via withholding iron from pathogens, we found no growth inhibition when the pathogens were exposed to purified ferribactin (Figure S2), the non-iron chelating precursor of pyoverdine.

To further validate that the antibacterial effect of pyoverdine is due to iron sequestration, we saturated pyoverdine s3b09 with varying concentrations of iron and repeated the dose-response experiment with *A. baumannii* (Figure S3). Here, we expect pyoverdine potency to decrease because iron is no longer a growth limiting factor. Indeed, we observed that the inhibition of *A. baumannii* was greatly reduced with intermediate levels of iron supplementation (40 μ M) and completely stalled with high levels of iron supplementation (200 μ M) (Figure S3).

Pyoverdines have low toxicity for mammalian cell lines, erythrocytes, and hosts

We investigated the potential toxicity of crude-purified pyoverdines against mouse neuroblastoma-spinal cord (NSC-34) hybrid cells and human embryonic kidney 293 (HEK-293) cells. NSC-34 are motor neuron-like cells, which do not divide in their differentiated state, thus mimicking established tissue. For these cells, we observed no adverse effects on cell viability at low and intermediate pyoverdine concentrations (Figure 4A), representing the doses that strongly inhibited the pathogens (Figure 3). A moderate decrease of cell viability was only observed at the highest pyoverdine concentrations (Figure 4A). Conversely, HEK-293 cells divide rapidly and inform us on whether pyoverdines interfere with cell proliferation. We indeed noticed a more pronounced effect on cell viability already at intermediate pyoverdine concentrations (Figure 4A), which suggests that pyoverdine chelates the iron required for cell proliferation.

To assess the haemolytic activity of pyoverdines, we exposed sheep erythrocytes to a concentration gradient of the crude-purified pyoverdines. We found that haemolysis was extremely low (<1%) relative to the positive control Triton X-100 surfactant for pyoverdine concentrations up to 0.5 (Figure 4B) and remained low even for the highest concentration (haemolysis rates for pyoverdines 3A06, 3G07 and s3b09 were 2.01%, 1.89% and 5.92%, respectively). This result shows that pyoverdines are unable to retrieve iron from haemoglobin.

Finally, we injected pyoverdine into the larvae of *G. mellonella* and followed host survival over time. We selected three pyoverdine concentrations (0.01, 0.05, and 0.1), which we later used for the infection experiments, and which were minimally toxic to cell lines (coloured dots in Figure 4A). We found that pyoverdine injections did not significantly reduce host survival at low and intermediate concentrations (Figure 4C and Table S1). In contrast, we observed a tendency towards higher larval mortality levels with the highest pyoverdine concentration administered, especially for 3G07, suggesting that pyoverdines can be mildly toxic for the host at high concentrations.

Pyoverdine treatment increases the survival of infected hosts

We then investigated whether pyoverdines can be used to treat infections in *G. mellonella*. We used *A. baumannii*, *K. pneumoniae*, and *P. aeruginosa* as pathogens. We did not include *S. aureus* for these experiments as it is literally avirulent for *G. mellonella* given the infection loads and time frames used⁴². We first determined the bacterial infection load needed to reduce larval survival by at least 50% (Figure S4) over 48 hours. We found this to be the case with 1.8×10^5 CFU/larvae of *A. baumannii* and 8.9×10^5 CFU/larvae of *K. pneumoniae*. For *P. aeruginosa*, all larvae died within 48 hours even with the lowest infection dose and we decided to use 56 CFU/larvae. We infected the




Figure 4.

Toxicity assays for pyoverdines from environmental *Pseudomonas* spp. against human cell lines, sheep erythrocytes and the host larvae of *G. mellonella*.

(A) We exposed mouse neuroblastoma-spinal cord (NSC-34) and human embryonic kidney 293 (HEK-293) cells to three crude-purified pyoverdines (3A06, 3G07, s3b09) that were among the most potent ones to inhibit bacterial growth. An MTT assay was used to assess the metabolic activity of cells as an indicator of cell viability and proliferation. Cell viability data are scaled relative to the pyoverdine-free treatment, whereby dots and error bars show means and standard error across three replicates, respectively. The absolute concentrations of the crude-purified pyoverdines are unknown and concentrations are therefore expressed relative to the highest one used. Coloured dots indicate pyoverdine dosages used for the *in vivo* experiments (pyoverdines/10 μ L). Dose-response curves were fitted using 5-parameter logistic regressions. (B) We evaluated the haemolytic activity of the pyoverdines by adding them to sheep erythrocytes along a concentration gradient (range 0.002 – 1). Triton X-100 and PBS served as positive and negative control, respectively. Haemolytic activity is scaled relative to the positive control, dots and error bars show means and standard error across 6 replicates from two independent experiments, respectively. (C) To assess the toxic effects of pyoverdine on the host, we injected pyoverdines (three relative concentrations, 0.01, 0.05, 0.1) into larvae 4 hours after a mock infection with PBS. The percentage of larval survival was tracked over 48 hours post-treatment. Data stem from three independent experiments with each 10 larvae per infection and treatment.

larvae with one of the three pathogens, and subsequently treated them with either pyoverdine (relative concentrations 0.01, 0.05 and 0.1) or a PBS control four hours post infection, followed by monitoring larval survival over time (**Figure 5** [↗](#)).

We found that the two pyoverdines 3A06 and 3G07 were effective treatments against *A. baumannii* and *K. pneumoniae* and significantly decreased the risk of death compared to untreated infections (**Figure 5** [↗](#), Table S2). Specifically, *A. baumannii* treated with 3A06 at a concentration of 0.05 and 3G07 at a concentration of 0.01 reduced the risk of death (hazard) by 57.9% ($z = -1.973$, $P = 0.0485$) and 59.4% ($z = -2.057$, $p = 0.0397$), respectively. For s3b09, the same tendency was observed ($z = -1.496$, $p = 0.135$) albeit not significant. For *K. pneumoniae*, the pyoverdine treatments 3A06 and 3G07 (concentration of 0.01) reduced the risk of death by 51.5% ($z = -2.027$, $p = 0.0426$) and 57.6% ($z = -2.341$, $p = 0.0192$), respectively. Lower doses of pyoverdine seemed to be more effective than the highest concentration, reinforcing our observation that pyoverdine can exhibit some toxicity at higher concentrations (**Figure 4C** [↗](#)). Infections with *P. aeruginosa* led to near 100% larval mortality regardless of pyoverdine treatment and dosage so that no significant treatment effect arose. Altogether, our infection experiments show that pyoverdines are effective against the moderately virulent pathogens *A. baumannii* and *K. pneumoniae*, while they are ineffective against the fast-killing highly virulent *P. aeruginosa*.

Pathogens show low levels of resistance evolution against pyoverdine treatment

We assessed the ability of *A. baumannii*, *K. pneumoniae*, *P. aeruginosa* and *S. aureus* to evolve resistance against the pyoverdine treatment by exposing all pathogens to both single pyoverdine treatments (3A06, 3G07, and s3b09) and combination treatments (double and triple combinations) over 16 days of experimental evolution. One exception was *K. pneumoniae*, which was only exposed to 3G07 as only this pyoverdine was inhibitory. We had six independent lineages (populations) per pathogen and treatment combination (see Table S2 for treatment concentrations). We further subjected all pathogens to the antibiotic ciprofloxacin as positive control in which resistance evolution is expected. Additionally, we let the pathogens grow in untreated medium to control for medium adaptation. Overall, we had 180 lineages that were transferred daily to fresh medium.

After experimental evolution, we observed that all four pathogens grew significantly better under ciprofloxacin treatment than the ancestor and beyond the level observed for medium adaptation (Table S3; **Figure 6A** [↗](#)). This provides strong evidence for pervasive resistance evolution against this conventional antibiotic. In contrast, we found that none of the four pathogens treated with pyoverdines experienced a significant increase in growth relative to the observed level of medium adaptation (Table S3, **Figure 6A** [↗](#)), suggesting low potentials of resistance evolution at the population level.

However, population screens can hide the presence of resistant clones within populations. We thus picked eight random clones from two populations for each of the pyoverdine single treatments and one combo treatment for each pathogen. With a total of 208 clones, we repeated the above growth assay and expressed clonal growth as fold-change relative to the growth of populations evolved in medium alone (**Figure 6B** [↗](#)). For *A. baumannii* and *K. pneumoniae*, the average growth fold-change across clones was 1.02 ± 0.004 and 0.84 ± 0.02 , respectively, and there was no sign of resistant clones being present in populations. This was different for *P. aeruginosa* and *S. aureus*, for which we observed high variation in growth fold-changes across clones, with many clones growing moderately better than the medium controls. The growth increase across clones was 1.11 ± 0.02 and 1.10 ± 0.02 , and significant in three and two population of *P. aeruginosa* and *S. aureus*, respectively. Thus, the clonal screen indicates some level of adaptation to pyoverdine treatment for these two pathogens.




Figure 5.

Pyoverdine treatments significantly increase survival of the host *G. mellonella* when infected with *A. baumannii* and *K. pneumoniae*.

Larvae of the greater wax moth were first infected with either *A. baumannii*, *K. pneumoniae*, or *P. aeruginosa* and then treated with one of three pyoverdines (3A06, 3G07, or s3b09) at four relative pyoverdine concentrations (0, 0.01, 0.05, 0.1). All panels show the larvae survival over 48 hours post-treatment. Data stem from three independent experiments with each 10 larvae per infection and treatment. Asterisks indicate significant differences in larval survival between treated and untreated infections based on Cox proportional hazard regressions ($p < 0.05$).




Figure 6.

Phenotypic and genotypic analysis of experimentally evolved pathogens reveal weak levels of resistance evolution against pyoverdine treatment.

(A) We exposed evolved and ancestral pathogen populations to the treatment in which they evolved in and quantified their growth (area under the curve). Growth values were scaled relative to the ancestor in untreated medium, and the panels show the scaled growth differences between evolved and ancestral populations. The shaded areas show the scaled growth difference between the ancestor and the populations evolved in untreated medium and is representative of medium adaptation. The blue and green dots represent the pathogen populations with the lowest (population 1) and highest (population 2) scaled growth difference, respectively, which were subsequently used to pick clones. The dots show mean values across two independent replicates and asterisks show significant growth increases relative to the medium-adapted control. Box plots show the median and the first and third quartiles across the 6 independently evolved populations. Whiskers represent the 1.5x interquartile range. (B) We repeated the above growth assays with 208 individual clones evolved under pyoverdine treatment and 24 populations evolved in medium alone (control). Each square represents a clone or a control population, and the number indicates the number of mutations identified based on whole-genome sequencing. The heatmap shows the fold-change in growth relative to the evolved control populations. Asterisks depict significant fold-increases in growth compared to control populations. (C) Relationship between the number of mutations and fold-change in growth across all sequenced clones and populations ($n = 68$). (D) Heatmap showing the number of mutations per pathogen and per functional gene categories across clones together with the respective fold-change in growth ($n = 47$).

To obtain insights into the genetic basis of putative resistance evolution, we sequenced the whole genomes of 52 evolved clones (16 per pathogen, except 4 for *K. pneumoniae*), 16 medium-evolved control populations and the 4 ancestors. We first mapped reads to the reference genomes and then to our ancestors. We called single nucleotide polymorphisms and small deletions/insertions that arose during experimental evolution (**Figure 6B**). We observed that growth peaks in clones with one mutation and decreases in clones with more mutations (**Figure 6C**, linear fit: $F_{1,66} = 7.51$, $p = 0.0079$), a pattern that suggests positive selection of a few specific advantageous mutations. Next, we filtered out mutations in genes that occurred at least twice in the control populations and identified 14 (*A. baumannii*), 0 (*K. pneumoniae*), 15 (*P. aeruginosa*), and 18 (*S. aureus*) mutations that exclusively arose under pyoverdine treatment and could thus be associated with resistance evolution. We classified the mutated genes in functional categories (**Figure 6D**, Table S4) and found that genes relevant for biofilm and motility formed the only category in which mutations were both frequent and consistently associated with increased growth in *P. aeruginosa* and *S. aureus* (**Figure 6D**).

Discussion

We investigated whether pyoverdines produced by environmental *Pseudomonas* spp. have antibacterial activities against human opportunistic pathogens through iron sequestration and withholding this trace element from pathogens. Our screen involving 320 environmental isolates and 12 pathogen strains revealed five top pyoverdine candidates with broad-spectrum activity. These candidates could be distinguished from non-inhibitory pyoverdines by their high CCS values (standing for larger and more complex molecules) and their higher iron complexation stability. Experiments with crude- and HPLC-purified pyoverdines showed that pyoverdines completely stall the growth of *A. baumannii* and *S. aureus*, while showing intermediate activity against *K. pneumoniae* and *P. aeruginosa*. When administered as treatment to infected *G. mellonella* larvae, we observed significantly increased host survival rates in infections with *A. baumannii* and *K. pneumoniae*, demonstrating that pyoverdine exhibits antibacterial activity in hosts. Furthermore, we found low toxicity of pyoverdines at effective concentrations and observed low potentials for resistance evolution. Overall, our results reveal pyoverdines from non-pathogenic *Pseudomonas* spp. as potent antibacterials against several human opportunistic pathogens.

Our novel treatment approach assumes that pyoverdines sequester iron and thereby induce iron starvation and growth arrest in pathogens. Several of our findings support this view. First, the growth-inhibiting effect of our top pyoverdine candidates exclusively occurred under iron-limited conditions, while pathogen growth was restored under iron-replete conditions (**Figure 1**, Figure S3). Second, non-inhibitory ferri-pyoverdines were more prone towards collision induced unfolding than inhibitory ferric-pyoverdines (**Figure 2D**), suggesting that inhibitory pyoverdines have a higher iron binding affinity. Third, the pyoverdine precursor ferribactin did not inhibit pathogen growth (Figure S2). The two molecules are identical apart from one ring in the chromophore core that is not cyclized and not oxidized in ferribactin, which compromises the binding of iron. These findings suggest that pyoverdine does not target the pathogens directly, but rather indirectly via the induction of iron starvation.

We observed that pyoverdines showed pathogen-specific efficacies in curbing bacterial growth (**Figure 3**). We propose that differences can be explained by the various strategies of these pathogens to cope with iron stress. *A. baumannii* and *S. aureus* produce siderophores that have a much simpler chemical structures than pyoverdines and are expected to bind iron with lower affinity (*A. baumannii*: acinetobactin (association constant: 10^{26} M^{-1}), fimsbactin (10^{27} M^{-1}) and baumannoferrin (unknown)^{43–45}, *S. aureus*: staphyloferrin A+B (unknown)⁴⁶) than pyoverdine (10^{32} M^{-1})²⁹. Hence, the growth suppression of *A. baumannii* and *S. aureus* most likely occurs because their siderophores are too weak to successfully compete with pyoverdine. The situation is different for *K. pneumoniae*, which produces enterobactin, the siderophore with

the highest known iron affinity (10^{52} M^{-1})³⁸. Consistent with the concept of iron competition, we observed that pyoverdine treatment is less effective against *K. pneumoniae*, and was only potent for one out of three pyoverdines (3G07, **Figure 3**). Finally, pyoverdine treatment is not expected to work well against *P. aeruginosa*, which itself produces a version of this siderophore. The two competing pyoverdines likely have similar iron affinities and the presence of pyoverdine receptors in the pathogen might foster the uptake of the supplemented pyoverdine. Our results indeed support the view that pyoverdine treatment is not effective against *P. aeruginosa*. Taken together, we predict that pyoverdine treatment could be highly potent against certain pathogens like *A. baumannii* and *S. aureus* that have less efficient siderophores and are unable to use the pyoverdine as iron source.

Not only bacteria but also host cells need iron. Consequently, pyoverdine treatment could negatively affect host iron homeostasis. Our toxicity assays reveal that there is a range of pyoverdine concentrations for which only mild adverse effects against cell lines and *G. mellonella* larvae are observed and no haemolysis occurs. Especially the finding that pyoverdines have no negative effects on the survival of non-replicating NSC-34 cells is encouraging, as this cell line is representative of intact host tissue, in which iron is bound to strong chelators such as transferrin, lactoferrin or ferritin^{47,48}. The result that pyoverdines are unable to obtain iron from haemoglobin is encouraging, too. Previous work showed that the pyoverdine of *P. aeruginosa* can retrieve iron from human transferrin when working in concert with proteases (proteolytically degrading transferrin)⁴⁹ or phenazines (spontaneously reducing ferric to ferrous iron)⁵⁰. However, these *P. aeruginosa* specific mechanisms require the presence of this pathogen⁵¹, which is not the case when purified pyoverdines are used as a treatment against other pathogens like *A. baumannii*. Nonetheless, we found that pyoverdines can have certain adverse effects, especially for the rapidly proliferating HEK293 cells (**Figure 4**). These cells represent a regenerating tissue and thus have high iron demands. Similarly, we observed mild adverse effects for some pyoverdines (3G07) when administered at high concentrations to *G. mellonella* larvae (**Figure 5**). These findings corroborate the results from previous studies on *Caenorhabditis elegans*, where pyoverdines were found to interfere with host iron homeostasis^{52,53}. These considerations highlight that finding the right pyoverdine concentration will be key to meet the fine line between maximizing pathogen inhibition and minimizing tissue damage. By contrast, the adverse host effects of pyoverdines might come with additional opportunities, for instance in cancer therapy, where siderophores reduce iron levels in tumours and slow tumour progression⁵⁴.

We propose that pyoverdine treatment could synergistically interact with the host innate immune system. Mammalian neutrophils produce siderocalin to sequester bacterial siderophore-iron complexes, rendering them non-functional⁴⁷. While this strategy is efficient against many siderophores, some bacteria have evolved so-called stealth siderophores that no longer bind to siderocalin. Pyoverdines are such stealth siderophores⁵⁵. Regarding pyoverdine treatment, this means that pathogens with regular siderophores will hit a double wall: their siderophores are immobilized by siderocalins, while pyoverdines lock away the remaining iron.

Another promising finding of our work is that the potential for resistance evolution against pyoverdines seems low. Conceptually, one reason for low resistance evolution could be that pyoverdines, unlike many antibiotics, are not internalized into bacterial cells. Thus, classic resistance mechanisms such as reduced drug influx, increased drug efflux, and intra-cellular target modification cannot operate⁵⁶. We initially expected that resistance evolution would involve mechanisms that improve the pathogen's iron acquisition. This could include the up-regulation of the pathogen's own siderophores or switching from siderophore-based ferric iron acquisition to the use of reductases that foster ferrous iron uptake⁵⁶. However, our sequencing analysis did not reveal mutations in genes associated with iron uptake systems. One explanation is that such mutations (albeit beneficial) might not reach high frequencies in populations because increased siderophore production and secretion would increase the iron acquisition of resistant

and non-resistant cells alike⁵⁶. We instead observed an overrepresentation of mutations in biofilm and motility genes in *P. aeruginosa* and *S. aureus*. While we do not know the phenotypes of these mutations, one possibility is that increased biofilm formation could offer some protection against pyoverdine treatment, for example by restricting pyoverdine from chelating free iron. Important to note is also that patterns of resistance evolution may differ between *in vitro* and *in vivo* settings^{4,57}. For example, our experimental evolution setup did not allow for horizontal transfer of siderophore receptors, a scenario that might occur in the context of polymicrobial infections and confer resistance to pyoverdine therapy.

In conclusion, our treatment approach using pyoverdine is based on the principle of paying like with like. Pathogens secrete siderophores to scavenge iron for their growth. We add a strong heterologous siderophore as treatment to withhold iron from pathogens to curb their growth. The idea to target bacterial iron up-take mechanisms has been highlighted in the literature for quite some time^{13,58–61}. While certain approaches focussed on the inhibition of siderophore synthesis⁵⁸, others were based on the administration of synthetic iron chelators¹³. However, synthetic iron chelators are often toxic and have lower iron affinities compared to bacterial siderophores. The advantage of our approach is that we capitalize on naturally evolved iron chelators from non-pathogenic species with extremely high iron affinity. We have shown that these natural iron chelators have a high potency to make iron inaccessible to pathogens, while showing low toxicity for hosts at treatment effective concentrations and low potentials for resistance evolution. They could become a new source of efficient antibacterials either as independent treatment or in combination with conventional antibiotics.

Material and methods

Bacterial strains

We used a strain collection of 320 natural isolates, originating from soil and pond habitats. The isolates originate from 16 independent soil and water samples (20 isolates per sample) collected on the Campus Irchel Park of the University of Zurich. In previous work, we have extensively studied and characterized all isolates^{17,30,32}. Details on the isolates can be found in the Supporting Information.

We used a collection of 14 strains of opportunistic human pathogens (**Table 1**), including *A. johnsonii*, *A. junii*, *C. sakazakii*, *K. michiganensis* (belonging to the *K. oxytoca* complex)⁶² and *Shigella* sp. (all provided by the laboratory of Leo Eberl, University of Zurich), and *B. cenocepacia* strains H111 and K56-2, *P. aeruginosa* strains PA14 and PAO1, *E. coli* K12 and *S. aureus* strains Cowan and JE2 from our own strain collection. *A. baumannii* (DSM 30007) and *K. pneumoniae* (DSM 30104) were purchased from the German Collection of Microorganisms and Cell Cultures GmbH.

Culturing conditions

Pathogen overnight cultures were grown in either 8 mL tryptic soy broth (TSB; for the two *S. aureus* strains) or 8 mL lysogeny broth (LB; all other pathogens) in 50 mL tubes at 37 °C and 220 rpm agitation. Following two washing steps with 0.8% NaCl, we adjusted the overnight cultures to an optical density at 600 nm (OD₆₀₀) of 1. For overnights of environmental isolates, we followed the same procedure when working with low sample sizes, with the only difference being that culturing occurred at 28 °C. For the large-scale screening experiments, we grew environmental isolates in 200 µL LB in 96-well plates shaken at 170 rpm. Cultures from plates were then directly used for experiments. For all main experiments (screen for bioactive pyoverdines and dose-response curves), we used casamino acid (CAA) medium (1% casamino acids, 5 mM K₂HPO₄ * 3H₂O, 1 mM MgSO₄ * 7 H₂O, 25 mM HEPES buffer). This medium contains low levels of iron, triggering pyoverdine production⁶³.

Supernatant screening assays

We created sterile supernatants from all the 320 *Pseudomonas* isolates. Specifically, we transferred 2 µL of overnight cultures (grown in 96-well plates in LB in four-fold replication for 24 hours as described above) to new plates containing 200 µL of CAA medium supplemented with 250 µM 2,2'-bipyridyl. Cultures were incubated for 24 hours and subsequently centrifuged at 2250g for 10 minutes. The supernatants were transferred to 0.2 µm membrane filter plates (PALL AcroPrep Advance), centrifuged a second time, and frozen at -20 °C.

For the supernatant screening assay, the pathogens were cultured and diluted as described above and added to 140 µL CAA medium in 96-well plates. We added 60 µL of thawed *Pseudomonas* supernatants (30%) or 0.8% NaCl as control and incubated the plates at 37 °C and 170 rpm. Growth was measured as OD₆₀₀ after 24 hours using a Tecan Infinite M-200 plate reader (Tecan Group, Männedorf, Switzerland). We then scaled the growth of pathogens in the supernatant relative to their growth in the control treatment. Relative growth values smaller and larger than one indicate growth inhibition and promotion by the supernatant, respectively.

To test whether pyoverdine is responsible for pathogen growth reduction, we repeated the screening assay but supplemented the CAA medium containing the *Pseudomonas* supernatants with 40 µM FeCl₃. In this iron-rich condition, pyoverdine should not be able to limit iron availability and thus pathogen growth should be restored.

Structure elucidation of pyoverdine and analyses of chemical properties

From the seven top supernatant candidates that inhibited all twelve pathogens, we extracted the pyoverdines and elucidated their structure. For this purpose, we have developed a new protocol that allows structure elucidation from low volume supernatants using UHPLC-HR-MS/MS. The methodological details are described in Rehm *et al.*^{28,34}. Afterwards, we assessed whether growth inhibiting pyoverdines can be distinguished from non-inhibiting pyoverdines by their chemical properties. We focussed on polarity, CCS values, and stability upon collision induced unfolding or fragmentation. Detailed protocols and instrument settings can be found in the Supporting Information.

Crude pyoverdine purification

To test the efficacy of the pyoverdines against the four focal human opportunistic pathogens (*A. baumannii*, *K. pneumoniae*, *P. aeruginosa* (PAO1), and *S. aureus* (JE2)), and to assess their cytotoxicity against human cell lines, we crude-purified three pyoverdines (3A06, 3G07, s3b09). The three pyoverdines were among the five most potent ones identified in our supernatant screening assay. The adapted method for pyoverdine purification from the previous studies^{17,35} are further described in the Supporting Information.

HPLC purification of pyoverdine and ferripectin

To further validate that pyoverdines are the compounds that induce pathogen growth inhibition, we purified the most common pyoverdine (s3b09) among the most potent pyoverdines from our screens using reversed-phase high-performance liquid chromatography (HPLC). Methodological details thereof are described in the Supporting Information.

Dose-response curves

To determine the inhibition potential of crude- and HPLC-purified pyoverdines, as well as ferripectin, we subjected *A. baumannii*, *K. pneumoniae*, *P. aeruginosa* and *S. aureus* to serially diluted extracts. For this, the extracts were dissolved in CAA medium, filter sterilized and diluted in the same medium. Bacterial overnight cultures were grown and prepared as described above

and added to a total of 200 μL medium on a 96-well plate with 4 replicates per extract dilution. As positive control where growth inhibition is expected, we subjected all three pathogens to a dilution series of the antibiotic ciprofloxacin (highest concentration 4 $\mu\text{g}/\text{mL}$), in the exact same way as for the extracts. We incubated the plates at 37 °C statically in the plate reader and measured the OD_{600} every 15 minutes for 24 hours. Before each measurement, plates were shaken for 15 seconds. We repeated the experiment up to five times with different initial extract starting concentrations. This was done to ensure that the full dose response to pyoverdine was captured. Thus, sample size per individual extract concentration varied between four and 20 replicates. We subtracted the blank values and the background values caused by the treatment of the respective extract concentration from the measured growth data and calculated the integral (area under the growth curve) using the R package *Growthcurver*⁶⁴. We then expressed growth relative to the control medium, where bacteria grew in CAA without extract or antibiotic addition. Finally, we plotted extract/antibiotic concentration vs. relative growth and fitted dose-response curves (see below).

Cytotoxicity assays

To determine the cytotoxicity of pyoverdine treatments against mammalian cells, we subjected human embryonic kidney 293 (HEK-293) and mouse motor neuron-like neuroblastoma-spinal cord hybrid (NSC-34) cells to the crude extract of the three most potent pyoverdines (3A06, 3G07, s3b09). Detailed culturing conditions can be found in the Supporting Information. Briefly, approximately 10^4 cells/well were plated on 96-well plates (Greiner Bio-One, 655090) and, after 48 hours, the culture medium was replaced and the pyoverdines were added covering a range of concentrations. After another 48 hours, the pyoverdine-containing culture medium was removed, and cells were incubated with 1 $\mu\text{g}/\text{mL}$ thiazolyl blue tetrazolium bromide (Sigma, D5655) in 100 μL of fresh medium for 1.5 hours (NSC-34) or 40 minutes (HEK-293). The reaction was stopped, and formazan crystals simultaneously solubilized with the addition of 100 μL of a solution containing 10% sodium dodecyl sulfate (SDS; Sigma-Aldrich, 05030) and 0.03% HCl (Supelco, 100319). Finally, the absorbance of cell debris and other contaminants at 630 nm were subtracted from the absorbance of the solubilized formazan at 570 nm.

Haemolysis assay

To determine the haemolytic activity of pyoverdines, we quantified sheep erythrocyte haemolysis at different pyoverdine concentrations. We first centrifuged 25 mL of fresh sheep blood (Chemie Brunschwig AG, Switzerland) in a 50 mL tube at 1500g for 5 minutes at 4 °C. Plasma was then gently aspirated, replaced by PBS and tubes were gently inverted to mix. This washing procedure was repeated three times. Pyoverdines were diluted in PBS to final relative concentrations of 1, 0.5, 0.25, 0.2, 0.02, and 0.002. PBS and 0.1% Triton X-100 (Sigma Aldrich, Switzerland) served as negative and positive control, respectively. 100 μL erythrocyte solution was mixed with either 100 μL pyoverdine solution or the controls in a 96-well plate in triplicates. Following an incubation at 37 °C and 120 rpm for 1 hour, the plate was centrifuged at 1036g for 5 minutes at 4 °C. Subsequently, 100 μL of the supernatant were transferred to a new 96-well plate. Haemoglobin release was measured at 570 nm in the plate reader and the percent haemolysis was calculated relative to the positive control (% haemolysis = $(\text{OD}_{570} \text{ of sample} - \text{negative control}) / (\text{positive control} - \text{negative control})$).

Infection model

We tested the efficacy of pyoverdine to treat infections in the host model *G. mellonella*. Detailed larvae handling and determination of LD_{50} (lethal dose, killing 50% of the larvae) are described in the Supporting Information. For the main experiments, we used the LD_{50} values for *A. baumannii* and *K. pneumoniae* (Figure S4). For *P. aeruginosa*, we use 56 CFU/larvae because all doses tested

killed 100% of the larvae (Figure S4). As negative control, we administered 10 μ L/larvae of sterile PBS. Larvae were incubated at 37 °C in the dark for 4 hours to allow pathogens to establish infections.

To treat infected larvae, pyoverdines 3A06, 3G07, and s3b09 were used at three concentrations that were non-toxic for the NSC-34 cells. For this, we prepared stocks of 60 mg/mL, 30 mg/mL, and 6 mg/mL of crude pyoverdines. Since the injected treatment volume was 10 μ L per larvae, the final pyoverdine concentrations were 0.6 mg, 0.3 mg, and 0.06 mg per larvae, respectively. When scaling these concentrations to the doses used for the *in vivo* dose-response curves (Figure 4), the relative pyoverdine concentrations were 0.1, 0.05 and 0.01. Pyoverdine treatments were administered through injections (see Supporting Information) with the larvae being immobilized on ice for 15 minutes. Following treatment, larvae were kept at 37°C in the dark. From the 12th hour post infection (hpi), their survival was regularly checked for a total of 48 hpi. Specifically, larvae were poked with a pipette tip and individuals not responding to the physical stimulation were considered dead. We conducted three independent experiments per pathogen and treatment at different days with larvae from different batches.

Experimental evolution experiment

We first determined the IC₅₀ concentrations of (crude) pyoverdines 3A06, 3G07, and s3b09, and the antibiotic ciprofloxacin against all four pathogens. For this, we fitted logistic regressions to the dose-response curves. We used these IC₅₀ values as treatment concentrations during experimental evolution (Table S2). We also mixed different pyoverdines in double and triple combo treatments. In combo treatments, pyoverdine concentrations were equivalent to those of the single treatments, meaning that we divided the single treatment concentrations by two or three. The experimental evolution experiment included (i) three single pyoverdine treatments (3A06, 3G07, s3b09), (ii) three double pyoverdine treatments (3A06 and 3G07, 3A06 and s3b09, 3G07 and s3b09), (iii) one triple pyoverdine treatment (3A06, 3G07 and s3b09), and (iv) one ciprofloxacin treatment for *A. baumannii*, *P. aeruginosa* and *S. aureus*. For *K. pneumoniae*, which was only affected by one pyoverdine (3G07), the experiment consisted of one single pyoverdine treatment and one ciprofloxacin treatment.

A detailed description of the experimental evolution setup is provided in the Supporting Information. Briefly, for each pathogen-treatment combination, we included six independently replicated lineages (populations) on 96-well plates. Each plate further contained control populations that evolved in plain CAA medium. At the start, we grew the pathogens overnight, washed and diluted them as described above, and transferred 2 μ L of the diluted pathogens into 200 μ L medium in 96-well plates. Subsequently, the plates were incubated at 37 °C for 22 hours, shaken at 170 rpm. After incubation, we measured growth (OD₆₀₀) of all evolving lineages with the plate reader. We then diluted the evolving cultures 1:1,000 in CAA medium and added them to a fresh treatment plate. After the transfer, the new plates were again incubated at 37 °C and 170 rpm agitation for 22 hours. We added glycerol to the old plates to a final concentration of 25% and stored them at -80 °C.

Phenotypic screening of evolved populations

We screened evolved populations for improved growth performance relative to the ancestor under pyoverdine treatment, which we took as a proxy for resistance evolution. For this purpose, we thawed the final plates (day 16) of the experimental evolution experiment and transferred 2 μ L of all populations to 200 μ L of LB or TSB (for *S. aureus*) to create overnight cultures in 96-well plates. We further included ancestor cultures of all four species. After incubation for 18 hours at 37 °C and 170 rpm, we diluted overnights 1:1,000 with 0.8% NaCl and subjected all evolved populations to the conditions they had evolved in. Ancestors were simultaneously exposed to the respective treatment conditions. The culturing volume and conditions matched those used during experimental evolution, but this time we incubated the plates in a plate reader, measuring OD₆₀₀

every 15 minutes to obtain growth kinetics. The population screen was repeated twice independently. From the growth kinetics, we first calculated the area under the curve (AUC) for each growth curve. AUC values were then scaled relative to the growth of the ancestor in the untreated CAA control medium. For each evolved population, we then calculated the difference in growth in comparison to the mean growth of the ancestor subjected to the same treatment.

Phenotypic screening of evolved clones

To screen for resistant clones within populations, we selected the pathogen populations with the lowest and highest growth difference for each single treatment and for one combination treatment and picked 8 random colonies. We transferred them to 200 μ L LB or TSB (*S. aureus*) in 96-well plates and incubated plates for 18 hours at 37 °C (170 rpm) to create overnight cultures. We followed the same protocol as above for quantifying the difference in growth of evolved and ancestral clones.

Genomic analyses of evolved populations and clones

We sequenced the genomes of the four ancestors, four control populations that had evolved in medium without pyoverdine treatment, and two random clones (out of the eight picked) per population that had received pyoverdine treatment and were used for clonal growth analysis (Figure 6B [↗](#)). The Supporting Information features a detailed description of the sequencing and data analysis protocols used. After variant calling, we screened for mutations that could be associated with resistance to pyoverdine treatment. For this analysis, we excluded variants with low coverage (< 30 reads), synonymous mutations, mutations already present in the ancestor, and mutations that occurred at least twice in evolved control populations.

Data analyses

All statistical analyses were performed in R 4.0.2 and RStudio version 1.3.1056. We conducted Shapiro-Wilk tests and consulted diagnostic plots to check whether our data (residuals) were normally distributed. For normally distributed data sets, we used linear models and two-sided tests for all analyses. For non-normally distributed data sets, we used non-parametric rank tests. We used ANOVA to test whether chemical properties differ between growth inhibiting and non-inhibiting pyoverdines, in their ferri-versus apo-pyoverdine state. For the *in vitro* dose-response and cytotoxicity experiments, we fitted dose-response curves with either 4- or 5-parameter logistic regressions using the `nplr` package⁶⁵ [↗](#). Survival analyses were performed using the Cox proportional hazards regression model or the log-rank test (adjusted for multiple comparisons using the Holm method) from the `survival` package⁶⁶ [↗](#). We used two-sided t-tests, Welch's t-tests, or Wilcoxon tests to compare growth differences between the ancestor and the evolved populations under treatment. To compare growth differences between the ancestor and the evolved clones under treatment, we used one- and two-way ANOVAs.

Acknowledgements

We thank Richard Allen, Dominik Bär, Johanna Giger, Kevin Schiefelbein and Julien Weber for laboratory assistance, and Markus Seeger for insightful discussion. This work was supported by funding from the Swiss National Science Foundation (grant numbers: 31003A_182499 to RK, PP00P3_144862 and PP00P3_176966 to MP) and a Candoc Grant (Forschungskredit) from the University of Zurich (to MPB).

Authors contributions

VV and RK conceived the project. VV, KR, CC and MPB performed the experiments and analysed experimental results. All authored interpreted the data. VV and RK wrote the manuscript with contributions from all authors.

Competing interests

Authors declare no competing interests.

References

1. Murray C. J. *et al.* (2022) **Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis** *The Lancet* **399**:629–655
2. WHO (2022) **WHO. 2021 antibacterial agents in clinical and preclinical development: an overview and analysis. (2022).** at <https://www.who.int/publications-detail-redirect/9789240047655>
3. Beyer P., Paulin S (2020) **The Antibacterial Research and Development Pipeline Needs Urgent Solutions** *ACS Infect. Dis* **6**:1289–1291
4. Bell G., MacLean C (2018) **The Search for ‘Evolution-Proof’ Antibiotics** *Trends Microbiol* **26**:471–483
5. Monserrat-Martinez A., Gambin Y., Sierrecki E (2019) **Thinking Outside the Bug: Molecular Targets and Strategies to Overcome Antibiotic Resistance** *Int. J. Mol. Sci* **20**
6. Rezzoagli C., Granato E. T., Kümmerli R (2020) **Harnessing bacterial interactions to manage infections: a review on the opportunistic pathogen *Pseudomonas aeruginosa* as a case example** *J. Med. Microbiol* **69**:147–161
7. Bhushan A., Peters E. E., Piel J (2017) **Entotheonella Bacteria as Source of Sponge-Derived Natural Products: Opportunities for Biotechnological Production**
8. Ghosh C., Sarkar P., Issa R., Haldar J (2019) **Alternatives to Conventional Antibiotics in the Era of Antimicrobial Resistance** *Trends Microbiol* **27**:323–338
9. Mei Y., Chen Y., Sivaccumar J. P., An Z., Xia N., Luo W (2022) **Research progress and applications of nanobody in human infectious diseases** *Front. Pharmacol* **13**
10. Rezzoagli C., Archetti M., Mignot I., Baumgartner M., Kümmerli R (2020) **Combining antibiotics with antivirulence compounds can have synergistic effects and reverse selection for antibiotic resistance in *Pseudomonas aeruginosa*** *PLOS Biol* **18**
11. Saeki E. K., Kobayashi R. K. T., Nakazato G (2020) **Quorum sensing system: Target to control the spread of bacterial infections** *Microb. Pathog* **142**
12. Wale N., Sim D. G., Jones M. J., Salathe R., Day T., Read A. F (2017) **Resource limitation prevents the emergence of drug resistance by intensifying within-host competition** *Proc. Natl. Acad. Sci* **114**:13774–13779
13. Hatcher H. C., Singh R. N., Torti F. M., Torti S. V (2009) **Synthetic and natural iron chelators: therapeutic potential and clinical use** *Future Med. Chem* **1**:1643–1670
14. Hug J. J., Krug D., Müller R (2020) **Bacteria as genetically programmable producers of bioactive natural products** *Nat. Rev. Chem* **4**:172–193
15. Imai Y. *et al.* (2019) **A new antibiotic selectively kills Gram-negative pathogens** *Nature* **576**:459–464

16. Ling L. L. *et al.* (2015) **A new antibiotic kills pathogens without detectable resistance** *Nature* **517**:455–459
17. Butaitė E., Baumgartner M., Wyder S., Kümmerli R (2017) **Siderophore cheating and cheating resistance shape competition for iron in soil and freshwater *Pseudomonas* communities** *Nat. Commun* **8**
18. Deveau A., Gross H., Palin B., Mehnaz S., Schnepf M., Leblond P., Dorrestein P. C., Aigle B (2016) **Role of secondary metabolites in the interaction between *Pseudomonas fluorescens* and soil microorganisms under iron-limited conditions** *FEMS Microbiol. Ecol* **92**
19. Hider R. C., Kong X. (2010) **Chemistry and biology of siderophores** *Nat. Prod. Rep.* **27**:637–657
20. Figueiredo A. R. T., Özkaya Ö., Kümmerli R., Kramer J (2022) **Siderophores drive invasion dynamics in bacterial communities through their dual role as public good versus public bad** *Ecol. Lett* **25**:138–150
21. Gu S. *et al.* (2020) **Competition for iron drives phytopathogen control by natural rhizosphere microbiomes** *Nat. Microbiol* **5**:1002–1010
22. Lee W., van Baalen M., Jansen V. A. A (2012) **An evolutionary mechanism for diversity in siderophore-producing bacteria** *Ecol. Lett* **15**:119–125
23. Niehus R., Picot A., Oliveira N. M., Mitri S., Foster K. R (2017) **The evolution of siderophore production as a competitive trait** *Evolution* **71**:1443–1455
24. Visca P., Imperi F., Lamont I. L (2007) **Pyoverdine siderophores: from biogenesis to biosignificance** *Trends Microbiol* **15**:22–30
25. Bonneau A., Roche B., Schalk I. J (2020) **Iron acquisition in *Pseudomonas aeruginosa* by the siderophore pyoverdine: an intricate interacting network including periplasmic and membrane proteins** *Sci. Rep* **10**
26. Cornelis P., Tahrioui A., Lesouhaitier O., Bouffartigues E., Feuilloley M., Baysse C., Chevalier S (2022) **High affinity iron uptake by pyoverdine in *Pseudomonas aeruginosa* involves multiple regulators besides Fur, PvdS, and FpvI** *BioMetals*
27. Meyer J.-M., Gruffaz C., Raharinosy V., Bezverbnaya I., Schäfer M., Budzikiewicz H (2008) **Siderotyping of fluorescent *Pseudomonas*: molecular mass determination by mass spectrometry as a powerful pyoverdine siderotyping method** *BioMetals* **21**:259–271
28. Rehm K., Vollenweider V., Kümmerli R., Bigler L (2022) **A comprehensive method to elucidate pyoverdines produced by fluorescent *Pseudomonas* spp. by UHPLC-HR-MS/MS** *Anal. Bioanal. Chem* **414**:2671–2685
29. Meyer, Abdallah M. A. (1978) **The Fluorescent Pigment of *Pseudomonas fluorescens*: Biosynthesis, Purification and Physicochemical Properties** *Microbiology* **107**:319–328
30. Butaitė E., Kramer J., Wyder S., Kümmerli R (2018) **Environmental determinants of pyoverdine production, exploitation and competition in natural *Pseudomonas* communities** *Environ. Microbiol* **20**:3629–3642

31. Butaitė E., Kramer J., Kümmerli R (2021) **Local adaptation, geographical distance and phylogenetic relatedness: Assessing the drivers of siderophore-mediated social interactions in natural bacterial communities** *J. Evol. Biol* **34**:1266–1278
32. Kramer J., López Carrasco M. Á., Kümmerli R (2020) **Positive linkage between bacterial social traits reveals that homogeneous rather than specialised behavioral repertoires prevail in natural *Pseudomonas* communities** *FEMS Microbiol. Ecol* **96**
33. Meyer J.-M (2000) **Pyoverdines: pigments, siderophores and potential taxonomic markers of fluorescent *Pseudomonas* species** *Arch. Microbiol* **174**:135–142
34. Rehm K., Vollenweider V., Kümmerli R., Bigler L (2023) **Rapid identification of pyoverdines of fluorescent *Pseudomonas* spp. by UHPLC-IM-MS** *BioMetals* **36**:19–34
35. Meyer J.-M., Stintzi A., De Vos D., Cornelis P., Tappe R., Taraz K., Budzikiewicz H. (1997) **1997. Use of Siderophores to Type *Pseudomonads*: The Three *Pseudomonas aeruginosa* Pyoverdine Systems** *Microbiology* **143**:35–43
36. Budzikiewicz H., Schäfer M., Fernández D. U., Matthijs S., Cornelis P (2007) **Characterization of the chromophores of pyoverdins and related siderophores by electrospray tandem mass spectrometry** *BioMetals* **20**:135–144
37. Nadal-Jimenez P., Koch G., Reis C. R., Muntendam R., Raj H., Jeronimus-Stratingh C. M., Cool R. H., Quax W. J (2014) **PvdP Is a Tyrosinase That Drives Maturation of the Pyoverdine Chromophore in *Pseudomonas aeruginosa*** *J. Bacteriol* **196**:2681–2690
38. Harris W. R., Carrano C. J., Cooper S. R., Sofen S. R., Avdeef A. E., McArdle J. V., Raymond K. N (1979) **Coordination chemistry of microbial iron transport compounds. 19. Stability constants and electrochemical behavior of ferric enterobactin and model complexes** *J. Am. Chem. Soc.* **101**:6097–6104
39. Ghysels B., Dieu B. T. M., Beatson S. A., Pirnay J.-P., Ochsner U. A., Vasil M. L., Cornelis P (2004) **2004. FpvB, an alternative type I ferripyoverdine receptor of *Pseudomonas aeruginosa*** *Microbiology* **150**:1671–1680
40. González J., Salvador M., Özkaya Ö., Spick M., Reid K., Costa C., Bailey M. J., Avignone Rossa C., Kümmerli R., Jiménez J. I (2021) **Loss of a pyoverdine secondary receptor in *Pseudomonas aeruginosa* results in a fitter strain suitable for population invasion** *ISME J* **15**:1330–1343
41. Sexton D. J., Glover R. C., Loper J. E., Schuster M (2017) ***Pseudomonas protegens* Pf-5 favours self-produced siderophore over free-loading in interspecies competition for iron** *Environ. Microbiol* **19**:3514–3525
42. Ménard G., Rouillon A., Ghukasyan G., Emily M., Felden B., Donnio P.-Y (2021) ***Galleria mellonella* Larvae as an Infection Model to Investigate sRNA-Mediated Pathogenesis in *Staphylococcus aureus*** *Front. Cell. Infect. Microbiol* **11**
43. Bohac T. J., Fang L., Giblin D. E., Wencewicz T. A (2019) **Fimsbactin and Acinetobactin Compete for the Periplasmic Siderophore Binding Protein BauB in Pathogenic *Acinetobacter baumannii*** *ACS Chem. Biol* **14**:674–687
44. Shapiro J. A., Wencewicz T. A (2016) **Acinetobactin Isomerization Enables Adaptive Iron Acquisition in *Acinetobacter baumannii* through pH-Triggered Siderophore Swapping** *ACS Infect. Dis* **2**:157–168

45. Sheldon J. R., Skaar E. P (2020) **Acinetobacter baumannii can use multiple siderophores for iron acquisition, but only acinetobactin is required for virulence** *PLOS Pathog* **16**
46. Hammer N. D., Skaar E. P (2011) **Molecular Mechanisms of Staphylococcus aureus Iron Acquisition** *Annu. Rev. Microbiol* **65**:129–147
47. Cassat J. E., Skaar E. P. (2013) **Iron in Infection and Immunity** *Cell Host Microbe* **13**:509–519
48. Skaar E. P (2010) **The Battle for Iron between Bacterial Pathogens and Their Vertebrate Hosts** *PLOS Pathog* **6**
49. Döring G., Pfestorf M., Botzenhart K., Abdallah M. A (1988) **Impact of proteases on iron uptake of Pseudomonas aeruginosa pyoverdine from transferrin and lactoferrin** *Infect. Immun* **56**:291–293
50. Cox C. D (1986) **Role of pyocyanin in the acquisition of iron from transferrin** *Infect. Immun* **52**:263–270
51. Wolz C., Hohloch K., Ocaktan A., Poole K., Evans R. W., Rochel N., Albrecht-Gary A. M., Abdallah M. A., Döring G (1994) **Iron release from transferrin by pyoverdine and elastase from Pseudomonas aeruginosa** *Infect. Immun* **62**:4021–4027
52. Kang D., Kirienko N. V (2021) **An In Vitro Cell Culture Model for Pyoverdine-Mediated Virulence** *Pathogens* **10**
53. Kirienko N. V., Kirienko D. R., Larkins-Ford J., Wählby C., Ruvkun G., Ausubel F. M (2013) **Pseudomonas aeruginosa Disrupts Caenorhabditis elegans Iron Homeostasis, Causing a Hypoxic Response and Death** *Cell Host Microbe* **13**:406–416
54. Pita-Grisanti V., Chasser K., Sobol T., Cruz-Monserrate Z (2022) **Understanding the Potential and Risk of Bacterial Siderophores in Cancer** *Front. Oncol* **12**
55. Peek M. E., Bhatnagar A., McCarty N. A., Zughaier S. M (2012) **Pyoverdine, the Major Siderophore in Pseudomonas aeruginosa, Evades NGAL Recognition** *Interdiscip. Perspect. Infect. Dis*
56. Rezzoagli C., Wilson D., Weigert M., Wyder S., Kümmerli R (2018) **Probing the evolutionary robustness of two repurposed drugs targeting iron uptake in Pseudomonas aeruginosa** *Evol. Med. Public Health* **2018**:246–259
57. Sommer M. O. A., Munck C., Toft-Kehler R. V., Andersson D. I (2017) **Prediction of antibiotic resistance: time for a new preclinical paradigm?** *Nat. Rev. Microbiol* **15**:689–696
58. Imperi F., Massai F., Facchini M., Frangipani E., Visaggio D., Leoni L., Bragonzi A., Visca P (2013) **Repurposing the antimycotic drug flucytosine for suppression of Pseudomonas aeruginosa pathogenicity** *Proc. Natl. Acad. Sci* **110**:7458–7463
59. Kelson A. B., Carnevali M., Truong-Le V (2013) **Gallium-based anti-infectives: targeting microbial iron-uptake mechanisms** *Curr. Opin. Pharmacol* **13**:707–716
60. Schalk I. J., Mislin G. L. A (2017) **Bacterial Iron Uptake Pathways: Gates for the Import of Bactericide Compounds** *J. Med. Chem* **60**:4573–4576

61. Thompson M. G., Corey B. W., Si Y., Craft D. W., Zurawski D. V (2012) **Antibacterial Activities of Iron Chelators against Common Nosocomial Pathogens** *Antimicrob. Agents Chemother* **56**:5419–5421
62. Yang J., Long H., Hu Y., Feng Y., McNally A., Zong Z (2021) **Klebsiella oxytoca Complex: Update on Taxonomy, Antimicrobial Resistance, and Virulence** *Clin. Microbiol. Rev* **35**
63. Cunrath O., Geoffroy V. A., Schalk I. J (2016) **Metallome of Pseudomonas aeruginosa: a role for siderophores** *Environ. Microbiol* **18**:3258–3267
64. Sprouffske K., Wagner A (2016) **Growthcurver: An R package for obtaining interpretable metrics from microbial growth curves** *BMC Bioinformatics* **17**
65. Commo F., Bot B. (2016) **nplr: N-Parameter Logistic Regression** *R package version 0*:1–7
66. Therneau T. (2020) **Therneau, T. A Package for Survival Analysis in R. (2020).** at <https://CRAN.R-project.org/package=survival>

Article and author information

Vera Vollenweider

Department of Quantitative Biomedicine, University of Zurich, Zurich, Switzerland
ORCID iD: [0000-0003-2920-3031](https://orcid.org/0000-0003-2920-3031)

Karoline Rehm

Department of Chemistry, University of Zurich, Zurich, Switzerland

Clara Chepkirui

Institute of Microbiology, Eidgenössische Technische Hochschule (ETH) Zurich, Zurich, Switzerland

Manuela Pérez-Berlanga

Department of Quantitative Biomedicine, University of Zurich, Zurich, Switzerland

Magdalini Polymenidou

Department of Quantitative Biomedicine, University of Zurich, Zurich, Switzerland

Jörn Piel

Institute of Microbiology, Eidgenössische Technische Hochschule (ETH) Zurich, Zurich, Switzerland

Laurent Bigler

Department of Chemistry, University of Zurich, Zurich, Switzerland

Rolf Kümmerli

Department of Quantitative Biomedicine, University of Zurich, Zurich, Switzerland

For correspondence: rolf.kuemmerli@uzh.ch

ORCID iD: [0000-0003-4084-6679](https://orcid.org/0000-0003-4084-6679)

Copyright

© 2024, Vollenweider et al.

This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Vaughn Cooper

University of Pittsburgh, Pittsburgh, United States of America

Senior Editor

Wendy Garrett

Harvard T.H. Chan School of Public Health, Boston, United States of America

Reviewer #1 (Public Review):

Summary:

In an era of increasing antibiotic resistance, there is a pressing need for the development of novel sustainable therapies to tackle problematic pathogens. In this study, the authors hypothesize that pyoverdines - metal-chelating compounds produced by fluorescent pseudomonads - can act as antibacterials by locking away iron, thereby arresting pathogen growth. Using biochemical, growth, and virulence assays on 12 opportunistic pathogens strains, the authors demonstrate that pyoverdines induce iron starvation, but this effect was highly context-dependent. This same effect has been demonstrated for plant pathogens, but not for human opportunistic pathogens exposed to natural siderophores. Only those pathogens lacking (1) a matching receptor to take up pyoverdine-bound iron and/or (2) the ability to produce strong iron chelators themselves experienced strong growth arrest. This would suggest that pyoverdines might not be effective against all pathogens, thereby potentially limiting the utility of pyoverdines as global antibacterials.

Strengths:

The work addresses an important and timely question - can pyoverdines be used as an alternative strategy to deal with opportunistic pathogens? In general, the work is well conducted with rigorous biochemical, growth, and virulence assays. The work is clearly written and the findings are supported by high-quality figures.

Weaknesses:

I do not think there are any 'weaknesses' as such. However, it is well known that siderophore production is highly plastic, typically being upregulated in response to metal limitation (as well as toxic metal stress). Did the authors quantify whether pyoverdine supplementation altered siderophore production in the focal pathogens (either through phenotypic assays / transcriptomics)? Could such a phenotypic plastic response result in an increased capacity to scavenge iron from the environment? Importantly, increased expression of siderophores has been shown to enhance pathogen virulence (e.g. Lear et al 2023: increased pyoverdine production is linked with increased virulence in *Pseudomonas aeruginosa*). I really appreciate the amount of work the authors have put into this study, but I would suggest expanding the discussion a bit to include a few sentences on (1) unintentional consequences of pyoverdine treatment (e.g. changes in gene expression and non-siderophore-related mutations (e.g. biofilm formation)) on disease dynamics/pathogen virulence, and (2) the efficacy of siderophore treatment under more natural conditions, i.e. when the pathogens

have to compete with other species in the resident community (i.e. any other effects than resistance evolution through HGT of pyoverdine receptors as mentioned).

- <https://doi.org/10.7554/eLife.92493.1.sa1>

Reviewer #2 (Public Review):

In this work, Vollenweider et al. examine the effectiveness of using natural products, specifically molecules that chelate iron, to treat infectious agents. Through the purification of 320 environmental isolates, 25 potential candidates were identified from natural products based on inhibition assays and were further screened. The structural information and chemical composition were determined.

The paper is well-structured and thorough; targeting virulence factors in this manner is a great idea. My enthusiasm is dampened by the mediocre effects of the compounds. The lack of a dose-response curve in the survivability assays suggests a limited scope for these molecules. While it is encouraging that the best survivability occurred at the lowest toxicity level, it opens questions as to how effective such molecules can be. Either the reduction in mortality was offset by using higher concentrations, which was not observed in the compound-alone test, or there is no dose-response curve. The latter would suggest to me that the variation in survivability is not due to the addition of siderophores.

I would also like to see how these molecules compare to other iron-chelating molecules. Desferoxamine is a bacteria-derived siderophore that is FDA-approved. However, it is not used to treat infections. Would the author consider comparing their candidate molecules to well-studied molecules? This also raises questions about the novelty of this work; I think the authors could rephrase the discussion to better reflect that bioprospecting for iron-chelating molecules has previously occurred and been successful.

Finally, I am concerned about the few mutations reported in the resistance study. Looking at the SI, it appears that very few mutations were seen. It is unclear what filtering the authors used to arrive at such a low number of mutations. Even filtering against mutations that were selected by adaptation to the media, it seems low that only a handful of clones had distinct mutations.

This paper has a lot of strengths. The workflow is logical and well-executed; the only significant weakness is the effect of the molecules and the lack of an explanation for a dose-response curve in the survivability assay, especially when compared to the data reported in Figure 3. As the authors describe in lines 214-217.

- <https://doi.org/10.7554/eLife.92493.1.sa0>