

mitoBK_{Ca} is functionally expressed in murine and human breast cancer cells and promotes metabolic reprogramming

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

[About eLife's process](#)

Reviewed preprint posted

December 11, 2023 (this version)

Posted to bioRxiv

October 4, 2023

Sent for peer review

September 21, 2023

Helmut Bischof, Selina Maier, Piotr Koprowski, Bogusz Kulawiak, Sandra Burgstaller, Joanna Jasińska, Kristian Serafimov, Dominic Gross, Werner Schroth, Lucas Matt, David Arturo Juarez Lopez, Irina Bonzheim, Florian A. Büttner, Falko Fend, Matthias Schwab, Andreas L. Birkenfeld, Roland Malli, Michael Lämmerhofer, Piotr Bednarczyk, Adam Szewczyk, Robert Lukowski 

Department of Pharmacology, Toxicology and Clinical Pharmacy, Institute of Pharmacy, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany • Dr Margarete Fischer-Bosch Institute of Clinical Pharmacology, Auerbachstraße 112, 70376 Stuttgart, Germany • Laboratory of Intracellular Ion Channels, Nencki Institute of Experimental Biology, Polish Academy of Sciences, 3 Pasteur Street, 02-093 Warsaw, Poland • NMI Natural and Medical Sciences Institute at the University of Tuebingen, Markwiesenstraße 55, Reutlingen 72770, Germany • Institute of Pharmaceutical Sciences, Pharmaceutical (Bio-)Analysis, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany • Medical Clinic IV, University Hospital Tuebingen, Otfried-Müller-Straße 10, 72076 Tuebingen, Germany • Institute of Pathology and Neuropathology, University Hospital Tuebingen, 72076 Tuebingen, Germany • Image-Guided and Functionally Instructed Tumor Therapies Cluster of Excellence (iFIT), University of Tuebingen, Roentgenweg 11, 72076 Tuebingen, Germany • Department of Clinical Pharmacology, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany • Department of Biochemistry and Pharmacy, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany • German Cancer Consortium (DKTK), German Cancer Research Center, Partner Site Tuebingen, Auf der Morgenstelle 15, 72076 Tuebingen, Germany • Gottfried Schatz Research Center, Molecular Biology and Biochemistry, Medical University of Graz, Neue Stiftingtalstraße 6, 8010 Graz, Austria • BioTechMed Graz, Mozartgasse 12, 8010 Graz, Austria • Department of Physics and Biophysics, Warsaw University of Life Sciences (SGGW), Nowoursynowska St. 159 02-776 Warsaw, Poland

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Summary

Alterations in the function of K⁺ channels such as the voltage- and Ca²⁺ activated K⁺ channel of large conductance (BK_{Ca}) reportedly promote breast cancer (BC) development and progression. Underlying molecular mechanisms remain, however, elusive. Here, we provide electrophysiological evidence for a BK_{Ca} splice variant localized to the inner mitochondrial membrane of murine and human BC cells (mitoBK_{Ca}). Through a combination of genetic knockdown and knockout along with cell permeable BK_{Ca} channel blocker, we show that mitoBK_{Ca} modulates overall cellular and mitochondrial energy production and mediates the metabolic rewiring referred to as the “Warburg effect”, thereby promoting BC cell proliferation in the presence and absence of oxygen. Additionally, we detect mitoBK_{Ca} and

BK_{Ca} transcripts in low or high abundance, respectively, in clinical BC specimens. Together, our results emphasize, that targeting mitoBK_{Ca}, combined with established anti-cancer approaches, could represent a novel treatment strategy for selected BC patients.

eLife assessment

The large-conductance Ca²⁺ activated K⁺ channel has been reported to promote breast cancer progression. The present study presents **convincing** evidence that an intracellular subpopulation of this channel reprograms breast cancer cells towards the Warburg phenotype, one of the metabolic hallmarks of cancer. This **important** finding advances the field of cancer cell metabolism and has potential therapeutic implications. However, additional experiments are needed to ascribe the metabolic reprogramming to BK channels located in mitochondria.

Introduction

Cancer represents a complex disease characterized by unconstrained cell proliferation and the spread of malignant cells in the body (Kalia, 2015 [↗](#); Seyfried and Shelton, 2010 [↗](#)). It is one of the leading causes of death worldwide, with millions of new cases diagnosed each year (Sung et al., 2021 [↗](#)). Globally, the most prevalent form of cancer represents breast cancer (BC) (Sung et al., 2021 [↗](#)). Despite many available anti-cancer treatments which largely depend on the steroid and epidermal growth factor (HER2) receptor status (Dunnwald et al., 2007 [↗](#)), cancer cells frequently escape from existing therapies due to adaptations (P. Wu et al., 2021 [↗](#)). Therefore, the identification of novel targets and therapeutic strategies that confer benefits for at least a subset of patients, whose cancer displays specific molecular or cellular features, is of utmost relevance.

Important factors that emerged as new cancer targets are ion channels (Li and Xiong, 2011 [↗](#)). Especially alterations in the expression levels and function of potassium ion (K⁺) channels are critically related to cancer malignancy and progression (Li et al., 2023 [↗](#), p. 0). One of these channels represents the calcium ion (Ca²⁺) and voltage-activated K⁺ channel of large conductance (BK_{Ca}) (Mohr et al., 2022 [↗](#)). Canonical BK_{Ca} channels usually localize in the plasma membrane (PM) of cells, and contribute to the regulation of the cytosolic K⁺ content, the PM potential ($\Delta\Psi_{PM}$), cell cycle, -volume, and -motility (Burgstaller et al., 2022a [↗](#)). Opening of BK_{Ca} channels results in K⁺ efflux, increasing the electrochemical driving force for Ca²⁺ entry into the cancer cell and affecting pathological cell growth and death (Ouadid-Ahidouch and Ahidouch, 2013 [↗](#)). Accordingly, in BC cells (BCCs), an upregulation of BK_{Ca} has been associated with increased malignancy (Huang and Jan, 2014 [↗](#); Mohr et al., 2022 [↗](#); Oeggerli et al., 2012 [↗](#)). However, besides their localization in the PM, several K⁺ channels, including BK_{Ca}, have also been identified in the inner mitochondrial membrane (IMM) (mitoBK_{Ca}), a topic which has been extensively reviewed recently (Checchetto et al., 2021 [↗](#); Kulawiak and Szewczyk, 2022 [↗](#); Szabo and Szewczyk, 2023 [↗](#); Szewczyk et al., 2009 [↗](#); Wrzosek et al., 2020). mitoBK_{Ca} was first described by Siemen et al. in a human glioma cell line more than 20 years ago (Siemen et al., 1999 [↗](#)). So far, mitoBK_{Ca} has further been found e.g. in bronchial epithelial cells (Dabrowska et al., 2022 [↗](#)), neurons (Douglas et al., 2006 [↗](#)), skeletal muscle cells (Skalska et al., 2008 [↗](#)), and in cardiac myocytes (Xu et al., 2002 [↗](#)). In the latter, a splice variant, the BK_{Ca}-DEC isoform containing a unique C-terminal exon of 50 amino acids forms the functional mitoBK_{Ca} channel at the IMM (Singh et al., 2013 [↗](#)). Little, however, is known about the molecular identity of mitoBK_{Ca} in other cell types, and mitochondrial localization of BK_{Ca} in BCCs has not been demonstrated so far.

Cancer cells show increased energy demands due to their high proliferation rates. Thus, tumor cells compensate for their elevated energy demand by increasing metabolic activities and adapting to nutrient-poor metabolic niches in the tumor microenvironment, allowing them to overcome oxygen (O_2)-dependent mitochondrial metabolism (Eales et al., 2016; Gross et al., 2022; Jang et al., 2013; Nazemi and Rainero, 2020). This metabolic switch from oxidative phosphorylation to glycolysis, frequently referred to as the “Warburg effect”, describes the phenomenon that cancer cells rather secrete lactate to the extracellular matrix (ECM), instead of utilizing pyruvate to fuel the TCA cycle (Warburg, 1924). This lactate secretion towards the ECM was shown to promote multiple microenvironmental cues causing tumor progression (de la Cruz-López et al., 2019). Interestingly, extracellular K^+ may also impair effector T-cell function, and, thus, the anti-tumor immune response (Eil et al., 2016), while, within the cancer cell, functions of several glycolytic enzymes rely on K^+ (Bischof et al., 2021; Gohara and Di Cera, 2016). Further, the presence of mitoBK_{Ca} contributes to the K^+ entry into the mitochondrial matrix to interfere with mitochondrial volume changes and the mitochondrial membrane potential ($\Delta\Psi_{mito}$) (Krabbendam et al., 2018). Consequently, (mito)/BK_{Ca}-dependent mechanisms may severely affect energy production pathways and the resulting energy supply of cancer cells.

To elucidate how BK_{Ca} contributes to increasing BCC malignancy (Oeggerli et al., 2012), we utilized BK_{Ca} pro- and deficient human BCC lines including MDA-MB-453 and MCF-7 cells, and primary murine BCCs derived from the mouse mammary tumor polyoma middle T-antigen (MMTV-PyMT)-induced wild type (WT) or BK_{Ca} knock-out (BK-KO) BC model (Mohr et al., 2022). In these cells, either isoforms of BK_{Ca} were expressed, or BK_{Ca} was pharmacologically blocked by paxilline or iberiotoxin, two frequently used inhibitors of BK_{Ca} that either penetrate the cell or act exclusively at PM localized channels, respectively (Candia et al., 1992; Zhou and Lingle, 2014). These approaches were complemented by patch-clamp experiments, and by measuring the cellular ion homeostasis and metabolism with fluorescence live-cell imaging, extracellular flux analysis and liquid chromatography-mass spectrometry (LC-MS)-based approaches.

Our results emphasize that the presence of BK_{Ca} in the IMM affects mitochondrial bioenergetics, thereby increasing BCC malignancy. This effect was specifically mediated by the DEC isoform of BK_{Ca}, i.e., mitoBK_{Ca}. Functional expression of BK_{Ca}-DEC was validated by single-channel patch-clamp recordings BCC-derived mitoplasts. Importantly, we also identified BK_{Ca}-DEC expression in a subset of BC patient biopsies using nanostring-based mRNA expression analysis. Finally, mitoBK_{Ca} crucially contributed to murine and human BCC proliferation and hypoxic resistance, and its activity increased lactate secretion resulting in higher extracellular acidification rates, even in the presence of O_2 .

Combined, our analyses provide, for the first time, a mechanistic link between functionally relevant mitochondrial BKCa isoforms in cancer cells and the promotion of the Warburg effect.

Results

Functional characterization of BKCa expression in BCCs

First, we assessed the functional expression of BK_{Ca} in established human BCC lines by performing whole-cell patch-clamp experiments. Primary BCCs obtained from transgenic, BC-bearing MMTV-PyMT WT or BK-KO mice were used as positive or negative controls, respectively (Mohr et al., 2022). Patch-clamp experiments revealed that depolarizing stimuli delivered in 20 mV increments induced K^+ outward currents that were larger in MMTV-PyMT WT compared to BK-KO cells (Figs. 1A and 1B, and Figs. S1A and S1B). To validate that these increased currents were elicited by BK_{Ca}, we pharmacologically inhibited the channel by using either paxilline or iberiotoxin (Candia et al., 1992; Zhou and Lingle, 2014). While the current remained unaffected by these treatments in MMTV-PyMT BK-KO cells (Fig. 1B and Fig. S1B), peak

currents ($I_{\max}$) were drastically reduced in MMTV-PyMT WT cells (Fig. 1A and Fig. S1A). Further, we analyzed the plasma membrane potential ($\Delta\Psi_{\text{PM}}$) in these cells using the voltage-sensitive fluorescent dye Dibac4(3) (Adams and Levin, 2012). The $\Delta\Psi_{\text{PM}}$ was more polarized in MMTV-PyMT WT compared to BK-KO cells (Fig. 1C), as expected, due to the presence of hyperpolarizing BK_{Ca} channels in WT cells (N'Gouemo, 2014). Similar current-voltage relationships were recorded using the human BCC lines MDA-MB-453 and MCF-7, expressing either high or low levels of BK_{Ca} mRNA transcripts, respectively (Mohr et al., 2022). Analysis of the outward currents activated by depolarization demonstrated a paxilline- and iberiotoxin-sensitive current in MDA-MB-453 cells (Fig. 1D and Fig. S1C), indicative for functional BK_{Ca} channels in their PM, but not in MCF-7 cells (Fig. 1E and Fig. S1D), which is well in line with previous studies (Mohr et al., 2022). Based on the almost non-existent level of BK_{Ca} -mediated PM currents in MCF-7 cells, we utilized these cells to express different BK_{Ca} isoforms fused to a red fluorescent protein (RFP). Therefore, we either used the recently identified mito BK_{Ca} isoform (Singh et al., 2013), namely $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$, or the same channel lacking the C-terminal amino acids, referred to as BK^{RFP} (Fig. 1F).

We hypothesized, that, in analogy to cardiac myocytes (Singh et al., 2013), the expression of $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$ in MCF-7 cells may yield a functional channel present in the IMM. To test this, MCF-7 cells were first transiently transfected either with RFP fused to a glycosylphosphatidylinositol (GPI)-anchor (RFP-GPI) directing RFP to the PM, or with a cytochrome c oxidase subunit 8 (COX8) mitochondrial leading sequence fusion protein (mtRFP) (Fig. S1E). Analysis of the colocalization of mtRFP with MitoGREEN, a dye that specifically stains mitochondria, resulted in high colocalization scores, while a low overlap was observed for RFP-GPI transfected MCF-7 cells (Fig. S1E). Subsequently, the same experiments were performed with MCF-7 cells expressing BK^{RFP} or $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$. Although the mitochondrial localization score was lower for $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$ compared to mtRFP, this BK_{Ca} variant showed a significantly higher overlap with the mitochondrial dye than $\text{BK}_{\text{Ca}}\text{RFP}$ (Fig. 1G). Nevertheless, as not all RFP signal in the $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$ transfected MCF-7 originated from mitochondria, we investigated the PM localization of the two channel isoforms by patch-clamp. Compared to native MCF-7 cells (Fig. 1E), both, BK^{RFP} and $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$, increased the PM outward current (Fig. 1H and Figs. S1F and S1G). Despite comparable expression levels of the RFP signals (Fig. S1H), presence of BK^{RFP} caused significantly bigger currents across the PM compared to $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$ (Fig. S1H), indicative either for i) higher intracellular abundance or ii) major functional differences of $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$. Importantly, the conductance of both channels was sensitive to the BK_{Ca} blockers paxilline and iberiotoxin, yielding comparable PM conductances to native MCF-7 cells (Figs. 1E and 1H and Figs. S1F and S1G).

BK_{Ca} modulates global and subcellular Ca^{2+} homeostasis in BCCs

As BK_{Ca} potentially affects cellular Ca^{2+} fluxes (Ouadid-Ahidouch and Ahidouch, 2013), we next investigated the cytosolic (Figs. 2A-C), endoplasmic reticulum (ER) (Figs. 2D-F), and mitochondrial (Figs. 2G-I) Ca^{2+} homeostasis in these cells. First, we assessed changes in the cytosolic Ca^{2+} concentration ($[\text{Ca}^{2+}]_{\text{cyto}}$) over-time in response to cell stimulation with the purinergic receptor agonist adenosine-5'- triphosphate (ATP) (Müller et al., 2020) using the fluorescent Ca^{2+} indicator Fura-2 (Grynkiewicz et al., 1985), either under control conditions (Fig. 2A) or in the presence of paxilline (Fig. S2A) or iberiotoxin (Fig. S2B). Analysis of the Fura-2 fluorescence emission ratio showed significantly elevated basal (Fig. 2A) and ATP-elicited maximal $[\text{Ca}^{2+}]_{\text{cyto}}$ responses (Fig. 2C) in MMTV-PyMT WT compared to BK-KO cells under control conditions. Interestingly, the elevated basal $[\text{Ca}^{2+}]_{\text{cyto}}$ of WT cells was reduced to BK-KO cell levels in response to paxilline (Fig. 2A and Fig. S2A), but not by iberiotoxin (Fig. 2A and Fig. S2B), which is a peptide-based pore-blocking toxin that cannot pass the PM (Candia et al., 1992). Subsequently, basal and ATP-elicited maximal $[\text{Ca}^{2+}]_{\text{cyto}}$ transients were recorded in the human BCC line MDA-MB-453, where both BK_{Ca} blockers reduced the basal $[\text{Ca}^{2+}]_{\text{cyto}}$ (Fig. 2B), while maximal $[\text{Ca}^{2+}]_{\text{cyto}}$ was not altered by these treatments (Fig. S2D). Eventually,




Figure 1

Characterization of BKCa channels in murine and human BCCs.

(A; B) I-V curves (left) and corresponding maximal currents (right) of MMTV-PyMT WT (A) and MMTV-PyMT BK-KO cells (B), either under control conditions, or in the presence of paxilline or iberiotoxin. Data represents average $\pm$ SEM. N (cells) = 15 WT ctrl, 17 WT + PAX, 17 WT + IBTX, 16 BK-KO ctrl, 17 BK-KO + PAX, 19 BK-KO + IBTX. *** $p \leq 0.001$, Brown-Forsythe and Welch ANOVA test followed by Games-Howell's multiple comparison test. $\dagger p \leq 0.001$ compared to respective WT condition, Welch's t-test. (C) Representative fluorescence images (left) and statistics (right) of MMTV-PyMT WT and BK-KO cells loaded with the $\Delta\Psi_{PM}$ sensitive dye DIBAC4(3). N = 6 independent experiments, ** $p \leq 0.01$, Unpaired t-test. (D) I-V curves (left) and maximal currents (right) of MDA-MB-453 cells, either under control conditions, or in the presence of paxilline or iberiotoxin. Data represents average $\pm$ SEM. N (cells) = 30 ctrl, 22 + PAX, 24 + IBTX. *** $p \leq 0.001$, Kruskal-Wallis test followed by Dunn's multiple comparison test. (E) I-V curves (left) and maximal currents (right) of MCF-7 cells, either under control conditions, or in the presence of paxilline or iberiotoxin. Data shows average $\pm$ SEM. N (cells) = 16 ctrl, 20 + PAX, 15 + IBTX. (F) Schematic representation of constructs used for over-expression in MCF-7 cells. The DEC exon is indicated in green. (G) Representative images (left) of MCF-7 cells either expressing BK^{RFP} (upper) or BK_{Ca}-DEC^{RFP} (lower), additionally stained with MitoGREEN. Average Pearson correlation $\pm$ SEM of MitoGREEN and RFP of BK_{Ca} or BK_{Ca}-DEC are shown. N (cells) = 18 BK_{Ca} - RFP, 23 BK_{Ca}-DEC^{RFP}. ** $p \leq 0.01$, Unpaired t-test. (H) I-V curves (left and middle) and corresponding maximal currents (right) of MCF-7 cells expressing BK^{RFP} (left) or BK_{Ca}-DEC^{RFP} (middle), respectively, either under control conditions, or in the presence of paxilline or iberiotoxin. Data represents average $\pm$ SEM. N (cells) = 18 BK^{RFP} ctrl, 14 BK^{RFP} + PAX, 19 BK^{RFP} + IBTX, 18 BK_{Ca}-DEC^{RFP} ctrl, 21 BK_{Ca}-DEC^{RFP} + PAX, 18 BK_{Ca}-DEC^{RFP} + IBTX. ** $p \leq 0.01$, *** $p \leq 0.001$, Brown-Forsythe and Welch ANOVA test followed by Games-Howell's multiple comparison test. $\dagger p \leq 0.01$ between ctrl conditions, Welch's t-test.

$[Ca^{2+}]_{cyto}$ was assessed in MCF-7 cells expressing or lacking the different BK_{Ca} isoforms (**Fig. 2C**). These experiments confirmed previous findings with the other BCC lines, as the expression of both splice variants increased the basal (**Fig. 2C**) and maximal $[Ca^{2+}]_{cyto}$ (**Fig. S2E**), although the effect was clearly more pronounced upon expression of $BK_{Ca}-DEC^{RFP}$.

Next, we visualized changes in the ER $[Ca^{2+}]$ ($[Ca^{2+}]_{ER}$). Therefore, BCCs were transfected with D1ER, an established genetically encoded, FRET-based Ca^{2+} sensor targeted to the lumen of the ER (Palmer et al., 2004). $[Ca^{2+}]_{ER}$ was depleted by extracellular Ca^{2+} removal and chelation by EGTA, followed by inhibition of the sarcoplasmic endoplasmic reticulum Ca^{2+} ATPase (SERCA) with BHQ and activation of inositol 1,4,5-trisphosphate (IP_3) receptors upon purinergic receptor stimulation with ATP (Lape et al., 2008; Müller et al., 2020; Salter and Hicks, 1995). Experiments were either performed under control conditions (**Figs. 2D-F**), or in the presence of paxilline or iberiotoxin for MMTV-PyMT (**Fig. 2D** and **Figs. S2F** and **S2G**), MDA-MB-453 (**Fig. 2E**), and MCF-7 cells expressing BK_{Ca}^{RFP} or $BK_{Ca}-DEC^{RFP}$ (**Fig. 2F**). Throughout all BCCs investigated, expression of BK_{Ca} reduced $[Ca^{2+}]_{ER}$, potentially indicating that the channel was i.) functional in this cell compartment and ii.) involved in regulating the Ca^{2+} homeostasis of the ER. In MMTV-PyMT WT cells, $[Ca^{2+}]_{ER}$ was restored by the cell-permeable BK_{Ca} inhibitor paxilline (**Fig. 2D** and **Fig. S2F**) but not by the cell-impermeable iberiotoxin (**Fig. 2D** and **Fig. S2G**), while both inhibitors restored the $[Ca^{2+}]_{ER}$ in MDA-MB-453 cells (**Fig. 2e**). Accordingly, overexpression of both RFP-tagged BK_{Ca} isoforms in MCF-7 cells depleted the $[Ca^{2+}]_{ER}$ (**Fig. 2F**).

$[Ca^{2+}]_{cyto}$ and $[Ca^{2+}]_{ER}$ reportedly affect the mitochondrial $[Ca^{2+}]$ ($[Ca^{2+}]_{mito}$) (Wacquier et al., 2019). Since an accumulation of K^+ within the mitochondrial matrix may oppose mitochondrial Ca^{2+} uptake (Checchetto et al., 2021), we assessed the effects of functional BK_{Ca} expression on $[Ca^{2+}]_{mito}$ utilizing 4mtD3cpV, a genetically encoded Ca^{2+} sensor targeted to the mitochondrial matrix (Palmer et al., 2006). Again, BCCs were treated with ATP to investigate the $[Ca^{2+}]_{mito}$ upon cell stimulation. In MMTV-PyMT WT (**Fig. 2G**) as well as MDA-MB-453 cells (**Fig. 2H**), functional expression of BK_{Ca} reduced basal $[Ca^{2+}]_{mito}$. Interestingly, in these two BCC types, basal and maximally elicited $[Ca^{2+}]_{mito}$ peaks increased in response to paxilline (**Figs. 2G** and **2H** and **Figs. S2H**, **S2J** and **S2K**), but not iberiotoxin treatment (**Figs. 2G** and **2H** and **Figs. S2I-K**). These findings confirm a role of intracellular located BK_{Ca} channels in modulating $[Ca^{2+}]_{mito}$ dynamics. Importantly, in MCF-7 cells, basal $[Ca^{2+}]_{mito}$ was only affected upon expression of the mitochondrially targeted $BK_{Ca}-DEC^{RFP}$, but not BK^{RFP} (**Fig. 2I**), whereas neither of the two isoforms affected the maximal $[Ca^{2+}]_{mito}$ (**Fig. S2L**). Presumably, by facilitating K^+ fluxes across the IMM, $BK_{Ca}-DEC^{RFP}$ expression reduces the driving force for Ca^{2+} uptake and thus the resulting Ca^{2+} signals in the mitochondrial matrix.

In total, our $[Ca^{2+}]$ imaging approaches emphasize that different BK_{Ca} isoforms at different localizations, i.e. the PM or intracellular organelles, may either amplify or weaken $[Ca^{2+}]_{cyto}$ and $[Ca^{2+}]_{ER}$ signals, respectively. These opposing effects are expected as BK_{Ca} -mediated uptake of K^+ into the ER should limit the Ca^{2+} uptake capacity of this subcellular compartment, while functional channel expression at the PM might increase the driving force for Ca^{2+} influx to fuel $[Ca^{2+}]_{cyto}$. $[Ca^{2+}]_{mito}$, however, seems to be exclusively and effectively suppressed by the activity of intracellularly located BK_{Ca} variants in MMTV-PyMT WT and MDA-MB-453 cells and solely by the expression of $BK_{Ca}-DEC^{RFP}$ in MCF-7 cells.

The metabolic activity of murine and human BCCs is modulated by intracellular BK_{Ca}

Given the involvement of Ca^{2+} and K^+ in regulating key features of cellular metabolism (Bischof et al., 2021; Dejos et al., 2020; Gohara and Di Cera, 2016), we proposed, that BK_{Ca} may alter metabolic activities of BCCs (Rossi et al., 2019). Hence, we analyzed the extracellular acidification rate (ECAR) as a measure of lactate secretion, i.e. glycolytic activity in MMTV-PyMT and MDA-MB-453 cells (Burgstaller et al., 2021a). ECAR measurements unveiled an increased




Figure 2

BkCa modulates the subcellular Ca^{2+} homeostasis in BCCs.

Cytosolic (A – C), endoplasmic reticulum (ER) (D – F), and mitochondrial Ca^{2+} dynamics (G – I) over-time of MMTV-PyMT WT and MMTV-PyMT BK-KO cells (A, D, G), MDA-MB-453 cells (B, E, H) or MCF-7 cells (C, F, I). All data represent average $\pm$ SEM. At time points indicated in the panels, cytosolic and mitochondrial Ca^{2+} alterations were evoked by extracellular stimulation with ATP (A – C, G – I), or by Ca^{2+} depletion of the ER using Ca^{2+} -free buffer containing the Ca^{2+} chelator EGTA, followed by administration of the SERCA inhibitor BHQ prior to ATP administration (D – F). MMTV-PyMT (A, D, G) and MDA-MB-453 (B, E, H) cells were either measured under control conditions, or in the presence of paxilline (+ PAX) or iberiotoxin (+ IBTX). MCF-7 cells (C, F, I) either expressed an RFP (ctrl), $\text{BK}_{\text{Ca}}^{\text{RFP}}$, or $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$. N (independent experiments) = A: 17 WT ctrl, 18 BK-KO ctrl, 6 WT + PAX and BK-KO + PAX, 5 WT + IBTX and BK-KO + IBTX, B: 4 for all, C: 5 for all, D: 14 WT ctrl, 13 BK-KO ctrl, 8 WT + PAX and BK-KO + PAX, 6 WT + IBTX and BK-KO + IBTX, E: 7 ctrl, 9 + PAX, 5 + IBTX, F: 4 for all, G: 11 WT ctrl, 12 BK-KO ctrl, 6 WT + PAX and BK-KO + PAX, 5 WT + IBTX, 4 BK-KO + IBTX, H: 8 ctrl and + PAX, 5 + IBTX, I: 5 ctrl, 4 + $\text{BK}_{\text{Ca}}^{\text{RFP}}$ and $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$. ** $p \leq 0.01$, *** $p \leq 0.001$. Kruskal-Wallis test followed by Dunn's MC test (A, B, C, D, I), One-Way ANOVA test followed by Tukey's MC test (E, F, G) or Brown-Forsythe ANOVA test followed by Games-Howell's MC test (H). # $p \leq 0.05$, # $p \leq 0.001$ compared to respective WT condition, Mann-Whitney test (A, D, ctrl in G), or Welch's t test (+ IBTX in G).

basal ECAR of MMTV-PyMT WT compared to BK- KO cells (**Figs. 3A** and **3B**). Subsequently, we performed a mitochondrial stress test by injection of Oligomycin-A, FCCP, and Antimycin-A, which increased ECARs in both cell types due to ATP-synthase inhibition, mitochondrial uncoupling and complex III blockade, respectively (**Fig. 3A**). Maximal ECAR, received upon FCCP injection, was elevated in WT compared to BK-KO cells (**Fig. 3C**). Besides this evidence derived from a gene-targeted BK_{Ca} channel-deficient model, we used paxilline or iberiotoxin as pharmacological BK_{Ca} modulators (**Figs. S3A** and **S3B**). Interestingly, only paxilline, but not iberiotoxin treatment equalized basal (**Fig. 3B**) and maximal ECARs (**Fig. 3C**) between WT and BK-KO, suggesting that intracellular BK_{Ca} channels stimulate the glycolytic activity of BCCs. Moreover, in MDA-MB-453 cells iberiotoxin treatment did neither affect basal (**Fig. 3D** and **3E**) nor maximal ECARs (**Fig. 3F**). Paxilline, however, increased basal and maximal ECAR in these cells (**Figs. 3E** and **3F**).

Next, we investigated the oxygen consumption rates (OCRs) (Burgstaller et al., 2021a). (**Fig. 3G**). Interestingly, the increased basal and maximal ECAR (**Figs. 3A-C**) in MMTV-PyMT WT cells correlated with a higher basal (**Fig. 3H**) and maximal (**Fig. 3I**) OCR compared to BCCs lacking BK_{Ca}. Paxilline treatment reduced the basal and maximal OCR of BK_{Ca} proficient MMTV-PyMT cells to the BK-KO level (**Figs. 3H** and **3I** and **Fig. S3C**), while iberiotoxin did not have any effect (**Figs. 3H** and **3I** and **Fig. S3D**) again suggesting that the latter toxin cannot reach the metabolism-relevant population of intracellular BK_{Ca} channels. Similar results were obtained in MDA-MB-453 cells, despite in these cells paxilline treatment increased the OCR (**Figs. 3J-L**), potentially due to the increased supply of oxidative phosphorylation with glycolytic substrates (**Figs. 3D-F**), while iberiotoxin had no effect.

To confirm that BK_{Ca} regulates the bioenergetic profile of BCCs, we subsequently applied LC-MS-based metabolomics according to the workflow presented in **Figure S3E**. We included typical analytes of glycolysis such as lactate (**Fig. 3M**) and pyruvate (**Fig. 3N**), as well as selected metabolites of mitochondrial metabolism including asparagine (**Fig. 3O**) and glutamine (**Fig. 3P**). Metabolomics data confirmed the metabolic differences between MMTV-PyMT WT and BK-KO cells and further demonstrated that BK_{Ca} inhibition by paxilline directly affected the concentrations of these metabolites (**Figs. 3M-P**). As observed before for the ECAR and OCR measurements, MMTV-PyMT and MDA-MB-453 cells responded, however, differently to paxilline treatment (**Figs. 3M-P**). In summary, our data strengthen the notion that intracellular BK_{Ca} modulates the cellular energy balance of murine and human BCCs.

BK_{Ca} alters the mitochondrial function of BCCs

To clarify how BK_{Ca} regulates BCC cell metabolic activities, we examined cellular bioenergetics in real time using single-cell fluorescence microscopy. First, we assessed the mitochondrial membrane potential ($\Delta\Psi_{\text{mito}}$) of MMTV-PyMT and MDA-MB-453 cells using TMRM under control and BK_{Ca} inhibitor-treated conditions, followed by depolarization of $\Delta\Psi_{\text{mito}}$ using the proton ionophore FCCP (Joshi and Bakowska, 2011). These measurements unveiled, however, a less polarized $\Delta\Psi_{\text{mito}}$ of MMTV-PyMT WT cells compared to BK-KO cells under control conditions (**Figs. 4A** and **4B**). Paxilline, but not iberiotoxin, equalized $\Delta\Psi_{\text{mito}}$ between MMTV-PyMT WT and BK-KO cells (**Fig. 4B**). Identical results were obtained in MDA-MB-453 cells (**Figs. 4C** and **4D**).

Interestingly, we found that $\Delta\Psi_{\text{mito}}$ of MMTV-PyMT cells was tightly dependent on the extracellular glucose concentration ($[\text{GLU}]_{\text{ex}}$) (**Fig. S4A**). Under high $[\text{GLU}]_{\text{ex}}$ conditions (25.0 mM), the difference in $\Delta\Psi_{\text{mito}}$ between MMTV-PyMT WT and BK-KO cells disappeared, as $\Delta\Psi_{\text{mito}}$ increased significantly in WT and decreased significantly in BK-KO cells compared to low $[\text{GLU}]_{\text{ex}}$ (2.0 mM). Because F₀F₁-ATP- synthase (ATP Synthase) may hydrolyze ATP in an attempt to




Figure 3

BKCa channels promote the metabolic activity of BCCs.

(A, D) Average ECAR over-time ± SEM of MMTV-PyMT WT (A, red) and BK-KO cells (A, black), or MDA-MB-453 cells (D) in response to administration of Oligomycin-A (+O), FCCP (+F) and Antimycin-A (+A) at time points indicated. (B, E) Average basal and (C, F) maximal ECAR ± SEM of MMTV-PyMT WT (B, C, left) and BK-KO cells (B, C, right), or MDA-MB-453 cells (E, F) under control conditions, or in the presence of paxilline or iberiotoxin. (G, J) Average OCR over-time ± SEM of MMTV-PyMT WT (G, red) and BK-KO cells (G, black), or MDA-MB-453 cells (J) in response to administration of Oligomycin-A (+O), FCCP (+F) and Antimycin-A (+A) at time points indicated. (H, K) Average basal and (I, L) maximal OCR ± SEM of MMTV-PyMT WT (H, I, left) and BK-KO cells (H, I, right), or MDA-MB-453 cells (K, L) under control conditions, or in the presence of paxilline or iberiotoxin. (M – P) LC-MS-based determination of major glycolytic and mitochondrial metabolites in particular Lactate (M, [Lactate]), Pyruvate (N, [Pyruvate]), Asparagine (O, [Asparagine]) and Glutamine (P, [Glutamine]) of MMTV-PyMT WT (left panels), BK- KO (middle panels) and MDA-MB-453 cells (right panels), either under control conditions or after cell cultivation with paxilline. N (independent experiments) = A, B, C, G, H, I: 7 WT ctrl and BK-KO, 3 for all others, D, E, F, J, K, L: 3 for all, M – P: 7 for BK-KO ctrl, 6 for all others. * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, Kruskal-Wallis test followed by Dunn's MC test (B, E, F, I), Brown-Forsythe and Welch ANOVA test followed by Games-Howell's MC test (C, H), One-Way ANOVA test followed by Tukey's MC test (K, L) or Mann-Whitney test (M – P). # $p \leq 0.05$, † $p \leq 0.01$, ‡ $p \leq 0.001$, to respective WT condition, Mann-Whitney test (B, C, + IBTX in I, M – P), Welch's t-test (ctrl in H and I) or Unpaired t-test (+ IBTX in H).




Figure 4

Expression of BKCa modulates mitochondrial function and glucose uptake of BCCs.

(A – D) Representative fluorescence images and -ratios ($F_{\text{mito}}/F_{\text{nuc}}$) over-time (A, C), and corresponding statistics $\pm$ SEM (B, D) representing $\Delta\Psi_{\text{mito}}$ of TMRM-loaded MMTV-PyMT WT and BK-KO (A, B) and MDA-MB-453 cells (C, D) under basal conditions (A, C, upper images) and upon administration of FCCP for mitochondrial depolarization (A, C lower images). (E – J) $[\text{ATP}]_{\text{mito}}$ dynamics $\pm$ SEM over-time of MMTV-PyMT WT and BK-KO cells (E), MDA-MB-453 cells (G) and MCF-7 cells (I) in response to extracellular glucose removal (left panels) or upon administration of Oligomycin-A (right panels). F, H and J show changes of $[\text{ATP}]_{\text{mito}}$ induced by glucose removal to Oligomycin-A administration $\pm$ SEM, under control conditions, or in the presence of paxilline or iberiotoxin (F, H), or upon expression of BK^{RFP} or BK-DEC^{RFP} (J). (K – M) Basal mitochondrial H₂O₂ concentrations $\pm$ SEM of MMTV-PyMT WT (K, left), BK-KO (K, right), MDA-MB-453 (L) and MCF-7 cells (M), either under control conditions, in the presence of paxilline or iberiotoxin (K, L), or upon expression of BK^{RFP} or BK-DEC^{RFP} (M). (N, P) Representative fluorescence wide-field images (left) and corresponding statistics $\pm$ SEM (right) of MMTV-PyMT WT (N, left images and red bars) and BK-KO cells (N, right images and black bars) or MDA-MB-453 cells (P) incubated with 2-NBDG, either in the absence (upper images) or presence of FCCP (lower images). (O, Q) Average $\pm$ SEM of FCCP induced change in 2-NBDG uptake of MMTV- PyMT WT (O, left) and BK-KO cells (O, right), or MDA-MB-453 cells (Q) either under control conditions, or in the presence of paxilline or iberiotoxin. Values above 1 indicate that mitochondria support glucose uptake, values below 1 that mitochondria prevent glucose uptake. N (independent experiments) = A – D, N – Q: 4 for all, E – J [- Glucose]: 8 WT ctrl, BK-KO ctrl and BK-KO + IBTX, 7 WT + IBTX and BK-KO + PAX and MCF-7 ctrl, 3 MDA-MB-453 + PAX, 5 for all others. E – J [+ Oligomycin-A]: 11 WT ctrl, 8 BK-KO ctrl and MDA-MB-453 + IBTX, 7 WT + PAX and WT + IBTX, 6 BK-KO + PAX, 3 MDA-MB-453 + PAX and 5 for all others. * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, Kruskal-Wallis test followed by Dunn's MC test (B, D, F, H, J, K, O), Brown-Forsythe and Welch ANOVA test followed by Games- Howell's MC test (L, M), Mann-Whitney test (N), Unpaired t-test (P) or Welch's t-test (Q). # $p \leq 0.05$, $\dagger p \leq 0.01$, $\ddagger p \leq 0.001$, to respective WT condition, Mann-Whitney test (B, F, + PAX and + IBTX in K, ctrl in O), Unpaired t-test (ctrl in K, + PAX and + IBTX in O).

maintain $\Delta\Psi_{\text{mito}}$, these results strongly suggest that the forward (ATP producing) vs reverse (ATP consuming) mode of the ATP synthase is affected by the BK_{Ca} status of the cells (Naguib et al., 2018).

To unravel the substrate dependency of these BCCs for maintaining their energy homeostasis in dependency of BK_{Ca} , we measured the mitochondrial ATP concentration ($[\text{ATP}]_{\text{mito}}$) using a genetically encoded ATP sensor targeted to the mitochondrial matrix, mtAT1.03 (Imamura et al., 2009). Mitochondrial ATP reportedly responds most dynamically to energy metabolism perturbations (Depaoli et al., 2018). To assess the sources of ATP in the cells, we either deprived the cells of $[\text{GLU}]_{\text{ex}}$, or administered Oligomycin-A to inhibit the ATP-synthase (Depaoli et al., 2018) (Figs. 4E-J and Figs. S4B-G). In MMTV-PyMT cells, glucose removal as well as ATP-synthase inhibition reduced $[\text{ATP}]_{\text{mito}}$, albeit the effect was more pronounced upon $[\text{GLU}]_{\text{ex}}$ depletion (Figs. 4E and 4F and Figs. S4B and S4C). Remarkably, paxilline, but not iberiotoxin treatment reduced the glucose, and increased ATP-synthase dependency of MMTV-PyMT WT cells for maintaining $[\text{ATP}]_{\text{mito}}$ (Figs. 4E and 4F and Figs. S4B and S4C). These observations were confirmed in MDA-MB-453 cells, even though i.) these cells showed an overall higher dependency on the ATP-synthase, and ii.) the paxilline-sensitivity of $[\text{ATP}]_{\text{mito}}$ maintenance was much less pronounced (Figs. 4G and 4H and Figs. S4D and S4E).

To validate these findings in a BK_{Ca} deficient model, these experiments were performed with MCF-7 cells either expressing exclusively RFP as control, BK^{RFP} , or $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$ (Fig. 4I). Expression of $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$, but not BK^{RFP} , resulted in a high dependency on $[\text{GLU}]_{\text{ex}}$ to maintain $[\text{ATP}]_{\text{mito}}$ while the $[\text{ATP}]_{\text{mito}}$ rundown in the presence of Oligomycin A was identical for both BK_{Ca} splice variants. This suggests that $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$ specifically triggers a high $[\text{GLU}]_{\text{ex}}$ sensitivity and simultaneously an independency on ATP derived from ATP-synthase for maintaining $[\text{ATP}]_{\text{mito}}$ as demonstrated by the ratio of the respective changes under these experimental conditions (Figs. 4I and 4J and Figs. S4F and S4G). These findings suggest that intracellular (mitochondrial) BK_{Ca} contributes to the metabolic reprogramming of BCCs.

In an extension to extracellular flux analyses, LC-MS metabolomics (Fig. 3), and live-cell imaging experiments (Figs. 4A-J), pointing to a BK_{Ca} -dependent “oncometabolic” phenotype, we assessed the concentrations of mitochondrial hydrogen peroxide ($[\text{H}_2\text{O}_2]_{\text{mito}}$) using mitoHyPer3 (Bilan et al., 2013), a genetically encoded fluorescent indicator for monitoring H_2O_2 in the mitochondrial matrix (Figs. 4K-M). These experiments demonstrated increased levels of $[\text{H}_2\text{O}_2]_{\text{mito}}$ in MMTV-PyMT WT (Fig. 4K) and MDA-MB-453 cells (Fig. 4L), or specifically upon $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$ expression in MCF-7 cells (Fig. 4M). Excessive reactive oxygen species (ROS) synthesis is caused by uncoupling of the respiratory chain and it is considered as an indicator of mitochondrial stress, which, among other reasons, promotes mutagenesis and BCC progression. In this regard, mito BK_{Ca} may serve as an “uncoupling” protein (Gałęcka et al., 2021), triggering ATP synthase to operate in reverse mode to consume instead of producing ATP, thereby counteracting the dissipation of the proton gradient and consequently the loss of $\Delta\Psi_{\text{mito}}$. Consequently, the lack of ATP must be compensated, e.g. by accelerating glycolysis. To investigate this assumption, we performed glucose uptake measurements using 2-NBDG, a fluorescent glucose analogue, which is taken up via glucose transporters (GLUTs), phosphorylated by hexokinase (HK) isoforms to generate 2-NBDG-6-phosphate and subsequently remains within the cell (Bischof et al., 2021) (Figs. S4H and S4I). Unexpectedly, we found that MMTV-PyMT BK-KO cells showed a higher rate of glucose uptake and phosphorylation compared to WT cells under basal conditions (Fig. 4N). To unravel the role of mitochondria in contributing to glucose uptake by supplying ATP to HKs, we next performed these experiments in the presence of FCCP, which disrupts mitochondrial ATP production (Losano et al., 2017). If ATP-synthase works in forward mode, FCCP treatment should reduce or even prevent oxidative ATP production, and, subsequently, ATP-dependent 2-NBDG phosphorylation (Fig. S4H). Contrary, if ATP-synthase works in reverse mode, it may compete with HKs for ATP, and FCCP treatment should abolish this competition, leading to increased 2-NBDG phosphorylation in BCCs (Fig. S4I). Interestingly, our experiments

unveiled increased 2-NBDG uptake in MMTV-PyMT WT and reduced uptake in BK-KO cells upon mitochondrial depolarization (Fig. 4N [↗](#) and Figs. S4J [↗](#) and S4K). To facilitate the interpretation, we calculated the FCCP-induced change in 2-NBDG uptake. These analyses revealed that mitochondria are less effective in assisting 2-NBDG uptake and phosphorylation in MMTV-PyMT WT cells, while they “support” these processes in BK-KO cells under control conditions (Fig. 4O [↗](#)). To test whether the changes in 2-NBDG uptake are sensitive to pharmacological modulation of BK_{Ca}, we performed a similar set of experiments in the presence of paxilline or iberiotoxin. While iberiotoxin treatment did not have any effect, paxilline treatment shifted the activity of the ATP-synthase to the ATP-supplying “forward” mode (Fig. 4O [↗](#) and Figs. S4J [↗](#) and S4K). Comparable effects were obtained in MDA-MB-453, despite paxilline treatment reduced basal accumulation of 2-NBDG in these cells (Figs. 4P [↗](#) and 4Q [↗](#) and Fig. S4L [↗](#)). Overall, the experiments performed so far suggest that mitochondria rather consume than generate ATP if BK_{Ca} was functionally expressed intracellularly in BCCs.

BK_{Ca} locates functionally in the IMM of murine and human BCCs

So far, our data suggest an important contribution of intracellularly located BK_{Ca}, possibly mitoBK_{Ca}, in reprogramming cancer cell metabolism. Thus, we applied an electrophysiological approach to provide functional evidence for endogenous mitoBK_{Ca}. Single-channel patch-clamp experiments were conducted using mitoplasts isolated from MMTV-PyMT WT and BK-KO, MDA-MB-453, and MCF-7 cells. In MDA-MB-453- and MMTV-PyMT WT-derived mitoplasts we indeed detected channels of large conductance (Fig. 5A [↗](#) and Fig. S5A [↗](#)). For these channels, the open probability did not significantly vary with the voltage (Fig. 5B [↗](#)). Only at very negative and positive voltages of approximately -150 mV and +150 mV differences in the open probabilities were observed (Fig. 5C [↗](#)). The bursts of single-channel openings showed an average conductance of 212 ± 2 pS (Fig. 5D [↗](#)). This large conductance and the sensitivity towards Ca²⁺ (Figs. 5E [↗](#) and 5F [↗](#) and Fig. S5B [↗](#)) and paxilline (Figs. 5E [↗](#) and 5F [↗](#) and Fig. S5C [↗](#)) pointed to mitoBK_{Ca} that exhibits the pharmacologic characteristics of canonical BK_{Ca} channels present at the PM. Overall, our electro-pharmacological experiments unveiled 3 different classes of channels with either small (≤ 100 pS), medium (≤ 150 pS) or large (~ 210 pS) conductance, where the latter conductivity corresponds to mitoBK_{Ca} (Liu et al., 1999 [↗](#)). MitoBK_{Ca} was detected at a frequency of 15% in 210 patches of MDA-MB-453 mitoplasts and at a lower frequency of 6% in 127 mitoplast patches of MMTV-PyMT WT cells. Importantly, this channel was absent in 76 mitoplast patches from MMTV-PyMT BK-KO and 58 mitoplast patches from MCF-7 cells (Fig. 5G [↗](#)). These findings provide evidence that the molecular entity for the K⁺ channel derives from the nuclear *Kcnma1* gene, which is ablated in the MMTV-PyMT BK-KO.

These experiments were further corroborated by immunoblotting experiments using whole-cell lysates and sub-cellular homogenates of different purity. We detected BK_{Ca}, the Na⁺/K⁺ ATPase as a marker of the PM, and cytochrome c oxidase subunit IV (COXIV) as a marker of mitochondria in these samples. A protein band corresponding to BK_{Ca} was identified not only in whole-cell lysates, but also in isolated mitochondria. Importantly, the Na⁺/K⁺ ATPase was absent in the latter protein fraction, confirming the purity of the mitochondrial preparation (Fig. S5D [↗](#)). These data confirm the presence of mitoBK_{Ca}, potentially BK_{Ca}-DEC, in the utilized BK_{Ca} proficient BCCs.

mitoBK_{Ca} promotes the Warburg effect, triggers cellular O₂ independency and stimulates BCC proliferation

Based on the observed influence of mitoBK_{Ca} on glycolysis and mitochondrial metabolism, we addressed, whether the channel contributes to the Warburg effect, commonly observed in cancer cells (Bischof et al., 2021 [↗](#); Warburg, 1924 [↗](#)). Therefore, we assessed the Warburg index (WI) by investigating the cytosolic lactate concentration ([LAC]_{cyto}) over-time using Laconic, a FRET-based lactate sensor (San Martín et al., 2013). [LAC]_{cyto} was followed in response to either inhibiting mitochondrial metabolism by administration of NaN₃, a complex IV inhibitor, to stop pyruvate




Figure 5

BKCa activity is present in the inner mitochondrial membrane (IMM) of BCCs.

(A) Representative BK_{Ca} single-channel recordings of the IMM of mitoplasts isolated from MDA-MB-453 cells using a symmetric 150/150 mM isotonic KCl solution containing 100 μM Ca^{2+} at voltages ranging from -80 to +80 mV as indicated in the panel. (B) Open probability analysis of $\text{mitoBK}_{\text{Ca}}$ at different voltages received from experiments as performed in (A). $N = 8$. (C) Single-channel currents of the IMM of mitoplasts isolated from MDA-MB-453 cells recorded using a voltage ramp protocol ranging from -150 to +150 mV. Above and below the ramp are enlarged excerpts of the records shown in rectangles. (D) Current-voltage (I-V) plot based on single-channel recordings of MDA-MB-453 cells as performed in a, using a symmetric 150/150 mM KCl isotonic solution containing 100 μM Ca^{2+} . $N = 11$. (E, F) Representative single-channel recordings of the IMM of mitoplasts isolated from MDA-MB-453 cells (E) and corresponding open probabilities at +40 mV in a symmetric 150/150 mM KCl isotonic solution under control conditions (100 μM Ca^{2+}), after reducing Ca^{2+} to 1 μM , re-addition of 100 μM Ca^{2+} and finally after application of 5 μM paxilline in the presence of 100 μM Ca^{2+} . Data in (F) show average $\pm$ SEM. * $p < 0.05$, ** $p < 0.01$ using Friedman test followed by Dunn's multiple comparison test, $n = 7$. (G) Pie chart displaying the percentage of $\text{mitoBK}_{\text{Ca}}$ channel currents (green) possessing a conductance of ~210 pS, versus the total number of patch-clamp experiments performed using mitoplasts isolated from MDA-MB-453 cells (upper left), MMTV-PyMT WT cells (upper right), MCF-7 cells (lower left) and MMTV-PyMT BK-KO cells (lower right). Black segments represent empty patches, bright- and dark grey fraction demonstrate percentage of channels possessing smaller conductances of ≤ 100 pS and ≤ 150 pS, respectively. All recordings were low-pass filtered at 1 kHz. "c" and "o" indicate the closed- and open state of the channel, respectively.

consumption (Leary et al., 1998 [↗](#)), or upon subsequent inhibition of lactate secretion towards the ECM via monocarboxylate transporter 1 (MCT-1) using BAY-8002 (Quanz et al., 2018 [↗](#)) (Figs. 6A [↗](#) and 6B [↗](#)). Indeed, MMTV-PyMT WT cells exhibited an increased WI compared to BK-KO cells under control conditions (Fig. 6C [↗](#)), indicating that the presence of BK_{Ca} favors lactate secretion rather than TCA-dependent utilization of pyruvate. Paxilline, but not iberiotoxin treatment reduced the WI in WT cells to the BK-KO level (Fig. 6C [↗](#)). The WI profiles of MDA-MB-453 (Fig. 6D [↗](#)) and MCF-7 cells (Fig. 6E [↗](#)) showed the same sensitivity towards paxilline, while in the latter the expression of BK_{Ca}-DEC^{RFP}, but not BK^{RFP}, stimulated the WI. To validate the contribution of the different BK isoforms, endogenous BK_{Ca} transcripts in MMTV-PyMT and MDA-MB-453 cells were targeted using specific siRNAs targeting either the major BK_{Ca} isoforms or specifically the DEC exon of BK_{Ca} (Fig. S6A [↗](#)). Cell treatment with the respective siRNAs reduced the expression of BK_{Ca} or BK_{Ca}-DEC (Fig. S6B [↗](#) and S6C). Interestingly, the knockdown of BK_{Ca} interfered with the WI in these cells (Figs. 6F [↗](#) and 6G [↗](#)). This effect was reproduced by specific silencing of the BK_{Ca}-DEC isoform (Figs. 6F [↗](#) and 6G [↗](#)), indicating that BK_{Ca}-DEC-derived mitoBK_{Ca} channels stimulate the Warburg effect in BCCs.

As glycolytic metabolites and ATP fuel cell proliferation, we investigated the proliferation rates of MMTV-PyMT WT, MMTV-PyMT BK-KO, and MDA-MB-453 cells over-time in the absence or presence of paxilline or iberiotoxin. Corroborating previous findings (Mohr et al., 2022 [↗](#)), these analyses revealed faster proliferation of WT compared to BK-KO cells under control conditions (Fig. 6H [↗](#)). Interestingly, paxilline, but not iberiotoxin treatment, reduced the proliferation of MMTV-PyMT WT to the level of BK_{Ca}-deficient cells (Fig. 6H [↗](#)). These findings were confirmed in MDA-MB-453 cells (Fig. 6I [↗](#)), suggesting mitoBK_{Ca} as key-player in mediating the metabolic rewiring. Next, we studied whether the increased WI affects the proliferation rate of these BCCs at low O₂ tension (Nazemi and Rainero, 2020 [↗](#)). Colony formation assays (CFAs) revealed an increased hypoxic resistance of MMTV-PyMT WT compared to BK-KO cells, as the number and size of colonies lacking BK_{Ca} was reduced upon O₂- deprivation (Fig. 6J [↗](#) and Fig. S6D [↗](#)). In sum, these data demonstrate a malignancy-promoting effect of mitoBK_{Ca} in BCC.

mitoBK_{Ca} is of potential clinical relevance

To determine the clinical relevance of our findings, we investigated whether BK_{Ca}-DEC transcripts are present in primary BC tissue. Therefore, the mRNA expression of BK_{Ca} and BK_{Ca}-DEC was analyzed by nanostring analysis in bulk tumor biopsies isolated from 551 BC patients. Remarkably, all 551 samples tested positive for BK_{Ca} mRNA expression, with 10 of the samples showing significant expression of BK_{Ca}-DEC above the log2 expression threshold of 5.5 (Figs. 6K [↗](#) and 6L [↗](#)). Importantly, if BK_{Ca}-DEC was expressed, the expression of BK_{Ca} well correlated with the expression of BK_{Ca}-DEC ($R^2 = 0.6060$, $p = 0.008$) (Fig. 6M [↗](#)).

In sum, our experiments emphasize the presence of BK_{Ca} in different intracellular organelles including the ER and vesicles of the secretory pathway, yielding its PM localization. At these sites, BK_{Ca} modulates the Ca²⁺ homeostasis and regulates $\Delta\Psi_{PM}$. K⁺ efflux across the PM may additionally affect glycolysis. Importantly, functionally relevant BK_{Ca} also locates in the IMM of BCCs, promoting, presumably by the K⁺ accumulation in the matrix following channel activation, ΔY_{mito} depolarization and consequently ATP- synthase activity in reverse mode as well as a depletion of [Ca²⁺]_{mito}. These profound ionic and bioenergetic changes ultimately trigger the proliferation of BCCs in a low oxygen environment, as found in solid tumors (Fig. 6N [↗](#)). Taken together, functional expression of mitoBK_{Ca} could possibly denote a prognostic or therapeutic marker for BC patients, and its pharmacologic modulation could represent a novel anti-cancer treatment strategy.




Figure 6

BKC_a-DEC expression contributes to the metabolic remodeling and growth of murine and human BCCs and is present in primary tumor samples.

(A) Schematic representation of the fate of glucose in glycolysis. The tricarboxylic acid (TCA) cycle or lactate secretion via monocarboxylate transporters (MCT) can be inhibited, either using NaN_3 or BAY-8002. GLUT: Glucose transporter, HKs: Hexokinases. (B) Representative cytosolic lactate concentration ($[\text{LAC}]_{\text{cyto}}$) of a MMTV-PyMT WT cell over-time in response to administration or removal of NaN_3 and BAY-8002 at time points indicated. Dashed lines indicate slopes taken for assessment of the "Warburg index. (C – G) Average Warburg indices $\pm$ SEM of MMTV-PyMT WT (C, F, left), MMTV-PyMT BK-KO (C, F, right), MDA-MB-453 cells (D, G) and MCF-7 cells (E) calculated from the experiments as shown in (B), either under control conditions, in the presence of paxilline or iberiotoxin (C, D), upon expression of BK_{RFP} or $\text{BK}_{\text{Ca}}\text{-DEC}^{\text{RFP}}$ (E), or upon cell treatment with a scrambled siRNA (siScrbl), or siRNA against a common BK_{Ca} sequence targeting all known splice variants (siBK), or a siRNA specifically designed to knockdown $\text{BK}_{\text{Ca}}\text{-DEC}$ (siDEC) (F, G). (H, I) Normalized MTT absorbance over-time of MMTV-PyMT WT (H, left) and BK-KO cells (H, right), and MDA-MB-453 cells (I), either under control conditions, or in the presence of paxilline or iberiotoxin. J, Representative images and corresponding statistics of colony formation assays using MMTV-PyMT WT or BK-KO cells in the presence or absence of O_2 . (K – N) mRNA expression of BK_{Ca} and $\text{BK}_{\text{Ca}}\text{-DEC}$ as performed by Nanostring analysis of 551 BC patient samples. (K) Log₂ expression counts of BK_{Ca} and $\text{BK}_{\text{Ca}}\text{-DEC}$. The threshold for positive expression level was set to $\log_2 = 5.5$ (dashed line). (L) Log₂ expression counts of $\text{BK}_{\text{Ca}}\text{-DEC}$ blotted over the log₂ expression counts of BK_{Ca} . 10 of the 551 patient samples showed expression of $\text{BK}_{\text{Ca}}\text{-DEC}$ above the threshold of $\log_2 = 5.5$ (dashed line), whereas 541 patient samples were $\text{BK}_{\text{Ca}}\text{-DEC}$ negative. (M) Correlation of the log₂ expression counts of $\text{BK}_{\text{Ca}}\text{-DEC}$ positive samples with the log₂ expression counts of BK_{Ca} in the primary human BC material. (N) Summarizing scheme of BK_{Ca} in cancer cell homeostasis. N (independent experiments) = A – E: 4 WT and BK-KO ctrl (A), 6 WT + PAX (C), 5 BK-KO + PAX (C), WT siScrbl and WT siBK (F) and MDA-MB-453 + PAX (D), MCF-7 + $\text{BK}_{\text{Ca}}\text{-RFP}$ (E), 4 WT and BK-KO + IBTX (C), 7 for all others. F, G: 4 WT siDEC, 5 for all others. H – J: 4 for all. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, Kruskal-Wallis test followed by Dunn's MC test (C, E, F, G, I), One-Way ANOVA test followed by Tukey's MC test (H) or Mann-Whitney test (D). † $p < 0.01$, ‡ $p < 0.001$ compared to respective WT condition, Unpaired t-test (ctrl, C, J) or Mann-Whitney test (+ IBTX, C, F).

Discussion

Here, we demonstrate for the first time that BK_{Ca}-DEC (mitoBK_{Ca}) is functionally expressed in BCCs. BK_{Ca}-DEC modulates BCC metabolism, stimulates the Warburg effect, and accelerates cell proliferation rates in the presence and absence of O₂. These tumor- and malignancy-promoting effects were sensitive to BK_{Ca} inhibition using the cell-permeable BK_{Ca} inhibitor paxillin e(Zhou and Lingle, 2014 [↗](#)), but not by the cell-impermeable blocker iberiotoxin (Candia et al., 1992 [↗](#)), indicating that intracellular BK_{Ca}, presumably mitoBK_{Ca}, mediates malignant BCC behavior and tumor development.

In line with recent single-cell RNA sequencing data of 26 primary breast tumors (S. Z. Wu et al., 2021 [↗](#)), we found high transcript levels for BK_{Ca} throughout the analyzed BC samples, in addition to BK_{Ca}-DEC, albeit in a much smaller subset of BC. Further, we confirmed functional BK_{Ca} expression in the PM of MMTV-PyMT WT and MDA-MB-453 cells, while MMTV-PyMT BK-KO and MCF-7 cells showed no or very low PM BK_{Ca} currents. Therefore, MCF-7 cells represented a suitable model to investigate the effects of BK_{Ca} over-/expression on the metabolic homeostasis of human BCC. Similarly to cardiac myocytes, expression of BK_{Ca}-DEC yielded mitochondrial localization of BK_{Ca}-DEC, although its abundance in the IMM appeared less pronounced compared to cardiomyocytes (Singh et al., 2013 [↗](#)). While a direct comparison is difficult as plasmid transfections may have caused unexpected effects, our finding that BK_{Ca}-DEC^{RFP} caused significantly reduced currents across the PM compared to BK confirm its increased intracellular abundance. ^{RFP} putatively

Based on the potential impact of BK_{Ca} on the cellular $\Delta\Psi_{PM}$ and ion balance (Burgstaller et al., 2022a [↗](#)), we conducted an in-depth investigation of the (sub)cellular Ca²⁺ homeostasis. Across the BCCs examined, we observed that functional BK_{Ca} expression modulated [Ca²⁺]_{cyto} dynamics. These alterations showed, however, differential sensitivities to BK_{Ca} inhibitors in MMTV-PyMT WT and MDA-MB-453 cells. While basal [Ca²⁺]_{cyto} in MDA-MB-453 cells was reduced by both, paxilline and iberiotoxin treatment, MMTV-PyMT WT cells were only sensitive to paxilline. Examination of [Ca²⁺]_{ER} confirmed the results from [Ca²⁺]_{cyto}, as [Ca²⁺]_{ER} levels increased with paxilline and iberiotoxin in MDA-MB-453, but only upon paxilline exposure in MMTV-PyMT WT cells, suggesting differential effects of intracellular and PM- localized BK_{Ca} on Ca²⁺ handling in these cells. Interestingly, in MCF-7 cells, both BK_{Ca} splice variants mediated the opposing effects on the basal [Ca²⁺]_{cyto} and [Ca²⁺]_{ER} levels, which either in- or decreased, respectively, upon transient BK ^{RFP} or BK -DEC^{RFP} expression. This is expected as the transitory hyperpolarization and the efflux of K⁺ due to the opening of PM BK_{Ca} provides the driving force for Ca²⁺ entry into cytoplasm, while a BK_{Ca}-mediated K⁺ increase within the ER, presumably through PM-directed channels crossing the ER membrane in the secretory pathway, would oppose the Ca²⁺ refilling (Burgstaller et al., 2022a [↗](#)). Moreover, these results are in line with the higher proliferative capability of BK_{Ca} proficient BCCs due to the manifold roles of Ca²⁺ as second messenger (Burgstaller et al., 2022a [↗](#)). Importantly, however, [Ca²⁺]_{mito} of BK_{Ca} proficient MMTV-PyMT WT and MDA-MB-453 cells was exclusively sensitive for paxilline, and it was specifically affected by the expression of BK_{Ca}-DEC^{RFP}, but not by BK ^{RFP}, in MCF-7 cells, suggesting that this Ca²⁺ pool is exclusively controlled by intracellular BK.

Subcellular Ca²⁺ alterations could reportedly alter cellular bioenergetics, as Ca²⁺ directly regulates metabolic enzymes and activities (Rossi et al., 2019 [↗](#)). Indeed, extracellular flux analysis, LC-MS-based metabolomics and fluorescence-based live-cell imaging confirmed, that the observed alteration in sub- cellular Ca²⁺ homeostasis caused by BK_{Ca}, especially mitoBK_{Ca}, has severe effects on cell metabolism. Our data further emphasize, that the presence of mitoBK_{Ca}, as confirmed by mitoplast patch-clamp and Western blot analysis of isolated mitochondria, depolarizes BCC mitochondria, which is in line with a previous study showing an impact of mitoBK_{Ca} activation on $\Delta\Psi_{mito}$ (Kicinska et al., 2016 [↗](#)). mitoBK_{Ca}- dependent depolarization of $\Delta\Psi_{mito}$ in turn triggers cellular glucose dependency and reverses the activity of the mitochondrial ATP-synthase to

consume ATP for restoring $\Delta\Psi_{\text{mito}}$. Finally, BK_{Ca}-DEC-derived mitoBK_{Ca} channels promote the Warburg effect and ultimately stimulated proliferation rates of BCCs. Our data fit earlier findings from glioma cells, where an O₂-sensitivity of mitoBK_{Ca} was observed, which probably increased the hypoxic resistance of these cancer cells (Gu et al., 2014 [DOI](#)). Excessive production and release of lactate, a hallmark of the Warburg effect, leads to extracellular acidification, subsequently creating a microenvironment that promotes tumorigenesis and metastasis as well as the resistance to anti-tumor immune responses and therapy (de la Cruz-López et al., 2019 [DOI](#); Nazemi and Rainero, 2020 [DOI](#); P. Wu et al., 2021 [DOI](#)). Extracellular K⁺ [K⁺]_{ex} in turn, accumulating within the necrotic core of solid tumors, was shown to interfere with effector T-cell function triggering immune escape of cancer cells (Eil et al., 2016 [DOI](#)). To elucidate whether mitoBK_{Ca} directly contributes to lactate-induced tumor aggressiveness or [K⁺]_{ex}, live-cell imaging of extracellular metabolites in 3D BCC models should be applied in future investigations (Burgstaller et al., 2022b [DOI](#), 2021b [DOI](#)).

Finally, our results demonstrate for the first time that BK_{Ca}-DEC transcripts are present in human BC biopsies. Although only a small proportion of patients was positive for BK_{Ca}-DEC expression, this finding could be of considerable clinical relevance considering the link between mitoBK_{Ca} function and BCC metabolism. Importantly, the design of our study likely underestimates the incidental number of BK_{Ca}-DEC positive BC, as i.) only hormone-receptor positive BC specimens were included, and ii.) bulk-tumor mRNA was analyzed, hampering the detection of low-abundant or tightly regulated transcripts against a strong background of non-cancer cells present in bulk tumor tissues. Finally, due to the small number of positive hits (N=10 positive *versus* N=541 BK_{Ca}-DEC negative specimens) and the lack of (sufficient) follow-up information in some of these cases, we are currently unable to correlate BK_{Ca}-DEC expression with, for example, treatment response or survival. Thus, future studies are warranted to show how the tumor's BK_{Ca}-DEC status can help to predict or therapeutically improve standard chemo-endocrine treatments. The rather low abundance of BK_{Ca}-DEC in the clinical samples, however, is in agreement with our mitoplast patch clamp experiments with mitoBK_{Ca}-mediated K⁺ currents being detected at frequencies between 6 to 15%, suggesting that either a small proportion of mitochondria express functional mitoBK_{Ca}, or that the abundance of the channel per mitochondrion is low, requiring sensitive mechanistic approaches for its detection.

In summary, our data highlight a potentially druggable mitoBK_{Ca} isoform in BCCs, whose molecular entity is mainly formed by the *Kcnma1* encoded BK_{Ca}-DEC splice variant. This channel promotes metabolic alterations in cancer cells, even under low-oxygen conditions, which may ultimately be of clinical interest for new anti-cancer therapies.

Material and Methods

Buffers and solutions

If not otherwise stated, all chemicals were purchased from Carl Roth GmbH, Karlsruhe, Germany. Buffers used in this study comprised the following:

Physiologic buffer for single cell live recordings contained (in mM): 138 NaCl, 5 KCl, 2 CaCl₂, 1 MgCl₂, 2 glucose, 10 HEPES, pH set to 7.4 with NaOH. No glucose was added during glucose removal experiments, while glucose was increased to 25 mM to investigate glucose dependency of the mitochondrial membrane potential. 0.1 mM Ethylene glycol bis(2-aminoethylether)-N, N, N', N'-tetra acetic acid (EGTA) (Sigma Aldrich Chemie GmbH, Taufkirchen, Germany) instead of 2.0 mM CaCl₂ was added to obtain Ca²⁺ free buffer. The following compounds were added to yield the following final concentration: 3 μM Oligomycin-A, 200 nM Carbonyl cyanide-p-trifluoromethoxyphenylhydrazone (FCCP), 5 μM paxilline (all from (Santa Cruz Biotechnology, Dallas, USA), 30 nM iberiotoxin (Selleckchem, Planegg, Germany), 5 mM NaN₃, 3 μM BAY-8002, 15

μM 2,5-Di-(*t*-butyl)-1,4- hydroquinone (BHQ) (all from Sigma Aldrich Chemie GmbH), 100 μM adenosine-5'-triphosphate (ATP). For H_2O insoluble compounds, final DMSO concentration in the buffer did not exceed 0.1%. For experiments of glutamine and pyruvate removal, 2 mM GlutaMAX and 1 mM sodium pyruvate (both Thermo Fisher Scientific, Waltham, USA) were added to the starting buffer.

Intracellular buffer used for whole-cell patch-clamp experiments contained (in mM): 130 K-Gluconate, 5 KCl, 2 Mg-ATP 0.1, CaCl_2 , 0.2 Na₂-GTP, 0.6 EGTA, 5 HEPES, pH = 7.2 with KOH.

Cell equilibration buffer fluorescence microscopy-based contained (in mM): 135 NaCl, 5 KCl, 2 CaCl_2 , 1 MgCl_2 , 2.6 NaHCO_3 , 0.44 KH_2PO_4 , 0.34 Na_2HPO_4 , 10 glucose, 10 HEPES, 2 GlutaMAX, 1 sodium pyruvate, with 1x MEM amino acids and 1x MEM vitamins added (both Thermo Fisher Scientific). pH was adjusted to 7.4 using NaOH.

Buffers used for mitochondrial isolation, mitoplast preparation, and single channel patch-clamp comprised the following: The preparation solution contained (in mM): 250 sucrose, 5 HEPES, pH = 7.2. The mitochondrial storage buffer contained (in mM): 150 KCl, 0.1 CaCl_2 , 20 HEPES, pH = 7.2. The hypotonic buffer contained (in mM) 0.1 CaCl_2 , 5 HEPES, pH = 7.2. The hypertonic buffer contained (in mM): 750 KCl, 0.1 CaCl_2 , 30 HEPES, pH 7.2. Low- Ca^{2+} solution (1 μM Ca^{2+}) contained (in mM): 150 KCl, 1 EGTA, 0.752 CaCl_2 , 10 HEPES, pH = 7.2.

Cell culture and transfection

Mouse mammary tumor virus polyoma middle T antigen (MMTV-PyMT) cells were isolated from tumors of MMTV-PyMT transgenic FVB/N WT or BK-KO mice. Tumor growth *in vivo* and biopsies were authorized by the local ethics *Committee for Animal Research* (Regierungspräsidium Tuebingen, Germany), and were performed in accordance with the *German Animal Welfare Act*. Animals were kept on a 12-hour light/ dark cycle under temperature- and humidity-controlled conditions with unlimited access to food (Altromin, Lage, Germany) and water. MMTV-PyMT cells used in this study were isolated from 3 – 7 different female breast-cancer bearing WT and 3 – 4 different female breast-cancer bearing BK-KO animals at an age of ~ 12 – 14 weeks. Upon dissection, tumors were carefully minced into pieces using atraumatic forceps, lysed by 1 mg mL^{-1} Collagenase-D (Roche, Basel, Switzerland) for 10 min, and cultured as follows: Cells were grown in modified improved minimal essential medium (IMEM) supplemented with 5% fetal bovine serum (FBS), 1 mM sodium pyruvate and 100 U mL^{-1} penicillin and 100 $\mu\text{g mL}^{-1}$ streptomycin (all purchased from Thermo Fisher Scientific) at 37°C and 5% CO_2 in a humidified incubator. Fibroblasts were removed by exposure of the cultures to 0.25% trypsin-EDTA in PBS (Thermo Fisher Scientific) and short incubation at 37°C (~ 1 minute). After gently tapping the plate, trypsin-EDTA with detached fibroblasts was removed and cells were further cultured in supplemented modified IMEM at 37°C and 5% CO_2 until subculturing.

MCF-7 and MDA-MB-453 cells were purchased from the Global Bioresource Center (ATCC). Cells were cultivated in Dulbecco's modified eagle's medium (DMEM) supplemented with 10% FBS, 1 mM sodium pyruvate and 100 U mL^{-1} penicillin and 100 $\mu\text{g mL}^{-1}$ streptomycin (Thermo Fisher Scientific) at 37°C and 5% CO_2 in a humidified incubator.

Subculturing of cells was performed when cells reached a confluency of 80 – 90%. Therefore, cell culture medium was removed, cells were washed 1x with PBS, and trypsin-EDTA at a final concentration of 0.25% trypsin-EDTA in PBS was added. Subsequently, cells were incubated at 37°C and 5% CO_2 in a humidified incubator until cell detachment occurred (~ 2-5 minutes). Trypsinization was stopped by adding supplemented cell culture media and cells were pelleted at 300xg for 5 minutes. The supernatant was removed, and cells were seeded to new cell culture dishes as required. For fluorescence microscopic live-cell imaging experiments, cells were either

seeded in 6-well plates containing 1.5 H 30 mm circular glass coverslips (Paul Marienfeld GmbH, Lauda-Königshofen, Germany). All other vessels and serological pipettes used for cell culture were ordered from Corning (Kaiserslautern, Germany).

Transfection of cells was performed according to manufacturer's instructions when cells showed a confluency of ~70%, either using PolyJET DNA transfection reagent (SignaGen Laboratories, Maryland, USA) for plasmid DNA transfection or Lipofectamine 2000 (Thermo Fisher Scientific) for siRNA transfection or co-transfection of siRNA with plasmid DNA. Plasmid DNA amount was reduced to 1/3 for transfection of mitochondrial-targeted probes to ensure proper mitochondrial localization of the probes. Plasmid transfections were performed 16 hours, siRNA transfections were performed 48 hours before the experiments. Paxilline or iberiotoxin were added 12 hours prior to the experiments to the respective cell culture medium. DMSO served as a control.

Whole-cell patch-clamp

For whole-cell patch-clamp experiments, 30,000 cells were seeded on the day before the experiment in 35 mm glass bottom μ Dishes (ibidi GmbH, Graefelfing, Germany) and cultivated in the respective supplemented cell culture medium over-night at 37°C and 5% CO₂ in a humidified incubator. The next day, cell culture medium was removed, and cells were washed 2x and maintained in prewarmed physiologic buffer. Subsequently, recordings were performed using borosilicate glass capillaries (0.86x1.5x100mm) (Science Products GmbH, Hofheim am Taunus, Germany), with a resistance of 4-6 MW, which were pulled using a model P-1000 flaming/ brown micropipette puller (Sutter Instruments, California, USA) and filled with intracellular buffer. A MP-225 micromanipulator served for pipette control (Sutter Instruments). Recordings were performed in whole-cell mode. Currents were evoked by 15 voltage square pulses (300 ms each) from the holding potential of -70 mV to voltages between -100 mV and +180 mV delivered in 20 mV increments. For amplifier control (EPC 10) and data acquisition, Patchmaster software (HEKA Elektronik GmbH, Lambrecht, Germany) was used. Voltages were corrected offline for the capacity. Data analysis was performed using Fitmaster software (HEKA Elektronik GmbH), Nest-o-Patch software (<http://sourceforge.net/projects/nestopatch> , written by Dr. V Nesterov), and Microsoft Excel (Microsoft, Washington, USA).

Cloning and plasmid preparation

Cloning was performed using conventional PCR-, restriction- and ligation-based procedures. BK_{Ca}-DEC was a gift from Michael J. Shipston and was N-terminally attached to an RFP (BK_{Ca}-DEC^{RFP}) using KpnI and BamHI restriction sites after PCR amplification (NEB Q5 High-Fidelity DNA-Polymerase, New England Biolabs (NEB), Ipswich, USA). For generation of BK^{RFP}, a PCR amplification of BK_{Ca}-DEC was performed, where the reverse primer omitted the last amino acids including the 50 amino acids encoding the DEC exon. Mitochondrial targeted TagRFP (mtRFP) was generated by fusing a double repeat of COX8 pre-sequence N-terminally to TagRFP. RFP-GPI was generated by fusing the membrane leading sequence (MLS) and the GPI-anchor signal of cadherin 13 N- and C-terminally to TagRFP, respectively. After PCR reactions, the DNA fragments were purified from gel electrophoresis using the Monarch DNA gel extraction kit (NEB), fragments and destination plasmid were digested using the respective restriction enzymes (NEB) and ligation (T4 DNA Ligase, NEB) and transformation (chemically competent NEB 5- α *E. coli*) were performed according to manufacturer's instructions. Plasmids were verified by sequencing (Microsynth AG, Balgach, Switzerland). DNA maxipreps were performed using the Nucleobond Xtra Maxi kit (Macherey Nagel GmbH & Co. KG, Düren, Germany). Purified DNA was stored at 4°C.

Apotome imaging

For apotome imaging of mtRFP, RFP-GPI, BK_{Ca}RFP or BK_{Ca}-DEC^{RFP} colocalization with mitochondria, MCF-7 cells were seeded on circular 18 mm glass coverslips (Marienfeld GmbH) in 12-well plates (Corning). Cells were transfected using PolyJet transfection reagent according to the

manufacturer's instructions. 16 hours after transfection medium was exchanged for fresh cell culture medium and cells were further cultivated for 24 hours. Subsequently, the medium was exchanged for fresh medium containing MitoGREEN (PromoCELL GmbH, Heidelberg, Germany) at a final concentration of 66.6 μM and cells were further incubated at 37°C and 5% CO_2 for 20 minutes. Subsequently, cells were washed 2x with cold PBS, paraformaldehyde (PFA) (Carl Roth GmbH) at a final concentration of 2% in PBS was added and cells were incubated for 30 minutes on ice. Next, cells were washed again 2x with PBS and mounted on glass slides (Th. Geyer GmbH & Co. KG, Renningen, Germany) with PermaFluor aqueous mounting medium (Thermo Fisher Scientific) and dried over-night in the dark at 4°C. Imaging was performed using a Zeiss axio imager Z1 equipped with an Apotome.2 (Carl Zeiss AG), a HBO 100 lamp and a 63x 1.4 plan apochromatic objective. Microscope control and image acquisition was performed using Zeiss zen 2.6 blue edition (Carl Zeiss AG). Image analysis was performed using the colocalization test in ImageJ with Fay randomization after cell selection by ROIs.

Fluorescence live-cell imaging

Cells were either analyzed using a Zeiss AXIO Observer Z1 or a Zeiss Axiovert 200m microscope (Carl Zeiss AG, Oberkochen, Germany). The Zeiss AXIO Observer Z1 was connected to a LEDHub high-power LED light engine (OMICRON Lasertechnik, Rodgau-Dudenhofen, Germany) and equipped with a EC Plan- Neofluar 40x/1.3 Oil DIC M27 objective (Carl Zeiss AG), an Optosplit II emission image splitter (Cairn Research Ltd, Faversham, UK), and a pco.panda 4.2 bi sCMOS camera (Excelitas PCO GmbH, Kelheim, Germany). The microscope possessed a BioPrecision2 automatic XY-Table (Ludl Electronic Products, Ltd., New York, USA). Optical filters included a 459/526/596 dichroic mirror and a 475/543/702 emission filter for FRET- and TMRM-based measurements, and a 409/493/573/652 dichroic mirror combined with a 514/605/730 emission filter for Dibac4(3), Fura-2 and 2-NBDG based measurements (all purchased from AHF Analysentechnik, Tübingen, Germany). The Optosplit II emission image splitter was equipped with a T505lpxr long-pass filter (AHF Analysentechnik). The LEDHub high-power LED light engine was equipped with a 340 nm, 385 nm, 455 nm, 470 nm and 505-600 nm LED, followed by the following emission filters, respectively: 340x, 380x, 427/10, 473/10 and 510/10 or 575/15 (AHF Analysentechnik). The Zeiss Axiovert 200m microscope was connected to a pe340^{fura} light source (CoolLED, Andover, UK), an Optosplit II emission image splitter (Cairn Research Ltd.) and a pco.panda 4.2 sCMOS camera (Excelitas PCO GmbH) and equipped with 340/26, 380/14 and switchable 427/10, 485/20 or 575/15 excitation filters (AHF Analysentechnik) in the light source, respectively, a 40x Fluor 1.30 oil immersion objective (Carl Zeiss AG), a 459/526/596 or 515LP dichroic mirror and a 475/543/702 or 525/15 emission filter (AHF Analysentechnik) in the microscope, and a T505lpxr (AHF Analysentechnik) in the Optosplit II. Image acquisition and control of both microscopes was performed using VisiView software (Visitron Systems GmbH, Puchheim, Germany). Perfusion of cells was performed using a PC30 perfusion chamber connected to a gravity-based perfusion system (NGFI GmbH, Graz, Austria) and a vacuum pump.

Fura-2 based Ca^{2+} measurements

For fura-2 based Ca^{2+} measurements, cells were taken from the humidified incubator at 37°C and 5% CO_2 , washed 1x with cell equilibration buffer and loaded with fura-2 AM (Biomol GmbH, Hamburg, Germany) at a final concentration of 3.3 μM in cell equilibration buffer for 45 minutes at room temperature.

Subsequently, cells were washed 2x with cell equilibration buffer and stored in equilibration buffer for additional 30 minutes prior to the measurements. Paxilline or iberitoxin treatment of the cells was performed 12 hours prior to the measurements and both inhibitors remained present during the fura-2 loading procedure and the measurement at concentrations of 5 μM and 30 nM, respectively. Imaging experiments were either performed on the Zeiss AXIO Observer Z1 or the Zeiss Axiovert 200m microscope (Carl Zeiss AG) in physiologic buffer using alternate excitations at 340 nm and 380 nm. Emissions were captured at roughly 514 nm (Zeiss AXIO

Observer Z1) or 525 nm (Zeiss Axiovert 200m). To evoke intracellular Ca^{2+} signals, cells were perfused with physiologic buffer containing ATP (Carl Roth GmbH) at a final concentration of 100 μM .

Genetically encoded sensor-based measurements

Cells were grown on 1.5 H 30 mm circular glass coverslips (Paul Marienfeld GmbH) for perfusion-based experiments. Cells were taken from the incubator, cell culture medium was removed, and cells were equilibrated for at least 30 minutes in cell equilibration buffer in ambient environment. Sensor plasmids used in this study comprised the following: D1ER (Palmer et al., 2004 [DOI](#)) (Addgene plasmid #36325) for measurement of $[\text{Ca}^{2+}]_{\text{ER}}$ and 4mtD3cpV (Palmer et al., 2006 [DOI](#)) (Addgene plasmid #36324) for measurement of $[\text{Ca}^{2+}]_{\text{mito}}$ were a gift from Amy Palmer & Roger Tsien, mtAT1.03 (Imamura et al., 2009 [DOI](#)) for measurement of $[\text{ATP}]_{\text{mito}}$ was a gift from Hiromi Imamura, Laconic (San Martín et al., 2013) (Addgene plasmid #44238) for measurement of $[\text{LAC}]_{\text{cyto}}$ and assessment of the Warburg index was a gift from Luis Felipe Barros, and mito-Hyper3 (Bilan et al., 2013 [DOI](#)) was a gift from Markus Waldeck-Weiermair. Experiments were either performed on the AXIO Observer Z1 or the Axiovert 200m microscope (Carl Zeiss AG). For experiments where paxilline (5 μM) or iberiotoxin (30 nM) were used, these compounds were added with the media change prior to the transfection with the PolyJet transfection reagent (SignaGen Laboratories) approximately 12 hours prior to the experiments. The compounds remained present throughout the experiment. FRET-based sensors were excited at 427/10 nm and emissions were collected simultaneously at roughly 475 and 543 nm. mito-Hyper3 was excited at 427/10 and 510/10 nm.

Extracellular flux analysis

Assessment of extracellular acidification rate (ECAR) and oxygen consumption rate (OCR) was performed using a Seahorse XFe24 analyzer (Agilent, Santa Clara, USA). The day before the assay, Seahorse XF24 cell culture microplates (Agilent) were coated with 0.5 mg mL^{-1} poly-L-lysine (Sigma Aldrich Chemie GmbH) for 30 minutes at 37°C, washed 2x with PBS followed by cell seeding of 50,000 MMTV-PyMT WT, 50,000 MMTV-PyMT BK-KO or 100,000 MDA-MB-453 cells per well of the 24-well plate in a final volume of 250 μL , either in the presence or absence of 5 μM paxilline, 30 nM iberiotoxin or an equivalent amount of DMSO. 4 wells contained medium without cells and served as blank. Cells were cultivated over-night at 37°C and 5% CO_2 in a humidified incubator. The Seahorse XFe24 sensor cartridges were equilibrated over-night at 37°C in Seahorse XF Calibrant solution according to manufacturer's instructions. The next day, cells were washed using Seahorse XF DMEM medium, pH 7.4 additionally supplemented with 5.5 mM glucose, 2 mM GlutaMAX and 1 mM sodium pyruvate (Thermo Fisher Scientific), with or without paxilline, iberiotoxin or DMSO according to manufacturer's instructions. With the last washing step, a final volume of 500 μL was adjusted per well and cells were incubated at 37°C in the absence of CO_2 for 30 minutes. Meanwhile the Seahorse XFe24 sensor cartridge was loaded with the following compounds: 55 μL of 20 μM Oligomycin-A, 62 μL of 3 μM FCCP and 69 μL of 25 μM Antimycin-A (Santa Cruz Biotechnology), yielding final concentrations of 2 μM , 300 nM and 2.5 μM upon injection, respectively. For analysis, ECAR and OCR rates were normalized for the protein concentration (μg) per well, which was assessed using the Pierce BCA protein assay kit (Thermo Fisher Scientific) according to manufacturer's instructions. Absorbance was measured at 540 nm using a TECAN multiplate reader (TECAN Group Ltd., Männedorf, Switzerland) and concentrations were assessed using a calibration curve.

LC-MS-based metabolomics

Formic acid, acetic acid, acetonitrile and methanol of Ultra LC-MS grade were supplied by Carl Roth (Karlsruhe, Germany). Ammonium hydroxide solution (Suprapur® quality 28.0 - 30.0% NH_3 basis) was purchased from Sigma-Aldrich (Merck, Taufkirchen, Germany). Deionized water was purified by a Purelab ultrapurification system (ELGA LabWater, Celle, Germany). Uniformly (^{13}C -labeled yeast extract of more than 2×10^9 *Pichia pastoris* cells (~15 mg; strain CBS 7435) was

obtained from ISOTopic Solutions (Vienna, Austria). All standards used were purchased from Sigma-Aldrich Chemie GmbH. Stock solutions of the individual calibrants were prepared at concentrations of 1 mg mL⁻¹ and used for further dilution. The individual stocks were stored at -80°C until use.

Targeted LC-MS analysis was performed using an Agilent 1290 Infinity II series UHPLC system from Agilent Technologies (Waldbronn, Germany) equipped with a binary pump, autosampler, thermostated column compartment and a QTrap 4500 mass spectrometer with a TurboIonSpray Source from SCIEX (Ontario, Canada). The samples were filled into homogenization tubes. Internal standards were added prior to sample preparation. For the extraction of the analytes, 1 mL of the ice-cold extraction solvent (50% methanol and 50% water) and 0.15 g of zirconia/glass beads were added to the cell pellets (which were slowly thawed on ice). The samples were homogenized (6,800 rpm, 1 min at 4 °C, 10 × 10 s, pause 30 s) with Cryolys Evolution using dry ice cooling (Bertin Technologies, France). The samples were then spun down for 5 min (16,100xg at 4 °C). The supernatant was carefully removed, transferred into fresh tubes and evaporated to dryness overnight under nitrogen using a high-performance evaporator (Genevac EZ-2) (Genevac, Ipswich, UK). The dry residue of the extract was reconstituted in 10 µL water and 90 µL acetonitrile, followed by 3 cycles of vortexing and sonication (30 seconds each). The samples were centrifuged at 18928xg for 5 min and the supernatant was used for further analysis.

Chromatographic separation was performed on a Waters (Eschborn, Germany) Premier BEH Amide column (150 x 2.1 mm, 1.7 µm). For metabolite analysis in ESI⁺ mode, mobile phases A and B were adjusted to a pH of 3.5 with formic acid and consisted of 20 mM ammonium formate in water and acetonitrile, respectively. In ESI⁻, the chromatographic conditions differed. Mobile phase A and B were adjusted to a pH of 7.5 with acetic acid and consisted of 20 mM ammonium acetate in water and acetonitrile, respectively. The gradient elution profile was the same for both positive and negative ionization modes (0.0 min, 100% B; 13 min 70% B; 15 min 70% B; 15.1 min 100% B; 20 min 100% B) and was carried out at a flow rate of 0.25 mL min⁻¹ and a constant column temperature of 35 °C. The injection volume was 5 µL. The autosampler was kept at 4°C. Ion source parameters were as follows: nebulizer gas (GS1, zero grade air) 50 psi, heater gas (GS2, zero grade air) 30 psi, curtain gas (CUR, nitrogen) 30 psi, source temperature (TEM) 450 °C, ion source voltage +5,500 V (positive mode) and - 4,500 V (negative mode). Due to the large number of transitions monitored simultaneously, the Scheduled-MRM function was enabled. A window of 30 seconds was set around the designated metabolite-specific retention time and the total cycle time was 1 s. Blank solvents (mobile phase A and B) followed by QCs were injected in the beginning of the chromatographic batch to ensure proper column and system equilibration.

Plasma- and mitochondrial membrane potential measurements

DY_{PM} was determined using Bis-(1,3-dibutylbarbituric acid)trimethine oxonol (Dibac4(3)) (Thermo Fisher Scientific). Cells were cultivated on 30 mm glass coverslips. On the day of analysis, cells were equilibrated in cell equilibration buffer containing Dibac4(3) at a concentration of 0.25 µg mL⁻¹ for 30 minutes and subsequently analyzed by fluorescence microscopy using 485/20 excitation at the Zeiss Axiovert 200m microscope. Emission was captured at roughly 525 nm.

DY_{mito} was assessed using Tetramethylrhodamin-Methylester (TMRM) (Thermo Fisher Scientific). After cell cultivation on 30 mm glass coverslips in the presence or absence of paxilline, iberiotoxin or DMSO for 12 hours, the cell culture medium was replaced with cell equilibration buffer containing 200 nM TMRM and cells were incubated in ambient environment for 30 minutes. Subsequently, cells were washed with physiologic buffer containing 2 mM or 25 mM glucose and 200 nM TMRM, and cells were equilibrated for further 30 minutes. The glass coverslips were transferred to the PC30 perfusion chamber and experiments were started using the gravity-based perfusion system (NGFI GmbH). Paxilline, iberiotoxin or DMSO (control) and 200 nM TMRM remained present throughout the experiment. Mitochondrial depolarization was induced by the perfusion of a buffer containing 200 nM FCCP. For analysis, the fluorescence emission at ≥600 nm

upon excitation at 575/15 nm of a region of interest (ROI) above mitochondria was divided by a ROI in the nucleus (mitochondria free area) and the ratio was plotted over-time, or basal values were given.

2-NBDG based glucose uptake measurements

Glucose uptake was assessed using 2-(N-(7-Nitrobenz-2-oxa-1,3-diazol-4-yl)Amino)-2-Deoxyglucose (2- NBDG) (Biomol GmbH). Therefore, cells were seeded on 30 mm glass coverslips (Marienfeld GmbH) in 6-well plates (Corning) and cultivated over-night at 37°C and 5% CO₂ in a humidified incubator. The next day, cells were taken from the incubator, cell culture medium was replaced for cell equilibration buffer and cells were equilibrated for 30 minutes at ambient environment. Subsequently, cells were washed 3x with glucose free physiologic buffer, and glucose free physiologic buffer containing 100 µM 2-NBDG, with or without paxilline and 200 nM FCCP or an equivalent amount of DMSO, was added to the cells, followed by incubation at 37°C for 30 minutes. Next, cells were washed 3x with glucose free physiologic buffer to remove any residual 2-NBDG and were analyzed by fluorescence microscopy using 485/20 excitation light. Emission was captured at roughly 525 nm (Zeiss Axiovert 200m).

Mitochondrial single-channel patch-clamp measurements

For mitoplast electrophysiology, mitochondria were isolated from BCC grown to confluency (>90%). Adherent cells were washed 2x with PBS, scraped from the dish, collected in a tube and centrifuged at 400xg for 5 minutes. Subsequently, the cell pellet was resuspended in preparation solution, followed by homogenization using a glass-glass homogenizer. Next, the homogenate was centrifuged at 9,200xg for 10 minutes. The resulting pellet was resuspended in preparation solution and centrifuged at 780xg for 10 minutes. The supernatant was collected, followed by centrifugation at 9,200xg for 10 minutes and resuspension of the mitochondrial fraction in the mitochondrial storage buffer. All procedures were performed at 4°C.

An osmotic swelling procedure was used for the preparation of mitoplasts from the isolated mitochondria. Therefore, mitochondria were added to the hypotonic buffer for ~1 minute to induce swelling and breakage of the outer mitochondrial membrane. Subsequently, isotonicity of the solution was restored by the addition of hypertonic buffer at a dilution of 1:5.

Patch-clamp experiments on mitoplasts were performed as previously described (Bednarczyk et al., 2013 [\[4\]](#)). The experiments were carried out in mitoplast-attached single-channel mode using borosilicate glass pipettes with a mean resistance of 10-15 MΩ. The patch-clamp glass pipette was filled with mitochondrial storage buffer. This isotonic solution was used as a control solution for all experiments. The size of the pipettes and the formation of the gigaseal were monitored by measuring electrical resistance. Connections were made with Ag/AgCl electrodes and an agar salt bridge (3 M KCl) for the ground electrode. The current was recorded using an Axopatch 200B patch-clamp amplifier (Molecular Devices, California, USA). To apply substances, a self- made perfusion system containing a holder with a glass pipe, a peristaltic pump, and Teflon tubing was used. All channel modulators were added as dilutions in the isotonic solution containing 100 µM CaCl₂.

The presented single-channel current-time recordings are representative for the most frequently observed conductances under the given conditions. The conductance was calculated from the current-voltage relationship. The probability of channel openings and the current amplitude were determined using the Single-Channel Search mode of the Clampfit 10.7 software (Molecular Devices).

Western blotting

Mitochondria for western blotting were prepared as described earlier (Pallotti and Lenaz, 2007 [↗](#)) with some modifications. Cells were washed and collected in PBS and centrifuged at 500 $\times$ g for 10 min. The pellet was frozen in liquid nitrogen and stored at -80°C. The next day, pellet was resuspended in ice-cold isolation buffer containing (in mM): 210 mannitol, 70 sucrose, 1 PMSF, 5 HEPES and bovine serum albumin (2.5 mg mL⁻¹), pH 7.2. Digitonin was added to a final concentration of 0.02–0.04% for additional membrane permeabilization. After 1 min of incubation, digitonin was diluted with isolation buffer and the cells were centrifuged at 3,000 $\times$ g for 5 min at 4°C. The pellet was resuspended in ice-cold isolation buffer. The cells were homogenized using a glass/glass homogenizer and centrifuged at 1,000 $\times$ g for 5 min at 4°C followed by another homogenization. The homogenates were centrifuged at 1,000 $\times$ g for 5 min at 4°C to remove cell remnants. The supernatants were collected and centrifuged for 60 min at 10,000 $\times$ g, at 4°C. Next, the pellet was resuspended in the isolation buffer without BSA and centrifuged at 10,000 $\times$ g for 30 min at 4°C, followed by resuspension in isolation buffer without BSA and centrifugation at 10,000 $\times$ g for 15 min at 4°C. This step was repeated once more. The pellet containing crude mitochondria was resuspended in 0.8 mL of 15% Percoll (in isolation buffer without BSA) and layered on top of a Percoll step gradient (23% and 40% Percoll layers). The suspension was centrifuged at 30,000 $\times$ g for 30 min at 4°C. Mitochondrial fraction located between the 23% and 40% Percoll layer was collected, diluted with isolation buffer without BSA and centrifuged at 10,000 $\times$ g for 15 min at 4°C. The final mitochondrial pellet was resuspended in storage buffer containing 500 mM sucrose and 5 mM HEPES, pH 7.2. Each mitochondrial isolation was performed using between 15–30 $\times$ 10⁶ cells.

A given amount of sample solubilized in Laemmli buffer (Bio-Rad) was separated by 10% Tris-tricine- SDS-PAGE and transferred onto polyvinylidene difluoride (PVDF) membranes (Bio-Rad). After protein transfer, the membranes were blocked with 10% nonfat dry milk solution in Tris-buffered saline with 0.2% Tween 20 and exposed to one of the following antibodies: anti-BK α antibody (NeuroMabs, USA, clone L6/60, diluted 1:200), anti-COXIV (Cell Signaling Technology, Leiden, Netherlands, no. 4844, 1:1,000), or anti- α 1 sodium potassium ATPase (Abcam, Berlin, Germany, no. ab7671, 1:1,000). This was followed by incubation with a secondary anti-rabbit (Thermo Fisher Scientific, no. 31460) or anti-mouse antibody (Thermo Fisher Scientific, no. SA1-100) coupled to horseradish peroxidase. The blots were developed using enhanced chemiluminescence solution (GE Healthcare). To estimate the molecular weight of the analyzed proteins PageRuler Prestained Protein Ladder (Thermo Fisher Scientific) was used.

qPCR analysis

mRNA was isolated from 6-well plates showing a cell confluency of ~90% using the Monarch Total RNA Miniprep kit (NEB). siRNA transfection was performed 48 hours prior to RNA isolation using the Lipofectamine 2000 transfection reagent (Thermo Fisher Scientific) according to manufacturer's instructions. qPCR reactions were performed using the GoTaq 1-Step RT-qPCR System (Promega GmbH, Walldorf, Germany). 100 ng of isolated mRNA were used per reaction. Primers were designed to span exon – exon junctions and to recognize both, human and murine BK_{Ca} sequences. Different primer pairs were used for human and murine β -tubulin. Primer and siRNA sequences are listed in **Supplementary Table 1** [↗](#) and **Supplementary Table 2** [↗](#). Primers were purchased from Thermo Fisher Scientific, siRNAs were purchased from Microsynth AG. qPCR reactions were run in a CFX connect real-time PCR instrument (Bio-Rad Laboratories GmbH, Feldkirchen, Germany). Sample (Ct) values were normalized to Ct's of housekeeper gene (β -Tubulin) and calculated as 2^{-Ct} (normalized).

Proliferation assays

Proliferation assays were performed using 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Thermo Fisher Scientific) assay. For the assay, 2,500 MMTV-PyMT WT and BK-KO cells or 7,500 MDA-MB-453 cells were seeded per well in flat bottom 96-well cell culture plates

(Corning) in their respective cell culture medium. Per run, condition, and time-point, 7 technical replicates were performed. One well served as a blank per time-point and condition and did not contain cells. The next day, medium was exchanged for fresh cell culture medium either containing 5 μM paxilline, 30 nM iberiotoxin or an equivalent amount of DMSO as a control and the MTT assay was started for time-point 24 hours directly thereafter. For each time-point (24, 48, 72 and 96 hours), MTT reagent was added with a medium change (10 μL + 100 μL / well) at a final concentration of 455 $\mu\text{g mL}^{-1}$ and cells were incubated for 4 hours. Subsequently, 85 μL of medium were removed from each well, 85 μL of DMSO (Carl Roth GmbH) was added and wells were mixed by pipetting thoroughly. 85 μL of this mixture were transferred to a new 96-well plate (Corning) and absorbance was measured at 540 nm using a TECAN microplate reader (TECAN Group Ltd.). Absorbances at the different time-points (48, 72 and 96 hours) were normalized to the absorbances after 24 hours.

Colony formation assays

For colony formation assays, cell suspensions containing 400 cells mL^{-1} of MMTV-PyMT WT or BK-KO cells were prepared. 1 mL of this suspension was spread drop-per-drop per well of a 6-well cell culture plate (Corning) already containing 1 mL of cell culture medium using 1 mL syringes and 23G needles. Culture plates were gently moved left/right and back/forth to equally distribute the cells across the well, plates were kept at room temperature for 20 minutes to ensure uniform cell settling, and cells were subsequently cultured over-night at 37°C and 5% CO_2 in a humidified incubator. The next day, medium was exchanged for 2 mL of fresh cell culture medium, and cells were kept in the presence or absence of O_2 at 37°C and 5% CO_2 in humidified incubators for 7 days. After 7 days, cells were washed carefully 2x with PBS and fixed for 30 minutes on ice using 2% PFA in PBS. Meanwhile, a solution of crystal violet (0.01% w/v) was prepared in ddH₂O, cells were washed 2x with PBS and incubated in the crystal violet solution (Sigma Aldrich Chemie GmbH) for 60 minutes after fixation, followed by excessive washing with ddH₂O until a clear background was obtained. Plates were dried at room temperature for at least 12 hours before images were captured using an Amersham Imager 600 (GE Healthcare UK Limited, Buckinghamshire, UK). Analysis was performed using ImageJ.

Breast cancer patient study

A subset of 551 primary tumor specimens with available archival tumor blocks were obtained from a prospective, observational multicenter adjuvant endocrine treatment study of 1286 post-menopausal HR- positive breast cancer patients recruited between 2005 and 2011 (German Clinical Trial Register DRKS 00000605, “IKP211” study). Study inclusion and exclusion criteria have been previously described (Schroth et al., 2020 [DOI](#)). Patients had received standard endocrine treatment, i.e. tamoxifen, aromatase inhibitors, or switch regimens between both. Ethics approval was obtained from the Medical Faculty of the University of Tuebingen and the local ethics committees of all participating centers in Germany. Informed patient consent was obtained from all participants as required by institutional review boards and research ethics committees. All patient data were de-identified prior inclusion in this study.

Nanostring nCounter gene expression analysis

Gene expression analysis was performed according to manufacturer’s protocol. In brief, total RNA from human primary tumors of the IKP211 study (Quick-DNA/RNA FFPE, ZymoResearch, Freiburg, Germany) was extracted from 10 μm sections of FFPE tissues with at least 20% tumor cell content. Samples were enriched for tumor tissue by microdissection. A total of 250 ng RNA was subsequently hybridized with target-specific capture and color-coded reporter probe sets in a pre-warmed thermal cycler at 65°C for 20 h. In the post-hybridization process, the total volume was increased to 32 μL with RNase- free water. Fluorescence count measurements were immediately conducted in the Nanostring nCounter System. Data were analyzed using nSolver 4.0 and

normalized to housekeeping genes. ABCF1, NRDE2, POLR2A, PUM1 and SF3A1 served as housekeepers. Probes used for Nanostring nCounter gene expression analysis are listed in **Supplementary Table 3** [↗](#).

Statistical analysis

Statistical analysis was performed using Prism8 software (GraphPad Software, Boston, USA). All data were tested for normal distribution using D'Agostino and Pearson omnibus normality test. A two-tailed Unpaired t-test or Welch's t-test were used for statistical comparison of normally distributed data, depending on whether the variances of the populations were comparable or significantly different as tested by an F-test. A two-tailed Mann-Whitney test was used for pairwise comparison of non-normally distributed data. Comparison of >2 data sets was done either using One-way ANOVA followed by Tukey's multiple comparison (MC) test or Brown-Forsythe and Welch ANOVA test followed by Games-Howell's MC test for normally distributed data, depending on whether the variances were comparable or significantly different. A Kruskal-Wallis test followed by Dunn's MC test was performed if data were not normally distributed. The statistical tests used are indicated in the figure legends. p-values of ≤ 0.05 were considered as significant, where * or # $p \leq 0.05$, ** or † $p \leq 0.01$ and *** or ‡ $p \leq 0.001$. No priori sample size estimation was performed.

Acknowledgements

This work was funded by the Deutsche Forschungsgemeinschaft (DFG) with individual grants to RL and LM. RL is a member of the GRK2381: "cGMP: From Bedside to Bench", DFG grant number 335549539. SM, WS, MS and RL acknowledge financial support from the ICEPHA Graduate Program "Membrane-associated Drug Targets in Personalized Cancer Medicine". HB is a fellow of the Austrian Science Fund (FWF) funded Erwin-Schrödinger-Program, project number J-4457. SB acknowledges financial support from the Fritz Thyssen Stiftung. The authors thank Clement Kabagema-Bilan, Michael Glaser, Monika Zochowska and Antoni Wrzosek for excellent technical support and Peter Ruth for valuable discussions. This work was partially supported by the Nencki Institute of Experimental Biology, the Polish National Science Centre grant no. 2019/34/A/NZ1/00352 (AS), and by the FWF, project number I-3716 (RM).

Author contributions

HB and RL initiated and designed this study ; HB, SM, PK, BK, SB, JJ, KS, WS and PB performed experiments and were involved in data curation; HB, SM, PK, BK, SB, JJ, KS and WS assisted in formal analysis; HB, SM, SB and PB visualized data; HB, SB, WS, LM, MS, PB, AS and RL acquired financial support; HB, SM, PK, BK, SB, KS, WS, ML and PB designed the methodology; HB and RL performed project administration; PK, WS, LM, MS, ALB, RM, ML, PB, AS and RL provided resources; RL supervised the project and the personnel; HB and RL wrote the original draft. All authors critically reviewed the manuscript and stated comments.


Data availability

Raw data for the graphs can be found in source data files.

Declaration of interests

The authors declare no competing interests.


Supplementary Information



Supplementary Figure 1

Representative whole-cell patch-clamp traces and colocalization analysis in breast cancer cells.


(A - D) Representative whole-cell patch-clamp traces of MMTV- PyMT WT (A), MMTV-PyMT BK-KO (B), MDA-MB-453 (C) and MCF-7 wild-type cells (D), either under control conditions (ctrl), or in the presence of 5 μ M paxilline (+ PAX) or 30 nM iberiotoxin (+ IBTX), respectively, as indicated in the panels. (E) Representative images (left) of MCF-7 cells either expressing a mitochondrial targeted red fluorescent protein (mtRFP, upper images, second column) or a red fluorescent protein fused to a glycosylphosphatidylinositol (GPI)- anchor (RFP-GPI, lower images, second column). Cells were additionally stained with MitoGREEN for visualization of mitochondria (first column). Merge of the channels and a zoom are demonstrated. Right panel shows average Pearson correlation $\pm$ SEM of MitoGREEN and RFP of mtRFP (left bar) or RFP-GPI (right bar). Grey dashed lines indicate average colocalization scores of MitoGREEN and RFP of mtRFP and RFP-GPI, which is also indicated in [Figure 1g](#). N (cells) = 24 for mtRFP and 12 for RFP-GPI. *** $p \leq 0.001$ Unpaired t-test. (F, G) Representative whole-cell patch-clamp traces of MCF-7 cells either expressing BK^{RFP} (F) or BK_{Ca}-DEC^{RFP} (G) under control conditions (ctrl, left panels), or in the presence of 5 μ M paxilline (+ PAX, middle panels) or 30 nM iberiotoxin (+ IBTX, right panels). (H) Global cellular RFP-fluorescence intensities of MCF-7 cells used for patch-clamp experiments. Cells either expressed BK^{RFP} (red bar) or BK_{Ca}-DEC^{RFP} (green bar). Data represents average $\pm$ SEM of 18 cells for both conditions.



Supplementary Figure 2

Cytosolic-, ER-, and mitochondrial Ca^{2+} homeostasis is altered by functional BKC expression in breast cancer cells.


(A, B) Fura-2 ratio signals (F_{340}/F_{380}) of MMTV-PyMT WT or BK-KO cells over-time in response to cell stimulation with 100 μM ATP. Experiments were either performed in the presence of 5 μM paxilline (A) or 30 nM iberiotoxin (B). (C – E) Maximal fura-2 ratio signals of MMTV-PyMT WT and BK-KO (C), MDA-MB-453 (D) or MCF-7 cells (E) upon stimulation with 100 μM ATP, under control conditions, in the presence of 5 μM paxilline or 30 nM iberiotoxin (C, D), or upon expression of BK^{RFP} or BK-DEC^{RFP} (E). (F, G) FRET-ratio signals over-time of MMTV-PyMT WT and BK-KO cells expressing D1ER, a FRET- based ER Ca^{2+} sensor. Throughout the experiments, either 5 μM paxilline (F) or 30 nM iberiotoxin (G) were present. (H, I) $[\text{Ca}^{2+}]_{\text{mito}}$ over-time of MMTV-PyMT WT or BK-KO cells in response to cell stimulation with 100 μM ATP. $[\text{Ca}^{2+}]_{\text{mito}}$ was assessed using 4mtD3cpV, a FRET-based Ca^{2+} indicator targeted to the mitochondrial matrix. Experiments were either performed in the presence of 5 μM paxilline (H) or 30 nM iberiotoxin (I). (J – L) Maximal FRET-ratio (FRET/CFP) signals received upon stimulation of MMTV-PyMT WT or BK-KO (J), MDA-MB-453 (K) or MCF-7 cells (L) with 100 μM ATP. Experiments were either under control conditions, in the presence of 5 μM paxilline or 30 nM iberiotoxin (J, K), or upon expression of BK^{RFP} or BK-DEC^{RFP} (L). All data represent average $\pm$ SEM. N (independent experiments) = A, B: 6 WT + PAX and BK-KO + PAX, 5 WT + IBTX and BK-KO + IBTX. C – E: 17 WT ctrl, 18 BK-KO ctrl, 6 WT + PAX and BK-KO + PAX, 5 WT + IBTX, BK-KO + IBTX and all MCF-7, 4 for all MDA-MB-453. F, G: 8 WT + PAX and BK-KO + PAX, 6 WT + IBTX and BK-KO + IBTX. H, I: 6 WT + PAX and BK-KO + PAX, 5 WT + IBTX, 4 BK-KO + IBTX. J – L: 11 WT ctrl, 12 BK-KO ctrl, 6 WT + PAX and BK-KO + PAX, 5 WT + IBTX, MDA-MB-453 + IBTX and all MCF-7, 4 BK-KO + IBTX, 8 MDA-MB-453 ctrl and + PAX. * $p \leq 0.05$, *** $p \leq 0.001$, Kruskal-Wallis test followed by Dunn's MC test (C), One- Way ANOVA test followed by Tukey's MC test (E, J) or Brown-Forsythe and Welch ANOVA test followed by Games-Howell's MC test (K). ‡ $p \leq 0.001$ compared to respective WT condition, Mann-Whitney test (ctrl in C, J), Welch's t-test (+ PAX in C). Unpaired t-test (+ IBTX in J).



Supplementary Figure 3

Paxilline and iberiotoxin differentially modulate ECAR and OCR in MMTV-PyMT WT and BK-KO cells.


(A, B) Average ECAR over-time ± SEM of MMTV-PyMT WT (bright red and salmon) and BK-KO cells (dark and bright grey) in response to administration of Oligomycin-A (+O), FCCP (+F) and Antimycin-A (+A) at time points indicated in the panel. Experiments were either performed in the presence of 5 μM paxilline (A) or 30 nM iberiotoxin (B). N = 3 for all. (C, D) Average OCR over-time ± SEM of MMTV-PyMT WT (bright red and salmon) and BK-KO cells (dark and bright grey) in response to administration of Oligomycin-A (+O), FCCP (+F) and Antimycin-A (+A) at time points indicated in the panel. Experiments were either performed in the presence of paxilline (C) or iberiotoxin (D). N = 3 for all. (E) Schematic representation of the workflow for LC-MS-based metabolomics. HILIC: Hydrophilic interaction liquid chromatography



Supplementary Figure 4

BKCa modulates cellular substrate dependency for maintaining [ATP]mito and reverses FOF1 ATP-synthase.


(A) Average fluorescence ratios ($F_{\text{mito}}/F_{\text{nuc}}$) of MMTV- PyMT WT and BK-KO cells, either in the presence of 2.0 mM or 25.0 mM extracellular glucose. $N = 5$ BK-KO 2.0 mM GLU, 6 for all others. $***p \leq 0.001$, Mann-Whitney test. $\dagger p \leq 0.01$, $\ddagger p \leq 0.001$ compared to 2.0 mM glucose condition of the respective cell type, Mann-Whitney test. (B – E) Average changes in FRET-ratio signals $\pm$ SEM induced either upon extracellular glucose removal (B, D) or upon administration of Oligomycin-A (C, E) to MMTV-PyMT WT, BK-KO (B, C) or MDA- MB-453 cells (D, E) expressing mtAT1.03, a FRET-based ATP sensor targeted to the mitochondrial matrix. Experiments were either performed under control conditions or in the presence of 5 μ M paxilline or 30 nM iberiotoxin. N 's for glucose removal (-Glucose) experiments = 8 WT and BK-KO ctrl, 6 WT and BK-KO + PAX, 7 WT + IBTX, 5 BK-KO + IBTX, 5 MDA-MB-453 ctrl, 3 MDA-MB-453 + PAX, 5 MDA-MB-453 + IBTX. N 's for Oligomycin-A administration (+Oligomycin-A) experiments = 11 WT ctrl, 8 BK-KO ctrl, 7 WT + PAX and WT + IBTX, 6 BK-KO + PAX, 8 BK-KO + IBTX, 5 MDA-MB-453 ctrl, 3 MDA-MB-453 + PAX and 8 MDA-MB-453 + IBTX. $*p \leq 0.05$, $**p \leq 0.01$, $***p \leq 0.001$, Kruskal-Wallis test followed by Dunn's MC test (B, C, D). $\#p \leq 0.05$, $\ddagger p \leq 0.001$ compared to respective WT condition, Mann-Whitney test (ctrl in B, all in C), or Welch's t-test (+ IBTX in B). (F, G) Average changes in FRET-ratio signals $\pm$ SEM induced either upon extracellular glucose removal (F) or upon administration of Oligomycin-A (G) to MCF-7 cells expressing mtAT1.03, either in combination with a red fluorescent protein as control, BK_{Ca} BK_{Ca}-DEC^{RFP}. $*p \leq 0.05$, $**p \leq 0.01$, Kruskal-Wallis test followed by Dunn's MC test. $N = 6$ MCF-7 ctrl – glucose, 5 for all others. (H) Schematic representation of processes involved in 2-NBDG uptake. 2-NBDG is taken up via glucose transporters (GLUTs, green), and ATP-dependently phosphorylated by hexokinase isoforms (HKs, red) to 2-NBDG-6-phosphate (2-NBDG-6P). Under basal conditions, if F_0F_1 ATP-synthase is running in forward mode (left panel), mitochondria contribute to 2-NBDG uptake by ATP generation and delivery to HKs via the adenine nucleotide transporter (ANT, yellow), and the voltage-dependent anion channel (VDAC, violet). Contrary, under basal conditions, if F_0F_1 ATP-synthase activity is reversed (right panel) it competes with HKs for ATP. Subsequent inhibition of mitochondria due to their depolarization with FCCP (red) stops ATP-synthase activity. Under these conditions, 2-NBDG uptake will decrease if F_0F_1 ATP-synthase operates in forward mode (left panel), but will increase if F_0F_1 ATP-synthase shows reversed activity (right panel). (J – L) Average fluorescence signal (a.u.) $\pm$ SEM of MMTV-PyMT WT and BK-KO cells (J, K), or MDA-MB-453 cells (L) incubated with 100 μ M 2-NBDG at 37°C for 30 minutes either under control conditions (J) or in the presence of 200 nM FCCP for mitochondrial depolarization (K, L), with or without 5 μ M paxilline or 30 nM iberiotoxin. $N = 4$ for all. $*p \leq 0.05$, $***p \leq 0.001$, Kruskal-Wallis test followed by Dunn's MC test (J, and WT in K), One-Way ANOVA test (BK-KO in K) or Welch's t-test (L). $\#p \leq 0.05$, $\ddagger p \leq 0.001$ compared to respective WT condition, Mann-Whitney test (ctrl and + IBTX in J, ctrl in K), or Unpaired t-test (+ PAX in J and + PAX in K).



Supplementary Figure 5

BKCa is present in the inner mitochondrial membrane of MMTV- PyMT WT and MDA-MB-453 cells.

(A – C) Graphs show representative BK_{Ca} single-channel recordings of the inner mitochondrial membrane of mitoplasts isolated from MMTV-PyMT WT cells using a symmetric 150/150 mM isotonic KCl solution, either containing 100 μ M Ca²⁺ (A), 1 μ M Ca²⁺ (B), or 5 μ M Paxilline in the presence of 100 μ M Ca²⁺ (C), at voltages ranging from -80 to +80 mV as indicated in the panels. The patch in **a** contained two BK_{Ca} channels. “c” indicates the closed-, “o” the open state of the channel. (D) Representative western blot of BK_{Ca} (upper), Na⁺/K⁺ ATPase as a plasma membrane marker (middle), and cytochrome c oxidase subunit IV (COXIV) as a mitochondrial marker (lower). Western blot was performed using whole-cell lysates (L), the homogenate (H), crude isolated mitochondria (C) and mitochondria after percoll gradient purification (P) of MDA-MB-453 cells. N = 3.



Supplementary Figure 6

siRNA treatment reduces expression of BKCa and BKCa-DEC, and BKCa-DEC decreases dependency on oxidative metabolism in breast cancer cells.

(A) Schematic representation of siRNAs used for subsequent silencing experiments. Either an siRNA targeting all known BK_{Ca} isoforms, referred to as siBK (orange), or an siRNA specifically designed against the DEC exon (green), referred to as siDEC (violet), were used. (B, C) Relative mRNA expression levels $\pm$ SEM of BK_{Ca} and BK_{Ca}-DEC in MMTV-PyMT WT (B) and MDA-MB-453 cells (C), as analysed by qPCR. Cells were either treated with a scrambled siRNA as a control (siScrbl), siRNA against all known BK_{Ca} isoforms (siBK) or siRNA specifically targeting BK_{Ca}-DEC (siDEC), respectively. N = 4 siBK MMTV-PyMT WT, 3 for all others. (D) Normalized colony sizes of colony formation assays performed using MMTV-PyMT WT and BK-KO cells. Cells were cultivated for 7 days either in the presence or absence of O₂. *p < 0.05, Mann-Whitney test. N = 4 independent experiments for all conditions.

Gene	Accession number	Target sequence (5' – 3')
ABCF1	NM_001090.2	TCCCGCCAAGCCATGTTAGAAAATGCATCTGACA TCAAGCTGGAGAAGTTCAGCATCTCCGCTCATGG CAAGGAGCTGTTTCGTCAATGCAGACCTGTACA
NRDE2	NM_017970.3	TGGAGGTCTTATGTACAGATTCAGAATAAGTCCC ACAGTGCCAGCAAAACCAGGAGATTTTTTGACA CAATCACCAGGTCTGCCAAACCCCTGGAGCCTT
POLR2A	NM_000937.2	TTCCAAGAAGCCAAAGACTCCTTCGCTTACTGTC TTCCTGTTGGGCCAGTCCGCTCGAGATGCTGAGA GAGCCAAGGATATTCTGTGCCGTCTGGAGCAT
PUM1	NM_001020658.1	CTGGGGAACATCAGATCATTTCAGTTTCCCAGCCA ATCATGGTGCAGAGAAGACCTGGTCAGAGTTTCC ATGTGAACAGTGAGGTCAATTCTGTACTGTCC
SF3A1	NM_005877.4	GATGATGAGGTGTACGCACCAGGTCTGGATATTG AGAGCAGCTTGAAGCAGTTGGCTGAGCGGCGTA CTGACATCTTCGGTGTAGAGGAAACAGCCATTG
KCNMA1	NM_001014797.2	CCGTGCGACAGCCGGGGCCAACGCATGTGGTGG GCTTTCTGGCCTCCTCCATGGTGACTTTCTTCGG GGGCCTCTTCATCATCTTGCTCTGGCGGACGC
KCNMA1-DEC	XM_024447988.2	AAACAGAATGCAACAAGGATGAATAGAATGGGC CAAGAAAAGAAATGGTTTACAGATGAACCGGAT AATGCCTATCCCAGAAACATTCAAATCAAGCCCA

Supplementary Table 1

Probes used for Nanostring nCounter gene expression analysis.

Primer name	Sequence (5' – 3')	Species	Amplicon size
BK _{Ca} -DEC for	CAAACAGAATGCAACAAGGATG	Human / mouse	124 bp
BK _{Ca} -DEC rev	GTTAGCCATGTGGGTACTC	Human / mouse	
BK _{Ca} for	CGCCTCTTCATGGTCTTC	Human / mouse	134 bp
BK _{Ca} rev	ATGTGCTTTCTTCCACTAAC	Human / mouse	
h b-tubulin for	GGCCAGATCTTTAGACCAGAC	Human	120 bp
h b-tubulin rev	CACATCCAGGACAGAATCAAC	Human	
m b-tubulin for	AGTGTGGCAACCAGATC	Mouse	114 bp
m b-tubulin rev	AGTAAACGCTGATCCTCTC	Mouse	

Supplementary Table 2

Primers used for qPCR analysis.

siRNA name	Sequence (5' – 3')	Targeted Species
siScrbl	UUCUCCGAACGUGUCACGU-dTdT	Human / mouse
siBK	UAGGAAACCGCAAGAAUA-dTdT	Human / mouse
siBK-DEC	CCAGAUCAACCAAUAUAAA-dTdT	Human / mouse

Supplementary Table 3

siRNAs used for silencing based experiments.

References

1. Adams DS, Levin M (2012) **Measuring Resting Membrane Potential Using the Fluorescent Voltage Reporters DiBAC 4 (3) and CC2-DMPE** *Cold Spring Harb Protoc* **2012** <https://doi.org/10.1101/pdb.prot067702>
2. Bednarczyk P, Wieckowski MR, Broszkiewicz M, Skowronek K, Siemen D, Szewczyk A (2013) **Putative Structural and Functional Coupling of the Mitochondrial BKCa Channel to the Respiratory Chain** *PLoS ONE* **8** <https://doi.org/10.1371/journal.pone.0068125>
3. Bilan DS, Pase L, Joosen L, Gorokhovatsky AY, Ermakova YG, Gadella TWJ, Grabher C, Schultz C, Lukyanov S, Belousov VV (2013) **HyPer-3: A Genetically Encoded H₂O₂ Probe with Improved Performance for Ratiometric and Fluorescence Lifetime Imaging** *ACS Chem Biol* **8**:535–542 <https://doi.org/10.1021/cb300625g>
4. Bischof H *et al.* (2021) **Potassium ions promote hexokinase-II dependent glycolysis** *iScience* **24** <https://doi.org/10.1016/j.isci.2021.102346>
5. Burgstaller S, Bischof H, Lukowski R, Graier WF, Malli R (2021) **Investigating the K⁺ sensitivity of cellular metabolism by extracellular flux analysis** *STAR Protoc* **2** <https://doi.org/10.1016/j.xpro.2021.100876>
6. Burgstaller S, Bischof H, Matt L, Lukowski R (2022) **Assessing K⁺ ions and K⁺ channel functions in cancer cell metabolism using fluorescent biosensors** *Free Radical Biology and Medicine* **181**:43–51 <https://doi.org/10.1016/j.freeradbiomed.2022.01.026>
7. Burgstaller S *et al.* (2021) **Immobilization of Recombinant Fluorescent Biosensors Permits Imaging of Extracellular Ion Signals** *ACS Sens* **6**:3994–4000 <https://doi.org/10.1021/acssensors.1c01369>
8. Burgstaller S *et al.* (2022) **Monitoring extracellular ion and metabolite dynamics with recombinant nanobody-fused biosensors** *iScience* **25** <https://doi.org/10.1016/j.isci.2022.104907>
9. Candia S, Garcia ML, Latorre R (1992) **Mode of action of iberiotoxin, a potent blocker of the large conductance Ca(2+)-activated K⁺ channel** *Biophys J* **63**:583–590 [https://doi.org/10.1016/S0006-3495\(92\)81630-2](https://doi.org/10.1016/S0006-3495(92)81630-2)
10. Checchetto V, Leanza L, De Stefani D, Rizzuto R, Gulbins E, Szabo I. (2021) **Mitochondrial K⁺ channels and their implications for disease mechanisms** *Pharmacology & Therapeutics* **227** <https://doi.org/10.1016/j.pharmthera.2021.107874>
11. Dabrowska A, Zajac M, Bednarczyk P, Lukasiak A (2022) **Effect of Quercetin on mitoBKCa Channel and Mitochondrial Function in Human Bronchial Epithelial Cells Exposed to Particulate Matter** *IJMS* **24** <https://doi.org/10.3390/ijms24010638>
12. de la Cruz-López KG, Castro-Muñoz LJ, Reyes-Hernández DO, García-Carrancá A, Manzo-Merino J. (2019) **Lactate in the Regulation of Tumor Microenvironment and Therapeutic Approaches** *Front Oncol* **9** <https://doi.org/10.3389/fonc.2019.01143>

13. Dejos C, Gkika D, Cantelmo AR (2020) **The Two-Way Relationship Between Calcium and Metabolism in Cancer** *Front Cell Dev Biol* **8** <https://doi.org/10.3389/fcell.2020.573747>
14. Depaoli MR *et al.* (2018) **Real-Time Imaging of Mitochondrial ATP Dynamics Reveals the Metabolic Setting of Single Cells** *Cell Rep* **25**:501–512 <https://doi.org/10.1016/j.celrep.2018.09.027>
15. Douglas RM, Lai JCK, Bian S, Cummins L, Moczydlowski E, Haddad GG (2006) **The calcium-sensitive large- conductance potassium channel (BK/MAXI K) is present in the inner mitochondrial membrane of rat brain** *Neuroscience* **139**:1249–1261 <https://doi.org/10.1016/j.neuroscience.2006.01.061>
16. Dunnwald LK, Rossing MA, Li CI (2007) **Hormone receptor status, tumor characteristics, and prognosis: a prospective cohort of breast cancer patients** *Breast Cancer Res* **9** <https://doi.org/10.1186/bcr1639>
17. Eales KL, Hollinshead KER, Tennant DA (2016) **Hypoxia and metabolic adaptation of cancer cells** *Oncogenesis* **5**:e190–e190 <https://doi.org/10.1038/oncsis.2015.50>
18. Eil R *et al.* (2016) **Ionic immune suppression within the tumour microenvironment limits T cell effector function** *Nature* **537**:539–543 <https://doi.org/10.1038/nature19364>
19. Gałęcka S, Kulawiak B, Bednarczyk P, Singh H, Szewczyk A (2021) **Single channel properties of mitochondrial large conductance potassium channel formed by BK-VEDEC splice variant** *Sci Rep* **11** <https://doi.org/10.1038/s41598-021-90465-3>
20. (2016) **Molecular Mechanisms of Enzyme Activation by Monovalent Cations** *Journal of Biological Chemistry* **291**:20840–20848 <https://doi.org/10.1074/jbc.R116.737833>
21. Gross D, Bischof H, Maier S, Sporbeck K, Birkenfeld AL, Malli R, Ruth P, Proikas-Cezanne T, Lukowski R (2022) **IKCa channels control breast cancer metabolism including AMPK-driven autophagy** *Cell Death Dis* **13** <https://doi.org/10.1038/s41419-022-05329-z>
22. Grynkiewicz G, Poenie M, Tsien RY (1985) **A new generation of Ca²⁺ indicators with greatly improved fluorescence properties** *J Biol Chem* **260**:3440–3450
23. Gu XQ, Pamerter ME, Siemen D, Sun X, Haddad GG (2014) **Mitochondrial but not plasmalemmal BK channels are hypoxia-sensitive in human glioma: Glioma BK Channels are not Hypoxia Sensitive** *Glia* **62**:504–513 <https://doi.org/10.1002/glia.22620>
24. Huang X, Jan LY (2014) **Targeting potassium channels in cancer** *Journal of Cell Biology* **206**:151–162 <https://doi.org/10.1083/jcb.201404136>
25. Imamura H, Huynh Nhat KP, Togawa H, Saito K, Iino R, Kato-Yamada Y, Nagai T, Noji H (2009) **Visualization of ATP levels inside single living cells with fluorescence resonance energy transfer-based genetically encoded indicators** *Proc Natl Acad Sci USA* **106**:15651–15656 <https://doi.org/10.1073/pnas.0904764106>
26. Jang M, Kim SS, Lee J (2013) **Cancer cell metabolism: implications for therapeutic targets** *Exp Mol Med* **45**:e45–e45 <https://doi.org/10.1038/emm.2013.85>
27. Joshi DC, Bakowska JC (2011) **Determination of Mitochondrial Membrane Potential and Reactive Oxygen Species in Live Rat Cortical Neurons** *JoVE* **2704** <https://doi.org/10.3791/2704>

28. Kalia M (2015) **Biomarkers for personalized oncology: recent advances and future challenges** *Metabolism* **64**:S16–S21 <https://doi.org/10.1016/j.metabol.2014.10.027>
29. Kicinska A, Augustynek B, Kulawiak B, Jarmuszkiewicz W, Szewczyk A, Bednarczyk P (2016) **A large-conductance calcium-regulated K⁺ channel in human dermal fibroblast mitochondria** *Biochemical Journal* **473**:4457–4471 <https://doi.org/10.1042/BCJ20160732>
30. Krabbendam IE, Honrath B, Culmsee C, Dolga AM (2018) **Mitochondrial Ca²⁺-activated K⁺ channels and their role in cell life and death pathways** *Cell Calcium* **69**:101–111 <https://doi.org/10.1016/j.ceca.2017.07.005>
31. Kulawiak B, Szewczyk A (2022) **Current Challenges of Mitochondrial Potassium Channel Research** *Front Physiol* **13** <https://doi.org/10.3389/fphys.2022.907015>
32. Lape M, Elam C, Versluis M, Kempton R, Paula S (2008) **Molecular determinants of sarco/endoplasmic reticulum calcium ATPase inhibition by hydroquinone-based compounds** *Proteins* **70**:639–649 <https://doi.org/10.1002/prot.21542>
33. Leary SC, Battersby BJ, Hansford RG, Moyes CD (1998) **Interactions between bioenergetics and mitochondrial biogenesis** *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1365**:522–530 [https://doi.org/10.1016/S0005-2728\(98\)00105-4](https://doi.org/10.1016/S0005-2728(98)00105-4)
34. Li M, Tian P, Zhao Q, Ma X, Zhang Y (2023) **Potassium channels: Novel targets for tumor diagnosis and chemoresistance** *Front Oncol* **12** <https://doi.org/10.3389/fonc.2022.1074469>
35. Li M, Xiong Z-G (2011) **Ion channels as targets for cancer therapy** *Int J Physiol Pathophysiol Pharmacol* **3**:156–166
36. Liu H, Moczydlowski E, Haddad GG (1999) **O₂ deprivation inhibits Ca²⁺-activated K⁺ channels via cytosolic factors in mice neocortical neurons** *J Clin Invest* **104**:577–588 <https://doi.org/10.1172/JCI7291>
37. Losano JDA, Padín JF, Méndez-López I, Angrimani DSR, García AG, Barnabe VH, Nichi M (2017) **The Stimulated Glycolytic Pathway Is Able to Maintain ATP Levels and Kinetic Patterns of Bovine Epididymal Sperm Subjected to Mitochondrial Uncoupling** *Oxidative Medicine and Cellular Longevity* **2017**:1–8 <https://doi.org/10.1155/2017/1682393>
38. Mohr CJ *et al.* (2022) **Subunits of BK channels promote breast cancer development and modulate responses to endocrine treatment in preclinical models** *British J Pharmacology* **179**:2906–2924 <https://doi.org/10.1111/bph.15147>
39. Müller CE, Baqi Y, Namasivayam V. (2020) **Agonists and Antagonists for Purinergic Receptors In: Pelegrín P, editor Purinergic Signaling, Methods in Molecular Biology.** New York, NY: Springer New York :45–64 https://doi.org/10.1007/978-1-4939-9717-6_3
40. Naguib A *et al.* (2018) **Mitochondrial Complex I Inhibitors Expose a Vulnerability for Selective Killing of Pten-Null Cells** *Cell Reports* **23**:58–67 <https://doi.org/10.1016/j.celrep.2018.03.032>
41. Nazemi M, Rainero E (2020) **Cross-Talk Between the Tumor Microenvironment, Extracellular Matrix, and Cell Metabolism in Cancer** *Front Oncol* **10** <https://doi.org/10.3389/fonc.2020.00239>

42. N'Gouemo P (2014) **BK Ca channel dysfunction in neurological diseases** *Front Physiol* **5** <https://doi.org/10.3389/fphys.2014.00373>
43. Oeggerli M *et al.* (2012) **Role of KCNMA1 in Breast Cancer** *PLoS ONE* **7** <https://doi.org/10.1371/journal.pone.0041664>
44. Ouadid-Ahidouch H, Ahidouch A (2013) **K⁺ channels and cell cycle progression in tumor cells** *Front Physiol* **4** <https://doi.org/10.3389/fphys.2013.00220>
45. Pallotti F, Lenaz G (2007) **Isolation and Subfractionation of Mitochondria from Animal Cells and Tissue Culture Lines** *Methods in Cell Biology* Elsevier :3–44 [https://doi.org/10.1016/S0091-679X\(06\)80001-4](https://doi.org/10.1016/S0091-679X(06)80001-4)
46. Palmer AE, Giacomello M, Kortemme T, Hires SA, Lev-Ram V, Baker D, Tsien RY (2006) **Ca²⁺ indicators based on computationally redesigned calmodulin-peptide pairs** *Chem Biol* **13**:521–530 <https://doi.org/10.1016/j.chembiol.2006.03.007>
47. Palmer AE, Jin C, Reed JC, Tsien RY (2004) **Bcl-2-mediated alterations in endoplasmic reticulum Ca²⁺ analyzed with an improved genetically encoded fluorescent sensor** *Proc Natl Acad Sci USA* **101**:17404–17409 <https://doi.org/10.1073/pnas.0408030101>
48. Quanz M *et al.* (2018) **Preclinical Efficacy of the Novel Monocarboxylate Transporter 1 Inhibitor BAY-8002 and Associated Markers of Resistance** *Molecular Cancer Therapeutics* **17**:2285–2296 <https://doi.org/10.1158/1535-7163.MCT-17-1253>
49. Rossi A, Pizzo P, Filadi R (2019) **Calcium, mitochondria and cell metabolism: A functional triangle in bioenergetics** *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* **1866**:1068–1078 <https://doi.org/10.1016/j.bbamcr.2018.10.016>
50. Salter M, Hicks J (1995) **ATP causes release of intracellular Ca²⁺ via the phospholipase C beta/IP3 pathway in astrocytes from the dorsal spinal cord** *J Neurosci* **15**:2961–2971 <https://doi.org/10.1523/JNEUROSCI.15-04-02961.1995>
51. Martín A San, Ceballo S, Ruminot I, Lerchundi R, Frommer WB, Barros LF (2013) **A Genetically Encoded FRET Lactate Sensor and Its Use To Detect the Warburg Effect in Single Cancer Cells** *PLoS ONE* **8** <https://doi.org/10.1371/journal.pone.0057712>
52. Schroth W *et al.* (2020) **Gene Expression Signatures of BRCAness and Tumor Inflammation Define Subgroups of Early-Stage Hormone Receptor-Positive Breast Cancer Patients** *Clinical Cancer Research* **26**:6523–6534 <https://doi.org/10.1158/1078-0432.CCR-20-1923>
53. Seyfried TN, Shelton LM (2010) **Cancer as a metabolic disease** *Nutr Metab (Lond)* **7** <https://doi.org/10.1186/1743-7075-7-7>
54. Siemen D, Loupatatzis C, Borecky J, Gulbins E, Lang F. (1999) **Ca²⁺-Activated K Channel of the BK-Type in the Inner Mitochondrial Membrane of a Human Glioma Cell Line** *Biochemical and Biophysical Research Communications* **257**:549–554 <https://doi.org/10.1006/bbrc.1999.0496>
55. Singh H, Lu R, Bopassa JC, Meredith AL, Stefani E, Toro L (2013) **mitoBK Ca is encoded by the Kcnma1 gene, and a splicing sequence defines its mitochondrial location** *Proc Natl Acad Sci USA* **110**:10836–10841 <https://doi.org/10.1073/pnas.1302028110>

56. Skalska J *et al.* (2008) **A novel potassium channel in skeletal muscle mitochondria** *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1777**:651–659 <https://doi.org/10.1016/j.bbabi.2008.05.007>
57. Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, Bray F (2021) **Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries** *CA A Cancer J Clin* **71**:209–249 <https://doi.org/10.3322/caac.21660>
58. Szabo I, Szewczyk A (2023) **Mitochondrial Ion Channels** *Annu Rev Biophys* **52**:229–254 <https://doi.org/10.1146/annurev-biophys-092622-094853>
59. Szewczyk A, Jarmuszkiewicz W, Kunz WS (2009) **Mitochondrial potassium channels** *IUBMB Life* **61**:134–143 <https://doi.org/10.1002/iub.155>
60. Wacquier B, Combettes L, Dupont G (2019) **Cytoplasmic and Mitochondrial Calcium Signaling: A Two-Way Relationship** *Cold Spring Harb Perspect Biol* **11** <https://doi.org/10.1101/cshperspect.a035139>
61. Warburg O (1924) **Warburg O. 1924. Über den Stoffwechsel der Carcinomzelle.** *Naturwissenschaften* **12**:1131–1137. doi:10.1007/BF01504608 *Naturwissenschaften* **12**:1131–1137 <https://doi.org/10.1007/BF01504608>
62. Wrzosek A, Augustynek B (2020) **Mitochondrial Potassium Channels as Druggable Targets** *Biomolecules* **10** <https://doi.org/10.3390/biom10081200>
63. Wu P, Gao W, Su M, Nice EC, Zhang W, Lin J, Xie N (2021) **Adaptive Mechanisms of Tumor Therapy Resistance Driven by Tumor Microenvironment** *Front Cell Dev Biol* **9** <https://doi.org/10.3389/fcell.2021.641469>
64. Wu SZ *et al.* (2021) **A single-cell and spatially resolved atlas of human breast cancers** *Nat Genet* **53**:1334–1347 <https://doi.org/10.1038/s41588-021-00911-1>
65. Xu W, Liu Y, Wang S, McDonald T, Van Eyk JE, Sidor A, O'Rourke B. (2002) **Cytoprotective Role of Ca²⁺ - Activated K⁺ Channels in the Cardiac Inner Mitochondrial Membrane** *Science* **298**:1029–1033 <https://doi.org/10.1126/science.1074360>
66. Zhou Y, Lingle CJ (2014) **Paxilline inhibits BK channels by an almost exclusively closed-channel block mechanism** *Journal of General Physiology* **144**:415–440 <https://doi.org/10.1085/jgp.201411259>

Article and author information

Helmut Bischof

Department of Pharmacology, Toxicology and Clinical Pharmacy, Institute of Pharmacy, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany
ORCID iD: [0000-0003-2380-600X](https://orcid.org/0000-0003-2380-600X)

Selina Maier

Department of Pharmacology, Toxicology and Clinical Pharmacy, Institute of Pharmacy, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany, Dr Margarete Fischer-Bosch Institute of Clinical Pharmacology, Auerbachstraße 112, 70376 Stuttgart, Germany

Piotr Koprowski

Laboratory of Intracellular Ion Channels, Nencki Institute of Experimental Biology, Polish Academy of Sciences, 3 Pasteur Street, 02-093 Warsaw, Poland

Bogusz Kulawiak

Laboratory of Intracellular Ion Channels, Nencki Institute of Experimental Biology, Polish Academy of Sciences, 3 Pasteur Street, 02-093 Warsaw, Poland

Sandra Burgstaller

Department of Pharmacology, Toxicology and Clinical Pharmacy, Institute of Pharmacy, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany, NMI Natural and Medical Sciences Institute at the University of Tuebingen, Markwiesenstraße 55, Reutlingen 72770, Germany

Joanna Jasińska

Laboratory of Intracellular Ion Channels, Nencki Institute of Experimental Biology, Polish Academy of Sciences, 3 Pasteur Street, 02-093 Warsaw, Poland

Kristian Serafimov

Institute of Pharmaceutical Sciences, Pharmaceutical (Bio-)Analysis, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany

Dominic Gross

Department of Pharmacology, Toxicology and Clinical Pharmacy, Institute of Pharmacy, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany

Werner Schroth

Dr Margarete Fischer-Bosch Institute of Clinical Pharmacology, Auerbachstraße 112, 70376 Stuttgart, Germany

Lucas Matt

Department of Pharmacology, Toxicology and Clinical Pharmacy, Institute of Pharmacy, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany

David Arturo Juarez Lopez

Medical Clinic IV, University Hospital Tuebingen, Otfried-Müller-Straße 10, 72076 Tuebingen, Germany

Irina Bonzheim

Institute of Pathology and Neuropathology, University Hospital Tuebingen, 72076 Tuebingen, Germany

Florian A. Büttner

Dr Margarete Fischer-Bosch Institute of Clinical Pharmacology, Auerbachstraße 112, 70376 Stuttgart, Germany

Falko Fend

Institute of Pathology and Neuropathology, University Hospital Tuebingen, 72076 Tuebingen, Germany

Matthias Schwab

Dr Margarete Fischer-Bosch Institute of Clinical Pharmacology, Auerbachstraße 112, 70376 Stuttgart, Germany, Institute of Pathology and Neuropathology, University Hospital Tuebingen, 72076 Tuebingen, Germany, Image-Guided and Functionally Instructed Tumor Therapies Cluster of Excellence (iFIT), University of Tuebingen, Roentgenweg 11, 72076 Tuebingen, Germany, Department of Clinical Pharmacology, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany, Department of Biochemistry and Pharmacy, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany, German Cancer Consortium (DKTK), German Cancer Research Center, Partner Site Tuebingen, Auf der Morgenstelle 15, 72076 Tuebingen, Germany

Andreas L. Birkenfeld

Medical Clinic IV, University Hospital Tuebingen, Otfried-Müller-Straße 10, 72076 Tuebingen, Germany

Roland Malli

Gottfried Schatz Research Center, Molecular Biology and Biochemistry, Medical University of Graz, Neue Stiftingtalstraße 6, 8010 Graz, Austria, BioTechMed Graz, Mozartgasse 12, 8010 Graz, Austria
ORCID iD: [0000-0001-6327-8729](https://orcid.org/0000-0001-6327-8729)

Michael Lämmerhofer

Institute of Pharmaceutical Sciences, Pharmaceutical (Bio-)Analysis, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany

Piotr Bednarczyk

Department of Physics and Biophysics, Warsaw University of Life Sciences (SGGW), Nowoursynowska St. 159 02-776 Warsaw, Poland

Adam Szewczyk

Laboratory of Intracellular Ion Channels, Nencki Institute of Experimental Biology, Polish Academy of Sciences, 3 Pasteur Street, 02-093 Warsaw, Poland

Robert Lukowski

Department of Pharmacology, Toxicology and Clinical Pharmacy, Institute of Pharmacy, University of Tuebingen, Auf der Morgenstelle 8, 72076 Tuebingen, Germany

For correspondence: robert.lukowski@uni-tuebingen.de

Copyright

© 2023, Bischof et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Alejandro San Martín

Centro de Estudios Científicos, Chile

Senior Editor

Caigang Liu

Shengjing Hospital of China Medical University, China

Reviewer #1 (Public Review):

Bischoff et al present a carefully prepared study on a very interesting and relevant topic: the role of ion channels (here a Ca^{2+} -activated K^{+} channel BK) in regulating mitochondrial metabolism in breast cancer cells. The potential impact of these and similar observations made in other tumor entities has only begun to be appreciated. That being said, the authors pursue in my view an innovative approach to understanding breast cancer cell metabolism.

Considering the following points would further strengthen the manuscript:

Methods:

1. The authors use an extracellular Ca^{2+} concentration (2 mM) in their Ringer's solutions that is almost twice as high as the physiologically free Ca^{2+} concentration (ln 473). Moreover, the free Ca^{2+} concentration of their pipette solution is not indicated (ln 487).
2. Ca^{2+} I measurements: The authors use ATP to elicit intracellular Ca^{2+} signals. Is this then a physiological stimulus for Ca^{2+} signaling in breast cancer? What is the rationale for using ATP? Moreover, it would be nice to see calibrated baseline values of Ca^{2+} i.
3. Membrane potential measurements: It would be nice to see a calibration of the potential measurements; this would allow us to correlate the IV relationship with membrane potential. Without calibration, it is hard to compare unless the identical uptake of the dye is shown.

Does paxilline or IbTx also induce depolarization?

4. Mito-potential measurements: Why did the authors use such a long time course and preincubate cells with channel blockers overnight? Why did they not perform paired experiments and record the immediate effect of the BK channel blockers in the mito potential?
5. MTT assays are also based on mitochondrial function - since modulation of mito function is at the core of this manuscript, an alternative method should be used.

Results:

1. Fig. 5G: The number of BK "positive" mitoplasts is surprisingly low - how does this affect the interpretation? Did the authors attempt to record mitoBK current in the "whole-mitoplast" mode? How does the mitoBK current density compare with that of the plasma membrane? Is it possible to theoretically predict the number of mitoBK channels per mitochondrion to elicit the observed effects? Can these results be correlated with the immuno-localization of mitoBK channels?
2. There are also reports about other mitoK channels (e.g. $\text{Kv}1.3$, $\text{KCa}3.1$, KATP) playing an important role in mitochondrial function. Did the authors observe them, too? Can the authors speculate on the relative importance of the different channels? Is it known whether they are expressed organ-/tumor-specifically?

Reviewer #2 (Public Review):

Summary:

The large-conductance Ca^{2+} activated K^{+} channel (BK) has been reported to promote breast cancer progression, but it is not clear how. The present study carried out in breast cancer cell

lines, concludes that BK located in mitochondria reprograms cells towards the Warburg phenotype, one of the metabolic hallmarks of cancer.

Strengths:

The use of a wide array of modern complementary techniques, including metabolic imaging, respirometry, metabolomics, and electrophysiology. On the whole, experiments are astute and well-designed and appear carefully done. The use of BK knock-out cells to control for the specificity of the pharmacological tools is a major strength. The manuscript is clearly written. There are many interesting original observations that may give birth to new studies.

Weaknesses:

The main conclusion regarding the role of a BK channel located in mitochondria appears is not sufficiently supported. Other perfectible aspects are the interpretation of co-localization experiments and the calibration of Ca^{2+} dyes. These points are discussed in more detail in the following paragraphs:

1. May the metabolic effects be ascribed to a BK located in mitochondria? Unfortunately not, at least with the available evidence. While it is clear these cells have a BK in mitochondria (characteristic K^{+} currents detected in mitoplasts) and it is also well substantiated that the metabolic effects in intact cells are explained by an intracellular BK (paxilline effects absent in the BK KO), it does not follow that both observations are linked. Given that ectopic BK-DEC appeared at the surface, a confounding factor is the likely expression of BK in other intracellular locations such as ER, Golgi, endosomes, etc. To their credit, authors acknowledge this limitation several times throughout the text ("...presumably mitoBK...") but not in other important places, particularly in the title and abstract.

2. MitoBK subcellular location. Pearson correlations of 0.6 and about zero were obtained between the locations of mitoGREEN on one side, and mRFP or RFP-GPI on the other (Figs. 1G and S1E). These are nice positive and negative controls. For BK-DECRFP however, the Pearson correlation was about 0.2. What is the Z resolution of apotome imaging? Assuming an optimum optical section of 600 nm, as obtained by a 1.4 NA objective with a confocal, that mitochondria are typically 100 nm in diameter and that BK-DECRFP appears to stain more structures than mitoGREEN, the positive correlation of 0.2 may not reflect colocalization. For instance, it could be that BK-DECRFP is not just in mitochondria but in a close underlying organelle e.g. the ER. Along the same line, why did BK-RFP also give a positive Pearson? Isn't that unexpected? Considering that BK-DEC was found by patch clamping at the plasma membrane, the subcellular targeting of the channel is suspect. Could it be that the endogenous BK-DEC does actually reside exclusively in mitochondria (a true mitoBK), but overflows to other membranes upon overexpression? Regarding immunodetection of BK in the mitochondrial Percoll preparation (Fig. S5), the absence of NKA demonstrates the absence of plasma membrane contamination but does not inform about contamination by other intracellular membranes.

3. Calibration of fluorescent probes. The conclusion that BK blockers or BK expression affects resting Ca^{2+} levels should be better supported. Fluorescent sensors and dyes provide signals or ratios that need to be calibrated if comparisons between different cell types or experimental conditions are to be made. This is implicitly acknowledged here when monitoring ER Ca^{2+} , with an elaborate protocol to deplete the organelle in order to achieve a reading at zero Ca^{2+} .

4. Line 203. "...solely by the expression of BKCa-DECRFP in MCF-7 cells". Granted, the effect of BKCa-DECRFP on the basal FRET ratio appears stronger than that of BK-RFP, but it appears that the latter had some effect. Please provide the statistics of the latter against the control group (after calibration, see above).

Reviewer #3 (Public Review):

The original research article, titled "mitoBKCa is functionally expressed in murine and human breast cancer cells and promotes metabolic reprogramming" by Bischof et al, has demonstrated the underlying molecular mechanisms of alterations in the function of Ca^{2+} activated K^{+} channel of large conductance (BKCa) in the development and progression of breast cancer. The authors also proposed that targeting mitoBKCa in combination with established anti-cancer approaches, could be considered as a novel treatment strategy in breast cancer treatment.

The paper is clearly written, and the reported results are interesting.

Strengths:

Rigorous biophysical experimental proof in support of the hypothesis.

Weaknesses:

A combinatorial synergistic study is missing.