

Plastic vasomotion entrainment

Daichi Sasaki, Ken Imai, Yoko Ikoma, Ko Matsui 

Super-network Brain Physiology, Graduate School of Life Sciences, Tohoku University, Sendai 980-8577 Japan

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

[About eLife's process](#)

Reviewed preprint version 1

February 7, 2024 (this version)

Posted to preprint server

November 21, 2023

Sent for peer review

November 20, 2023

 https://en.wikipedia.org/wiki/Open_access Copyright information

Abstract

The presence of global synchronization of vasomotion induced by oscillating visual stimuli was identified in the mouse brain. Endogenous autofluorescence was used and the vessel “shadow” was quantified to evaluate the magnitude of the frequency-locked vasomotion. This method allows vasomotion to be easily quantified in non-transgenic wild-type mice using either the wide-field macro-zoom microscopy or the deep-brain fiber photometry methods. Vertical stripes horizontally oscillating at a low temporal frequency (0.25 Hz) were presented to the awake mouse and oscillatory vasomotion locked to the temporal frequency of the visual stimulation was induced not only in the primary visual cortex but across a wide surface area of the cortex and the cerebellum. The visually induced vasomotion adapted to a wide range of stimulation parameters. Repeated trials of the visual stimulus presentations resulted in the entrainment of the amplitude of the vasomotion. Horizontally oscillating visual stimulus is known to induce horizontal optokinetic response (HOKR). The amplitude of the eye movement is known to increase with repeated training sessions and the flocculus region of the cerebellum is known to be essential for this learning to occur. Here, we show a strong correlation between the average HOKR performance gain and the vasomotion entrainment magnitude in the cerebellar flocculus. Therefore, the plasticity of vasomotion and neuronal circuits appeared to occur in parallel. Efficient energy delivery by the entrained vasomotion may contribute to meeting the energy demand for increased coordinated neuronal activity and the subsequent neuronal circuit reorganization.

eLife assessment

This manuscript presents potentially **valuable** results indicating a plastic enhancement in the vasomotion response of pial cortical arterioles to external stimulation in awake mice using a wide range of external visual stimulation paradigms. The evidence for this interesting effect, with broad potential applications, is, however, **incomplete** because it is unclear whether the effect is a modulation of vasomotion rather than stimulus-driven hyperaemia and whether it is reversible. These results are relevant for scientists and clinicians interested in the regulation of blood flow in the brain.

Introduction

Although the energy consumption in the brain for information processing is super-efficient compared to modern computers, the neuronal circuit still requires an adequate energy supply for encoding information and for inducing plastic changes. Energy (i.e. oxygen and glucose) is delivered to the brain via the blood vessels. Vasodilation and vasoconstriction occurring at seconds time scale have been reported (Davis et al., 2008 [↗](#); Fujii et al., 1990 [↗](#); Hundley et al., 1988 [↗](#); Jones TW, 1853). The temporal pattern of such vasomotion is distinct from the frequency of heartbeat, which in mice is ~10 Hz. The slow vasomotion is likely beneficial for the efficient delivery of blood in small vessels as has been also suggested by simulations of micro-fluid dynamics (Meyer et al., 2002 [↗](#)). Vasomotion has also been suggested to facilitate the clearance of unwanted substances such as A β (Aldea et al., 2019 [↗](#); van Veluw et al., 2020 [↗](#)).

In this study, vasomotion dynamics in the cerebral cortex were studied through intact crania using a macro-zoom microscope. Interestingly, the global brain vasomotion temporal dynamics perfectly locked to the oscillatory visual stimulus with repeated presentations. Vasomotion response to sensory stimulus has been observed previously; however, plastic entrainment of vasomotion has never been described. Horizontally oscillating visual pattern is known to induce optokinetic response (HOKR). Repeated exposure to such a pattern results in an increased amplitude of the HOKR eye movement (Ito, 1982 [↗](#); Kanaya et al., 2023 [↗](#); Waespe and Henn, 1977 [↗](#)). Neuronal circuit plasticity required for the HOKR learning takes place in the flocculus region of the cerebellum. An optical fiber was inserted close to the flocculus and fiber photometry was performed to study the vasomotion dynamics in the deep brain region (Ikoma et al., 2023a [↗](#), 2023b [↗](#)). A concomitant increase in the HOKR performance and in the magnitude of the frequency-locked vasomotion was observed. These results suggest the plastic change of vasomotion dynamics may support and interact with the neuronal circuit plasticity. Future therapeutic strategies may be directed at promoting efficient entrainment of vasomotion, which may lead to enhanced cognitive functions.

Results

Presence of spontaneous vasomotion

Blood vessels in the brain can change their diameter through the actions of the smooth muscle cells and the pericytes (Davis et al., 2008 [↗](#); Peppiatt et al., 2006 [↗](#)). Here, we observed spontaneous vasomotion *in vivo* in unanesthetized awake mice. We sought to understand how vasomotion is regulated by sensory stimuli, how vasomotion can be entrained, and the functional significance of vasomotion plasticity in information processing in the brain.

Since anesthesia or the creation of an invasive cranial window could potentially perturb naturally occurring vasomotion, first, we studied the presence of spontaneous vasomotion in the brain of awake wild-type mice through the intact skull using a macro-zoom microscope. To visualize the blood vessels, a fluorescence dye, TexasRed dextran, was administrated through the tail vein. As shown later, blood vessels on the surface of the brain could be visualized through an intact skull when the skull is coated with a transparent resin; however, the images were inevitably blurred. Therefore, in initial experiments, the thinned-skull cranial window was created (Figure 1A [↗](#)). Spontaneous vasodilation/vasoconstriction oscillations were observed. To quantify the vasomotion dynamics, we attempted to measure the vessel diameter; however, even through the thinned-skull cranial window, the exact boundary between the vessel and the brain parenchyma was difficult to determine. Therefore, the TexasRed fluorescence intensity profile perpendicular to the axis of blood flow in the vessel was measured, and the full width at 10% maximum of the TexasRed fluorescence was taken as an indication of vessel diameter (Figure 1B-C [↗](#); 45.7 ± 9.6 (S.D.) μm , $n =$

5 vessels observations). The diameter of the selected vessels suggests that mostly the pial arteries or the arterioles were observed for the stereomicroscope studies. Later in the manuscript, we will be dealing with vasomotion dynamics observed with the optical fiber photometry methods, in which the blood vessel type under the detection of the fiber could not be identified. Spontaneous vasomotion dynamic differences between different classes of blood vessels would be of interest to study using a more sophisticated *in vivo* two-photon microscope which we do not own. Here, we focused on the more bulk dynamics observed with our limited resolution of the microscope and fiber photometry.

Spontaneous vessel diameter dynamics could be revealed; however, the quantification was pixelated due to the resolution limit of the microscope (**Figure 1E**, top). Therefore, the peak TexasRed fluorescence within the blood vessels was also quantified. The vessel diameter and the peak TexasRed fluorescence correlated well; i.e. a larger vessel diameter resulted in a higher peak fluorescence (**Figure 1F**). Higher resolution of the vasomotion dynamics could be examined with the peak fluorescence measurement method (**Figure 1E**, bottom). Spontaneous vasodilation events were observed with the near-instantaneous rise and an apparent exponential decay. The cumulative frequency of the inter-event interval, the peak amplitude of the dilated vessel diameter, and the half-width of the duration of individual events were summarized (**Figure 1G**). The vasodilation events occurred relatively periodically (15.8 ± 4.8 (S.D.) seconds), were relatively fixed in amplitude (10.4 ± 3.7 (S.D.)%), and the half-width of duration was also relatively constant (1.6 ± 0.4 (S.D.) seconds; $n = 9$ sessions from 5 vessels observations).

It is possible that even the creation of the thinned skull may perturb the naturally occurring vasomotion, as heat or vibration created by the drilling was reported to influence the surface brain tissue (Augustinaite and Kuhn, 2020). Thus, the blood vessels were observed through the intact skull coated with a transparent resin. The blood vessels could be surprisingly well resolved with this method without the use of a two-photon microscope; however, compared to the thin skull observation, the boundary between the vessel and the brain parenchyma was more vague (**Figure 2A-C**). Texas Red fluorescence intensity profile perpendicular to the vessel was often shallower than that observed with the thinned skull; however, the correlation between the full width at 10% maximum of the profile and the peak intensity was preserved (**Figure 2D**). As the intensity profile was shallow with intact skull imaging, the full width at 10% maximum of the profile is less reliable as an indicator of vessel diameter and measurements of the TexasRed intensity fluctuations would better represent the vessel volume changes stemming from vasomotion.

During these macro-zoom microscopy recordings, we noticed that an intense autofluorescence from the brain tissue could be detected with ~ 430 nm excitation and ~ 540 nm emission. Most of the autofluorescence appeared to be coming from the brain parenchyma and the inside of the blood vessels were mostly devoid of this autofluorescence. Therefore, with the simultaneous TexasRed and autofluorescence imaging, the blood vessels and the brain parenchyma could be separately visualized (**Figure 2E, F**). Dynamic fluorescence intensity changes of the TexasRed and autofluorescence were simultaneously observed (**Figure 2G**). The two fluorescence traces nearly completely mirrored each other (**Figure 2H, I**). Therefore, the inverted autofluorescence dynamics would also reflect the vasomotion dynamics. In later experiments, $\Phi 400$ μm optical fibers will be used to measure the vasomotion dynamics in the deep brain areas. To simulate the situation using images from the brain surface, $\Phi 500$ μm circular region of interest (ROI) was drawn and the average TexasRed fluorescence dynamics were plotted (**Figure 2J, K**). As apparent from the spectrogram of the trace, several cycles of oscillatory fluctuations at low frequencies (0.1-0.5 Hz) were spontaneously observed. The frequency of the oscillatory fluctuation within a cluster was not strictly fixed at a constant value. The occurrence of clusters was also not periodic, and the duration of a cluster also varied. The presence of similar low-frequency vascular oscillations has been reported in previous studies, but in those studies, the cycle of fluctuation




Figure 1

Spontaneous vasomotion detection

(A) A fluorescence image of a mouse brain through the thinned-skull cranial window using a single-photon wide-field macro-zoom microscope. TexasRed was intravenously injected and the blood vessels on the brain surface could be readily resolved. (B) Magnified image of a single blood vessel. TexasRed fluorescence intensity profile was calculated at the cross-section indicated by the blue line. (C) An example of a single-line intensity profile. (D) TexasRed intensity profiles averaged for multiple lines. The profile during the basal, undilated phase (solid line shown on the left, the same profile shown as a dotted line on the right) and the dilated phase (solid line on the right, dotted line on the left). Peak amplitudes of the intensity profiles (red cross) and full-width at 10% maximum (horizontal lines). (E) An example of a time series data of the calculated vessel diameter and TexasRed intensity. (F) Correlation of the peak amplitude of the calculated vessel diameter with the TexasRed peak intensity. (G) Characteristics of the vessel dilation events across multiple vessels were summarized. Inter-event interval (left), the peak amplitude of the vessel diameter (middle), and the half-width of the duration of individual events (right) are shown for individual sessions collected from multiple vessels (gray lines, $n = 9$ sessions from 5 vessels of 3 animals; duration of each session was from 1.5 to 11 min). Averaged cumulative histograms are shown as blue lines.




Figure 2

Autofluorescence signals could be used for vasomotion detection through the intact skull

(A) A fluorescence image of a mouse brain through the intact skull. Although with less resolution, individual blood vessels containing TexasRed could be visualized. (B) Magnified image of a single blood vessel. (C) A single-line intensity profile. (D) Averaged TexasRed intensity profile at the basal (left) and the dilated (right) states. The intensity profiles were shallower compared to the profiles analyzed with thinned-skull images. Therefore, rather than the vessel diameter estimated by the full width at 10% maximum of the intensity profile, the TexasRed intensity would likely better reflect the volume changes of the blood vessels. (E) A merged image of the TexasRed signal in the blood vessel (magenta) and the autofluorescence signal likely coming exclusively from the brain parenchyma (green). (F) Spatial intensity profile along the cross-section of TexasRed and autofluorescence signals in (E). (G) The temporal transitions of the TexasRed and the autofluorescence intensity signals at an ROI encompassing the vessel are shown in (E). The two signals mostly mirrored each other, suggesting that vessel dilation appears as the TexasRed intensity increase and the autofluorescence intensity decrease. (H) An enlarged segment of the traces is shown in (G). The inverted autofluorescence trace (light green) resembles the TexasRed intensity trace. (I) Autofluorescence intensity (AI) plotted against the TexasRed intensity (TI) for the duration of the traces shown in (G). (J) An image of the mouse brain through the intact skull. $\Phi 500\ \mu\text{m}$ circular ROI is shown (dotted yellow line). This diameter corresponds closely to the diameter ($\Phi 400\ \mu\text{m}$) of the optical fiber used in fiber photometry experiments. (K) Spontaneous TexasRed fluctuation (top) was detected through the intact skull within the ROI shown in (J). The spectrum below shows that the low temporal frequency vasomotion does not continuously occur and short phases of vascular oscillation occur with frequency not locked to a narrow specific range.

appeared to be constant and the periodic cycles continued without much interruption (Davis et al., 2008 [↗](#); Jones TW, 1853). The reason why the vasomotion detected in our system through the intact skull in awake *in vivo* mice was less periodic was unknown.

Entrainment of vasomotion with visual stimuli presentation

Horizontally oscillating visual stimuli are known to induce eye movement which follows the stimuli in head-fixed mice. This response to the stimuli is termed, horizontal optokinetic response (HOKR). Repeated presentation of the stimuli also induces a gradual increase in the amplitude of the eye movement and the small flocculus region of the cerebellum has been shown to be responsible for this learning (Aziz et al., 2014 [↗](#); Ito, 1982 [↗](#)). Immediately after TexasRed was tail vein injected, using the whole brain surface imaging with the macro-zoom microscope, TexasRed fluorescence fluctuation in response to the HOKR stimuli was studied (**Figure 3A-B** [↗](#)).




Figure 3

Vasomotion entrainment with the oscillating visual stimulus

(A) Fluorescence images were taken from the whole brain surface through the intact skull using the macro-zoom microscope in awake mice. Horizontally oscillating (at 0.25 Hz temporal frequency) vertical stripes were shown as the visual stimuli on the two perpendicular monitors placed in front of the mouse. (B) Representative image of the near-whole brain surface after intravenous injection of TexasRed. The location of ROI in V1 is shown with the yellow dotted line. (C) TexasRed signals at V1-ROI during visual stimulation are shown. After repeated sessions, the vasomotion gradually became entrained to the frequency of the visual stimulus. At the 1st training session (Novice, left), no specific peak was observed in the amplitude spectrum (bottom) of the TexRed signals (middle). After several trainings (Trained, middle), a 0.25 Hz peak (arrowhead) in the amplitude spectrum appeared. After saturated training (Expert, right), the 0.25 Hz oscillation of the TexRed signal was apparent in the raw traces (middle). (D) (upper) An example data from an Expert mouse. Immediately after the onset of a visual stimulation session, the 0.25 Hz band appeared in the TexasRed trace (left). The 0.25 Hz band dissipated immediately upon the cessation of the visual stimulation (middle). The 0.25 Hz band lasted throughout the 15-minute session in this particular mouse, albeit the amplitude became smaller toward the end of the session (right). (lower) The amplitude values at 0.25 Hz in the spectrogram were plotted against time for three different animals. The left graph is from the same Expert animal and the same session as shown in the upper graph. The two other graphs (middle, right) are from different sessions. The amplitudes at 0.25 Hz during the visual stimulation were mostly higher than those before the stimulation (horizontal gray line); however, the amplitude did not stay constant in most cases. The vasomotion appears to come in and out of the frequency-lock to the visual stimulus. (E) The peak amplitude at 0.25 Hz over the mean amplitude in the 0.1-1 Hz range (PR0.25) was calculated for each pixel in the whole brain during the visual stimulation for Novice, Trained, and Expert mice. PR0.25 increase with repeated sessions was not restricted to the visual cortex but a near-uniform increase was observed throughout the whole brain surface. The color bar indicates PR0.25. (F) The color scale for the Expert example shown in (E) is saturated. Therefore, a different color scale (right) was applied for the same data. (G) The averaged PR0.25 in the whole brain surface ROI gradually increased as the training progressed ($n = 5$ animals). Novice vs. Trained, $p = 0.039$ * $p < 0.05$; Novice vs. Expert, $p = 0.0030$ ** $p < 0.005$; paired t-test with Holm correction for multiple test. Data from individual animals are shown in gray and the averaged data across all animals are shown in red. Data are presented as mean $\pm$ SEM, as error bars. (H) TexasRed signals in ROIs at V1, V2, S1, and cerebellum (Cb). Signals in different locations fluctuated with similar frequency with similar phases.

Horizontally oscillating vertical black and white stripes were presented to the head-fixed mouse with a temporal frequency of 0.25 Hz and with a maximal visual angle (amplitude) of 17 degrees for 15 minutes per session (**Figure 3A**). In response to the first visual training session, fixed frequency oscillation of the TexasRed intensity at the primary visual cortex (V1) was not observed. TexasRed signals in these “Novice” mice, fluctuated with non-fixed low frequencies which consisted mainly of <0.3 Hz (**Figure 3C**, left). After a couple of more training sessions and just before the 4th training session, TexasRed was administered again. In these “Trained” mice, the TexasRed signal in the cortex fluctuated periodically. Fourier transform of the TexasRed signal at V1 during the 15 minutes’ training session revealed a clear sharp peak in the amplitude spectrum at 0.25 Hz, which was not observed in the Novice mice (**Figure 3C**, middle). This 0.25 Hz is the temporal frequency of the HOKR oscillatory visual stimuli. With further repeated training sessions, the TexasRed signal fluctuation further becomes extremely tuned to the frequency of the visual stimuli (“Expert”, **Figure 3C**, right). In this Expert mouse, the 0.25 Hz fluctuation in the TexasRed signal becomes visible even in the raw traces. Using this Expert mouse, the time course of the vasomotion frequency locked to the visual stimulus was examined (**Figure 3D**). Before the visual stimulus, the amplitude at 0.25 Hz was negligible; however, as soon as the visual stimulus started, a sharp rise in 0.25 Hz frequency amplitude increased. The amplitude gradually declined but stayed significant until the cessation of the visual stimulus and the vasomotion amplitude locked to the visual stimulus frequency quickly dissipated (**Figure 3D**, upper row and far left of the lower row). This characteristic of an instant onset and a steady increase was not always shared

in all animals. In many cases, the frequency locked to the 0.25 Hz came in and out throughout the visual stimulation session; however, mostly throughout the session, the vasomotion amplitude at 0.25 Hz frequency was above the baseline measured before the visual stimulation (examples in **Figure 3D** [↗](#), lower row, middle and right panel).

Visual stimulus affecting vasomotion in the V1 was more or less expected; however, does it affect other regions of the cortex? To examine this, data from the near-whole brain surface observation with a macro-zoom microscope was studied. First, the TexasRed fluorescence images were Gaussian filtered in the spatial XY dimension to take out the random noise presumably created within the imaging system. Then, the TexasRed signal fluctuation during the 15-minute session was examined for every pixel imaged. The peak of the amplitude of the TexasRed signal at 0.25 Hz temporal frequency was calculated and was divided with the mean amplitude at 0.1-1 Hz range (Peak ratio of 0.25 Hz; PR0.25). This normalization process was necessary because each region had a difference in the baseline fluctuation level of the fluorescence signal. Color-coded map of the PR0.25 was mostly uniform throughout the cortex and gradually increased from Novice, Trained, to Expert (**Figure 3E** [↗](#)). Averaged PR0.25 in the whole brain surface ROI gradually increased as the training progressed (**Figure 3G** [↗](#)). The increase in the PR0.25 was so great in the Expert that the color map was mostly saturated in **Figure 3E** [↗](#), right panel. Therefore, the color code was scaled down in **Figure 3F** [↗](#). PR0.25 in V1 and V2 was high but not particularly high compared to the primary sensory cortex (S1) or the cerebellum (Cb) or, in fact, anywhere else that could be observed. An example of the actual TexasRed signal fluctuation trace in various ROI is shown in **Figure 3H** [↗](#). The 0.25 Hz signal fluctuations were apparent in all ROI and, in addition, the phase of the signal fluctuation was also nearly synchronized ($n = 5$ animals; data not shown). These data show that frequency and phase-locked vasomotion are not restricted to the visual-related area, and enhancement and entrainment of vasomotion are observed widely throughout the surface of the brain.

The use of autofluorescence signal for examination of vasomotion

The small cerebellar flocculus region deep and away from the cranial surface is responsible for the cerebellum HOKR motor learning. To examine the vasomotion in such a deep brain region, optical fiber was implanted into the cerebellum (**Figure 4C** [↗](#)). The eye movement amplitude should gradually increase in response to repeated sessions of HOKR visual stimulation training; however, we found that, if TexasRed had to be injected before every training session, the mouse did not learn very well. As an alternative to measuring the vessel volume dynamics directly with TexasRed fluorescence, the use of fluorescence in the brain parenchyma was attempted.




Figure 4

Autofluorescence signals could be used to quantify the vasomotion entrainment

(A) Images of the TexasRed fluorescence in the blood vessels and the direct YFP fluorescence expressed in astrocytes of a transgenic mouse. The blood vessels appear as dark “shadows” in the dYFP fluorescence image. (B) An optical fiber was implanted in the cerebellum close to the flocculus and another optical fiber (highlighted in green) from the fiber photometry apparatus was connected to this implanted fiber. Horizontally oscillating visual stimulation was presented on the monitors. A hot mirror reflected the IR image of the eye and the eye location was monitored using a camera. (C) Location of the optical fiber implantation relative to the cerebellar flocculus. (D) Simultaneously recording of the TexasRed and the dYFP signals from the optical fiber implanted in the flocculus during the HOKR visual stimulation session. The dYFP mirrored the TexasRed signal fluctuations. Fluorescence signal data were filtered at 150-350 mHz. (E) 20 5-second segments of the TexasRed and the dYFP signal traces were extracted and averaged. The two traces nearly completely mirrored each other. The temporal frequency of the visual stimulation presented to the transgenic mouse was 0.20 Hz. (F) Cross-correlation between the two traces had a negative peak with a lag time of Δt of 0 sec. (G) The phase shift between TexasRed and dYFP signals was calculated for $n = 13$ segments with 5-minute durations each. (H) In a separate set of experiments, the TexasRed signal and the endogenous autofluorescence signal were simultaneously recorded in wild-type mice. 20 4-second segments were extracted and averaged. The traces were not completely inverted and a certain amount of phase shift was observed. The temporal frequency of the visual stimulation presented to the wild-type mouse was 0.25 Hz. (I) Cross-correlation of TexasRed and Autofluorescence signals. Δt of 0.8 sec was observed in this particular example. (J) Phase shifts in TexasRed and autofluorescence signals were calculated for $n = 28$ segments. (K) Averaged data of 35 traces of the extracted 4-second segments of the TexasRed and the autofluorescence signals during the 1st training (Novice) and the 4th training (Trained). Data was filtered with 150-350 mHz. An increase in the amplitudes was observed from both signals. (L) Amplitude spectrums for TexasRed and autofluorescence during the 5 minutes of the 1st training (Novice) and that of the 4th training (Trained). Signal data were high-passed filtered at 50 mHz. (M) PR0.25 of the TexasRed signals and the autofluorescence signals in the 5 minutes during the visual stimulation were plotted against each other. A clear correlation was observed indicating that the autofluorescence signal can be used to estimate the magnitude of the vasomotion frequency-locked to the visual stimulus.

As shown in **Figure 4A**, the blood vessels are devoid of the fluorescence from the YFP expressed in astrocytes, and thus the blood vessels appear as dark regions in the images. Therefore, dilation or constriction would appear as a decrease and increase in the fluorescence signal of the YFP. Such concept of “shadow” imaging of the dynamics of brain blood volume (BBV) has been demonstrated using the fiber photometry method previously (Ikoma et al., 2023a, 2023b). Here, we first used a transgenic mouse, *Mlc1-tTA::tetO-YC_{nan050}*, which expresses FRET-based CFP and YFP conjugated sensor protein specifically in astrocytes, including Bergmann glial cells (Kanamaru et al., 2014; Tanaka et al., 2012). Instead of exciting the CFP with ~ 430 nm, YFP was directly excited by delivering ~ 505 nm light through the implanted optical fiber in the cerebellum (dYFP). The dYFP fluorescence would not be affected by the Ca^{2+} , and would only be affected by the BBV and possibly the cytosolic pH, as YFP protein is sensitive to changes in the pH and YFP fluorescence can be quenched by acidification. Both TexasRed and dYFP signals simultaneously monitored with fiber photometry fluctuated with the 0.25 Hz temporal frequency directly matching the visual stimulus, but mirrored in the direction of change (**Figure 4D**). The TexasRed signal oscillations were locked to the temporal frequency of the visual stimulus. This shows that, not only in the cortex, but also deep in the cerebellum, the vasomotion can be induced with visual stimulation. Cross-correlation of the TexasRed and dYFP signals shows that a negative peak with ~ 0 sec lag time (Δt ; **Figure 4E, F**). The phase shift between TexasRed and dYFP was determined by using the lag time (Δt) and divided by the presented visual stimulation oscillation cycle time and multiplied by 360° and added 180° . The average phase shift of dYFP relative to TexasRed was $184.4^\circ \pm 17.6^\circ$ (S.D.) (**Figure 4G**; $n = 13$ segments with 5-minute durations each), indicating that these signals were nearly completely inverted. This shows that dYFP signals can be used for

assessing the vasomotion dynamics. The cytosolic pH may change; however, even if there are effects of the cytosolic pH fluctuations, pH changes appear to be in phase with the oscillatory vasomotion.

Unfortunately, robust HOKR learning could not be reliably induced in this line of transgenic mice for unknown reasons. Therefore, C57BL/6Njcl wild-type mice were used for the experiments described in the rest of the manuscript. As introduced in **Figure 2E**, autofluorescence excited with 430 nm light could be detected. Using the images taken with the macro-zoom microscope and limiting the ROI in a small region surrounding the blood vessels, mirrored traces of the TexasRed and autofluorescence were observed (**Figure 2H**). However, when the signals collected using the $\Phi 400\ \mu\text{m}$ fiber photometry were analyzed, TexasRed fluorescence and autofluorescence did not mirror completely and a significant shift in the phase was observed (**Figure 4H-J**; $246.9^\circ \pm 49.9^\circ$ (S.D.), $n = 28$ segments). The endogenous substance in the brain parenchyma that produces autofluorescence was not identified but it could be flavin or other related substances (Croce and Bottiroli, 2015; Huang et al., 2002; Islam et al., 2013). Production and degradation of flavin and other metabolites may be induced by the fluctuation in the blood vessel diameter with a fixed delay time. The phase shift in the autofluorescence could be due to the additive effect of “shadow” imaging of the vessel and to the concentration fluctuation of the autofluorescent metabolite.

The TexasRed fluorescence and the autofluorescence signals are shown for the initial sessions (Novice, **Figure 4K**) and after several training sessions (Trained). The amplitude of the 0.25 Hz temporal frequency component of both the TexasRed and the autofluorescence signals increased with training (**Figure 4K, L**). PR0.25 of autofluorescence was plotted against that of the TexasRed for multiple training sessions and a strong correlation between these two parameters was found (**Figure 4M**). Therefore, although the autofluorescence dynamics result from the fluctuations reflecting the “shadow” of vasomotion and an unknown fluctuation source, the PR0.25 apparently reflects the amplitude of the vasomotion locked to the visual stimulation cycle. In the remainder of the manuscript, the autofluorescence fluctuations were mainly examined to understand the dynamics of vasomotion.

Vasomotion followed the visual stimulation dynamics over a wide parameter range

Optimal visual stimulation properties for inducing frequency-locked vasomotion were searched. The visual stimulation used in all other sections of this study was the following. The temporal frequency of horizontal oscillation was 0.25 Hz; the spatial amplitude of the horizontal oscillation was 17° in the visual angle of the mouse; the horizontal spatial cycle of the vertical stripes was 6.4° in the visual angle. First, the temporal frequency was changed from 0.25 Hz to 0.15 Hz or 0.50 Hz while keeping other parameters fixed. Interestingly, the autofluorescence fluctuations followed the changes in the temporal frequency, and a clear sharp amplitude at the frequency locked to the presented visual stimulation was evident in the amplitude spectrum in all conditions (**Figure 5A**). The peak ratio of the autofluorescence was examined for temporal frequency ranging from 0.15 to 0.50 Hz and normalized to that of 0.25 Hz (PR0.25; **Figure 5D**, left panel). No significant difference was found, showing that vasomotion can lock to any temporal frequency within the range examined.

Similarly, the effect of changing the spatial amplitude of the horizontal oscillations was examined, while keeping other parameters fixed. A large amplitude would require the mouse to move its eyes largely within their eye sockets. The temporal frequency-locked vasomotion was apparent in the low 4° amplitude as well as in the high 25° amplitude (**Figure 5B**). Relative peak ratio normalized to that of 17° amplitude showed that the vasomotion was substantially induced in response to any amplitude between 4° and 25° (**Figure 5D**, middle panel). Lastly, the horizontal spatial cycle of the vertical stripes was varied (**Figure 5C**). Again, the vasomotion could follow




Figure 5

Vasomotion entrainment over a wide range of visual stimulation parameters

(A-C) autofluorescence signals at the flocculus in response to visual stimulation with various parameters. In the original condition, the Temporal freq (the temporal frequency of horizontal oscillation) was 0.25 Hz, the Amplitude (the spatial amplitude of the horizontal oscillation) was 17° in the visual angle of the mouse, and the Spatial cycle (the horizontal spatial cycle of the vertical stripes) was 6.4°. All the fluorescence signal data were filtered with 50-1000 mHz. (A) The temporal frequency of visual oscillation was varied. The location of the peaks in the autofluorescence signal amplitude spectrogram exactly matched the frequency of the visual stimulus in almost all cases (arrowhead). (B) The amplitude of the visual oscillation was varied. (C) The spatial cycle of the stripe patterns was varied. (D) The relative peak ratio of the autofluorescence signal at the temporal frequency matching that of the presented visual stimulus was calculated. This relative peak ratio was normalized to that of the original condition. Varying the Temporal freq, the Amplitude, or the Spatial cycle almost had no effect on the relative peak ratio. This suggests that vasomotion can tune in to any visual stimulation within the parameter range that we tested. Significant differences were shown only when the Spatial cycle was varied (Spatial cycle, 6.4° (original) vs. 9°, $p = 0.0028$ * $p < 0.0125$; 6.4° (original) vs. 18°, $p = 0.015$ * $p < 0.017$; one-sample t-test with Holm correction for multiple tests). Individual mice's data (as an average from the 3 sessions of the 5-minute test) are shown in pale green, and the average data from the examined mice ($n = 5$ mice) are shown in dark green.

the visual stimuli irrespective of the fineness of the presented stripes (**Figure 5D** [↗](#), right panel). The combination of parameters of the visual stimuli used in the rest of the manuscript was well within the adequate range that would induce maximal visually induced vasomotion.

Vasomotion is not the consequence of eye movements

We have shown above that the bulk dynamics of vasomotion under the optical fiber could be quantified with autofluorescence intensity fluctuations using fiber photometry. HOKR visual stimulation induces both eye movement and frequency-locked vasomotion. Here, we examined whether the eye movement itself is the cause of vasomotion.

First, the right eye trace in response to the horizontally oscillating visual stimulus was examined (**Figure 6A** [↗](#)). The eye tracked the visual stimulation well and there was virtually no phase shift in the eye trace relative to the visual stimulation cycle (**Figure 6B** [↗](#)). The TexasRed signal fluctuation in the flocculus in the right hemisphere was oscillatory with the visual stimulation; however, the phase of the TexasRed signal was either 0° or 180° relative to the visual stimulation, depending on the animal (**Figure 6C** [↗](#)). This indicates that in the 0° phase-shifted animals, the maximum vasodilation and vasoconstriction occur when the right eye is at the maximal nasal and temporal locations, respectively. Conversely, in the 180° phase-shift animals, it is the other way around. In either mouse, the maximum vasodilation and vasoconstriction occur when the speed of the eye movement reaches zero. The reason why the phase of the vasomotion becomes fixed to 0° or 180°, depending on the animal, is unresolved.

As the vasomotion apparently followed the visual stimuli unbelievably well, we suspected several artifacts affecting the fluorescence recording. First, we suspected that the visual stimulation itself is penetrating the back of the eye, through the brain, and to the flocculus of the cerebellum. Thus, the photo intensity detected with the optical fiber could be coming directly from the oscillatory visual stimulus. To address this possibility, the mouse was anesthetized by isoflurane (**Figure 6D** [↗](#)). Under anesthesia, the eye fails to follow the screen. The vasomotion was detected with the “shadow” imaging method by studying the dYFP fluctuations. No oscillatory signals were detected in the dYFP. Next, in the TexasRed injected mouse, the HOKR visual stimulation was presented, which induced a nice eye movement following the visual stimulus (**Figure 6E** [↗](#)). However, in this case, we did not send the excitation light through the optical fiber but kept the PMT active. If the visual stimulus on the presented screen is somehow reaching the optical fiber, then it should be detected in this condition, but it was not.

Finally, we investigated the possibility that the fluorescence signal fluctuation is due to the motion artifact. The physical eye movement could produce a movement in the brain tissue and the optical fiber location relative to the blood vessels could be moved, leading to the fluctuation of the fluorescence signals. When the mouse is presented with visual stimulation, in the trained mouse, the eye usually follows the visual stimulus completely. However, even during a single session, the frequency-locked TexasRed signal was not always apparent. In an example shown in **Figure 6F** [↗](#), the eye trace was completely in sync with the visual stimulation either in the left panel segment or in the right. However, TexasRed signal fluctuation with an amplitude spectrogram of 0.25 Hz was not apparent in the left panel segment but was significant in the right panel segment. This observation argues against the possibility that the detected fluorescence fluctuation is the direct result of eye movement. The amplitude of the oscillatory eye movement was plotted against the PR0.25 of the TexasRed during a single visual session (**Figure 6G** [↗](#)). The cycle of visual stimulation producing either a large or small eye movement did not always result in a large or small amplitude fluctuation of the vasomotion, respectively, and no correlation was found.




Figure 6

Fluorescence fluctuations are not the results of eye movement artifact

(A) An example of the visual stimulation time course, the eye trace, the TexasRed signal, and the autofluorescence signal fluctuations are shown. The data was filtered with 50–500 mHz. (B) The phase of the eye trace relative to the visual stimulus. The visual stimulus was presented with a 4-second cycle (0.25 Hz), thus, the eye trace data was segmented with 4-second durations during the first 5 minutes of each training. The segmented data were aligned, averaged, and fitted with a sine curve. The phase difference between the visual stimulus and the fit to the averaged eye trace was calculated. The mean of the phase in each animal across multiple sessions is shown as a vertical black bar. The average phase lag was $2.3 \pm 1.0^\circ$ in 5 animals, indicating that the eye trace synchronize nearly completely with the visual stimulus. (C) The phase of the TexasRed signal fluctuations relative to the visual stimulus. The average TexasRed signal fluctuation with a 4-second cycle was calculated from segments extracted from several sessions, each with 1–6 minutes in duration. The phase lag of the TexasRed signal relative to the visual stimulus was clustered either at 0° or 180° . This shows that the local brain blood volume in the right flocculus becomes the largest when the right eye is either in the most nasal position or the temporal position. (D) Under anesthesia with isoflurane, the eye movement was not induced. The dYFP signal from the YCnano50 expressed in astrocytes also did not oscillate. (E) With no excitation light sent to the optical fiber, the PMT detected no signal (0 V). The presented visual stimulus did not reach the fiber optics to directly induce oscillatory signals. (F) The amplitude of the eye movement was normally nearly constant during a single session; however, slightly strong (left) and weak (right) eye movements were observed. The TexasRed signal fluctuation at the frequency locked to the visual stimulus was not constant during the session and PR0.25 of the TexasRed signal could be seen in some (left) but not the other (right) segments of the session. (G) PR0.25 of TexasRed signal was plotted against the oscillatory eye movement amplitude in 40-second segments within a single visual stimulation session (15 min). The first segment was excluded ($n = 21$ segments total). No correlation was found between these two factors ($R^2 = 0.12$).

Possibility of the contribution of vasomotion entrainment to the HOKR performance

Repeated HOKR trainings usually result in a larger amplitude of the eye oscillations, which reaches close to the actual visual presentation amplitude. We have shown above that the frequency-locked vasomotion in the cortex also got entrained with repeated training sessions (**Figure 3**). It has been shown that HOKR performance increase relies on the neuronal and glial activity in the cerebellar flocculus (Kanaya et al., 2023). We next evaluated whether the plasticity of the vasomotion dynamics in the flocculus correlates with the HOKR performance increase.

HOKR-spaced training paradigm was used. Briefly, 15-minute training sessions were repeated with 1-hour intervals 4 times on Day 1 (i.e. the training day; **Figure 7A**). Averaged amplitudes of the eye movement during the first and last 3 minutes of each training session were calculated. The calculated averaged eye amplitudes were normalized to the initial amplitude measured at the first 3 minutes of the first training session. This value was termed the “relative gain”. A brief 2-minute tests were done on Day 2 and Day 5 to evaluate the long-term changes in the performance. Along with the HOKR eye movement measurements, autofluorescence signals were measured with the optical fiber implanted close to the flocculus. The amplitude of the autofluorescence signal at a temporal frequency locked to the visual stimulation frequency (0.25 Hz) was analyzed. As above, this amplitude was divided by the mean amplitude in the 0.1-1 Hz range to derive the PR0.25. As the autofluorescence fluctuations were often noisy, the autofluorescence signals during the first and last 5 minutes of each training session were used for the PR0.25 calculation. The long-term memory tests done on Day 2 and Day 5 were only 2 minutes in duration, thus, the full 2 minutes were used for the PR0.25 calculations.

As described previously, HOKR performance increased with repeated training sessions during Day 1 (**Figure 5B-D**). With the current experimental conditions, the average HOKR performance tested on Day 2 and Day 5 was nearly the same as the performance on the last session of Day 1; however, long-term changes in the performance were sometimes apparent in individual cases (**Figure 7C**). The PR0.25 of the autofluorescence signal changes with the spaced training protocol basically followed the plastic changes of the eye movement (**Figure 7E**). PR0.25 tended to increase with repeated training sessions.

The HOKR relative gain was calculated for each test (i.e. 10 tests total; the early and the late time tests of each of the 4 sessions on Day 1, and tests on Day 2 and Day 5), and the HOKR relative gain was averaged across all animals examined ($n = 15$ mice). Similarly, the PR0.25 in autofluorescence fluctuations were averaged across animals in 10 tests. The average HOKR relative gains were plotted against the average PR0.25 values and a strong correlation was found (**Figure 7F**; each data point represents an average score of $n = 15$ mice and 10 data points are shown for the 10 tests). In addition, it should be pointed out that a large variance was found both in the HOKR relative gain and the PR0.25 of the autofluorescence fluctuations across animals (X and Y error bars indicate S.D. in **Figure 7F**). Therefore, in individual animal recordings, the strong correlation between the HOKR relative gain and the PR0.25 of the autofluorescence fluctuations may not always be apparent. It is possible that the plasticity governing HOKR and vasomotion increases could be induced by a parallel process.

Correlations between HOKR and vasomotion across individual animals were searched. HOKR relative gain on Day 5 normalized to that for the 4th training on Day 1 was plotted against the PR0.25 of the autofluorescence fluctuations during the last 5 minutes of the 4th training session on Day 1 (**Figure 7G**). Interestingly, a moderate but significant correlation between these parameters was found ($n = 15$ mice). This may suggest that the level of visually-induced




Figure 7

A concomitant increase in the HOKR performance and vasomotion entrainment

(A) Schematic schedule of the HOKR spaced training paradigm. The 15-minute training sessions were repeated at 1-hour intervals for 4 times on Day 1 (the training day). The average amplitudes of the eye movement during the first and last 3 minutes of each training session were analyzed (during the period indicated by the blue box in the schematics). The 2-minute tests measuring long-term memory were done on Day 2 and Day 5. (B) Representative traces of the visual stimulus, the eye trace, and the autofluorescence during the 1st and the 4th training on Day 1 and the test done on Day 2. The amplitude of the eye movement and the vasomotion gradually increased in amplitude with repeated sessions. (C) The eye traces and the autofluorescence traces were segmented in 4-second durations, corresponding to the visual stimulation frequency of 0.25 Hz, and aligned and averaged. (D) The amplitude of the eye movement relative to that of the start of the 1st training session on Day 1 was refined as the HOKR relative gain. Average HOKR relative gain across $n = 15$ animals is plotted (mean $\pm$ SEM). (E) PR0.25 of the autofluorescence signals was averaged across $n = 15$ animals and plotted against sessions. (F) Averaged HOKR relative gain was plotted against the averaged PR0.25 of autofluorescence signals. Error bars indicated S.D. ($n = 15$ animals). (G) HOKR relative gain on Day 5 normalized to that on the 4th training on Day 1 was plotted against the PR0.25 of autofluorescence signals at the end of the 4th training on Day 1 ($n = 15$ animals).

vasomotion contributes to the consolidation of HOKR performance in later days. These results imply a complex and intimate relationship between vasomotion and neuronal circuit plasticity, learning, and memory.

Discussion

Vasomotion, defined as the rhythmical contraction and relaxation of the blood vessel, was first described in the vein of the bat wing (Jones, 1853 [\[1\]](#)) and was subsequently observed mainly in the arteries and arterioles of numerous species (Fuji et al., 1990 [\[2\]](#); Hundley et al., 1988 [\[3\]](#)). These vasomotions are not the secondary effects from the pulsation of blood pressure induced by the heartbeat, which in mice is a lot faster (~10 Hz; Ho et al., 2011 [\[4\]](#); Kramer et al., 1993 [\[5\]](#)) compared with the vasomotion that was observed in this study (~0.1 – 0.5 Hz).

Spontaneous vasomotion requires coordinated activity of the vascular smooth muscle cells (SMCs). Oscillatory release of calcium ions from the sarcoplasmic reticulum results in oscillations in the membrane potential of SMCs. Via the gap junctions between SMCs, synchronized oscillations are created (Haddock et al., 2006 [\[6\]](#); Haddock and Hill, 2002 [\[7\]](#); Hill et al., 1999 [\[8\]](#); Peng et al., 2001 [\[9\]](#); Schuster et al., 2000 [\[10\]](#)). Blood flow in the central nervous system was assumed to be regulated primarily by the SMCs in precapillary arterioles; however, capillaries lacking smooth muscle have been shown to be controlled by pericytes (Peppiatt et al., 2006 [\[11\]](#)). The spontaneous vasomotion observed in our study could be the result of constriction and dilation of arterioles or capillaries controlled by SMCs or pericytes, respectively. Astrocytes also constitute the neurovascular unit; however, whether astrocytic calcium is involved in controlling local cerebral blood flow is in question (Ozawa et al., 2023 [\[12\]](#)). It is still possible that astrocytes could regulate vasomotor amplitudes via stretch-mediated feedback (Bazargani and Attwell, 2016 [\[13\]](#); Haidey et al., 2021 [\[14\]](#)).

Spontaneous vasomotion has been suggested to be driven by the periodic neural activity in the resting-state (Martuzzi et al., 2009 [\[15\]](#); Mateo et al., 2017 [\[16\]](#)). Neuronal projections from the neuronal nuclei projecting to the whole brain may be controlling and synchronizing vasomotion via neurovascular coupling (Korte et al., 2023 [\[17\]](#)). It is also possible that ultra-slow vasomotion could be driven not only by the neuronal rhythms, but also by the oscillatory systemic changes, such as blood pressure and carbon dioxide in the blood (Panerai et al., 1998 [\[18\]](#); Sassaroli et al., 2012 [\[19\]](#); Wise et al., 2004 [\[20\]](#)).

In addition to spontaneous vasomotion, functional hyperemia has been shown to be induced by visual stimulation (Heeger et al., 2000 [\[21\]](#); Kayser, 2004 [\[22\]](#); Niessing et al., 2005 [\[23\]](#); Rivadulla et al., 2011 [\[24\]](#); Shaw et al., 2021 [\[25\]](#); Stickland et al., 2019 [\[26\]](#)). Visual stimulation evokes neuronal action potential firing and through the neurovascular coupling mechanisms, vasomotion is affected. If a pulsatile visual stimulus is periodically presented, that would result in vasomotion fluctuation with the same frequency as the presented visual stimulus (Kayser, 2004 [\[22\]](#); van Veluw et al., 2020 [\[27\]](#)). Previously, such visual stimulus-evoked vasomotion was reported to be restricted to the visual cortex (van Veluw et al., 2020 [\[27\]](#)). However, in our study, we show that vasomotion entrainment occurs with repeated HOKR visual stimulation, and the coordinated vasomotion, which is frequency-locked to the visual stimulus, was observed not only in the visual cortex but also widely throughout the surface of the cortex and the cerebellum, and also deep in the flocculus region of the cerebellum. It is possible that vasomotion entrainment is induced in other areas as well, including the limbic system and the brain stem.

We observed in our study that repeated visual stimulus results in vasomotion entrainment. Repeated visual stimulus may induce synaptic plasticity at the neuronal projections from the retina to a central nucleus that controls a brain-wide vasomotion. The neuronal projections from the central nucleus to the rest of the brain could also be plastically changed. It is also possible that

plasticity may occur not in the neuronal circuits but in the neurovascular coupling mechanisms. The constriction and dilation mechanisms in the SMCs and pericytes could be the actual target of plasticity leading to vasomotion entrainment.

Vasomotion could be beneficial for regulating the blood flow and the vascular surface area, which would affect the supply of nutrients (e.g. glucose) and oxygen in the brain. Mathematical modeling has shown that effective vascular resistance decreases with vasomotion (Meyer et al., 2002 [\[1\]](#)). Low-frequency vasomotion has been shown to be beneficial for tissue oxygenation, especially in regions where perfusion is restricted (Davis et al., 2008 [\[2\]](#); Nilsson and Aalkjaer, 2003 [\[3\]](#)). As astrocyte processes surround the blood vessels, the coordinated activity of astrocytes is presumed to be important for the efficient delivery of the energy source to neurons for information coding and plasticity. Furthermore, it was recently reported that vasomotion could also drive paravascular clearance or intramural periarterial drainage (IPAD) (van Veluw et al., 2020 [\[4\]](#)). Initially, arterial pulsation from the heart was considered to be the driving force for the IPAD; however, simulation modeling suggested that vasomotion was required for the efficient IPAD (Aldea et al., 2019 [\[5\]](#)).

40-Hz sensory stimulation has been proposed to be potentially beneficial for the treatment of Alzheimer's disease (AD) (Adaikkan et al., 2019 [\[6\]](#); Iaccarino et al., 2018 [\[7\]](#); Martorell et al., 2019 [\[8\]](#)). However, it was recently claimed that the 40-Hz flickering light does not entrain gamma oscillations and suppress amyloid beta (A β) accumulation in the brains of the AD model mouse (Soula et al., 2023 [\[9\]](#)). In another study, it was shown that flashing visual stimuli could increase the amplitude of vasomotion, which resulted in an increase in the paravascular clearance rates in the visual cortex of awake mice (van Veluw et al., 2020 [\[4\]](#)). In mice with cerebral amyloid angiopathy, impaired paravascular clearance and A β accumulation occur (Charidimou et al., 2017 [\[10\]](#); Peng et al., 2016 [\[11\]](#)). It was also suggested that low-frequency arteriolar oscillations could drive the drainage of soluble waste products (van Veluw et al., 2020 [\[4\]](#)).

In our study, we show that the HOKR-driving oscillatory visual stimulus, which is much slower (0.25 Hz) than the 40 Hz, is effective in producing a brain-wide oscillatory vasomotion. In addition, repeated presentation with the spaced training protocol was effective in driving the plastic vasomotion entrainment. Paravascular clearance and drainage of waste products, such as A β , may be enhanced with the vasomotion entrainment. We show that synaptic plasticity change related to cerebellar motor learning is enhanced in parallel with the vasomotion entrainment. It is possible that vasomotion enhancement may increase the meta-plasticity and provide a basis for efficient neuronal plasticity to occur. In the cerebellum, neuron-to-astrocyte (or Bergmann glia) (Matsui et al., 2005 [\[12\]](#); Matsui and Jahr, 2004 [\[13\]](#), 2003 [\[14\]](#)) and astrocyte-to-neuron (Beppu et al., 2021 [\[15\]](#), 2014 [\[16\]](#); Sasaki et al., 2012 [\[17\]](#)) signaling pathways have been identified and such neuron-astrocyte interaction has been shown to be essential for the online HOKR learning to occur (Kanaya et al., 2023 [\[18\]](#)). Late offline HOKR learning was dependent in part on the phagocytosis activity of astrocytes (Morizawa et al., 2022 [\[19\]](#)). It is possible that the energy required for the long-term plastic changes in the brain is supported by the cooperative actions of the astrocytes and the entrained vasomotion. A β has been shown to constrict capillaries and the resulting slight hypoxia and neuronal damage may actually be the cause of dementia in AD (Nortley et al., 2019 [\[20\]](#)). The slow sensory stimulation may possibly be effective for AD treatment; however, evidence of enhanced A β drainage and cognitive decline resistance with such sensory stimulus using AD model mice would be required in the future to support such a claim.

Materials and Methods

Animals

All experiments were performed in accordance with the recommendations of the Regulations for Animal Experiments and Related Activities at Tohoku University. Efforts were made to minimize animal suffering and to reduce the number of animals used. Mice were housed individually for more than 7 days under standard laboratory conditions with food and water *ad libitum*.

C57BL/6NJcl male mice (2–4 months old) were used in this study in order to reduce the variance of HOKR performance among animals (Aziz et al., 2014 [DOI](#)). Mlc1-tTA::tetO-YC_{nano50} double transgenic mice were also used, which express the tetracycline transactivator (tTA) under the astrocyte-specific Mlc1 promoter (Tanaka et al., 2012 [DOI](#)). The expressed tTA activates the tetO promoter and the FRET-based Ca²⁺ sensor YC_{nano50} is expressed specifically in astrocytes, including the cerebellar Bergmann glial cells (Kanemaru et al., 2014 [DOI](#)). Astrocyte Ca²⁺ dynamics associated with vasomotion were of potential interest to study; however, HOKR learning could not be reliably induced in Mlc1-tTA::tetO-YC_{nano50} transgenic mice for unknown reasons. Thus, we concentrated on the vasomotion dynamics resolved by the autofluorescence detection for most of the manuscript. Only the YFP part of the CFP-YFP conjugated protein of YC_{nano50} was used to assess the “shadow” imaging principle for detecting vasomotion dynamics.

Wide-field macro-zoom microscope

For surgical preparation, mice were anesthetized under ~5% isoflurane for induction and 0.5–2.0% isoflurane mixed with the air for maintenance. The scalp was cut open and the periosteum was gently removed to expose the crania. A stainless steel chamber frame (CF-10; Narishige, Tokyo, Japan) was used as a holding jig to place the head in place under the microscope. The chamber frame was attached to the edges of the exposed crania of the mouse to avoid obstruction of the view of the cortex and the cerebellum using a synthetic resin (Super bond C & B; Sun Medical Company, Shiga, Japan) mixed with charcoal particles. In the thinned skull imaging, the skull was gently thinned with a drill with precautions not to penetrate the skull. In both thinned skull and intact skull imaging, the crania were covered with a UV-curing resin (Jelly Nail; IML Inc., Tokyo, Japan) to prevent desiccation of the bone. The resin greatly improved the transparency of the skull and it could protect the field of view for over weeks. The resin and the short UV exposure are commonly applied to human fingernails without health concerns. Even after the short UV exposure for curing resin, fluorescence proteins in the mouse brain were not photobleached. A tube made of black aluminum was attached to the top of the mouse's head so as not to obstruct the view of the microscope. All gaps between the head and this custom-made tube were sealed with the Jelly Nail resin with charcoal particles to prevent light from the video monitor used for HOKR visual stimulus from leaking in. The mice were allowed to recover from the surgery for at least 3 days.

Prior to setting the mouse under the microscope, the mouse was intravenously injected via the tail vein with TexasRed dextran (70,000 MW, Invitrogen; prepared 25 mg/kg in saline at 0.1 ml). Immediately after the injection, the mouse was mounted on a chamber holder (MAG-1 and MAG-A; Narishige) using the chamber frame fixed to its skull. The mouse's body was loosely restrained with a plastic cylinder. The head-fixed mouse was set under a wide-field macro-zoom microscope (MVX10; Olympus, Tokyo, Japan) equipped with an image splitting optics (W-VIEW GEMINI A12801-01; Hamamatsu) and a digital CMOS camera (ORCA-Flash4.0 V3; Hamamatsu Photonics, Shizuoka, Japan). An inverted cone made of a black opaque sheet was attached to the objective lens to prevent the outside light from leaking in and to prevent the fluorescence excitation light from the microscope from reaching the mouse's vision and startling the mouse. The inverted cone was capped by the tube attached to the mice's head, thus, the light insulation was complete.

The excitation light to the microscope was provided by an LED-based light source (Niji; Blue Box Optics, Blackwood, UK). For excitation of the TexasRed, excitation of the autofluorescence, and direct excitation of the YFP (dYFP) of the YC_{nano50}, were accomplished with 560 nm LED light source with a bandpass filter of ET580/25x (Chroma, VT, USA), 440 nm LED light source with a bandpass filter ET430/24x (Chroma), and 490 nm LED light source with a bandpass filter ET505/20x (Chroma), respectively. A multi-band beamsplitter (69008bs; Chroma) was used to send multiple excitation wavelength lights to the mouse while allowing the emission lights of the TexasRed, the autofluorescence, and the dYFP to pass through. When necessary, the emission light of the TexasRed and that of the autofluorescence or the dYFP were split using a dichroic mirror (T590lpxr, Chroma). The wavelength of the emission lights was restricted using a custom bandpass filter (passband: 473±12 nm and 625±15 nm, Chroma), a bandpass filter ET540/40m (Chroma), or a bandpass filter ET540/30m (Chroma) for the detection of TexasRed, autofluorescence, or dYFP, respectively.

The frame exposure time of the camera was 89 msec and frames were recorded at 5 Hz sampling frequency. The magnification of the microscope was x0.8 or x6.3. All settings of the camera were controlled by the HCImageLive software (Hamamatsu). The power of excitation light at the wavelengths of 560 nm and 440 nm delivered from Niji were 1.7 mW and 3.2 mW in x0.8 magnification, respectively. The power of excitation light with the wavelength of 560 nm, 440 nm, and 490 nm in x6.3 magnification were 8 mW, 13 mW, and 4 mW, respectively. In all of the recording configurations, each excitation light was sequentially delivered in 91 msec pulses with 109 msec intervals.

Fiber photometry

Prior to surgical implantation of the optical fiber, the mouse was anesthetized with three types of mixed anesthesia, consisting of 0.75 mg/kg of medetomidine hydrochloride (Domitol; Nippon Zenyaku Kogyo Co., Ltd, Fukushima, Japan), 4 mg/kg of midazolam (Midazolam, Sandoz Inc., Japan), and 5 mg/kg of butorphanol tartrate (Vetorphale; Meiji Seika Pharma Co., Ltd., Tokyo, Japan). A screw (6 mm head diameter and 10 mm length) was attached to the cranial bone with its head down between bregma and lambda using a synthetic resin (Super bond C & B; Sun Medical Company). This screw was used as a holding jig to keep the mouse's head fixed toward the visual stimulus presenting monitor. A glass optical fiber (Core Φ400 μm, 0.39 NA; FT400UMT; Thorlabs, NJ, USA) was implanted in the cerebellum close to the flocculus (coordinates X = 2.0 mm, Y = -5.65 mm, Z = 2.9 mm, α = 10°). This optical fiber was fixed using a synthetic resin (Super bond C&B) mixed with charcoal particles to prevent external lights from penetrating the resin. For recovery from three types of mixed anesthesia, 0.75 mg/kg of atipamezole hydrochloride (Antisedan; Zenyaku Kogyo Co., Ltd, Tokyo, Japan) was administered. The mice were allowed to recover from surgery for at least 5 days. When necessary, the mice were intravenously injected via the tail vein with TexasRed dextran prior to recording. Each mouse was head-restrained by tightening the screw on the head of the mouse to the custom-made apparatus and its body was loosely restrained in an upward-sloping plastic cylinder.

A custom-made optical apparatus for fiber photometry was used (Lucir, Tsukuba, Japan). Excitation lights were delivered from the SPECTRA X Light Engine (Lumencor, OR, USA). An optical fiber from the fiber photometry apparatus was connected to the optical fiber implanted in the mouse with a ferrule-to-ferrule connection using a split sleeve. Excitation light was sent to this optical fiber and the power of excitation light with the wavelength of 575 nm, 510 nm, and 440 nm were 3 mW, 11 μW, and 2.2 mW, respectively, at the tip of the implanted optical fiber. The emitted light came back through the optical fiber, reached the fiber photometry apparatus, and was detected using the two photomultiplier tubes (PMTs) (PMTH-S1-CR316-02; Zolix Instruments Co., Ltd., China) with current-to-voltage amplifiers (HVC1800; Zolix). The electronic signals from the PMTs were digitized with Micro1401-3 plus ADC12 top box (CED, Cambridge, UK) and recorded using the Spike2 software (CED) at a 5 kHz sampling frequency. In all of the recording

configurations, each excitation light was delivered in 20 msec pulses with 180 msec intervals. The signals were averaged within the 20 msec pulse, resulting in an effective sampling frequency of 5 Hz.

For excitation of TexasRed and dYFP fluorescence, lights with the wavelength of 575 nm and 510 nm restricted with a multi-bandpass filter 69008x (Chroma) were used. A multi-band beamsplitter (ZT445/514/594rpc; Chroma) was used to send multiple excitation lights to the mice and to pass through the emission lights of TexasRed or dYFP. The emission lights of TexasRed and dYFP were split by a dichroic mirror, FF605-Di02 (Semrock, IL, USA). The wavelength of the emission lights was restricted with a bandpass filter ET640/30x (Chroma) for TexasRed and with ET539/21x (Chroma) for dYFP detection.

For excitation of TexasRed and autofluorescence, lights with the wavelength of 575 nm and 440 nm restricted by a multi-bandpass filter 69008x (Chroma) were used. A multi-band beamsplitter 69008bs (Chroma) was used to send multiple excitation lights to the mice and to pass through the emission lights of TexasRed and autofluorescence. The emission lights of TexasRed and autofluorescence were split by a dichroic mirror FF605-Di02 (Semrock). The wavelength of the emission lights was restricted with a bandpass filter ET645/75m (Chroma) for TexasRed and with ET537/29m (Chroma) for autofluorescence detection.

For excitation of only the autofluorescence, light with the wavelength of 440 nm restricted with a bandpass filter ET436/20x (Chroma) was used. A dichroic mirror T495lpxr (Chroma) was used to send the excitation light to the mice and to pass through the emission light of the autofluorescence. The wavelength of the emission light was directly restricted with a bandpass filter FF01-550/88 (Semrock) for autofluorescence detection.

Visual stimulus presentation

For visual stimulation, sinusoidal oscillations of vertical stripes displayed on two PC monitors arranged orthogonally were presented to the animal. The visual stimuli were created and controlled by the ImageJ/Fiji software (National Institute of Health, MD, USA) or by the custom program written in MATLAB (Mathworks, Natick, MA, USA), for use with the macro-zoom microscope or the fiber photometry, respectively. In experiments other than the ones shown in **Figure 5** [↗](#), the amplitude of the oscillations was 17° in the visual angle of the mouse, the temporal frequency of the oscillations was 0.25 Hz, and the spatial cycle of the displayed stripe pattern was 6.4°.

Analysis of the eye movement for the HOKR experiments

In fiber photometry experiments, the eye movements were simultaneously recorded. The method of recording and analyzing eye movement was previously described in detail (Kanaya et al., 2023 [↗](#)). During the HOKR experiments, the mouse was exposed to white noise sounds, which helped keep the mouse alert. An angled hot mirror, which reflects only the IR light, was placed between the mouse's right eye and the monitor, thus the visual field of the mouse was unobstructed. The monitored eye (the right eye) was illuminated with two infrared (IR) LEDs. The corneal reflections of the IR LEDs were used as the reference point. The reflected IR image of the eye was captured using a CCD camera (DN3V-30BU; Shodensha Inc., Osaka, Japan) with the IR-cut filter removed.

The captured video taken at 30 fps was analyzed using DeepLabCut (Mathis et al., 2018 [↗](#)), and the center of the pupil was tracked (see Kanaya et al., 2023 [↗](#) for details). The angle of the eye direction was calculated based on the pupil and camera position using a previously introduced method (Sakatani and Isa, 2004 [↗](#)). When the difference in the center eye position between two consecutive frames (1/30 sec) was more than 1° in visual angle, it was assumed that a saccadic eye movement had occurred. These incidents of saccadic movements were removed. Drifts in the eye

movement were eliminated by a band-pass filter with a 100-500 mHz frequency. The amplitudes of the eye were calculated by the difference between the averaged maximum peak and minimum peak. HOKR relative gain was calculated by normalizing the amplitude of the eye movement at the start of the first training session.

Vessel diameter and vasomotion analysis with macro-zoom microscope images

Images acquired with the macro-zoom microscope were analyzed with ImageJ/Fiji and AxoGraph (Axograph Scientific, New South Wales, Australia) software. The blood vessels focused and studied were likely to be arteries or arterioles found on the surface of the brain which were filled with TexasRed fluorophore via intravenous injection. The blood vessel in focus was rotated so that the flow of the blood would be aligned with the vertical axis. The stack of images taken as time series was cropped with the blood vessel in the center of the image, and this stack is named the “original stack”. The background intensity was measured within the region of interest (ROI) on the left and right of the vessel and an average value was calculated for each image of the stack. A new stack with the same XY pixel dimensions as the “(1) original stack” was created and each image of the stack had a single uniform value of the background intensity calculated above. This background stack was subtracted from the “(1) original stack” to create the “(2) background subtracted stack”. Due to the dilution of the TexasRed fluorophore and the photobleaching effect, the fluorescence in the blood vessels tends to fade in time. Thus, a normalization procedure to counter the fading effect was developed. First, using the above “(2) background subtracted stack”, an ROI was set well within the blood vessels and a time series data of the average intensity within the ROI was calculated. This data was strongly low-pass Gaussian filtered at 5 mHz, and the transient fluctuations of intensity due to vasomotion were totally eliminated leaving only the baseline decline of intensity due to the fading effect was left. Again, a new stack with the same XY pixel dimensions as the “(1) original stack” was created and each image of the stack had a single uniform value of the baseline intensity calculated above. The “(2) background subtracted stack” was divided by this baseline stack to create the “(3) baseline normalized stack”. With this procedure, the images are normalized to the basal TexasRed intensity in the blood vessels. TexasRed intensity differs among vessels and trials, thus this procedure is effective in the normalization of multiple experiments. The “(3) baseline normalized stack” was resliced with a line vertical to the vessel using the ImageJ/Fiji software and the temporal change of the TexasRed profile along this line was created. Multiple lines vertical to the vessel were averaged allowing a better signal-to-noise ratio. This data was low-pass Gaussian filtered (~ 0.05 pixel) in each slice. The left side and the right side of the vessel often had different background intensity levels. Therefore, these cross-section profiles of the blood vessels were processed with a flat-sloping baseline function in AxoGraph to eliminate this horizontal slope. The cross-section profile of an undilated vessel in the basal phase was referred and 10% of the maximum value in the profile was calculated. The full width of the cross-section profile at this threshold value was defined as the vessel diameter. The calculated vessel diameter at each time point was normalized to the vessel diameter in the undilated phase.

The vasodilation events were detected using the event detection function in AxoGraph. Thresholds were determined arbitrarily as the amplitudes of the vasodilation varied across the vessels. Inter-event interval and peak amplitudes were calculated. Unit dilation time was defined as the full-width time at the half-maximum of the amplitude of the vasodilation. Inter-event interval, peak amplitude, and unit dilation time were transformed into probability density function and the cumulative relative frequency was calculated.

Frequency analysis

Essentially, the same procedure was used for analyzing the temporal frequency profile of the fluorescence signal data recorded from either the macro-zoom microscopy or the fiber photometry. First, the fluorescence fade was corrected and the baseline value of the fluorescence

signals was normalized. Using MATLAB, the original fluorescence signal data taken with a sampling frequency of 5 Hz were divided by their own low-pass filtered data (the passband was 50 mHz and the stopband was 10 mHz). The result was subtracted by 1 and multiplied by 100 to transform the data into a percentage display. 5 minutes or 15 minutes of data taken during visual stimulus presentation were processed by fast Fourier transform (FFT). The amplitude spectrum and spectrogram were created by executing FFT or short-term Fourier transform (STFT), respectively, by custom programs written in MATLAB. For the spectrogram; the window size was 120 sec, 60 sec, or 40 sec, and the step size was 10 sec, 4 sec, or 2 sec.

The magnitude of the fluorescence signal frequency locked to the presented temporal frequency of the visual stimulus was expressed as the peak ratio of 0.25 Hz (PR0.25). For deriving the PR0.25, first, the peak amplitude of the fluorescence signal at 0.25 Hz was calculated. Next, the mean amplitude of the signal in the 0.1-1 Hz range was calculated. PR0.25 is the signal amplitude at 0.25 Hz divided by the mean at 0.1-1 Hz.

The whole brain surface map of the PR0.25 magnitude presented in **Figure 3E** [↗](#) was created using the following methods. The raw stacked image data taken during the visual stimulus presentation were first processed by using a Gaussian filter with $\sigma = 165 \mu\text{m}$ in the x and y dimensions. The time series data of each pixel were normalized using the method described above. The time series data of each pixel was then processed using FFT and the PR0.25 during the 15 minutes of visual stimulation was calculated and the magnitude of the PR0.25 was color-coded and mapped.

Phase analysis

The phase shifts between the periodic oscillatory signals of the TexasRed and the dYFP signals, or between the TexasRed and the autofluorescence signals, were evaluated using the cross-correlation analysis. The lag time of the negative peak around 0 sec in the cross-correlation was defined as the Δt . The temporal cycle of the horizontally oscillating visual stimuli was 4 sec. A negative peak was usually found within the ± 4 -sec range in the cross-correlation graph in almost all sessions. Data with no negative peak within this range were excluded from further analysis.

The phase shifts between the visual stimuli and the eye movement were calculated using the following method. The eye movement data during the first 5 minutes of each training was divided into segments of 4 sec in duration, resulting in a series of 75 traces. These traces were averaged and fitted with a sinusoidal curve. The phase of the eye movement was calculated using this fitted curve to the averaged eye movement trace. The phase shifts between the visual stimuli and the TexasRed signal were analyzed similarly. However, as shown in **Figure 6F** [↗](#), although the eye movement reliably followed the visual stimulation throughout the session, vasomotion apparently came in and out of the frequency locked to the visual stimulation. Therefore, PR0.25 was calculated for the TexasRed traces and only when the PR0.25 was higher than 3, that segment of the TexasRed trace was extracted. Using these selected segments of the TexasRed signals, the phase shift between the visual stimulation and the TexasRed signal was calculated.

Statistical analysis

All statistical tests were done with Origin Pro 8.6 software (OriginLab, MA, United States). Data are presented as means $\pm$ SEM, except in **Figure 7F** [↗](#). The number of samples is indicated in the figure legend. Statistical analyses were performed with the paired t-test (two-sided), one-sample t-test, and Holm correction for multiple tests.

Author Contributions

D.S. and K.M. conceived the project and designed the study. D.S. performed the experiments and analysis. K.I. contributed to the data analysis. Y. I. contributed to the development of the experimental methods and tools and supervised the initial surgical and experimental procedures. D.S. and K.M. wrote the manuscript. All authors discussed the results and commented on the manuscript.

Acknowledgements

We are thankful to all members of the Ko Matsui laboratory for their invaluable assistance at every stage of the experiments. This work was supported by JSPS Fellowships 22KJ0262 (to D.S.), Grant-in-Aid for Early-Career Scientists 22K15218 (to Y.I.), Grant-in-Aid for Transformative Research Areas (A) “Glial Decoding”: 20H05896 (to K.M.), “Biology of Behavior Change”: 23H04659 (to K.M.), Grant-in-Aid for Scientific Research on Innovative Areas “Brain Information Dynamics”: 18H05110, 20H05046 (to K.M.), Grant-in-Aid for Scientific Research (B): 19H03338, 22H02713 (to K.M.), Research Foundation for Opto-Science and Technology (to K.M.), Takeda Science Foundation (to K.M.), the NOVARTIS Foundation for the Promotion of Science (to K.M.), and the Uehara Memorial Foundation (to K.M.).

Author Information

Daichi Sasaki

Super-network Brain Physiology, Graduate School of Life Sciences, Tohoku University, Sendai 980-8577 Japan

Ken Imai

Super-network Brain Physiology, Graduate School of Life Sciences, Tohoku University, Sendai 980-8577 Japan

Yoko Ikoma

Super-network Brain Physiology, Graduate School of Life Sciences, Tohoku University, Sendai 980-8577 Japan

Ko Matsui

Super-network Brain Physiology, Graduate School of Life Sciences, Tohoku University, Sendai 980-8577 Japan

ORCID iD: 0000-0003-1068-9705

For correspondence: matsui@med.tohoku.ac.jp

References

1. Adaikkan C *et al.* (2019) **Gamma Entrainment Binds Higher-Order Brain Regions and Offers Neuroprotection** *Neuron* **102**:929–943 <https://doi.org/10.1016/j.neuron.2019.04.011>
2. Aldea R, Weller RO, Wilcock DM, Carare RO, Richardson G (2019) **Cerebrovascular smooth muscle cells as the drivers of intramural periarterial drainage of the brain** *Front Aging Neurosci* **11**:1–17 <https://doi.org/10.3389/fnagi.2019.00001>
3. Augustinaite S, Kuhn B (2020) **Chronic Cranial Window for Imaging Cortical Activity in Head-Fixed Mice** *STAR Protoc* **1** <https://doi.org/10.1016/j.xpro.2020.100194>
4. Aziz W, Wang W, Kesaf S, Mohamed AA, Fukazawa Y, Shigemoto R (2014) **Distinct kinetics of synaptic structural plasticity, memory formation, and memory decay in massed and spaced learning** *Proc Natl Acad Sci U S A* **111** <https://doi.org/10.1073/pnas.1303317110>
5. Bazargani N, Attwell D (2016) **Astrocyte calcium signaling: The third wave** *Nat Neurosci* **19**:182–189 <https://doi.org/10.1038/nn.4201>
6. Beppu K, Kubo N, Matsui K (2021) **Glial amplification of synaptic signals** *J Physiol* **599**:2085–2102 <https://doi.org/10.1113/jP280857>
7. Beppu K, Sasaki T, Tanaka KF, Yamanaka A, Fukazawa Y, Shigemoto R, Matsui K (2014) **Optogenetic countering of glial acidosis suppresses glial glutamate release and ischemic brain damage** *Neuron* **81**:314–320 <https://doi.org/10.1016/j.neuron.2013.11.011>
8. Charidimou A, Boulouis G, Gurol ME, Ayata C, Bacskai BJ, Frosch MP, Viswanathan A, Greenberg SM (2017) **Emerging concepts in sporadic cerebral amyloid angiopathy** *Brain* **140**:1829–1850 <https://doi.org/10.1093/brain/awx047>
9. Croce AC, Bottiroli G (2015) **New light in flavin autofluorescence** *Eur J Histochem* **59**:317–318 <https://doi.org/10.4081/ejh.2015.2576>
10. Davis MJ, Hill MA, Kuo L (2008) **Local Regulation of Microvascular Perfusion** *Microcirculation* :161–284 <https://doi.org/10.1016/B978-0-12-374530-9.00006-1>
11. Fujii K, Heistad DD, Faraci FM (1990) **Vasomotion of basilar arteries in vivo** *Am J Physiol - Hear Circ Physiol* **258** <https://doi.org/10.1152/ajpheart.1990.258.6.h1829>
12. Haddock RE, Grayson TH, Brackenbury TD, Meaney KR, Neylon CB, Sandow SL, Hill CE (2006) **Endothelial coordination of cerebral vasomotion via myoendothelial gap junctions containing connexins 37 and 40** *Am J Physiol - Hear Circ Physiol* **291**:2047–2056 <https://doi.org/10.1152/ajpheart.00484.2006>
13. Haddock RE, Hill CE (2002) **Differential activation of ion channels by inositol 1,4,5-trisphosphate (IP3)- and ryanodine-sensitive calcium stores in rat basilar artery vasomotion** *J Physiol* **545**:615–627 <https://doi.org/10.1113/jphysiol.2002.027904>
14. Haidey JN *et al.* (2021) **Astrocytes regulate ultra-slow arteriole oscillations via stretch-mediated TRPV4-COX-1 feedback** *Cell Rep* **36** <https://doi.org/10.1016/j.celrep.2021.109405>

15. Heeger DJ, Huk AC, Geisler WS, Albrecht DG (2000) **Spikes versus BOLD: What does neuroimaging tell us about neuronal activity?** *Nat Neurosci* **3**:631–633 <https://doi.org/10.1038/76572>
16. Hill CE, Eade J, Sandow SL (1999) **Mechanisms underlying spontaneous rhythmical contractions in irideal arterioles of the rat** *J Physiol* **521**:507–516 <https://doi.org/10.1111/j.1469-7793.1999.00507.x>
17. Ho D, Zhao X, Gao S, Hong C, Vatner DE, Vatner SF (2011) **Heart Rate and Electrocardiography Monitoring in Mice** *Curr Protoc Mouse Biol* **1**:123–139 <https://doi.org/10.1002/9780470942390.mo100159>
18. Huang S, Heikal AA, Webb WW (2002) **Two-photon fluorescence spectroscopy and microscopy of NAD(P)H and flavoprotein** *Biophys J* **82**:2811–2825 [https://doi.org/10.1016/S0006-3495\(02\)75621-X](https://doi.org/10.1016/S0006-3495(02)75621-X)
19. Hundley WG, Renaldo GJ, Levasseur JE, Kontos HA (1988) **Vasomotion in cerebral microcirculation of awake rabbits** *Am J Physiol - Hear Circ Physiol* **254** <https://doi.org/10.1152/ajpheart.1988.254.1.h67>
20. Iaccarino HF *et al.* (2018) **Author correction: Gamma frequency entrainment attenuates amyloid load and modifies microglia (Nature, (2016), 540, 7632, (230–235), 10.1038/nature20587)** *Nature* **562** <https://doi.org/10.1038/s41586-018-0351-4>
21. Ikoma Y, Sasaki D, Matsui K (2023) **Local brain environment changes associated with epileptogenesis** *Brain* **146**:576–586 <https://doi.org/10.1093/brain/awac355>
22. Ikoma Y, Takahashi Y, Sasaki D, Matsui K (2023) **Properties of REM sleep alterations with epilepsy** *Brain* **146**:2431–2442 <https://doi.org/10.1093/brain/awac499>
23. Islam MS, Honma M, Nakabayashi T, Kinjo M, Ohta N (2013) **pH dependence of the fluorescence lifetime of FAD in solution and in cells** *Int J Mol Sci* **14**:1952–1963 <https://doi.org/10.3390/ijms14011952>
24. Ito M (1982) **Cerebellar control of the vestibulo-ocular reflex--around the flocculus hypothesis** *Annu Rev Neurosci* **5**:275–296 <https://doi.org/10.1146/annurev.ne.05.030182.001423>
25. Jones TW (1853) **Discovery that veins of the bat's wing (which are furnished with valves) are endowed with rhythmical contractility and that the onward flow of blood is accelerated by each contraction** *Edinb Med Surg J* **79**:367–373 <https://doi.org/10.1098/rspl.1850.0048>
26. Kanaya T, Ito R, Morizawa YM, Sasaki D, Yamao H, Ishikane H, Hiraoka Y, Tanaka K, Matsui K (2023) **Glial modulation of the parallel memory formation** *Glia* **71**:2401–2417 <https://doi.org/10.1002/glia.24431>
27. Kanemaru K *et al.* (2014) **In Vivo visualization of subtle, transient, and local activity of astrocytes using an ultrasensitive Ca²⁺ indicator** *Cell Rep* **8**:311–318 <https://doi.org/10.1016/j.celrep.2014.05.056>
28. Kayser C (2004) **A Comparison of Hemodynamic and Neural Responses in Cat Visual Cortex Using Complex Stimuli** *Cereb Cortex* **14**:881–891 <https://doi.org/10.1093/cercor/bhh047>

29. Korte N, James G, You H, Hirunpattarasilp C, Christie I, Sethi H, Attwell D (2023) **Noradrenaline released from locus coeruleus axons contracts cerebral capillary pericytes via α_2 adrenergic receptors** *J Cereb Blood Flow Metab* **43**:1142–1152 <https://doi.org/10.1177/0271678X231152549>
30. Kramer K, van Acker SA, Voss HP, Grimbergen JA, van der Vijgh WJ, Bast A (1993) **Use of telemetry to record electrocardiogram and heart rate in freely moving mice** *J Pharmacol Toxicol Methods* **30**:209–215
31. Martorell AJ *et al.* (2019) **Multi-sensory Gamma Stimulation Ameliorates Alzheimer's-Associated Pathology and Improves Cognition** *Cell* **177**:256–271 <https://doi.org/10.1016/j.cell.2019.02.014>
32. Martuzzi R, Murray MM, Meuli RA, Thiran JP, Maeder PP, Michel CM, Grave De Peralta Menendez R, Gonzalez Andino SL (2009) **Methods for determining frequency- and region-dependent relationships between estimated LFPs and BOLD responses in humans** *J Neurophysiol* **101**:491–502 <https://doi.org/10.1152/jn.90335.2008>
33. Mateo C, Knutsen PM, Tsai PS, Shih AY, Kleinfeld D (2017) **Entrainment of Arteriole Vasomotor Fluctuations by Neural Activity Is a Basis of Blood-Oxygenation-Level-Dependent “Resting-State” Connectivity** *Neuron* **96**:936–948 <https://doi.org/10.1016/j.neuron.2017.10.012>
34. Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M (2018) **DeepLabCut: markerless pose estimation of user-defined body parts with deep learning** *Nat Neurosci* **21**:1281–1289 <https://doi.org/10.1038/s41593-018-0209-y>
35. Matsui K, Jahr CE (2004) **Differential control of synaptic and ectopic vesicular release of glutamate** *J Neurosci* **24**:8932–8939 <https://doi.org/10.1523/JNEUROSCI.2650-04.2004>
36. Matsui K, Jahr CE (2003) **Ectopic release of synaptic vesicles** *Neuron* **40**:1173–1183 [https://doi.org/10.1016/S0896-6273\(03\)00788-8](https://doi.org/10.1016/S0896-6273(03)00788-8)
37. Matsui K, Jahr CE, Rubio ME (2005) **High-concentration rapid transients of glutamate mediate neural-glia communication via ectopic release** *J Neurosci* **25**:7538–7547 <https://doi.org/10.1523/JNEUROSCI.1927-05.2005>
38. Meyer C, Vries G de, Davidge ST, Mayes DC (2002) **Reassessing the Mathematical Modeling of the Contribution of Vasomotion to Vascular Resistance** *J Appl Physiol* **92**:888–9 <https://doi.org/10.1152/jappl.2002.92.2.888>
39. Morizawa YM *et al.* (2022) **Synaptic pruning through glial synapse engulfment upon motor learning** *Nat Neurosci* **25**:1458–1469 <https://doi.org/10.1038/s41593-022-01184-5>
40. Niessing J, Ebisch B, Schmidt KE, Niessing M, Singer W, Galuske RAW (2005) **Hemodynamic signals correlate tightly with synchronized gamma oscillations** *Science (80-)* **309**:948–951 <https://doi.org/10.1126/science.1110948>
41. Nilsson H, Aalkjaer C (2003) **Vasomotion: mechanisms and physiological importance** *Mol Interv* **3**:79–89 <https://doi.org/10.1124/mi.3.2.79>
42. Nortley R *et al.* (2019) **Amyloid β oligomers constrict human capillaries in Alzheimer's disease via signaling to pericytes** *Science (80-)* **365** <https://doi.org/10.1126/science.aav9518>

43. Ozawa K, Nagao M, Konno A, Iwai Y, Vittani M, Kusk P, Mishima T, Hirai H, Nedergaard M, Hirase H (2023) **Astrocytic GPCR-Induced Ca²⁺ Signaling Is Not Causally Related to Local Cerebral Blood Flow Changes** *Int J Mol Sci* **24** <https://doi.org/10.3390/ijms241713590>
44. Panerai RB, Rennie JM, Kelsall AWR, Evans DH (1998) **Frequency-domain analysis of cerebral autoregulation from spontaneous fluctuations in arterial blood pressure** *Med Biol Eng Comput* **36**:315–322 <https://doi.org/10.1007/BF02522477>
45. Peng H, Matchkov V, Ivarsen A, Aalkjær C, Nilsson H (2001) **Hypothesis for the initiation of vasomotion** *Circ Res* **88**:810–815 <https://doi.org/10.1161/hh0801.089603>
46. Peng W *et al.* (2016) **Suppression of glymphatic fluid transport in a mouse model of Alzheimer's disease** *Neurobiol Dis* **93**:215–225 <https://doi.org/10.1016/j.nbd.2016.05.015>
47. Peppiatt CM, Howarth C, Mobbs P, Attwell D (2006) **Bidirectional control of CNS capillary diameter by pericytes** *Nature* **443**:700–704 <https://doi.org/10.1038/nature05193>
48. Rivadulla C, de Labra C, Grieve KL, Cudeiro J (2011) **Vasomotion and neurovascular coupling in the visual thalamus in vivo** *PLoS One* **6** <https://doi.org/10.1371/journal.pone.0028746>
49. Sakatani T, Isa T (2004) **PC-based high-speed video-oculography for measuring rapid eye movements in mice** *Neurosci Res* **49**:123–131 <https://doi.org/10.1016/j.neures.2004.02.002>
50. Sasaki T, Beppu K, Tanaka KF, Fukazawa Y, Shigemoto R, Matsui K (2012) **Application of an optogenetic byway for perturbing neuronal activity via glial photostimulation** *Proc Natl Acad Sci U S A* **109**:20720–20725 <https://doi.org/10.1073/pnas.1213458109>
51. Sassaroli A, Pierro M, Bergethon PR, Fantini S (2012) **Low-frequency spontaneous oscillations of cerebral hemodynamics investigated with near-infrared spectroscopy: A review** *IEEE J Sel Top Quantum Electron* **18**:1478–1492 <https://doi.org/10.1109/JSTQE.2012.2183581>
52. Schuster A, Hirotaka O, Beny JL, Stergiopoulos N, Meister JJ (2000) **Simultaneous arterial calcium dynamics and diameter measurements: application to myoendothelial communication** *Ann Biomed Eng* **28**:1088–1096
53. Shaw K, Bell L, Boyd K, Grijseels DM, Clarke D, Bonnar O, Crombag HS, Hall CN (2021) **Neurovascular coupling and oxygenation are decreased in hippocampus compared to neocortex because of microvascular differences** *Nat Commun* **12**:1–16 <https://doi.org/10.1038/s41467-021-23508-y>
54. Soula M, Martín-Ávila A, Zhang Y, Dhingra A, Nitzan N, Sadowski MJ, Gan WB, Buzsáki G (2023) **Forty-hertz light stimulation does not entrain native gamma oscillations in Alzheimer's disease model mice** *Nat Neurosci* **26**:570–578 <https://doi.org/10.1038/s41593-023-01270-2>
55. Stickland R, Allen M, Magazzini L, Singh KD, Wise RG, Tomassini V (2019) **Neurovascular Coupling During Visual Stimulation in Multiple Sclerosis: A MEG-fMRI Study** *Neuroscience* **403**:54–69 <https://doi.org/10.1016/j.neuroscience.2018.03.018>
56. Tanaka KF *et al.* (2012) **Expanding the Repertoire of Optogenetically Targeted Cells with an Enhanced Gene Expression System** *Cell Rep* **2**:397–406 <https://doi.org/10.1016/j.celrep.2012.06.011>

- 57. van Veluw SJ, Hou SS, Calvo-Rodriguez M, Arbel-Ornath M, Snyder AC, Frosch MP, Greenberg SM, Bacsikai BJ (2020) **Vasomotion as a Driving Force for Paravascular Clearance in the Awake Mouse Brain** *Neuron* **105**:549–561 <https://doi.org/10.1016/j.neuron.2019.10.033>
- 58. Waespe W, Henn V (1977) **Neuronal activity in the vestibular nuclei of the alert monkey during vestibular and optokinetic stimulation** *Exp Brain Res* **27**:523–538 <https://doi.org/10.1007/BF00239041>
- 59. Wise RG, Ide K, Poulin MJ, Tracey I (2004) **Resting fluctuations in arterial carbon dioxide induce significant low frequency variations in BOLD signal** *Neuroimage* **21**:1652–1664 <https://doi.org/10.1016/j.neuroimage.2003.11.025>

Article and author information

Daichi Sasaki

Super-network Brain Physiology, Graduate School of Life Sciences, Tohoku University, Sendai 980-8577 Japan

Ken Imai

Super-network Brain Physiology, Graduate School of Life Sciences, Tohoku University, Sendai 980-8577 Japan

Yoko Ikoma

Super-network Brain Physiology, Graduate School of Life Sciences, Tohoku University, Sendai 980-8577 Japan

Ko Matsui

Super-network Brain Physiology, Graduate School of Life Sciences, Tohoku University, Sendai 980-8577 Japan

For correspondence: matsui@med.tohoku.ac.jp

ORCID iD: [0000-0003-1068-9705](https://orcid.org/0000-0003-1068-9705)

Copyright

© 2024, Sasaki et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Brice Bathellier

CNRS, Paris, France

Senior Editor

Laura Colgin

University of Texas at Austin, Austin, United States of America

Reviewer #1 (Public Review):

This manuscript presents the first evidence for a plastic enhancement in the response of pial cortical arterioles to external stimulation. Specifically, they show (p8; Figure 3A-C) that repeated application of a visual stimulus at 0.25 Hz, at the upper edge of the vasomotor response, leads to a greater change in the diameter of pial arterioles at that frequency. This adds to the earlier, referenced work of Mateo et al (2017) that showed locking - or entrainment of pial arteriole vasomotion - by stimuli at different (0.0 to 0.3 Hz) frequencies.

The manuscript has a major flaw. Much as there is plasticity that leads to an increase in the amplitude of vasomotion at the drive frequency, the authors need to show reversibility. This could possibly be accomplished by driving the visual system at a different frequency, say 0.15 Hz, and observing if the 0.25 Hz response is then diminished. The authors could then test if their observation is repeatable by again driving at 0.25 Hz. Unless I missed the presentation on this point, there is no evidence for reversibility.

Drew, P. J., A. Y. Shih, J. D. Driscoll, P. M. Knutsen, D. Davalos, P. Blinder, K. Akassoglou, P. S. Tsai, and D. Kleinfeld. 2010. 'Chronic optical access through a polished and reinforced thinned skull', *Nature Methods*, 7: 981-84.

Morii, S., A. C. Ngai, and H. R. Winn. 1986. 'Reactivity of rat pial arterioles and venules to adenosine and carbon dioxide: With detailed description of the closed cranial window technique in rats', *Journal of Cerebral Blood Flow & Metabolism*, 6: 34-41.

<https://doi.org/10.7554/eLife.93721.1.sa2>

Reviewer #2 (Public Review):

Sasaki et al. investigated methods to entrain vasomotion in awake wild-type mice across multiple regions of the brain using a horizontally oscillating visual pattern which induces an optokinetic response (HOKR) eye movement. They found that spontaneous vasomotion could be detected in individual vessels of their wild-type mice through either a thinned cranial window or intact skull preparation using a widefield macro-zoom microscope. They showed that low-resolution autofluorescence signals coming from the brain parenchyma could be used to capture vasomotion activity using a macro-zoom microscope or optical fibre, as this signal correlates well with the intensity profile of fluorescently-labelled single vessels. They show that vasomotion can also be entrained across the cortical surface using an oscillating visual stimulus with a range of parameters (with varying temporal frequencies, amplitudes, or spatial cycles), and that the amplitude spectrum of the detected vasomotion frequency increases with repeated training sessions. The authors include some control experiments to rule out fluorescence fluctuations being due to artifacts of eye movement or screen luminance and attempt to demonstrate some functional benefit of vasomotion entraining as HOKR performance improves after repeat training. These data add in an interesting way to the current knowledge base on vasomotion, as the authors demonstrate the ability to entrain vasomotion across multiple brain areas and show some functional significance to vasomotion with regards to information processing as HOKR task performance correlates well with vascular oscillation amplitudes.

The aims of the paper are mostly well supported by the data, but some streamlining of the data presentation would improve overall clarity. The third aim to establish the functional significance of vasomotion in relation to plasticity in information processing could be better supported by the inclusion of some additional control experiments. Specifically:

1. The clarity and comprehensibility of the paper could be significantly enhanced by incorporating additional details in both the introduction and discussion sections. In the introduction, a succinct definition of the frequency range of vasomotion should be provided, as well as a better description of the horizontal optokinetic response (i.e. as they have in the results section in the first paragraph below the 'Entrainment of vasomotion with visual stimuli presentation' sub-heading). The discussion would benefit from the inclusion of a clear summary of the results presented at the start, and the inclusion of stronger justification (i.e. more citations) with regards to the speculation about vasomotion and neuronal plasticity (e.g. paragraph 5 includes no citations).
2. The novel methods for detecting vasomotion using low-resolution imaging techniques are discussed across the first four figures, but this gets a little bit confusing to follow as the authors jump back and forth between the different imaging and analysis techniques they have employed to capture vasomotion. The data presentation could be better streamlined - for instance by presenting only the methods most relevant for the functional dataset (in Figures 5-7), with the additional information regarding the various controls to establish the use of autofluorescence intensity imaging as a valid method for capturing vasomotion reduced to fewer figure panels, or moved to supplementary figures so as to not detract from the main novel findings contributed in this study.
3. The authors heavily rely on representative traces from individual vessels to illustrate their findings, particularly evident in Figures 1-4. While these traces offer a valuable visualization, augmenting their approach by presenting individual data points across the entire dataset, encompassing all animals and vessels, would significantly enhance the robustness of their claims. For instance, in Figures 1 and 2, where average basal and dilated traces are depicted for a representative vessel, supplementing these with graphs showcasing peak values across all measured vessels would enable the authors to convey a more holistic representation of their data. Or in Figure 3, where the amplitude spectrum is presented for individual Texas red fluorescence intensity changes in V1 across novice, trained, and expert mice, incorporating a summary graph featuring the amplitude spectrum value at 0.25Hz for each individual trace (across animals/imaging sessions), followed by statistical analysis, would fortify the strength of their assertions. Moreover, providing explicit details on sample sizes for each individual figure panel (where not a representative trace), including the number of animals or vessels/imaging sessions, would contribute to transparency and aid readers in assessing the generalisability of the findings.
4. In the experiments where mice are classed as "novice", "trained" or "expert", the inclusion of the specific range of the number of training sessions for each category would improve replicability.
5. The authors don't state whether mice were habituated to the imaging set-up prior to the first data collection, as head-fixation and restraint can be stress-inducing for animals, especially upon first exposure, which could impact their neurovascular coupling responses differentially in "novice" versus "trained" imaging sessions (e.g. see Han et al., 2020, DOI: <https://doi.org/10.1523/JNEUROSCI.1553-20.2020>). The stress associated with a tail vein injection prior to imaging could also partially explain why mice didn't learn very well if Texas Red was injected before the training session. If no habituation was conducted in these experiments, the study would benefit from the inclusion of some control experiments where "novice" responses were compared between habituated and non-habituated animals.

6. The experiments regarding the brain-wide vasomotion entrainment across the cortical surface would benefit from some additional information about how brain regions were identified (e.g. particularly how V1 and V2 were distinguished given how close together they are).
7. Whilst the authors show that HOKR task performance and vasomotion amplitude are increased with repeated training to provide some support to their aim of investigating the functional significance of vasomotion with regards to information processing plasticity, the inclusion of some additional control experiments would provide stronger evidence to address this aim. For instance, if vasomotion signalling is blocked or reduced (e.g. using optogenetics or in an AD mouse model where arteriole amyloid load restricts vasomotion capacity), does flocculus-dependent task performance (e.g. HOKR eye movements) still improve with repeated exposure to the external stimulus.

<https://doi.org/10.7554/eLife.93721.1.sa1>

Reviewer #3 (Public Review):

Summary:

Here the authors show global synchronization of cerebral blood flow (CBF) induced by oscillating visual stimuli in the mouse brain. The study validates the use of endogenous autofluorescence to quantify the vessel "shadow" to assess the magnitude of frequency-locked cerebral blood flow changes. This approach enables straightforward estimation of artery diameter fluctuations in wild-type mice, employing either low magnification wide-field microscopy or deep-brain fibre photometry. For the visual stimuli, awake mice were exposed to vertically oscillating stripes at a low temporal frequency (0.25 Hz), resulting in oscillatory changes in artery diameter synchronized to the visual stimulation frequency. This phenomenon occurred not only in the primary visual cortex but also across a broad cortical and cerebellar surface. The induced CBF changes adapted to various stimulation parameters, and interestingly, repeated trials led to plastic entrainment. The authors control for different artefacts that may have confounded the measurements such as light contamination and eye movements but found no influence of these variables. The study also tested horizontally oscillating visual stimuli, which induce the horizontal optokinetic response (HOKR). The amplitude of eye movement, known to increase with repeated training sessions, showed a strong correlation with CBF entrainment magnitude in the cerebellar flocculus. The authors suggest that parallel plasticity in CBF and neuronal circuits is occurring. Overall, the study proposes that entrained "vasomotion" contributes to meeting the increased energy demand associated with coordinated neuronal activity and subsequent neuronal circuit reorganization.

Strengths:

- The paper describes a simple and useful method for tracking vasomotion in awake mice through an intact skull.
- The work controls for artefacts in their primary measurements.
- There are some interesting observations, including the nearly brain-wide synchronization of cerebral blood flow oscillations to visual stimuli and that this process only occurs after mice are trained in a visual task.
- This topic is interesting to many in the CBF, functional imaging, and dementia fields.

Weaknesses:

- I have concerns with the main concepts put forward, regarding whether the authors are actually studying vasomotion as they state, as opposed to functional hyperemia which is sensory-induced changes in blood flow, which is what they are actually doing. I recommend

several additional experiments/analyses for them to explore. This is mostly further characterizing their effect which will benefit the interpretations.

-Neuronal calcium imaging would also benefit the study and improve the interpretations.

-The plastic effects in vasomotion synchronization that occur with training are interesting but they could use an additional control for stress. Is this really a plastic effect, or is it caused by progressively decreasing stress as trials and progress? I recommend a habituation control experiment.

Appraisal

I think the authors have an interesting effect that requires further characterization and controls. Their interpretations are likely sound and additional experiments will continue to support the main hypothesis. If brain-wide synchrony of blood flow can be trained and entrained by external stimuli, this may have interesting therapeutic potential to help clear out toxic proteins from the brain as seen in several neurodegenerative diseases.

<https://doi.org/10.7554/eLife.93721.1.sa0>

Author Response

Reviewer #1 (Public Review):

This manuscript presents the first evidence for a plastic enhancement in the response of pial cortical arterioles to external stimulation. Specifically, they show (p8; Figure 3A-C) that repeated application of a visual stimulus at 0.25 Hz, at the upper edge of the vasomotor response, leads to a greater change in the diameter of pial arterioles at that frequency. This adds to the earlier, referenced work of Mateo et al (2017) that showed locking - or entrainment of pial arteriole vasomotion - by stimuli at different (0.0 to 0.3 Hz) frequencies.

We thank the reviewer for positively identifying the value of our manuscript.

The manuscript has a major flaw. Much as there is plasticity that leads to an increase in the amplitude of vasomotion at the drive frequency, the authors need to show reversibility. This could possibly be accomplished by driving the visual system at a different frequency, say 0.15 Hz, and observing if the 0.25 Hz response is then diminished. The authors could then test if their observation is repeatable by again driving at 0.25 Hz. Unless I missed the presentation on this point, there is no evidence for reversibility.

The reviewer has raised a very important point of view. In our experiments, the visually induced vasomotion (or visual stimulus-triggered vasomotion) was always entrained by repeated trials of the 0.25 Hz temporal frequency stimuli. When the visual stimulation stops, the vasomotion frequency lock to 0.25 Hz quickly dissipates. After saturated training with this stimulus, the parameters of the visual stimulus were switched, for example to 0.15 Hz. The animal quickly adapted to this new stimulus paradigm and the vasomotion was frequency-locked to 0.15 Hz. The adaptation to this new paradigm occurred well within 5 minutes. In Fig. 5, various paradigms were randomly tested. In some of the trials, 0.25 Hz stimulus was tested after 0.15 Hz. The vasomotion also quickly adapted back to the 0.25 Hz. We agree with the reviewer that this reversibility could have been explicitly documented in the manuscript.

Drew, P. J., A. Y. Shih, J. D. Driscoll, P. M. Knutsen, D. Davalos, P. Blinder, K. Akassoglou, P. S. Tsai, and D. Kleinfeld. 2010. 'Chronic optical access through a polished and reinforced thinned skull', *Nature Methods*, 7: 981-84.

Morii, S., A. C. Ngai, and H. R. Winn. 1986. 'Reactivity of rat pial arterioles and venules to adenosine and carbon dioxide: With detailed description of the closed cranial window technique in rats', *Journal of Cerebral Blood Flow & Metabolism*, 6: 34-41.

Reviewer #2 (Public Review):

Sasaki et al. investigated methods to entrain vasomotion in awake wild-type mice across multiple regions of the brain using a horizontally oscillating visual pattern which induces an optokinetic response (HOKR) eye movement. They found that spontaneous vasomotion could be detected in individual vessels of their wild-type mice through either a thinned cranial window or intact skull preparation using a widefield macro-zoom microscope. They showed that low-resolution autofluorescence signals coming from the brain parenchyma could be used to capture vasomotion activity using a macro-zoom microscope or optical fibre, as this signal correlates well with the intensity profile of fluorescently-labelled single vessels. They show that vasomotion can also be entrained across the cortical surface using an oscillating visual stimulus with a range of parameters (with varying temporal frequencies, amplitudes, or spatial cycles), and that the amplitude spectrum of the detected vasomotion frequency increases with repeated training sessions. The authors include some control experiments to rule out fluorescence fluctuations being due to artifacts of eye movement or screen luminance and attempt to demonstrate some functional benefit of vasomotion entraining as HOKR performance improves after repeat training. These data add in an interesting way to the current knowledge base on vasomotion, as the authors demonstrate the ability to entrain vasomotion across multiple brain areas and show some functional significance to vasomotion with regards to information processing as HOKR task performance correlates well with vascular oscillation amplitudes.

We thank the reviewer for summarizing the value of our study and recognizing its significance.

The aims of the paper are mostly well supported by the data, but some streamlining of the data presentation would improve overall clarity. The third aim to establish the functional significance of vasomotion in relation to plasticity in information processing could be better supported by the inclusion of some additional control experiments.

We thank the reviewer for recognizing our vast amount of data supporting our findings. We agree that better data presentation could have improved the clarity of the manuscript.

Specifically:

- 1. The clarity and comprehensibility of the paper could be significantly enhanced by incorporating additional details in both the introduction and discussion sections. In the introduction, a succinct definition of the frequency range of vasomotion should be provided, as well as a better description of the horizontal optokinetic response (i.e. as they have in the results section in the first paragraph below the 'Entrainment of vasomotion with visual stimuli presentation' sub-heading). The discussion would benefit from the inclusion of a clear summary of the results presented at the start, and the inclusion of stronger justification (i.e. more citations) with regards to the speculation about vasomotion and neuronal plasticity (e.g. paragraph 5 includes no citations).*

We agree that a better description of vasomotion and horizontal optokinetic response could have been provided in the introduction. As the reviewer suggests, the discussion could also have started with the following summary of the results.

“We show that visually induced vasomotion can be frequency-locked to the visual stimulus and can be entrained with repeated trials. The initial drive for the vasomotion, or the sensory-evoked hyperemia, must be coming from the neuronal activity in the visual system. The vasomotion is likely triggered by activation of the neurovascular interaction (Kayser, 2004; van Veluw et al., 2020). Surprisingly, the entrained vasomotion was observed not only in the visual cortex but also widely throughout the surface of the brain and deep in the cerebellar flocculus. The global entrainment could be realized through separate mechanisms from the local neurovascular coupling. What is also unknown is where the plasticity occurs. The neuronal visual response in the primary visual cortex could potentially decrease with repeated visual stimulation presentation as the adaptive movement of the eye should decrease the retinal slip. With repeated training sessions, a more static projection of the presented image will likely be shown to the retina. The neurovascular coupling could be enhanced with increased responsiveness of the vasculature and vascular-to-vascular coupling could also be potentiated.”

1. The novel methods for detecting vasomotion using low-resolution imaging techniques are discussed across the first four figures, but this gets a little bit confusing to follow as the authors jump back and forth between the different imaging and analysis techniques they have employed to capture vasomotion. The data presentation could be better streamlined - for instance by presenting only the methods most relevant for the functional dataset (in Figures 5-7), with the additional information regarding the various controls to establish the use of autofluorescence intensity imaging as a valid method for capturing vasomotion reduced to fewer figure panels, or moved to supplementary figures so as to not detract from the main novel findings contributed in this study.

We apologize for the confusing presentation of the data. Many of the initial figures were technical; however, we feel that following these steps was necessary to logically conclude that shadow imaging of the autofluorescence could be used as an indicator of vasomotion. We do agree with the reviewer that going back and forth between different techniques can be confusing. We could have added separate supplementary figures to introduce the various methods used upfront before going into the findings.

1. The authors heavily rely on representative traces from individual vessels to illustrate their findings, particularly evident in Figures 1-4. While these traces offer a valuable visualization, augmenting their approach by presenting individual data points across the entire dataset, encompassing all animals and vessels, would significantly enhance the robustness of their claims. For instance, in Figures 1 and 2, where average basal and dilated traces are depicted for a representative vessel, supplementing these with graphs showcasing peak values across all measured vessels would enable the authors to convey a more holistic representation of their data. Or in Figure 3, where the amplitude spectrum is presented for individual Texas red fluorescence intensity changes in V1 across novice, trained, and expert mice, incorporating a summary graph featuring the amplitude spectrum value at 0.25Hz for each individual trace (across animals/imaging sessions), followed by statistical analysis, would fortify the strength of their assertions. Moreover, providing explicit details on sample sizes for each individual figure panel (where not a representative trace), including the number of animals or vessels/imaging sessions, would contribute to transparency and aid readers in assessing the generalisability of the findings.

We agree with the reviewer that summarization of the data across a number of vessels/imaging sessions would lead to more generalization of the findings. However, contrary to what the reviewer described, we did summarize the vessel diameter expansion events across multiple vessel observations in Fig. 1F, G. The vasomotion parameters were not

summarized for observation in intact skull shown in Fig. 2. However, this figure was intended just to show that vessel boundary cannot be well defined in intact skull imaging and Texas Red intensity or autofluorescence intensity fluctuation would give a better indication of vessel diameter fluctuation. In Fig. 3G, the peak ratio of 0.25 Hz was calculated for individual animals at Novice, Trained, and Expert levels and summarized for $n = 5$ animals. Statistical analysis was also done. The variability between imaging sessions within individual animals was not analyzed; thus, this could have been indicated.

1. *In the experiments where mice are classed as "novice", "trained" or "expert", the inclusion of the specific range of the number of training sessions for each category would improve replicability.*

We agree with the reviewer that classification on the level of training should have been explicitly indicated. Mice experiencing the first visual training session were defined as “Novice”. The mice that have experienced 3 training sessions are the “Trained” mice and the performance of the “Trained” mice during the 4th training session was evaluated. Mice that experienced 8 to 11 rounds of visual training sessions are the “Expert” mice.

1. *The authors don't state whether mice were habituated to the imaging set-up prior to the first data collection, as head-fixation and restraint can be stress-inducing for animals, especially upon first exposure, which could impact their neurovascular coupling responses differentially in "novice" versus "trained" imaging sessions (e.g. see Han et al., 2020, DOI: <https://doi.org/10.1523/JNEUROSCI.1553-20.2020>). The stress associated with a tail vein injection prior to imaging could also partially explain why mice didn't learn very well if Texas Red was injected before the training session. If no habituation was conducted in these experiments, the study would benefit from the inclusion of some control experiments where "novice" responses were compared between habituated and non-habituated animals.*

We agree with the reviewer that stress could well affect spontaneous vasomotion as well as visually induced vasomotion (or visual stimulus-triggered vasomotion). As the reviewer suggested, we could have compared the habituated and non-habituated mice to the initial visually induced vasomotion response. In addition, whether the experimentally induced increase in stress would interfere with the vasomotion or not could also be studied. With the Texas Red experiments, we observed that tail-vein injection stress appeared to interfere with the HOKR learning process. In the experiments presented in Fig. 3, Texas Red was injected before session 1. Vasomotion entrainment likely progressed with sessions 2 and 3 training. Before session 4, Texas Red was injected again to visualize the vasomotion. The vasomotion was clearly observed in session 4, indicating that the stress induced by tail-vein injection could not interfere with the generation of visually induced vasomotion.

1. *The experiments regarding the brain-wide vasomotion entrainment across the cortical surface would benefit from some additional information about how brain regions were identified (e.g. particularly how V1 and V2 were distinguished given how close together they are).*

The brain regions were identified by referring to the Mouse Brain Atlas. As the skull was intact, the location of bregma, lambda, and midline was clearly visible. We agree with the reviewer that strict separation of V1 and V2 could be difficult if we rely on the brain atlas alone. However, what we wanted to emphasize was that there was no specific localization of the vasomotion entrainment effect.

1. Whilst the authors show that HOKR task performance and vasomotion amplitude are increased with repeated training to provide some support to their aim of investigating the functional significance of vasomotion with regards to information processing plasticity, the inclusion of some additional control experiments would provide stronger evidence to address this aim. For instance, if vasomotion signalling is blocked or reduced (e.g. using optogenetics or in an AD mouse model where arteriole amyloid load restricts vasomotion capacity), does flocculus-dependent task performance (e.g. HOKR eye movements) still improve with repeated exposure to the external stimulus.

We agree that experimental intervention to vasomotion is ideal to test the functional significance of vasomotion. As pharmacological intervention lacks specificity, we are currently exploring the optogenetic approach. We have never thought of using the AD mouse as a model of restricted vasomotion by amyloid, and we agree this would be an interesting model to study. However, the AD mouse model would also have deficits other than the restricted vasomotion. On the other hand, we could test whether the repeated presentation of slowly oscillating visual stimuli can have beneficial effects in improving the cognitive abilities of AD model mice.

Reviewer #3 (Public Review):

Summary:

Here the authors show global synchronization of cerebral blood flow (CBF) induced by oscillating visual stimuli in the mouse brain. The study validates the use of endogenous autofluorescence to quantify the vessel "shadow" to assess the magnitude of frequency-locked cerebral blood flow changes. This approach enables straightforward estimation of artery diameter fluctuations in wild-type mice, employing either low magnification wide-field microscopy or deep-brain fibre photometry. For the visual stimuli, awake mice were exposed to vertically oscillating stripes at a low temporal frequency (0.25 Hz), resulting in oscillatory changes in artery diameter synchronized to the visual stimulation frequency. This phenomenon occurred not only in the primary visual cortex but also across a broad cortical and cerebellar surface. The induced CBF changes adapted to various stimulation parameters, and interestingly, repeated trials led to plastic entrainment. The authors control for different artefacts that may have confounded the measurements such as light contamination and eye movements but found no influence of these variables. The study also tested horizontally oscillating visual stimuli, which induce the horizontal optokinetic response (HOKR). The amplitude of eye movement, known to increase with repeated training sessions, showed a strong correlation with CBF entrainment magnitude in the cerebellar flocculus. The authors suggest that parallel plasticity in CBF and neuronal circuits is occurring. Overall, the study proposes that entrained "vasomotion" contributes to meeting the increased energy demand associated with coordinated neuronal activity and subsequent neuronal circuit reorganization.

We thank the reviewer for providing a thorough summarization of our manuscript.

Strengths:

- *The paper describes a simple and useful method for tracking vasomotion in awake mice through an intact skull.*
- *The work controls for artefacts in their primary measurements.*

- *There are some interesting observations, including the nearly brain-wide synchronization of cerebral blood flow oscillations to visual stimuli and that this process only occurs after mice are trained in a visual task.*
- *This topic is interesting to many in the CBF, functional imaging, and dementia fields.*

We thank the reviewer for positively recognizing the strength of the paper.

Weaknesses:

- *I have concerns with the main concepts put forward, regarding whether the authors are actually studying vasomotion as they state, as opposed to functional hyperemia which is sensory-induced changes in blood flow, which is what they are actually doing. I recommend several additional experiments/analyses for them to explore. This is mostly further characterizing their effect which will benefit the interpretations.*

We recognized that the terminology used in our paper was not explicitly explained. Traditionally, “vasomotion” is defined as the dilation and constriction of the blood vessels that occurs spontaneously at low frequencies in the 0.1 Hz range without any apparent external stimuli. Sensory-induced changes in the blood flow are usually called “hyperemia”. However, in our paper, we used the term, vasomotion, literally, to indicate both forms of “vascular” “motion”. Therefore, the traditional vasomotion was called “spontaneous vasomotion” and the hyperemia induced with slow oscillating visual stimuli was called “visually induced vasomotion”.

Using our newly devised methods, we show the presence of “spontaneous vasomotion”. However, this spontaneous vasomotion was often fragmented and did not last long at a specific frequency. With visual stimuli that slowly oscillated at temporal frequencies close to the frequency of spontaneous vasomotion, oscillating hyperemia, or “visually induced vasomotion” was observed.

- *Neuronal calcium imaging would also benefit the study and improve the interpretations.*

In our paper, we mainly studied the visually induced vasomotion (or visual stimulus-triggered vasomotion). Therefore, visual stimulation must first activate the neurons and, through neurovascular coupling, the initial drive for vasomotion is likely triggered. However, visually induced vasomotion is not observed in novice animals. Therefore, the visually induced vasomotion is not a simple sensory reaction of the vascular in response to neuronal activity in the primary visual cortex. We also do not know how the synchronized vasomotion can spread throughout the whole brain. Where the plasticity for vasomotion entrainment occurs is also unknown. To identify the extent of the neuronal contribution to the vasomotion triggering, whole brain synchronization, and vasomotion entrainment, simultaneous neuronal calcium imaging would be ideal. However, due to the fact that fluorescent Ca²⁺ indicators expressed in neurons would also be distorted by the “shadow” effect from the vasomotion, exquisite imaging techniques would be required.

- *The plastic effects in vasomotion synchronization that occur with training are interesting but they could use an additional control for stress. Is this really a*

plastic effect, or is it caused by progressively decreasing stress as trials and progress? I recommend a habituation control experiment.

As also pointed out by reviewer #2, we agree that, whether stress would affect visually induced vasomotion or not could be studied. Studying the visually induced vasomotion in mice well-habituated to the experimental apparatus would give an idea of whether stress could truly be a profounding factor affecting vasomotion. On the other hand, whether acutely induced stress can interfere with the already entrained vasomotion could also be studied. In the experiments presented in Fig. 3, Texas Red was injected via the tail vein, which would be quite stressful for the mouse. However, in the trained mouse, visually induced vasomotion could be observed regardless of the stress. It is likely that stress can interfere with the acquisition of vasomotion entrainment, but the already acquired entrainment will not be canceled with acute stress induced by tail-vein injection. We agree that further relationship between stress and vasomotion and plasticity related to vasomotion entrainment could be investigated.

Appraisal

I think the authors have an interesting effect that requires further characterization and controls. Their interpretations are likely sound and additional experiments will continue to support the main hypothesis. If brain-wide synchrony of blood flow can be trained and entrained by external stimuli, this may have interesting therapeutic potential to help clear out toxic proteins from the brain as seen in several neurodegenerative diseases.

We thank the reviewer for the positive evaluation of our manuscript. Strong entrainment of visually induced vasomotion was observed with a simple presentation of slowly oscillating visual stimuli for several days. This is a totally non-invasive method to train the vasomotion capacity. As the reviewer recognizes, potential benefits for the treatment of dementia and neurodegenerative diseases could be evaluated with further studies.