

Dual engagement of the nucleosomal acidic patches is essential for deposition of histone H2A.Z by SWR1C

Reviewed Preprint

Revised by authors after peer review.

About eLife's process

Reviewed preprint version 2

May 1, 2024 (this version)

Reviewed preprint version 1

February 28, 2024

Sent for peer review

December 21, 2023

Posted to preprint server

November 26, 2023

Alexander S. Baier, Nathan Gioacchini, Priit Eek, Erik M. Leith, Song Tan, Craig L. Peterson 

Program in Molecular Medicine, University of Massachusetts Chan Medical School, Worcester, MA, 01605 • Medical Scientist Training Program, T.H. Chan School of Medicine, University of Massachusetts • Interdisciplinary Graduate Program, Morningside Graduate School of Biomedical Sciences, University of Massachusetts Chan Medical School, Worcester, MA, 01605 • Center for Eukaryotic Gene Regulation, Department of Biochemistry and Molecular Biology, Pennsylvania State University, University Park, PA 16802, USA • Department of Chemistry and Biotechnology, Tallinn University of Technology, 12618 Tallinn, Estonia

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

The SWR1C chromatin remodeling enzyme catalyzes the ATP-dependent exchange of nucleosomal histone H2A for the histone variant H2A.Z, a key variant involved in a multitude of nuclear functions. How the 14-subunit SWR1C engages the nucleosomal substrate remains largely unknown. Studies on the ISWI, CHD1, and SWI/SNF families of chromatin remodeling enzymes have demonstrated key roles for the nucleosomal acidic patch for remodeling activity, however a role for this nucleosomal epitope in nucleosome editing by SWR1C has not been tested. Here, we employ a variety of biochemical assays to demonstrate an essential role for the acidic patch in the H2A.Z exchange reaction. Utilizing asymmetrically assembled nucleosomes, we demonstrate that the acidic patches on each face of the nucleosome are required for SWR1C-mediated dimer exchange, suggesting SWR1C engages the nucleosome in a “pincer-like” conformation, engaging both patches simultaneously. Loss of a single acidic patch results in loss of high affinity nucleosome binding and nucleosomal stimulation of ATPase activity. We identify a conserved arginine-rich motif within the Swc5 subunit that binds the acidic patch and is key for dimer exchange activity. In addition, our cryoEM structure of a Swc5-nucleosome complex suggests that promoter proximal, histone H2B ubiquitinylation may regulate H2A.Z deposition. Together these findings provide new insights into how SWR1C engages its nucleosomal substrate to promote efficient H2A.Z deposition.

eLife assessment

This manuscript presents an **important** analysis of the role that the nucleosome acidic patch plays in SWR1-catalyzed histone exchange. This manuscript contains **convincing** data which significantly expands our understanding of the complex process of H2A.Z deposition by SWR1 and therefore would be of interest to a broad readership.

Introduction

Eukaryotic genomes regulate access to information stored in their genetic code through the dynamic nucleoprotein structure known as chromatin. The basic unit of chromatin, the nucleosome, sequesters ~147 base pairs (bp) of DNA around a histone octamer consisting of a H3-H4 tetramer flanked by two H2A-H2B (AB) hetero-dimers¹. Rather than a static structure, chromatin is highly dynamic, due to the coordinated impact of histone post translational modifications, ATP-dependent chromatin remodeling enzymes, and the incorporation of histone variants². Disruption of these processes can have deleterious consequences for all nuclear events, including transcription, DNA replication, and genome stability pathways.

H2A.Z, a variant of the core histone H2A, is enriched within nucleosomes adjacent to genes transcribed by RNA polymerase II, replication origins, centromeres, and at sites of DNA double strand break repair³⁻⁷. In yeast, H2A.Z enrichment correlates with higher rates of nucleosome turnover that facilitates transcription, as well as antagonizing the spread of heterochromatic regions from telomeres and the silent mating type loci⁸⁻¹⁰. Recently, we found that H2A.Z plays a global, positive transcriptional role in yeast strains lacking the nuclear RNA exosome¹¹. While yeast H2A.Z is not essential for viability, H2A.Z is essential in metazoans where it is associated with both transcriptional activation and repression¹². For instance, in embryonic stem cells, deposition and function of H2A.Z is interdependent with the polycomb repressive complex 2 (PRC2), and loss of either H2A.Z or PRC2 leads to transcriptional de-repression of a host genes and a subsequent failure in lineage commitment¹³.

The site-specific deposition of H2A.Z is catalyzed by ATP-dependent chromatin remodeling enzymes related to the 14-subunit, yeast SWR1C complex^{14,15}. Mammals contain two enzymes related to SWR1C, the SRCAP and Tip60/p400 complexes, and subunits of the mammalian enzymes are highly conserved with those of SWR1C^{16,17}. Each of these enzymes are members of the larger INO80 subfamily of chromatin remodeling enzymes, distinguished by a large insertion domain between the two ATPase lobes of the catalytic ATPase (Swr1 in SWR1C)^{2,18,19}. This insertion domain serves as a docking site for several key subunits of SWR1C-like complexes, including the Rvb1/Rvb2 heterohexameric ring (RUVB1/RUVB2 in humans) which acts as a further scaffold for organizing additional subunits^{20,21}. In addition to Rvb1/Rvb2, previous studies have demonstrated that the deposition of H2A.Z by SWR1C requires several key subunits, including Swc2, Swc4, Yaf9, and Swc5^{20,22}. The Swc2 subunit and its mammalian ortholog, Yl-1, promotes nucleosome binding and assists SWR1C in the recognition of nucleosome free regions adjacent to gene promoter regions^{20,23,24}. Swc4 (DMAP1 in mammals) is also part of the yeast NuA4 histone acetyltransferase complex, and it appears to promote DNA and histone binding^{25,26}. Swc5 (Cfdp1 in mammals) is essential for the ATPase activity of SWR1C, but how Swc5 promotes SWR1C activity is poorly understood²⁷.

The subunit complexity of chromatin remodelers varies from the single subunit, yeast CHD1, to multi-subunit ~1MDa enzymes such as RSC, SWI/SNF, INO80C, and SWR1C. Despite differences in function and complexity, recent studies have shown that the activity of many remodelers require that they engage a solvent exposed, acidic surface on the nucleosome²⁸⁻³³. This “acidic patch” is composed of eight residues from histones H2B and H2A that are known to provide a docking interface for many proteins. For instance, PRC1, RCC1, and Orc1 bind to the nucleosome acidic patch through a common binding motif consisting of a loop region with an arginine residue that inserts directly into the acidic pocket³⁴⁻³⁶. Though these regions are primarily unstructured and basic, nucleosomal docking requires the specific side chain structure of an arginine, which cannot be substituted by lysine, resulting in these regions being referred to as “arginine anchors”^{37,38}. In the context of remodelers, ISWI uses an arginine anchor within its catalytic ATPase subunit to relieve an autoinhibitory mechanism following nucleosome binding²⁹.

Members of the SWI/SNF family of enzymes also use arginine-rich motifs within both the conserved Snf5-like and Snf2-like subunits to sandwich the complex to the octamer during remodeling activity^{39–41}. A recent cryoEM structure of INO80C also shows key interactions with the nucleosomal acidic patch⁴², and this remodeler is unable to remodel nucleosomes that lack this nucleosomal epitope²⁹.

Here we investigate the role of the nucleosome acidic patch in dimer exchange by SWR1C. Through Förster-resonance energy transfer (FRET), fluorescence intensity (FI), and fluorescence polarization (FP) assays, we interrogate how SWR1C is influenced by H2A/H2B acidic patch alterations, and we specifically probe how each acidic patch on the two faces of the nucleosome contribute to SWR1C activity. Surprisingly, we find that loss of even a single acidic patch results in the loss of dimer exchange activity, suggesting SWR1C must engage both acidic patches simultaneously in a “pincer-like” fashion to carry out dimer exchange. In addition, we found that both linker DNA and the contralateral incorporation of an H2A.Z/H2B dimer stimulated H2A.Z deposition. Efforts to identify potential SWR1C subunits responsible for acidic patch interactions led us to identify an arginine rich motif within the Swc5 subunit of SWR1C, and we show that this domain is essential for dimer exchange *in vitro*, the function of SWR1C *in vivo*, and Swc5 binding to the nucleosome. A low-resolution cryoEM structure of the Swc5-nucleosome complex confirms an acidic patch interaction, as well as contacts with the histone H4 N-terminal tail and the H2B C-terminal helix. These results indicate that SWR1C activity relies on communication between the Swc5 arginine motif and the nucleosome acidic patch to functionally engage nucleosomes for dimer exchange activity.

Results

Dimer exchange by SWR1C requires the nucleosomal acidic patch

To investigate the role of the nucleosomal acidic patch on dimer exchange by SWR1C, a fluorescence-based assay was initially employed. Recombinant nucleosomes were assembled with octamers containing wildtype *Saccharomyces cerevisiae* H2A/H2B (AB) dimers and *Xenopus laevis* H3/H4 tetramers or with *Saccharomyces cerevisiae* H2A/H2B dimers containing 8 alanine substitutions for residues that comprise the nucleosomal acidic patch (AB-apm, see method section for residue list). The DNA template for nucleosome reconstitutions was a 224-bp fragment containing a “601” nucleosome positioning sequence and an asymmetric, 77 bp linker DNA to mimic the structure of a promoter-proximal nucleosome located next to a nucleosome depleted region (77N0)⁴³. These substrates contain a Cy3 fluorophore conjugated to the linker-distal end of the nucleosomal DNA, and a Cy5 fluorophore attached to an engineered cysteine residue (H2A-119) within the histone H2A C-terminal domain. The Cy3 and Cy5 fluorophores are within an appropriate distance to function as a FRET pair, such that excitation of the Cy3 donor leads to efficient energy transfer to the Cy5 acceptor, as evidenced by the fluorescence emission peak at 670 nm. The dimer exchange activity of SWR1C is monitored by following the decrease in the 670 nm FRET signal due to eviction of the Cy5-labeled AB-Cy5 dimer (**Figure 1A**).

Reactions to measure dimer exchange by SWR1C were performed under single turnover conditions (excess enzyme to nucleosomal substrate) and contained free H2A.Z/H2B (ZB) dimers which act as an essential co-substrate. Addition of SWR1C to a wildtype nucleosome led to a rapid drop in FRET, showing the biphasic kinetics consistent with the sequential exchange of the two nucleosomal H2A/H2B (AB) dimers^{44,45}. Incubation of SWR1C with a nucleosome that lacks an intact acidic patch (AB-apm/AB-apm nucleosome) showed little decrease in the FRET signal, indicating that an intact nucleosomal acidic patch is essential for SWR1C to catalyze H2A/H2B eviction (**Figure 1B**).




Figure 1

H2A.Z deposition requires each nucleosomal acidic patch.

(A) Schematic of SWR1C mediated dimer exchange. Cylinder is representative of a nucleosome; light grey area of cylinder represents H3/H4 tetramer, dark grey area of cylinder represents AB heterodimer, yellow star represents Cy5, pink star represents Cy3, and solid black line represents DNA. The nucleosome undergoes two rounds of SWR1C mediated dimer eviction where ZB heterodimers (orange) replace AB heterodimers. **(B)** 77N0-Cy3 nucleosomes were remodeled by SWR1C under single turnover conditions (30 nM SWR1C, 10 nM 77N0-Cy3 nucleosomes, 70 nM ZB dimers, 1mM ATP or AMP-PNP). Dimer eviction is monitored by measuring Cy5 emission at 670 nm. SWR1C was able to perform dimer exchange on an AB-Cy5/AB-Cy5 nucleosome (red line) but was unable to exchange dimers on an AB-apm-Cy5/AB-apm-Cy5 nucleosome (black line). The AMP-PNP reaction contained the AB-apm-Cy5/AB-apm-Cy5 nucleosomal substrate. **(C)** Asymmetrically assembled 77N0-Cy3 and Cy3-0N77 nucleosomes were remodeled by SWR1C under single turnover conditions (50 nM SWR1C, 10 nM nucleosomes, 50 nM ZB dimers, 1mM ATP) and one-phase eviction rates were calculated in Prism 9 and plotted. The dimer labels are listed in linker proximal (LP), linker distal (LD) order (see inset for template nucleosome represented as a cylinder with H3/H4 tetramer in light grey, LP dimer in green, LD dimer in dark blue, DNA as a solid black line with Cy3 (pink star) on the 0bp-linker side). Nucleosomes with a ZB dimer contralateral to the dimer being evicted had significantly faster rates for both linker distal (first two bars) and linker proximal (second two bars). Linker distal eviction was also significantly faster than linker proximal eviction. Rates for nucleosomes containing an AB-apm dimer contralateral to the dimer being evicted (5th and 6th bars) were not significantly different from no enzyme controls (last bar) in either orientation. SWR1C was also unable to evict a dimer with an acidic patch mutant (7th bar). **(D)** Asymmetrically assembled 0N0-Cy3 nucleosomes were remodeled by SWR1C under single turnover conditions (50 nM SWR1C, 10 nM nucleosomes, 50 nM ZB dimers, 1mM ATP) and one-phase eviction rates were calculated in Prism 9 and plotted. Nucleosomes contained a single AB-Cy5 dimer and a contralateral AB or ZB dimer (see inset for template nucleosome represented as a cylinder with H3/H4 tetramer in light grey, AB-Cy5 dimer in dark grey, contralateral AB or ZB dimer represented by dark grey to orange gradient, 0N0-Cy3 DNA as a solid black line with Cy3 (pink star) label on one side). AB-Cy5/ZB 0N0-Cy3 nucleosomes had significantly faster eviction than AB-Cy5/AB 0N0-Cy3 nucleosomes. At least 3 independent nucleosome preparations were used for substrate, and error bars reflect 95% confidence intervals from at least 3 replicates.

Since the FRET assay is limited to monitoring the eviction of nucleosomal H2A/H2B dimers, a gel-based assay that monitors deposition of H2A.Z was employed to confirm these findings.¹⁹ In this assay, an unlabeled, 77N0 nucleosome was incubated with SWR1C, ATP, and free H2A.Z/H2B dimers in which H2A.Z contains a 3xFLAG tag at its C-terminus. Reaction products are separated on native PAGE, and formation of the heterotypic (AB/ZB) and homotypic (ZB/ZB) nucleosomal products are detected by their reduced gel migration due to the 3xFLAG tag on H2A.Z. Similar to the FRET-based assay, the SWR1C-catalyzed deposition of H2A.Z was largely complete by 30-60 minutes for the AB/AB nucleosomal substrate, but addition of SWR1C to the AB-apm/AB-apm nucleosome led to 10-fold reduction in H2A.Z deposition after 60 minutes (Supplemental Figure 1A, B, D). Together, these assays indicate that SWR1C requires an intact nucleosomal acidic patch for H2A.Z deposition.

To further interrogate the role for each of the two nucleosomal acidic patches in dimer exchange by SWR1C, we performed dimer exchange reactions with asymmetrically assembled nucleosomal substrates. Such asymmetric nucleosomes allow the fluorescent labelling of only a single H2A/H2B dimer, such that we could monitor which dimer was targeted for replacement. We generated these substrates by leveraging the asymmetric features of the 601 nucleosome positioning sequence and the spontaneous incorporation of dimers into purified hexasomes⁴⁶ (Supplemental Figure 2A). Hexasomes were generated by varying the ratio of *Xenopus laevis* H3/H4 tetramers to yeast H2A/H2B heterodimers on 77N0 or ON0 DNA that was labeled with Cy3 (Supplemental Figure 2B). After hexasome purification, nucleosomes of different compositions were reconstituted by titration of the appropriate yeast heterodimer (Supplemental Figure 2C, D). For nucleosomes with linker DNA, we list the asymmetric dimers in linker proximal, linker distal order throughout (e.g. AB/AB-apm for a nucleosome with a linker distal H2A/H2B dimer harboring the acidic patch alteration).

First, we monitored H2A.Z deposition with nucleosomes harboring an unlabeled H2A/H2B heterodimer with a disrupted acidic patch (AB-apm) assembled on the nucleosomal face opposite to the Cy5-labelled heterodimer that will be monitored for H2A.Z deposition (Supplemental Figure 3G,H). Strikingly, H2A.Z deposition was crippled, irrespective of whether the linker distal or linker proximal H2A/H2B heterodimer was the target for exchange (**Figure 1C**, AB-apm/AB-Cy5 and AB-Cy5/AB-apm substrates). We then tested if the Cy5-labelled heterodimer that is targeted for replacement also requires an intact acidic patch (Supplemental Figure 3I). In this case as well, H2A.Z deposition was not greater than the no enzyme control (**Figure 1C**, AB/AB-apm-Cy5 substrate). Thus, SWR1C requires that both faces of the nucleosome contain an intact acidic patch, irrespective of which H2A/H2B heterodimer is undergoing replacement.

Previously, we and others reported that SWR1C preferentially exchanges the linker distal H2A/H2B dimer, as monitored by either ensemble or single molecule FRET analyses.^{44,47,48} The asymmetric nucleosome substrates, harboring only a single, labeled H2A/H2B dimer were leveraged to further investigate this preferential exchange. Consistent with previous results, a substrate with a labeled, linker distal heterodimer is exchanged at a ~3-fold faster rate by SWR1C compared to the linker proximal dimer (**Figure 1C**, compare AB/AB-Cy5 to AB-Cy5/AB; Supplemental Figure 3A,B). To rule out the possibility that this preferential exchange was influenced by the unlabeled H2A/H2B dimer on the opposing nucleosomal face, nucleosomes harboring an unlabeled ZB dimer were also reconstituted. In these cases as well, the linker distal H2A/H2B dimer was exchanged at a more rapid rate, compared to the linker proximal heterodimer (**Figure 1C**, compare ZB/AB-Cy5 to AB-Cy5/ZB; Supplemental Figure 4 D,E). Surprisingly, incorporation of the ZB heterodimer on the contralateral (opposite to the dimer being evicted) nucleosome face stimulated H2A.Z deposition, irrespective of linker orientation (**Figure 1C**; purple bars). The stimulatory effect of contralateral ZB heterodimer placement remained even after elimination of the linker DNA (**Figure 1D**, Supplemental Figure 4C,F). Interestingly, these assays also confirmed that linker DNA has a stimulatory impact on the H2A.Z deposition reaction, with both linker distal and linker proximal eviction occurring at a faster rate than

eviction from a nucleosome core particle²³. To test whether the stimulation by linker DNA was due to differences in nucleosome binding affinity, we ensured the experiment was done under saturating enzyme levels by repeating it at a four-fold increase in the concentration of SWR1C. At this higher concentration of SWR1C, the rates were unchanged, and the stimulatory impact of linker DNA remained (Supplemental Figure 4A). Thus, both linker DNA and nucleosomal ZB dimers appear to stimulate H2A.Z deposition.

Intact nucleosomal acidic patches are required for SWR1C ATPase activity

The ATPase activity of SWR1C is stimulated by both its nucleosomal substrate and free ZB dimers¹⁹. Notably, the ATPase activity of SWR1C is not stimulated by a ZB/ZB nucleosome, implying that SWR1C can distinguish product from substrate¹⁹. To investigate the role of the nucleosomal acidic patch in the stimulation of SWR1C ATPase activity, SWR1C ATPase activity was measured in the presence or absence of different 77N0-Cy3 nucleosomes. Consistent with previous results¹⁹, SWR1C ATPase activity was stimulated by AB/AB nucleosomes as compared to the basal level. However, ATPase activity was not stimulated by either of the asymmetric acidic patch mutant nucleosomes (AB-apm/AB, AB/AB-apm) or by a nucleosome lacking both acidic patches (AB-apm/AB-apm), demonstrating the essential role of dual acidic patch engagement by SWR1C for stimulation of ATPase activity (Supplemental Figure 4B).

The nucleosomal acidic patch is a key driver of SWR1C binding affinity

The inability of APM nucleosomes to stimulate the ATPase activity of SWR1C could be due to a defect in binding or the inability to form an active complex post-binding. To explore whether disruption of each acidic patch might impact the nucleosome binding affinity of SWR1C for nucleosomes, we leveraged a fluorescence polarization assay. Using a Cy3-labeled 77N0 nucleosome, we determined a K_d of 13.6 nM for an AB/AB nucleosome (Figure 2A), similar to prior studies⁴⁴. However, disruption of even a single nucleosomal acidic patch severely weakened SWR1C binding, such that a K_d was unable to be determined (Figure 2B-D). Given the range of SWR1C concentrations that could be used in this assay, we estimate that alteration of one or both nucleosomal acidic patches decreases binding affinity by at least 10-fold. These data indicate that the inability of APM nucleosomes to stimulate SWR1C ATPase activity is most likely due to a defect at the nucleosome binding step (K_m), rather than K_{cat} , though an impact at this latter step cannot be eliminated as yet.

Swc5 interacts with the nucleosomal acidic patch

Recently, a potential arginine anchor domain, the archetypal motif for proteins interacting with the nucleosomal acidic patch, was identified within the N-terminal, acidic domain of the SWR1C subunit, Swc2. However, removal of this domain (Swc2-ZN) has only a modest impact on the dimer exchange activity of SWR1C, and this data, combined with the essential nature of the nucleosomal acidic patches for SWR1C activity, suggest that additional subunits must contribute to acidic patch recognition⁴⁹. The Swc5 subunit is one candidate, as loss of Swc5 eliminates H2A.Z deposition and nucleosome-stimulated ATPase activity²⁰ (See Figure 3B, C). Furthermore, removal of the Swc5 subunit from SWR1C crippled nucleosome binding, as measured by fluorescence polarization (Figure 3D). Previous studies of Swc5 identified an acidic N-terminal domain that interacts preferentially with AB dimers, and a conserved C-terminal domain, termed BCNT, that is essential for dimer exchange by SWR1C^{27,50}. An alignment of Swc5 homologs revealed a region adjacent to the BCNT domain, containing multiple arginine residues in a conserved basic region (Figure 3A).




Figure 2

Nucleosomal acidic patches are required for SWR1C nucleosome binding.

Asymmetrically assembled 77N0-Cy3 nucleosomes were incubated with SWR1C serially diluted in concentration from approximately 1 μ M to 1 nM. Fluorescence polarization values were collected and plotted by concentration for each nucleosome type. Each graph has a representative nucleosome at the top left (represented by a cylinder with H3/H4 tetramer in light grey, AB dimer in dark grey, AB-apm dimer in light blue, 77N0-Cy3 DNA as a solid black line with Cy3 (pink star) label on the 0bp-linker side). FP data was normalized between experiments by translation, which does not effect the binding curve fit so that plots can be compared at a glance. **(A)** AB/AB 77N0-Cy3 nucleosomes showed a K_d as determined by a fit to the Morrison equation (blue line). The K_d calculated for each individual replicate was 13.55, 13.32, 13.94, and 13.75. The average of those individual values is 13.64. The standard deviation is 0.27. A K_d was unable to be calculated for nucleosomes with a linker proximal acidic patch mutant **(B)**, a linker distal acidic patch mutant **(C)**, or both acidic patches mutated **(D)** at the range of concentrations of SWR1C tested suggesting at least a 10-fold reduction in binding affinity. Error bars reflect 95% confidence intervals from at least 3 replicates.




Figure 3

Swc5 contains a key arginine-rich motif.

(A) Alignment of SWR1C subunit Swc5 homologs reveal a conserved arginine in a region we named the RRKR motif in *S. cerevisiae* adjacent to the essential BCNT region and conserved LDW residues (asterisks are used to denote residues mutated in RRKR-4A and LDW-3A mutants). A linear schematic of this gene with the acidic N-terminal domain in orange, gene body in grey, RRKR motif in red and BCNT region in purple shows relative locations of these domains. **(B)** Swc5 and Swc5 derivatives were reconstituted into SWR1C^{Swc5Δ} complexes to measure their activity on dimer exchange using FRET at single turnover conditions (30 nM SWR1C, 10 nM 77N0-Cy3 AB-Cy5/AB-Cy5 nucleosomes, 70 nM ZB dimers, 1 mM ATP). The RFU for each reaction was normalized and plotted as a function of time. SWR1C^{Swc5Δ} was unable to carry out dimer exchange (black triangles), and recombinant Swc5 (rSwc5) rescued activity (red circles). SWR1C^{Swc5Δ} reconstituted with Swc5 containing mutations in the LDW (rSwc5^{LDW-3A}) or RRKR motifs (rSwc5^{RRKR-4A}) showed no dimer eviction activity (purple and blue circles respectively), while an N-terminal truncation of Swc5 (rSwc5⁷⁹⁻³⁰³) had a reduction in activity compared to full length Swc5 (yellow triangles). **(C)** Asymmetrically assembled 77N0-Cy3 nucleosomes were incubated with SWR1C^{Swc5Δ} serially diluted in concentration from approximately 1 μM to 1 nM. Fluorescence polarization values were collected and plotted by concentration. A representative nucleosome is shown at the top left (represented by a cylinder with H3/H4 tetramer in light grey, AB dimer in dark grey, 77N0-Cy3 DNA as a solid black line with Cy3 (pink star) label on the 0bp-linker side). **(D)** Nucleosomal stimulation of SWR1C ATPase activity with (purple bars) or without (green bars) ZB dimers added was measured for complexes containing SWR1C, SWR1C^{Swc5Δ}, SWR1C^{Swc5Δ} + rSwc5, and SWR1C^{Swc5Δ} + rSwc5^{RRKR-4A} using a phosphate sensor assay. Calculated rates were normalized to basal SWR1C activity (blue bar). Stimulation of ATPase activity was lost in the SWR1C^{Swc5Δ} complex and reduced in the SWR1C^{Swc5Δ} + rSwc5^{RRKR-4A} complex. Error bars reflect 95% confidence intervals from at least 3 replicates.

To investigate the role of this Swc5 arginine-rich domain, alanine substitutions were created in four basic residues (RRKR), and this derivative was recombinantly expressed and reconstituted into a SWR1C complex purified from a *swc5* deletion strain (SWR1C^{swc5Δ}). In addition, SWR1C was reconstituted with wildtype Swc5, a Swc5 derivative lacking the acidic N-terminal domain (Swc5^{79–303}), and a Swc5 derivative harboring alanine substitutions in the essential BCNT domain (Swc5^{LDW-3A}). In all cases, derivatives were incorporated into SWR1C complexes with equal efficiencies (Supplemental Figure 5). Consistent with previous studies, reconstitution with wildtype Swc5 fully restored dimer exchange activity to SWR1C^{swc5Δ} (**Figure 3B** [↗](#), red curve). In contrast, SWR1C harboring Swc5^{LDW-3A} had no detectable exchange activity, and the complex that lacks the N-terminal, acidic domain of Swc5 (Swc5^{79–303}) exhibited a modest defect, as expected from previous studies (**Figure 3B** [↗](#))²⁷. Strikingly, SWR1C that contained the Swc5^{RRKR-4A} derivative showed minimal activity in the FRET-based exchange assay (**Figure 3B** [↗](#)), and this complex exhibited only ~10% the activity of wildtype SWR1C in the gel-based H2A.Z deposition assay (Supplemental Figure 1A, C, D). Dimer exchange activity for each SWR1C complex was also mirrored by their ATPase activity (**Figure 3D** [↗](#)). SWR1C^{swc5Δ} lost nucleosome-stimulated ATPase activity, with or without free ZB dimers, as compared to wild-type. As expected, reconstitution of SWR1C^{swc5Δ} with recombinant Swc5 rescued ATPase activity. Underscoring the importance of the arginine-rich domain, reconstitution of SWR1C^{swc5Δ} with Swc5^{RRKR-4A} showed a large defect in nucleosome-stimulated ATPase activity, revealing its importance in both dimer exchange and ATPase activity. Additionally, we reinforced prior findings that the ATPase activity of the Swc5^{LDW-3A} complex was not stimulated by either nucleosomes or free ZB dimers, while the complex harboring Swc5^{79–303} had wildtype levels of ATPase activity (Supplemental Figure 4C).

To further investigate whether the arginine-rich domain within Swc5 binds to the nucleosomal acidic patch, *in vitro* binding assays were performed with nucleosome core particles (ON0) and recombinant Swc5 (**Figure 4A–F** [↗](#)). Increasing amounts of Swc5 were incubated with nucleosomes, and binding was visualized by native PAGE. Addition of wildtype Swc5 to nucleosomes led to formation of discrete complexes with an apparent K_d of 125.7 nM (**Figure 4A** [↗](#)). Likewise, binding of Swc5^{LDW-3A}, which has a disrupted BCNT domain, bound to nucleosomes with nearly identical affinity, compared to wildtype Swc5 (**Figure 4B** [↗](#)). Importantly, high affinity binding of Swc5 to nucleosomes required an intact acidic patch, with binding reduced ~3-fold when an APM nucleosome was used as the substrate (**Figure 4C** [↗](#)). Furthermore, the Swc5^{RRKR-4A} derivative also showed an ~3-fold loss in affinity, with an apparent K_d of 379.1 nM (**Figure 4D** [↗](#)). Together these data demonstrate that Swc5 has nucleosome binding activity, and the data are consistent with direct interactions between the Swc5 arginine-rich domain and the nucleosomal acidic patch.

A time-resolved FRET assay (TR-FRET) was employed (**Figure 5A** [↗](#)) as an additional solution approach to monitor the binding of Swc5 to nucleosomes⁵¹. In this assay, increasing concentrations of recombinant, 6His-tagged Swc5 were incubated with a 31N1 nucleosome. The 6His-Swc5 protein were labelled with a ULIGHT alpha-6xHIS acceptor antibody, and biotinylated nucleosomal DNA was labelled with a Eu-streptavidin donor fluorophore (**Figure 5A** [↗](#)). The binding of Swc5 to the nucleosome was monitored by quantitative measurements of the time-resolved FRET signal between the ULIGHT alpha acceptor and the Eu fluorophore. Binding curves for wildtype Swc5 yielded an apparent K_d of 133 ± 12 nM, a value that agrees well with the K_d measured by the nucleosome gel shift assay. Binding experiments performed in parallel with the Swc5^{RRKR-4A} derivative yielded a K_d of 593 ± 28 nM, confirming that the arginine-rich motif within Swc5 plays a key role in nucleosome recognition (**Figure 5B** [↗](#)).

While the gel shift and TR-FRET assays establish that Swc5 can bind to nucleosomes, data from these assays can only indirectly link the nucleosomal acidic patch to the Swc5 arginine-rich domain. To probe for more direct interactions, a fluorescence quenching assay was employed. These assays exploit nucleosomes harboring a site-specific, Oregon Green fluorophore whose emission is sensitive to the chemical environment whereby fluorescence is quenched by protein




Figure 4

Swc5 is a nucleosome binding subunit.

Gel mobility shift assays were performed with 5 nM ONO nucleosomes and recombinant Swc5 ranging in concentration from 0 to 500 nM, in 50 nM increments. Swc5 was able to bind AB/AB nucleosomes (**A**) but showed reduced binding of AB-apm/AB-apm nucleosomes (**B**). Swc5^{LDW-3A} bound AB/AB nucleosomes (**C**) whereas Swc5^{RRKR-4A} showed reduced binding (**D**). Percent of nucleosomes bound for each condition was calculated using ImageQuant and plotted by Swc5 concentration (**E**) and specific binding Kd values were predicted along with 95% confidence intervals (CI) predicted in Prism 9 (**F**).




Figure 5

TR-FRET assay confirms Swc5 nucleosome binding activity.

(A) Schematic of the TR-FRET assay. Recombinant 6His-Swc5 is labelled with a ULight alpha 6His acceptor antibody, and biotinylated nucleosomal DNA is labelled with an Eu-streptavidin acceptor fluorophore. The FRET signal increases as 6His-Swc5 binds to the nucleosome. **(B)** TR-FRET assay was performed with 2nM 31N1 nucleosomes and recombinant Swc5 or Swc5^{RRKR-4A} ranging in concentration from 0.6-6000 nM. Recombinant wildtype Swc5 bound with an apparent Kd of ~133nM, while the Swc5^{RRKR-4A} derivative bound with an apparent Kd of ~592nM. Note that the Swc5^{RRKR-4A} assays did not reach full saturation, and thus the measured Kd is likely an under-estimate of the reported value. Kd values were determined from triplicate titrations of each Swc5 variant and are reported as means ± standard error of the mean.

binding^{35,52}. Nucleosome core particles were reconstituted that contained an Oregon Green fluorophore covalently attached to either the histone H4 N-terminal domain (H4-tail) (**Figure 6A**, top) or to a residue directly adjacent to the nucleosomal acidic patch (**Figure 6A**, bottom). Titration of wildtype Swc5 led to the concentration-dependent quenching of the Oregon Green fluorophore positioned at the acidic patch, but little quenching was observed for the fluorophore located on the H4-tail (**Figure 6B**) demonstrating the sensitivity of this assay. Importantly, Swc5 did not quench the acidic patch probe when this substrate also harbored alanine substitutions within the acidic patch (**Figure 6B**). Examined in the same titration of concentrations, we found that the Swc5^{LDW-3A} derivative also showed specific quenching of the acidic patch probe (**Figure 6C**), but the Swc5^{RRKR-4A} derivative was defective for quenching the acidic patch probe (**Figure 6D**). Together these results are consistent with a direct interaction between the arginine-rich domain of Swc5 and the nucleosomal acidic patch, an interaction that is essential for dimer exchange by SWR1C.

Structure of a Swc5-nucleosome complex

We analyzed the structure of Swc5⁷⁹⁻³⁰³ in complex with a 147 bp nucleosome core particle using cryo-EM (**Figure 6**). We collected over 13,000 micrographs of the cross-linked sample and were able to easily resolve the nucleosome structure to 3-Å resolution (Supplemental Figure 6L-O). On the other hand, the moiety corresponding to Swc5 displayed severe structural heterogeneity, hindering our ability to build an atomic model for Swc5. The protein appears to be flexible on the nucleosome, with continuous conformational dynamics apparent (**Figure 7A-D**, Supplemental Movie 1). Nevertheless, reconstructions of particle subsets obtained by local 3D classification show that Swc5 forms multivalent interactions with the nucleosome via the nucleosome acidic patch, the H4 N-terminal histone tail, and the C-terminal helix of H2B (**Figure 7E-G**). In the majority of 3D reconstructions, we observe additional density at the nucleosome acidic patch attributable to an arginine residue, like in other complexes with arginine anchor interactions^{34,37}. Unfortunately, the low local resolution precludes the determination of the precise Swc5 arginine residue involved in this interaction (Supplemental Figure 6D-G). In contrast, the 3-Å resolution of the nucleosome core allows the identification of H2B-K123 as an interacting residue with Swc5. In most 3D classes, Swc5 forms either a partial or a complete bridge between H2B C-terminal helix including K123 and the N-terminal tail of H4.

The Swc5 arginine-rich domain is required for SWR1C function *in vivo*

Previous studies have shown that yeast lacking functional SWR1C grow poorly on media containing formamide^{14,27}. To investigate the functional role of the Swc5 arginine-rich domain *in vivo*, growth assays were performed with isogenic *swc5Δ* strains that harbor low copy vectors that express different Swc5 derivatives (**Figure 8**). As expected, the *swc5Δ* strain grew well on rich media, but was impaired on formamide media (vector), while cells expressing wildtype Swc5 grew well on both media (SWC5). Consistent with a previous study, a strain expressing the derivative with a disrupted BCNT domain showed slow growth on formamide media (Swc5^{LDW-3A}), and the Swc5⁷⁹⁻³⁰³ derivative that lacks the acidic N-terminal domain showed a moderate growth defect²⁷. Importantly, a strain expressing the Swc5^{RRKR-4A} derivative had a severe growth defect on formamide media, consistent with an important role of the arginine rich region for SWR1C function.




Figure 6

Swc5 interacts with the nucleosome acidic patch.

(A) Schematic of Oregon green based fluorescence quenching assay. Top shows a nucleosome (represented as a cylinder with H3/H4 tetramer in light grey, AB dimers in dark grey, and ON0 DNA as a solid black line) with an Oregon green label (green star) on H4. Oregon green is exquisitely sensitive to changes in its local environment. Binding of an acidic patch interacting protein (pink) does not interact with Oregon green label on H4, resulting in no change in fluorescence intensity. A label on H2B near the acidic patch (bottom) is affected by an acidic patch interacting protein, resulting in decreased fluorescence intensity. **(B)** Swc5 was added to 10 nM H4-OG AB/AB ON0 nucleosomes (orange squares), H2B-OG AB/AB ON0 nucleosomes (blue circles) and H2B-OG AB-apm/AB-apm ON0 nucleosomes (black circles) resulting in quenching of the AB/AB H2B-OG ON0 nucleosomes but not the H4-OG AB/AB ON0 nucleosomes or the H2B-OG AB-apm/AB-apm ON0 nucleosomes. **(C)** Swc5^{LDW-3A} similarly quenched H2B-OG AB/AB ON0 nucleosomes with no change in H4-OG AB/AB ON0 nucleosomes while Swc5^{RRKR-4A} **(D)** did not show significant quenching in either condition. Error bars reflect standard deviations from at least 3 replicates.




Figure 7

Cryo-EM structure of Swc5/nucleosome complex.

(A-D) Reconstructions of selected particle subsets obtained from 3D classification that was focused on the Swc5 moiety (purple) illustrate the dynamic nature of Swc5 binding to the nucleosome. Three interactions persist among the 3D classes: (E) Swc5 interaction with the N-terminal tail of histone H4 (green), (F) Swc5 interaction with the C-terminal helix of histone H2B including K123, and (G) an Arginine anchor bound into the acidic patch of the histone core. The gray circles and arrows denote the reconstructions and locations from where the cut-out panels originate.




Figure 8

Swc5 arginine-rich region is key for SWR1C activity in vivo.

Genetic complementation using a *swc5Δ* strain (*MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 swc5!1::kanMX4*) transformed with an empty pRS416 CEN/ARS *URA3* vector (top row in both panels) or the same vector with various alleles of *SWC5*. Cultures were grown to an OD of 1 and spotted with serial dilution onto plates with synthetic complete media lacking uracil without (left) or with (right) 2% formamide.

Discussion

The nucleosomal acidic patch has emerged as a key binding pocket for nearly all ATP-dependent chromatin remodeling complexes. Here, leveraging asymmetrically constructed nucleosomes, we have found that both acidic patches are essential for the ATP-dependent deposition of H2A.Z by SWR1C. Each acidic patch is key for high affinity nucleosome binding, and this defect in binding appears to result in an inability of APM nucleosomes to stimulate SWR1C ATPase activity. The requirement for both acidic patches suggests that SWR1C adopts a pincer conformation, contacting both acidic patches and leading to a conformation that stimulates the ATPase activity of SWR1C from basal levels. Like others have noted⁵³, the thermodynamic landscape of the nucleosome provides a much higher barrier to dimer exchange than sliding, and we suspect this is likely responsible for the greater effect of an asymmetric acidic patch mutant nucleosome on SWR1C as compared to other remodelers. Additionally, while the acidic patch plays a key role in the activity of SWR1C and other chromatin remodeling enzymes, it also serves as a docking point for other non-chromatin remodeling nuclear proteins^{38,54,55}. From our data, we can conclude that other acidic patch interacting proteins may consequently inhibit the activity of SWR1C even if only one acidic patch is sterically occluded.

Recently, the nucleosomal acidic patch has been shown to differentially impact the behavior of chromatin remodeling enzymes in an asymmetric manner^{31,33,46}. Studies on the sliding activity of the ISWI chromatin remodeling enzyme from both humans and *Drosophila melanogaster* show that the linker proximal acidic patch (the DNA entry side) is more important than the linker distal acidic patch for maintaining appropriate sliding activity⁴⁶. In contrast, budding yeast Chd1 shows a small reduction in sliding rate when either acidic patch is disrupted. The nucleosomal acidic patch is also essential for remodeling by SWI/SNF family members, and in this case the two acidic patches on the nucleosome are contacted by two different subunits – Snf5 (human SMARCA1) and the Snf2 (human BRG1) ATPase^{39–41,56}. Cramer and colleagues suggested that the RSC complex may use both acidic patches to sandwich the histone octamer⁴⁰, however, loss of the Snf5 (SMARCA1) arginine anchor has only a small impact on remodeling, whereas the key domain within the Snf2 subunit (the SnAC domain) is essential for activity^{41,57,58}. Thus, unlike SWR1C, both acidic patch surfaces are not essential for the activity of other remodeling enzymes.

The cryoEM structure of a SWR1C-nucleosome complex suggests that the enzyme has two major nucleosomal contacts – the Swr1 ATPase lobes interact with nucleosomal DNA ~2 helical turns from the nucleosomal dyad (SHL2.0), and the Swc6/Arp6 module interacts with DNA at the nucleosomal edge²¹. These contacts encompass the DNA gyre that wraps the AB dimer that is destined to be exchanged in the first round of catalysis (the linker distal dimer⁴⁵). The structure also indicated that each of the two nucleosomal acidic patches may make contacts with SWR1C subunits, although in both cases the unambiguous identification of the amino acid sequence was restricted by the resolution of cryoEM maps. It seems likely that the Swc2 subunit may contact the acidic patch of the H2A/H2B dimer that is not targeted for replacement. Unfortunately, the Swc5 subunit was not visualized in a previous cryoEM structure²¹, but here we find that Swc5 can make an arginine anchor-like interaction with the nucleosomal acidic patch. SWR1C that lacks Swc5 shows the same spectrum of defects as disruption of the nucleosome acidic patch, consistent with Swc5 playing a key role in anchoring SWR1C to the nucleosome. Based on the existing cryoEM map of the SWR1C-nucleosome complex, we propose that Swc5 contacts the H2A/H2B dimer that is targeted for eviction, positioned between the Arp6/Swc6 complex and the Swr1 ATPase. While some observed density at that nucleosome acidic patch was tentatively identified as Swc6 in the cryoEM map, the authors noted that the resolution in this region was insufficient to make a

putative sequence determination⁵⁹. Notably, this location places Swc5 close to its interaction surfaces on Swr1^{60,61}, as well as positioning its acidic N-terminal domain in an ideal location to capture the evicted AB dimer, as previously suggested⁵⁰.

One advantage of the hexosome strategy for reconstitution of asymmetric nucleosomal substrates is that we were able to reconstitute homogeneous, heterotypic Z/A nucleosomes. This nucleosome mimics the product of the first round of H2A.Z deposition, a potential intermediate towards the formation of a homotypic Z/Z nucleosome. Surprisingly, we found that incorporation of one H2A.Z/H2B dimer stimulated the second round of dimer exchange. This stimulation was observed for both linker proximal and linker distal positions, as well as for a nucleosome core particle. The heterotypic Z/A nucleosome (with linker) reproducibly stimulated deposition rate by ~20%, values which are unlikely to be detected in prior studies that used a standard A/A nucleosomal substrate and reaction conditions where nucleosomes were in large excess¹⁹. How incorporation of one H2A.Z/H2B dimer stimulates the second cycle of dimer exchange is not clear, but one possibility is that re-orientation or re-binding of SWR1C may be facilitated if the enzyme encounters an H2A.Z surface on the initial binding event.

Previous studies have investigated the impact of Swc5 on the nucleosome binding activity of SWR1C, concluding that this subunit does not play a role in nucleosome recognition. For instance, initial work from Wu and colleagues employed a nucleosomal array pulldown assay²³, while subsequent studies from Wigley and colleagues used a native gel, electrophoresis shift assay⁶⁰. In both cases, loss of Swc5 from SWR1C had no detectable impact on nucleosome binding. Furthermore, recent live cell imaging from the Wu group showed that Swc5 is not required for general chromatin association by SWR1C⁶². Together with our results, these data suggest that SWR1C may make strong nonspecific interactions with chromatin, most likely through the Swc2 subunit which is known to bind DNA²³. Indeed, deletion of Swc2 results in a substantial reduction in the localization of SWR1C to chromatin *in vivo*⁶². Furthermore, stabilization of a SWR1C-nucleosome complex in a gel matrix may mask the defects in binding that we observe here in a solution-based assay. Notably, our fluorescence polarization assay is unable to detect nucleosome binding by SWR1C if the affinity is reduced by more than 10-fold, and consequently our work indicates that Swc5 is essential for high affinity interactions that are required for subsequent H2A.Z deposition. Whether Swc5 plays roles subsequent to nucleosome recognition is currently unclear.

In addition to contacting the nucleosomal acidic patch, our Swc5-nucleosome structure indicates that Swc5 makes contacts with the H4 N-terminal tail and the H2B C-terminal helix. Previous work has shown that acetylation of the H4 tail can influence both SWR1C recruitment and activity, raising the possibility that the Swc5-H4 interaction may be functionally important⁶³. Interestingly, we find that Swc5 interacts with the H2B C-terminal helix via contact with H2B-K123 which is subject to transcription-associated modification with ubiquitin by the Bre1-Rad6 ubiquitin ligase complex. Furthermore, H2B-K123ub is enriched at the +1 nucleosome, the same nucleosome targeted by SWR1C for H2A.Z deposition. Strikingly, a ChIP-exo study found that H2B-K123ub is enriched at the NFR proximal face of the +1, while H2A.Z is enriched at the NFR distal surface⁶⁴. The anti-correlation between H2B-K123ub and H2A.Z is consistent with the idea that H2B ubiquitylation disrupts the Swc5-nucleosome interaction and regulates H2A.Z deposition. Interestingly, H2B-K123ub is also known to regulate the Chd1 remodeler, but in this case H2B-K123ub does not appear to contact Chd1, but rather it interacts with unwrapped nucleosomal DNA, stimulating sliding activity⁶⁵.

The human homolog of Swc5 is Craniofacial development protein 1 (Cfdp1), although there is currently a lack of information on the role of Cfdp1 in development. Mutations in a human homolog of SWR1C, SRCAP is implicated in Floating Harbor syndrome, which results in skeletal and facial defects⁶⁶. Complementary to this observation, additional evidence suggests Cfdp1 has an essential role in bone development⁶⁷. Interestingly, a survey of Cfdp1 mutations identified in

the TCGA transCancer atlas reveals many missense substitutions within the C-terminal BCNT domain, as well as a cluster of substitutions directly adjacent to the arginine-rich domain. The majority of these Cfdp1 alterations are associated with uterine endometrial carcinomas, suggesting that Cfdp1, and likely the SRCAP remodeler, are key for suppressing cancer initiation or progression in this cell type.

Methods

Yeast strains and culture conditions

The yeast strain *W1588-4C swr1::SWR1– 3xFLAG-P-kanMx-P htz1Δ::natMX* (yEL190) was a gift from Ed Luk (SUNY Stony Brook). Deletion of *SWC5* was constructed by standard gene replacement with a Hygromycin B cassette (pAG32)⁶⁸, and used to make *W1588-4C swr1::SWR1– 3xFLAG-P-kanMx-P htz1Δ::natMX swc5Δ::hphMX* (CY2535). The *swc5Δ* strain *MATa his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 YBR231c::kanMX4* (Y03371) used for spot assays was purchased from Euroscarf.

Spot assays were performed with the *swc5Δ* strain (Y03371) transformed with an empty CEN/ARS/URA3 vector pRS416, or ones containing WT and mutant *SWC5*. Cultures were diluted to an OD₆₀₀ of 1 and then serially diluted 1:10. A dilution (7 μL) was spotted onto synthetic complete media lacking uracil with and without 2% formamide. Plates were incubated for 3 days before imaging.

Plasmid Construction

The vector (pQE80L) was used for recombinant Swc5 expression. The *SWC5* gene and *SWC5*^{79–303} were inserted in frame with the N-terminal 6x histidine tag through Gibson assembly⁶⁹ after digesting the vector with BamHI and SphI. Once WT and truncated Swc5 were in frame, the alanine mutants [*swc5 (RRKR-4A)* and *swc5 (LDW-3A)*] were generated using QuickChange site directed mutagenesis. The pRS416 CEN/ARS/URA3 vector was digested with XbaI and SacI to allow for insertion of the *SWC5* gene with 500bp upstream of the start site and 500bp downstream of the stop. This region was initially PCR amplified from a W303 strain where the primers had homology to the XbaI and SacI cut sites of the vector. Gibson assembly was used to ligate the vector and insert (CP1566)⁶⁹. Once the WT *SWC5* gene was inserted, site directed mutagenesis was used on CP1566 to make *swc5 RRRK-4A* (CP1579) and *swc5 LDW-3A* (CP1580). *swc5 79-303* construct was generated using CP1566 to PCR amplify the 500bp upstream stretch of DNA and start site of *SWC5* and PCR amplifying another DNA fragment starting from Swc5 K79 to 500bp downstream of the stop. Primers were designed so each PCR fragment had homology to each other and the pRS416 vector at XbaI and SphI so that the two fragments and the pRS416 vector could be ligated together by Gibson assembly. Site directed mutagenesis was used to generate cysteine mutants used for Oregon Green labeling, as well as for single amino acid substitution used for assembly of acidic patch mutant (AB-apm) dimers – H2A-apm (H2A-E63A, H2A-E65A, H2A-D74A, H2A-D92A, H2A-D93A, H2A-E94A) and H2B-apm (H2B-E109A, H2B-E117A).

Protein purification for *in vitro* assays

Histones

Histones were generated as described previously^{45,70}. For experiments in this paper, we produced histones H2A, H2B, H2A-K119C, H2A-apm, H2B-apm, H2B-S115C, H2B-apm-S115C, H2A.Z, and H2A.Z-3xFLAG from *Saccharomyces cerevisiae*, as well as histones H3, H4, and H4-Q27C from *Xenopus laevis*. Fluorescent labels for histones were conjugated on an as needed basis prior to nucleosome or hexasome reconstitution as described previously⁷¹.

Swr1 c

Swr1-3xFlag yeast strains were grown in 6x2L of YEPD, supplemented with adenine at 30°C until reaching an OD₆₀₀ of 3. Cultures were centrifuged and pellets were noodled into liquid nitrogen and stored at -80°C. SWR1C was purified as previously described^{19,43,45} with minor modifications. A PM 100 cryomill was used to lyse the harvested yeast noodles with 6 × 1 min cycles at 400 rpm then stored at -80°C. Lysed cell powder (~200 mL) were resuspended in 200 mL 1x HC lysis buffer ([25 mM HEPES-KOH pH 7.6, 10% glycerol, 1 mM EDTA, 300 mM KCl, 0.01% NP40, 10 mM β-glycerophosphate, 1 mM Na-butyrate, 0.5 mM NaF, 1mM DTT plus 1× protease inhibitors [PI 1mM PMSF, 1mM benzamidine, 0.1 mg/mL pepstatin A, 0.1 mg/mL leupeptin, 0.1 mg/mL chymostatin)] in a 1L beaker with a stir bar at 4°C. Thawed cells were centrifuged for 2 hours at 35000rpm in a Ti45 rotor at 4°C then whole cell extract was transferred to a 250 mL falcon tube along with anti-Flag M2 agarose resin (1 mL bed volume) and nutated at 4°C for 4 hours. Resin and extract mixture was transferred in 25 mL increments to a 25 mL gravity column. After flow through of extract, the resin was washed with 4x 25 mL B-0.5 buffer (25 mM HEPES, pH 7.6, 1 mM EDTA, 2 mM MgCl₂, 10 mM β-glycerophosphate, 1 mM Na-butyrate, 0.5 mM NaF, 500 mM KCl, 10% glycerol, 0.01% NP40 and 1x PI) followed by 3x 10 mL B-0.1 buffer (25 mM HEPES, pH 7.6, 1 mM EDTA, 2 mM MgCl₂, 10 mM β-glycerophosphate, 1 mM Na-butyrate, 0.5 mM NaF, 100 mM KCl, 10% glycerol, 0.01% NP40 and 1x PI). For purification of SWR1C^{Swc5Δ} where recombinant Swc5 is added back, rSwc5 would be diluted to 5 μM in 2 mL of B-0.1 and added to resin for 15 min at 4°C followed by 2x5 mL B-0.5 washes then 2x5 mL B-0.1 washes. SWR1C was eluted by nutating resin in 1 mL B-0.1 with 0.5 mg/mL recombinant 3xFlag peptide (Genscript) for 1 hour twice in series. Collected eluant was combined and concentrated using a 100 kDa cutoff Amicon Ultra-0.5 mL centrifugal filter (Millipore). After concentrating down to ~150 μL, 3x Flag peptide is removed by serial dilution with fresh B-0.1 four more times in the concentrator. The subsequently concentrated SWR1C was aliquoted, flash frozen, and stored at -80°C. SWR1C concentration was determined by SDS-PAGE using a BSA (NEB) standard titration followed by SYPRO Ruby (Thermo Fisher Scientific) staining and quantification using ImageQuant 1D gel analysis.

Recombinant Swc5

Swc5 and variants were first cloned into a pQE80L expression vector in frame with a six histidine N-Terminal tag through Gibson assembly. Plasmids were transformed into Rosetta 2 DE3 competent cells (Novagen). 1L cultures of 2x YT (Yeast extract, tryptone) were grown at 37°C to OD₆₀₀ of 0.5-0.7 before adding 0.4 mM IPTG. After adding IPTG, cells continued to shake at room temperature for 3 hours. Cells were harvested by centrifugation at 4°C and 1L of Swc5 bacterial pellet was resuspended in 20 mL lysis buffer (10 mM Imidazole, 50 mM Tris pH 8.0, 300 mM NaCl, 1 mM PMSF, 1 mM DTT) before being flash frozen in liquid nitrogen and stored at -80°C. Pellets were thawed in a 37°C water bath then fresh 0.5 mM PMSF and 1 mM Benzamidine were added. Cells were sonicated 5x 30 seconds at 15% power, incubating on ice for 1 min between sonication intervals. Lysate was then centrifuged at 14000 rpm in a JA-17 rotor at 4°C for 25 min. Whole cell extract was incubated with 800 μL of HisPur Ni-NTA resin slurry, that had been prewashed with lysis buffer, for 2 hours at 4°C on nutator. The resin and whole cell extract mixture was centrifuged at 750 rcf for 5 minutes. The supernatant was aspirated, and the resin was placed in a 25 mL gravity column and washed with 5x 10 mL wash buffer (500 mM NaCl, 10 mM Imidazole, 50 mM Tris pH 8.0). Swc5 was eluted by capping the gravity column and adding 400 μL elution buffer (500 mM NaCl, 300 mM Imidazole, 50 mM Tris pH 8.0) to resin and incubating at 4°C for 10 min then collecting the flow through. The Swc5 eluate was dialyzed against 2x 250 mL of storage buffer (50 mM NaCl, 50 mM Tris pH 7.4, 10% glycerol) then flash frozen and stored at -80°C. Concentration was determined by BSA standard on a SDS PAGE gel.

Nucleosome reconstitution

These experiments utilized a variety of nucleosomes, both in terms of histone variants, labels, and mutations. Broadly, all nucleosomes contained a tetramer of *Xenopus laevis* H3 and H4, along with *Saccharomyces cerevisiae* H2A/H2B heterodimers, with mutations or variant substitutions as appropriate, reconstituted on DNA containing the Widom 601 positioning sequence with or without linkers and/or fluorescent labels. We refer to the Widom 601 sequence as a 147bp element, though structural studies have indicated that this sequence assembles a 145bp nucleosome core particle⁷². We decided to use *Xenopus laevis* H3/H4 tetramers due to slightly increased stability of nucleosomal product in solution. These histones show high sequence conservation with *Saccharomyces cerevisiae* H3 and H4 and we have previously demonstrated there is no difference in SWR1C eviction activity on full yeast or hybrid yeast-xenopus nucleosomes⁴⁷. Furthermore, the H4 tail region that interacts with Swc5 in our cryo-EM models is virtually identical in the orthologs.

For experiments using nucleosomes with symmetrical dimers, hybrid octamers were generated as previously described and nucleosomes were reconstituted via salt dialysis deposition on a Widom 601 positioning sequence containing DNA template^{43,73}.

Symmetrical nucleosomes produced include AB/AB and AB-apm/AB-apm mutant nucleosomes used for Swc5 quenching and gel shift assays contained yH2B-S115C or xH4-Q27C labeled with Oregon GreenTM 488 Maleimide (ThermoFisher) on a ON0 DNA template⁴³. Additionally, AB/AB and AB-apm/AB-apm containing nucleosomes in which all H2A dimers contain the label Cy5 were assembled on 77N0-Cy3 and ON0-Cy3 templates.

Asymmetrical nucleosomes were produced by first reconstituting *Xenopus laevis* H3/H4 tetramers as well as *Saccharomyces cerevisiae* dimers of interest separately instead of generating octamers. Tetramers, a single dimer type, and template DNA were mixed in a 1.4:1.8:1.0 ratio followed by reconstitution via salt dialysis identically to regular nucleosome reconstitution. This ratio was found to reliably produce a mixture of hexasomes and nucleosomes, and in the case of hexasomes the dimer is known to be deposited on the “strong” side of the Widom position sequence³¹. For asymmetric nucleosomes containing a single dimer labeled with Cy5, the template DNA was varied such that the labeled dimer is deposited on the strong side of the 601 positioning sequence. Note that our previous study demonstrated that the rates of H2A.Z deposition by SWR1C are nearly identical for either the weak or the strong side of a 601 mononucleosome⁴³. Hexasomes are then purified using a Bio-Rad MiniPrep Cell apparatus, a method by which proteins are collected via fraction collector after passage through a native-PAGE tube gel (60:1 Acrylamide/Bis), which is sufficient to separate hexasomes from nucleosomes^{31,46,74}. After checking fractions on a native-PAGE gel (29:1 Acrylamide/Bis) stained with SYBR Gold, the hexasome containing fractions were pooled and concentrated, followed by buffer exchange into remodeling buffer (25 mM HEPES, pH 7.6, 0.2 mM EDTA, 5 mM MgCl₂, 70 mM KCl, 1 mM DTT) with 20% glycerol, aliquoting, and flash-freezing. The introduction of a *Saccharomyces cerevisiae* heterodimer to hexasomes in solution leads to spontaneous incorporation and formation of a nucleosome, allowing for production of asymmetric nucleosomes with known orientation.

Six types of asymmetric nucleosomes for dimer exchange assays were produced in this manner containing linker-proximal or distal H2A-Cy5/H2B (AB-Cy5) heterodimers with placement of unlabeled AB, ZB, or AB-apm dimers on the contralateral side on a 77N0-Cy3 template DNA. Additionally, a linker distal AB-apm-Cy5 and linker proximal AB nucleosome, as well as AB/AB- Cy5 and ZB/AB-Cy5 core particles (ON0 template DNA) were produced.

For fluorescence polarization and ATPase studies, unlabeled dimers were used to produce hexasomes containing linker-distal AB or AB-apm dimers on 77N0-Cy3. These hexasomes were then used to generate nucleosomes with dimers (listed here and throughout the manuscript in

linker proximal/linker distal order) AB/AB, AB/AB-apm, AB-apm/AB, or AB-apm/AB-apm, all on the 77N0-Cy3 template.

Nucleosome dimer exchange assays

FRET based dimer exchanges assays were performed as previously described⁴⁵ using an ISS PC1 spectrofluorometer, a Tecan Infinite M1000 PRO microplate reader, or a Tecan Spark microplate reader. The nucleosome remodeling reactions (50-100 μ L) were performed in remodeling buffer at room temperature with 30-60 nM SWR1C, 10 nM 77N0-Cy3 and at least one dimer-Cy5 containing nucleosome, 60-70 nM ZB dimers, and 1 mM ATP (to start reaction). Dimer exchange was observed by exciting the Cy3 fluorophore at 530nm and monitoring the Cy5 emission 670. Data for nucleosomes with both dimers labeled was fit to the two-phase decay equation (Equation 1); where Y denotes RFU, Y_i the initial RFU, Y_p the final RFU, P_f the fast phase expressed as a fraction, P_s the slow phase expressed as a fraction, K_f the rate constant of the fast phase, K_s the rate constant of the slow phase, and t is time) while data for reactions with only one dimer labeled were fit to the one-phase decay equation (Equation 2; where Y denotes RFU, Y_i the initial RFU, Y_p the final RFU, k is the rate constant, and t is time).

(Equation 1)

$$Y = (P_f(Y_i - Y_p))^{-K_f t} + (P_s(Y_i - Y_p))^{-K_s t} + Y_p$$

(Equation 2)

$$Y = (Y_i - Y_p)^{-k t} + Y_p$$

Gel based dimer exchange assays used a modified protocol from a previous study.^{15,23} A 150 μ L dimer exchange reaction contained 30 nM SWR1C, 10 nM 77N0 nucleosomes (WT or APM), and 70nM H2A.Z^{flag}/H2B in remodeling buffer with 0.1 mg/mL BSA. The reaction was performed at room temperature and 20 μ L of the reaction was taken at each time point (time 0 is before 1 mM ATP is added) and quenched with 1 μ g of plasmid DNA to separate SWR1C and dimers from the nucleosome. Each time point was stored on ice until last sample was quenched. Each time point was loaded onto a 6% 0.5x TBE Native PAGE gel (29:1 Acrylamide/Bis) and electrophoresed for 90 min at 120V. Gels were stained with SYBR gold and imaged on a GE Typhoon.

Fluorescence polarization assay

Fluorescence polarization assays were performed as previously described⁷⁵ using a CLARIOstar microplate reader. The affinity of SWR1C for various nucleosomal substrates was tested by mixing 10 nM 77N0-Cy3 nucleosomes with SWR1C serially diluted in concentration from approximately 1 μ M to 1 nM in remodeling buffer. The dimensionless value of polarization was plotted for each concentration of SWR1C and the Morrison equation (Equation 3; where Bmax is maximum observed binding, [F] is the concentration of the labeled nucleosomes, [E] is the concentration of SWR1C, and $K_{d,app}$ is the apparent dissociation constant) was used to determine the binding affinity for substrates where binding was observed.

(Equation 3)

$$Y = Bmax \left(\frac{([F] + [E] + K_{d,app}) - \sqrt{([F] + [E] + K_{d,app})^2 - 4[F][E]}}{2[F]} \right)$$

ATPase assay

ATPase activity was measured using a phosphate binding protein reporter assay. For the phosphate sensor assay,^{19,76} 50 μ L reactions were prepared with 8 nM SWR1C mixed with 0.5 μ M phosphate sensor (fluorophore-conjugated phosphate binding protein, ThermoFisher) in remodeling buffer. Reactions containing nucleosome and ZB dimers had concentrations of 20 nM and 30 nM respectively. Reactions were initiated by adding 100 μ M ATP and monitored by the

measuring the fluorescence of the MDCC fluorophore on a Tecan Spark by exciting at 430 nm and setting the emission filter to 450 nm. All reagents and the microplate were pre-treated with a “Pi Mop” to remove free phosphate as previously described^{19,76}. Fluorescence intensity was monitored over time and the linear range of intensity change was plotted to calculate the rate in reciprocal seconds.

Swc5 binding assays

Gel shifts with recombinant WT or Swc5 derivatives were performed in 15 μ L reactions. Reactions contained varying amounts of Swc5 in a total volume of 5 μ L of storage buffer (10% glycerol, 50 mM Tris pH 7.4, 50 mM NaCl), 0.25 μ L 300nM ONO nucleosomes, 6.15 μ L water, 0.6 μ L 50% glycerol, and 3 μ L 5x binding buffer (125 mM HEPES pH 7.3, 250 mM NaCl, 25 mM MgCl₂, 0.05% Tween 20, 0.5 mg/mL BSA, 5 mM DTT). Reaction incubated in the dark at room temp for 30 min. Samples were loaded onto 6% Native PAGE gels (29:1 Acrylamide/Bis) and electrophoresed at 230V for 1 hour. Gels were imaged on a GE typhoon imager and binding was calculated by measuring the disappearance of the free nucleosome band. GraphPad Prism 8 was used to generate binding curves.

Swc5 fluorescent quenching assays were assembled similar to the gel shift assays but in 40 μ L reactions with 10 nM ONO Oregon Green nucleosomes (8 μ L 5x binding buffer, 1.3 μ L 300nM nucleosomes, 11.7 μ L water, 19 μ L Swc5/storage buffer). Master mixes of each reaction incubated for 30min at room temp in the dark before being loaded into 384 well flat black plate (PerkinElmer) and scanning each well 20 times on a Tecan Spark plate reader at 488 nm excitation and 530 nm emission. Quenching was calculated by taking the average relative fluorescence units (RFU) of each Swc5 containing well and dividing RFU average of the nucleosome only wells. The normalized fluorescence was plotted using Prism 8.

Time-Resolved FRET nucleosome binding assays were performed as described by Wesley et al⁵¹. Acceptor mixtures were prepared by mixing ULIGHT alpha-6xHIS acceptor antibody (Perkin Elmer) with 6xHIS-tagged Swc5 variants at a ratio of 1:20 and serially diluting across 13 concentrations in H66 buffer (20 mM HEPES pH 7.5, 66 mM NaCl, 5 mM MgCl₂, 5 mM DTT, 5% glycerol, 0.01% NP-40, 0.01% CHAPS, and 100 μ g/mL BSA). 2x donor mixtures were prepared by mixing 4 nM streptavidin-Eu (Perkin Elmer) with or without 2 nM yeast nucleosomes containing 177 bp of Widom 601 DNA (31+145+1) with a 5' biotin group on the 31 bp extension and incubating for 30 min at room temperature. Samples were prepared in 384-well plates by mixing 5 μ L of 2x donor mixtures with 5 μ L of acceptor mixtures at each dilution. Fluorescence signals were acquired at room temperature in a Victor Nivo multimode fluorescent plate reader (Perkin Elmer) using an excitation filter at 320 nm and emission filters at 615 and 665 nm. Emission signals at 615 and 665 nm were measured simultaneously following a 100 μ s delay. Kd values were determined from triplicate titrations of each Swc5 variant and are reported as means $\pm$ standard error of the mean.

Cryo-EM sample preparation

Swc5^{79–303} was mixed with 147 bp Widom 601 sequence-containing nucleosomes at a 2.5:1 molar ratio in H50 buffer (10 mM HEPES pH 7.5, 50 mM NaCl, 1 mM DTT). Precipitated material was removed by centrifugation at 16,000 x g for 10 min. The sample was chemically cross-linked using the GraFix method in a 10-30% glycerol gradient⁷⁷, prepared with a Gradient Master instrument (Biocomp) from top (H50 + 10% v/v glycerol) and bottom (H50 + 30% v/v glycerol, 0.15% v/v glutaraldehyde) solutions. The sample was applied on top of the gradient and centrifuged in a SW40Ti rotor (Beckman-Coulter) at 35,000 rpm, 4 $^{\circ}$ C for 16 h. Fractions were collected manually by pipetting from the top, and the reaction was quenched by adding Tris pH 7.5 to each fraction to 50 mM. Fractions were analyzed using a 6% native gel (29:1 Acrylamide/Bis) stained with ethidium bromide and negative-stain EM with uranyl formate to select for monodisperse particles of the complex. Selected fractions were dialyzed into H50 buffer supplemented with 0.1 mM PMSF and concentrated to $\sim$ 1 mg/mL.

Cryo-EM data collection and analysis

Swc5/nucleosome complex was prepared at 0.9 mg/mL. Quantifoil R1.2/1.3 holey carbon, 300 mesh, copper grids (Electron Microscopy Sciences) were glow discharged for 45 s at 15 mA with an easiGlow device (PELCO). After glow discharge, 3 μ L of sample was pipetted onto the grid and Grade 595 filter paper (Ted Pella) was used to blot the grid for 3 seconds, at 4 °C and 100% humidity. The grid was plunge-frozen in liquid ethane cooled to the temperature of liquid nitrogen with a Vitrobot Mark IV instrument (Thermo Scientific). A Krios G3i microscope (Thermo Scientific) operating at 300 kV was used in conjunction with a K3 direct electron detector camera (Gatan) to collect data in super-resolution mode at the Pacific Northwest Cryo-EM Center. 13,855 movies were collected at a nominal magnification of 29,000x with a physical pixel size of 0.788 Å (super-resolution, 0.394 Å). The total dose per exposure was 50 e⁻/Å² at a dose rate of ~18.5 e⁻/pix/s and each exposure was fractionated into 50 subframes. Data was recorded in the defocus range of -0.8 to -2.8 μ m.

Data was processed with the cryoSPARC v4 suite (Supplemental Figure 6)⁷⁸. Super-resolution movies were motion-corrected with “Patch Motion Correction” and binned using 2x Fourier cropping. CTF values were estimated with “Patch CTF Estimation”. Initial particles were picked with “Blob Picker” from a subset of micrographs, then used for *ab-initio* 3D reconstruction, and the best obtained model was used to generate templates for particle picking with “Template Picker”. Template-picked particles were refined into 3 classes using “Heterogeneous Refinement” with the previously generated *ab initio* reconstructions as initial volumes to remove junk particles. 1,846,495 nucleosomal particles were further filtered to remove low-quality particles using the “random-phase 3D classification” strategy⁷⁹. Briefly, particles were sequentially subjected to multiple rounds of Heterogeneous Refinement against a low-pass filtered reconstruction (40, 30, 25, 20, 15, 10, 8 Å low-pass) of the output volume from the previous round. This strategy yielded two classes of high-quality particles with the nucleosomal DNA either 1) completely wrapped around the core (585,573 particles, Non-uniform Refinement to 3.0-Å resolution), or 2) partially peeling off (392,348 particles, 3.1 Å). The Swc5 moiety is extremely noisy in these reconstructions due to apparent structural heterogeneity. We tried employing various methods including “3D Variability Analysis”⁸⁰ to either further classify the particles into more homogenous subsets or to describe the intrinsic conformational dynamics, but with limited success (data not shown). We obtained the most informative results with cryoSPARC’s algorithm of “3D Classification” without alignment. The 585k particles of fully wrapped nucleosome were classified into 10 classes using a mask focused on the sides of the histone core in the regions with Swc5 moiety apparent. The obtained particle classes were refined to ~3.1 Å using “Non-uniform Refinement”⁸¹. Modeling was performed with UCSF ChimeraX⁸².

Data availability

The data that support this study are available from the corresponding author upon reasonable request. Source data are provided with this paper. The cryo-EM maps depicted in **Figure 6A-D** have been deposited to the Electron Microscopy Data Bank under accession codes EMD-41852, EMD-41839, EMD-41853, and EMD-41851, respectively. The raw micrographs, extracted particles, and final 3D classification subsets are available from the Electron Microscopy Public Image Archive with the accession code EMPIAR-11681.

Code availability

Not applicable

Acknowledgements

We thank Ed Luk (SUNY Stony Brook) for the kind gift of the Swr1-3xFLAG strain; Jessica Feldman for vector pQE80L, and members of the Peterson lab for helpful discussions. We thank Joseph Cho for preparing cryo-EM grids and collecting preliminary data at the Penn State cryo-EM facility, and we thank Jose M. Espinola-Lopez for assistance with the TR-FRET assays. This work was supported by the National Institutes of Health [R35-GM122519 to C.L.P.] and [R35GM127034] to S.T., and by the Estonian Research Council [PUTJD906 to P.E.]. A portion of this research was supported by NIH grant U24GM129547 and performed at the PNCC at OHSU and accessed through EMSL (grid.436923.9), a DOE Office of Science User Facility sponsored by the Office of Biological and Environmental Research. This project is funded, in part, under a grant from the Pennsylvania Department of Health using Tobacco CURE Funds. The Department specifically disclaims responsibility for any analyses, interpretations, or conclusions. Molecular graphics figures were generated using UCSF ChimeraX. UCSF ChimeraX is developed by the Resource for Biocomputing, Visualization, and Informatics at the University of California, San Francisco, with support from NIH R01-GM129325 and the Office of Cyber Infrastructure and Computational Biology, National Institute of Allergy and Infectious Diseases.

Author contributions

A.S.B and N.G. performed all experimental studies, with the exception of the cryoEM analyses which were performed by P.E. and the TR-FRET assays which were performed by E.M.L. Data analysis and manuscript preparation was performed by A.S.B, N.G., P.E., S.T., and C.L.P.

Competing interests

The authors declare no competing interests.

References

1. Luger K., Mäder A. W., Richmond R. K., Sargent D. F., Richmond T. J (1997) **Crystal structure of the nucleosome core particle at 2.8 Å resolution** *Nature* **389**:251–260
2. Clapier C. R., Iwasa J., Cairns B. R., Peterson C. L (2017) **Mechanisms of action and regulation of ATP-dependent chromatin-remodelling complexes** *Nat. Rev. Mol. Cell Biol* **18**:407–422
3. Albert I., et al. (2007) **Translational and rotational settings of H2A.Z nucleosomes across the *Saccharomyces cerevisiae* genome** *Nature* **446**:572–576
4. Barski A., et al. (2007) **High-resolution profiling of histone methylations in the human genome** *Cell* **129**:823–837
5. Hartley P. D., Madhani H. D (2009) **Mechanisms that specify promoter nucleosome location and identity** *Cell* **137**:445–458
6. Boyarchuk E., Montes de Oca R., Almouzni G. (2011) **Cell cycle dynamics of histone variants at the centromere, a model for chromosomal landmarks** *Curr. Opin. Cell Biol* **23**:266–276
7. Xu Y., et al. (2012) **Histone H2A.Z controls a critical chromatin remodeling step required for DNA double-strand break repair** *Mol. Cell* **48**:723–733
8. Meneghini M. D., Wu M., Madhani H. D (2003) **Conserved histone variant H2A.Z protects euchromatin from the ectopic spread of silent heterochromatin** *Cell* **112**:725–736
9. Guillemette B., et al. (2005) **Variant histone H2A.Z is globally localized to the promoters of inactive yeast genes and regulates nucleosome positioning** *PLoS Biol* **3**:1–11
10. Dion M. F., et al. (2007) **Dynamics of replication-independent histone turnover in budding yeast** *Science* **315**:1405–1408
11. Bryll A. R., Peterson C. L (2022) **Functional interaction between the RNA exosome and the sirtuin deacetylase Hst3 maintains transcriptional homeostasis** *Genes Dev* **36**:17–22
12. Raisner R. M., et al. (2005) **Histone variant H2A.Z marks the 5' ends of both active and inactive genes in euchromatin** *Cell* **123**:233–248
13. Creighton M. P., et al. (2008) **H2AZ is enriched at polycomb complex target genes in ES cells and is necessary for lineage commitment** *Cell* **135**:649–661
14. Kobor M. S., et al. (2004) **A protein complex containing the conserved Swi2/Snf2-related ATPase Swr1p deposits histone variant H2A.Z into euchromatin** *PLoS Biol* **2**
15. Mizuguchi G., et al. (2004) **ATP-Driven Exchange of Histone H2AZ Variant Catalyzed by SWR1 Chromatin Remodeling Complex** *Science (80-.)* **303**:343–348
16. Johnston H., Kneer J., Chackalaparampil I., Yaciuk P., Chrivia J (1999) **Identification of a novel SNF2/SWI2 protein family member, SRCAP, which interacts with CREB-binding protein** *J. Biol. Chem* **274**:16370–16376

17. Fuchs M., et al. (2001) **The p400 complex is an essential E1A transformation target** *Cell* **106**:297–307
18. Ruhl D. D., et al. (2006) **Purification of a human SRCAP complex that remodels chromatin by incorporating the histone variant H2A.Z into nucleosomes** *Biochemistry* **45**:5671–5677
19. Luk E., et al. (2010) **Stepwise histone replacement by SWR1 requires dual activation with histone H2A.Z and canonical nucleosome** *Cell* **143**:725–736
20. Wu W. H., et al. (2005) **Swc2 is a widely conserved H2AZ-binding module essential for ATP-dependent histone exchange** *Nat. Struct. Mol. Biol* **12**:1064–1071
21. Willhoft O., et al. (2018) **Structure and dynamics of the yeast SWR1-nucleosome complex** *Science* (80-.) **362**
22. Watanabe S., et al. (2015) **Structural analyses of the chromatin remodelling enzymes INO80-C and SWR-C** *Nat. Commun* **6**
23. Ranjan A., et al. (2013) **Nucleosome-free region dominates histone acetylation in targeting SWR1 to promoters for H2A.Z replacement** *Cell* **154**
24. Liang X., et al. (2016) **Structural basis of H2A.Z recognition by SRCAP chromatin-remodeling subunit YL1** *Nat. Struct. Mol. Biol* **23**:317–323
25. Lu P. Y. T., Lévesque N., Kobor M. S (2009) **NuA4 and SWR1-C: two chromatin-modifying complexes with overlapping functions and components** *Biochem. Cell Biol* **87**:799–815
26. Gómez-Zambrano Á., et al. (2018) **Arabidopsis SWC4 Binds DNA and Recruits the SWR1 Complex to Modulate Histone H2A.Z Deposition at Key Regulatory Genes** *Mol. Plant* **11**:815–832
27. Sun L., Luk E (2017) **Dual function of Swc5 in SWR remodeling ATPase activation and histone H2A eviction** *Nucleic Acids Res* **45**:9931–9946
28. Dann G. P., et al. (2017) **ISWI chromatin remodellers sense nucleosome modifications to determine substrate preference** *Nature* **548**:607–611
29. Gamarra N., Johnson S. L., Trnka M. J., Burlingame A. L., Narlikar G. J (2018) **The nucleosomal acidic patch relieves auto-inhibition by the ISWI remodeler SNF2h** *Elife* **7**
30. Dao H. T., Dul B. E., Dann G. P., Liszczak G. P., Muir T. W (2020) **A basic motif anchoring ISWI to nucleosome acidic patch regulates nucleosome spacing** *Nat. Chem. Biol* **16**:134–142
31. Levendosky R. F., Sabantsev A., Deindl S., Bowman G. D (2016) **The Chd1 chromatin remodeler shifts hexasomes unidirectionally** *Elife* **5**
32. Eustermann S., et al. (2018) **Structural basis for ATP-dependent chromatin remodelling by the INO80 complex** *Nature* **556**:386–390
33. Lehmann L. C., et al. (2020) **Mechanistic Insights into Regulation of the ALC1 Remodeler by the Nucleosome Acidic Patch** *Cell Rep* **33**
34. Makde R. D., England J. R., Yennawar H. P., Tan S (2010) **Structure of RCC1 chromatin factor bound to the nucleosome core particle** *Nature* **467**:562–566

35. McGinty R. K., Henrici R. C., Tan S (2014) **Crystal structure of the PRC1 ubiquitylation module bound to the nucleosome** *Nature* **514**:591–596
36. De Ioannes P., et al. (2019) **Structure and function of the Orc1 BAH-nucleosome complex** *Nat. Commun* **10**
37. McGinty R. K., Tan S. (2015) **Nucleosome structure and function** *Chem. Rev* **115**:2255–2273
38. McGinty R. K., Tan S (2021) **Principles of nucleosome recognition by chromatin factors and enzymes** *Curr. Opin. Struct. Biol* **71**:16–26
39. Han Y., Reyes A. A., Malik S., He Y (2020) **Cryo-EM structure of SWI/SNF complex bound to a nucleosome** *Nature* **579**:452–455
40. Wagner F. R., et al. (2020) **Structure of SWI/SNF chromatin remodeller RSC bound to a nucleosome** *Nature* **579**
41. Ye Y., et al. (2019) **Structure of the RSC complex bound to the nucleosome** *Science (80-.)* **366**
42. Eustermann S., et al. (2018) **Structural basis for ATP-dependent chromatin remodelling by the INO80 complex** *Nature* **556**:386–390
43. Lowary P. T., Widom J (1998) **New DNA sequence rules for high affinity binding to histone octamer and sequence-directed nucleosome positioning** *J. Mol. Biol* **276**:19–42
44. Fan J., Moreno A. T., Baier A. S., Loparo J. J., Peterson C. L (2022) **H2A.Z deposition by SWR1C involves multiple ATP-dependent steps** *Nat. Commun* **13**
45. Singh R. K., et al. (2019) **Transient Kinetic Analysis of SWR1C-Catalyzed H2A.Z Deposition Unravels the Impact of Nucleosome Dynamics and the Asymmetry of Histone Exchange** *Cell Rep* **27**:374–386
46. Levandosky R. F., Bowman G. D (2019) **Asymmetry between the two acidic patches dictates the direction of nucleosome sliding by the ISWI chromatin remodeler** *Elife* **8**
47. Singh R. K., et al. (2019) **Transient Kinetic Analysis of SWR1C-Catalyzed H2A.Z Deposition Unravels the Impact of Nucleosome Dynamics and the Asymmetry of Histone Exchange** *Cell Rep* **27**:374–386
48. Poyton M. F., et al. (2022) **Coordinated DNA and histone dynamics drive accurate histone H2A.Z exchange** *Sci. Adv* **8**
49. Dai L., et al. (2021) **Recognition of the inherently unstable H2A nucleosome by Swc2 is a major determinant for unidirectional H2A.Z exchange** *Cell Rep* **35**
50. Huang Y., et al. (2020) **Role of a DEF/Y motif in histone H2A-H2B recognition and nucleosome editing** *Proc. Natl. Acad. Sci. U. S. A* **117**:3543–3550
51. Wesley N. A., et al. (2022) **Time Resolved-Fluorescence Resonance Energy Transfer platform for quantitative nucleosome binding and footprinting** *Protein Sci* **31**
52. Winkler D. D., Luger K., Hieb A. R (2012) **Quantifying chromatin-associated interactions: the HI-FI system** *Methods Enzymol* **512**:243–274

53. Clapier C. R (2021) **Sophisticated Conversations between Chromatin and Chromatin Remodelers, and Dissonances in Cancer** *Int. J. Mol. Sci* **22**
54. Kalashnikova A. A., Porter-Goff M. E., Muthurajan U. M., Luger K., Hansen J. C (2013) **The role of the nucleosome acidic patch in modulating higher order chromatin structure** *J. R. Soc. Interface* **10**
55. Skrajna A., et al. (2020) **Comprehensive nucleosome interactome screen establishes fundamental principles of nucleosome binding** *Nucleic Acids Res* **48**
56. Mashtalir N., et al. (2020) **A Structural Model of the Endogenous Human BAF Complex Informs Disease Mechanisms** *Cell* **183**:802–817
57. Sen P., Ghosh S., Pugh B. F., Bartholomew B (2011) **A new, highly conserved domain in Swi2/Snf2 is required for SWI/SNF remodeling** *Nucleic Acids Res* **39**
58. Mulvihill T. S., Nelson M. L., Verma N., Jones K. B., Cairns B. R. (2021) **A system for CBAF reconstitution reveals roles for BAF47 domains and BCL7 in nucleosome ejection** *bioRxiv* <https://doi.org/10.1101/2021.10.26.465931>
59. Willhoft O., et al. (2018) **Structure and dynamics of the yeast SWR1-nucleosome complex** *Science* **362**
60. Lin C. L., et al. (2017) **Functional characterization and architecture of recombinant yeast SWR1 histone exchange complex** *Nucleic Acids Res* **45**:7249–7260
61. Wu W. H., et al. (2009) **N terminus of Swr1 binds to histone H2AZ and provides a platform for subunit assembly in the chromatin remodeling complex** *J. Biol. Chem* **284**:6200–6207
62. Ranjan A., et al. (2020) **Live-cell single particle imaging reveals the role of RNA polymerase II in histone H2A.Z eviction** *Elife* **9**
63. Altaf M., et al. (2010) **NuA4-dependent acetylation of nucleosomal histones H4 and H2A directly stimulates incorporation of H2A.Z by the SWR1 complex** *J. Biol. Chem* **285**
64. Rhee H. S., Bataille A. R., Zhang L., Pugh B. F (2014) **Subnucleosomal structures and nucleosome asymmetry across a genome** *Cell* **159**:1377–1388
65. Sundaramoorthy R., et al. (2018) **Structure of the chromatin remodelling enzyme Chd1 bound to a ubiquitylated nucleosome** *Elife* **7**
66. Messina G., Attarrato M. T., Dimitri P (2016) **When chromatin organisation floats astray: the Srcap gene and Floating-Harbor syndrome** *J. Med. Genet* **53**:793–797
67. Celauro E., Carra S., Rodriguez A., Cotelli F., Dimitri P (2017) **Functional analysis of the cfdp1 gene in zebrafish provides evidence for its crucial role in craniofacial development and osteogenesis** *Exp. Cell Res* **361**:236–245
68. Goldstein A. L., McCusker J. H (1999) **Three new dominant drug resistance cassettes for gene disruption in *Saccharomyces cerevisiae*** *Yeast* **15**:1541–1553
69. Gibson D. G., et al. (2009) **Enzymatic assembly of DNA molecules up to several hundred kilobases** *Nat. Methods* **6**:343–345

70. Luger K., Rechsteiner T. J., Richmond T. J (1999) **Expression and purification of recombinant histones and nucleosome reconstitution** *Methods Mol. Biol* **119**:1–16
71. Zhou C. Y., Narlikar G. J (2016) **Analysis of Nucleosome Sliding by ATP-Dependent Chromatin Remodeling Enzymes** *Methods Enzymol* **573**:119–135
72. Vasudevan D., Chua E. Y. D., Davey C. A (2010) **Crystal Structures of Nucleosome Core Particles Containing the ‘601’ Strong Positioning Sequence** *J. Mol. Biol* **403**:1–10
73. Luger K., Rechsteiner T. J., Richmond T. J (1999) **Preparation of nucleosome core particle from recombinant histones** *Methods Enzymol* **304**:3–19
74. Nodelman I. M., Patel A., Levendosky R. F., Bowman G. D (2020) **Reconstitution and purification of nucleosomes with recombinant histones and purified DNA** *Curr. Protoc. Mol. Biol* **133**
75. Baier A. S., Peterson C. L (2022) **Fluorescence approaches for biochemical analysis of ATP-dependent chromatin remodeling enzymes** *Methods Enzymol* **673**:1–17
76. Brune M., Hunter J. L., Corrie J. E. T., Webb M. R (1994) **Direct, real-time measurement of rapid inorganic phosphate release using a novel fluorescent probe and its application to actomyosin subfragment 1 ATPase** *Biochemistry* **33**:8262–8271
77. Stark H (2010) **GraFix: Stabilization of Fragile Macromolecular Complexes for Single Particle Cryo-EM** *Methods Enzymol* **481**:109–126
78. Punjani A., Rubinstein J. L., Fleet D. J., Brubaker M. A (2017) **CryoSPARC: Algorithms for rapid unsupervised cryo-EM structure determination** *Nat. Methods* **14**
79. Gong X., et al. (2016) **Structural insights into the Niemann-Pick C1 (NPC1)-mediated cholesterol transfer and ebola infection** *Cell* **165**
80. Punjani A., Fleet D. J (2021) **3D variability analysis: Resolving continuous flexibility and discrete heterogeneity from single particle cryo-EM** *J. Struct. Biol* **213**
81. Punjani A., Zhang H., Fleet D. J (2020) **Non-uniform refinement: adaptive regularization improves single-particle cryo-EM reconstruction** *Nat. Methods* **17**
82. Pettersen E. F., et al. (2021) **UCSF ChimeraX: Structure visualization for researchers, educators, and developers** *Protein Sci* **30**

Article and author information

Alexander S. Baier

Program in Molecular Medicine, University of Massachusetts Chan Medical School, Worcester, MA, 01605, Medical Scientist Training Program, T.H. Chan School of Medicine, University of Massachusetts

Nathan Gioacchini

Program in Molecular Medicine, University of Massachusetts Chan Medical School, Worcester, MA, 01605, Interdisciplinary Graduate Program, Morningside Graduate School of Biomedical Sciences, University of Massachusetts Chan Medical School, Worcester, MA, 01605

Priit Eek

Center for Eukaryotic Gene Regulation, Department of Biochemistry and Molecular Biology, Pennsylvania State University, University Park, PA 16802, USA, Department of Chemistry and Biotechnology, Tallinn University of Technology, 12618 Tallinn, Estonia

Erik M. Leith

Center for Eukaryotic Gene Regulation, Department of Biochemistry and Molecular Biology, Pennsylvania State University, University Park, PA 16802, USA

Song Tan

Center for Eukaryotic Gene Regulation, Department of Biochemistry and Molecular Biology, Pennsylvania State University, University Park, PA 16802, USA

Craig L. Peterson

Program in Molecular Medicine, University of Massachusetts Chan Medical School, Worcester, MA, 01605

For correspondence: craig.peterson@umassmed.edu

Copyright

© 2024, Baier et al.

This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Yamini Dalal

National Cancer Institute, Bethesda, United States of America

Senior Editor

Amy Andreotti

Iowa State University, Ames, United States of America

Reviewer #1 (Public Review):

This manuscript presents an extremely exciting and very timely analysis of the role that the nucleosome acidic patch plays in SWR1-catalyzed histone exchange. Intriguingly, SWR1 loses activity almost completely if any of the acidic patches are absent. To my knowledge, this makes SWR1 the first remodeler with such a unique and pronounced requirement for the acidic patch. The authors demonstrate that SWR1 affinity is dramatically reduced if at least one of the acidic patches is absent, pointing to a key role of the acidic patch in SWR1 binding to the nucleosome. The authors also pinpoint a specific subunit - Swc5 - that can bind nucleosomes and engage the acidic patch and obtain a cryo-EM structure of Swc5 bound to a nucleosome. They also identify a conserved arginine-rich motif in this subunit that is critical for nucleosome binding and histone exchange *in vitro* and for SWR1 function *in vivo*. The authors provide evidence that suggests a direct interaction between this motif and the acidic patch.

Strengths:

The manuscript is well-written and the experimental data are of outstanding quality and importance for the field. This manuscript significantly expands our understanding of the fundamentally important and complex process of H2A.Z deposition by SWR1 and would be of great interest for a broad readership.

<https://doi.org/10.7554/eLife.94869.2.sa0>

Author response:

The following is the authors' response to the original reviews.

The reviewer comments have been helpful, and we have revised the manuscript to address the concerns of reviewer 2. In addition to text changes, we also added a negative control to Figure 1 to address concerns about photobleaching or DNA unwrapping.

Reviewer #1:

This manuscript presents an extremely exciting and very timely analysis of the role that the nucleosome acidic patch plays in SWR1-catalyzed histone exchange. Intriguingly, SWR1 loses activity almost completely if any of the acidic patches are absent. To my knowledge, this makes SWR1 the first remodeler with such a unique and pronounced requirement for the acidic patch. The authors demonstrate that SWR1 affinity is dramatically reduced if at least one of the acidic patches is absent, pointing to a key role of the acidic patch in SWR1 binding to the nucleosome. The authors also pinpoint a specific subunit - Swc5 - that can bind nucleosomes, engage the acidic patch, and obtain a cryo-EM structure of Swc5 bound to a nucleosome. They also identify a conserved arginine-rich motif in this subunit that is critical for nucleosome binding and histone exchange in vitro and for SWR1 function in vivo. The authors provide evidence that suggests a direct interaction between this motif and the acidic patch.

Strengths:

The manuscript is well-written and the experimental data are of outstanding quality and importance for the field. This manuscript significantly expands our understanding of the fundamentally important and complex process of H2A.Z deposition by SWR1 and would be of great interest to a broad readership.

We thank the reviewer for their enthusiastic and positive comments on our work.

Reviewer #2:

Summary:

In this study, Baier et al. investigated the mechanism by which SWR1C recognizes nucleosomal substrates for the deposition of H2A.Z. Their data convincingly demonstrate that the nucleosome's acidic patch plays a crucial role in the substrate recognition by SWR1C. The authors presented clear evidence showing that Swc5 is a pivotal subunit involved in the interaction between SWR1C and the acidic patch. They pared down the specific region within Swc5 responsible for this interaction. However, two central assertions of the paper are less convincing. First, the data supporting the claim that the insertion of one Z-B dimer into the canonical nucleosome can stimulate SWR1C to insert the second Z-B dimer is somewhat questionable (see below). Given that this claim contradicts previous observations made by other groups, this hypothesis needs further testing to eliminate potential artifacts. Secondly, the claim that SWR1C simultaneously recognizes the acidic patch on both sides of the nucleosome also needs further

investigation, as the assay used to establish this claim lacks the sensitivity necessary to distinguish any difference between nucleosomal substrates containing one or two intact acidic patches.

Strengths:

As mentioned in the summary, the authors presented clear evidence demonstrating the role of Swc5 in recognition of the nucleosome acidic patch. The identification of the specific region in Swc5 responsible for this interaction is important.

We thank the reviewer for their careful critique of our work. Below we address each major concern.

Major comments: (1) Figure 1B: It is unclear how much of the decrease in FRET is caused by the bleaching of fluorophores. The authors should include a negative control in which Z-B dimers are omitted from the reaction. In the absence of ZB dimers, SWR1C will not exchange histones. Therefore, any decrease in FRET should represent the bleaching of fluorophores on the nucleosomal substrate, allowing normalization of the FRET signal related to A-B eviction.

In this manuscript, as well as in our two previous publications (Singh et al., 2019; Fan et al., 2022), we have presented the results of no enzyme controls, +/- ZB dimers, no ATP controls, or AMP-PNP controls for our FRET-based, H2A.Z deposition assay (see also Figure S3). We do not observe significant levels of photobleaching in this assay, either during ensemble measurements or in an smFRET experiment. To aid the reader, we have added the AMP-PNP data for the experiment shown in Figure 1B. The results show there is less than a 10% decrease in FRET over 30', and the signal from the double acidic patch disrupted nucleosome is identical to this negative control.

(2) Figure S3: The authors use the decrease in FRET signal as a metric of histone eviction. However, Figure S3 suggests that the FRET signal decrease could be due to DNA unwrapping. Histone exchange should not occur when SWR1C is incubated with AMP-PNP, as histone exchange requires ATP hydrolysis (10.7554/eLife.77352). And since the insertion of Z-B dimer and the eviction of A-B dimer are coupled, the decrease of FRET in the presence of AMP-PNP is unlikely due to histone eviction or exchange. Instead, the FRET decrease is likely due to DNA unwrapping (10.7554/eLife.77352). The authors should explicitly state what the loss of FRET means.

We agree with the reviewer, that loss of FRET can be due to DNA unwrapping from the nucleosome. We have previously demonstrated this activity by SWR1C in our smFRET study (Fan et al., 2022). However, DNA unwrapping is highly reversible and has a time duration of only 1-3 seconds. We and others have not observed stable unwrapping of nucleosomes by SWR1C, but rather the stable loss of FRET reports on dimer eviction. We assume the reviewer is concerned about the rather large decrease in FRET signal shown in the AMP-PNP controls for Figure S3, panels A and D. For the other 7 panels, the decrease in FRET with AMP-PNP are minimal. In fact, if we average all of the AMP-PNP data points, the rate of FRET loss is not statistically different from no enzyme control reactions (nucleosome plus ZB dimers).

Data for panels A and D used a 77NO nucleosomal substrate, with Cy3 labeling the linker distal dimer. This is our standard DNA fragment, and it was used in Figure 1B. The only difference between data sets is that the data shown in Fig 1B used nucleosome reconstituted with a Cy5-labelled histone octamer, rather than the hexasome assembly method used for Fig S3. Three points are important. First, for all of these substrates, we assembled 3 independent nucleosomes, and the results are highly reproducible. Two, we performed a total of 6 experiments for the 77NO-Cy5 substrates to ensure that the rates were accurate (+/-ATP).

Third, and most important, we do not see this decrease in FRET signal in the absence of SWR1C (no enzyme control). This data was included in the data source file. Thus, it appears that there is significant SWR1C-induced nucleosome instability for these two hexasome-assembled substrates. We now note this in the legend to Figure S3. Key for this work, however, is that there is a large increase in the rate of FRET loss in the presence of ATP, and this rate is faster when a ZB dimer was present at the linker proximal location. In response to the last point, we state in the first paragraph of the results: “The dimer exchange activity of SWR1C is monitored by following the decrease in the 670 nm FRET signal due to eviction of the Cy5-labeled AB-Cy5 dimer (Figure 1A).”

(3) Related to point 2. One way to distinguish nucleosomal DNA unwrapping from histone dimer eviction is that unwrapping is reversible, whereas A-B eviction is not. Therefore, if the authors remove AMP-PNP from the reaction chamber and a FRET signal reappears, then the initial loss of FRET was due to reversible DNA unwrapping. However, if the removal of AMP-PNP did not regain FRET, it means that the loss of FRET was likely due to A-B eviction. The authors should perform an AMP-PNP and/or ATP removal experiment to make sure the interpretation of the data is correct.

See response to item 2 above

(4) The nature of the error bars in Figure 1C is undefined; therefore, the statistical significance of the data is not interpretable.

We apologize for not making this more explicit for each figure. The error bars report on 95% confidence intervals from at least 3 sets of experiments. This statement has been added to the legend.

*(5) The authors claim that the SWR1C requires intact acidic patches on both sides of the nucleosomes to exchange histone. This claim was based on the experiment in Figure 1C where they showed mutation of one of two acidic patches in the nucleosomal substrate is sufficient to inhibit SWR1C-mediated histone exchange activity. However, one could argue that the sensitivity of this assay is too low to distinguish any difference between nucleosomes with one (i.e., AB/AB-*apm*) versus two mutated acidic patches (i.e., AB-*apm*/AB-*apm*). The lack of sensitivity of the eviction assay can be seen when Figure 1B is taken into consideration. In the gel-shift assay, the AB-*apm*/AB-*apm* nucleosome exhibited a 10% SWR1C-mediated histone exchange activity compared to WT. However, in the eviction assay, the single AB/AB-*apm* mutant has no detectable activity. Therefore, to test their hypothesis, the authors should use the more sensitive in-gel histone exchange assay to see if the single AB/AB-*apm* mutant is more or equally active compared to the double AB-*apm*/AB-*apm* mutant.*

Our pincher model is based on three, independent sets of data, not just Figure 1C. First, as noted by the reviewer, we find that disruption of either acidic patch cripples the dimer exchange activity of SWR1C in the FRET-based assay. Whether the defect is identical to that of the double APM mutant nucleosome does not seem pertinent to the model. In a second set of assays, we used fluorescence polarization to quantify the binding affinity of SWR1C for wildtype nucleosomes, a double APM nucleosome, or each single APM nucleosome. Consistent with the pincher model, each single APM disruption decreases binding affinity at least 10-fold (below the sensitivity of the assay). Finally, we monitored the ability of different nucleosomes to stimulate the ATPase activity of SWR1C. Consistent with the pincher model, a single APM disruption was sufficient to eliminate nucleosome stimulation.

(6) The authors claim that the AZ nucleosome is a better substrate than the AA nucleosome. This is a surprising result as previous studies showed that the two insertion

steps of the two Z-B dimers are not cooperative (10.7554/eLife.77352 and 10.1016/J.CELREP.2019.12.006). The authors' claim was based on the eviction assay shown in Fig 1C. However, I am not sure how much variation in the eviction assay is contributed by different preparations of nucleosomes. The authors should use the in-gel assay to independently test this hypothesis.

For all data shown in our manuscript, at least three different nucleosome preparations were used. The impact of a ZB dimer on the rates of dimer exchange was highly reproducible among different nucleosome preparations and experiments. We also see reproducible ZB stimulation for three different substrates – with ZB on the linker proximal side, the linker distal side, and on one side of a core particle. We do not believe that our data are inconsistent with previous studies. First, the previous work referenced by the reviewer performed dimer exchange reactions with a large excess of nucleosomes to SWR1C (catalytic conditions), whereas we used single turnover reactions. Secondly, our study is the first to use a homogenous, ZA heterotypic nucleosome as a substrate for SWR1C. All previous studies used a standard AA nucleosome, following the first and second rounds of dimer exchange that occur sequentially. And finally, we observe only a 20-30% increase in rate by a ZB dimer (e.g. 77N0 substrates), and such an increase was unlikely to have been detected by previous gel-based assays.

Minor comments:

(1) Abstract line 4: To say 'Numerous' studies have shown acidic patch impact chromatin remodeling enzymes activity may be too strong.

Removed

(2) Page 15, line 15: The authors claim that swc5Δ was inviable on formamide media. However, the data in Figure 8 shows cell growth in column 1 of swc5Δ.

The term 'inviable' has been replaced with 'poor' or 'slow growth'

(3) The authors should use standard yeast nomenclature when describing yeast genes and proteins. For example, for Figure 8 and legend, Swc5Δ was used to describe the yeast strain BY4741; MATa; his3Δ1; leu2Δ0; met15Δ0; ura3Δ0; YBR231c::kanMX4. Instead, the authors should describe the swc5Δ mutant strain as BY4741 MAT a his3Δ1 leu2Δ0 met15Δ0 ura3Δ0 swc5Δ::kanMX4. Exogenous plasmid should also be indicated in italics and inside brackets, such as [SWC5-URA3] or [swc5(R219A)-URA3].

We apologize for missing this mistake in the Figure 8 legend. We had inadvertently copied this from the euroscarf entry and forgot to edit the entry. We decided not to add all the plasmid names to the figure, as it was too cluttered. We state in the figure legend that the panels show growth of swc5 deletion strains harboring the indicated swc5 alleles on CEN/ARS plasmids.

(4) According to Lin et al. 2017 NAR (doi: 10.1093/nar/gkx414), there is only one Swc5 subunit per SWR1C. Therefore, the pincher model proposed by the authors would suggest that there is a missing subunit that recognizes the second acidic patch. The authors should point out this fact in the discussion. However, as mentioned in Major comment 6, I am not sure if the pincher model is substantiated.

In our discussion, we had noted that the published cryoEM structure had suggested that the Swc2 subunit likely interacts with the acidic patch on the dimer that is not targeted for

replacement, and we proposed that Swc5 interacts with the acidic patch on the exchanging H2A/H2B dimer. We have now made this more clear in the text.