

Metabolic disruption impairs ribosomal protein levels, resulting in enhanced aminoglycoside tolerance

Reviewed Preprint

v2 • June 5, 2024

Revised by authors

Reviewed Preprint

v1 • February 14, 2024

Rauf Shiraliyev, Mehmet A. Orman 

William A. Brookshire Department of Chemical and Biomolecular Engineering, University of Houston, Houston, TX, 77204

 https://en.wikipedia.org/wiki/Open_access
 Copyright information

Abstract

Aminoglycoside antibiotics display broad-spectrum activity against Gram-negative and Gram-positive bacteria by targeting their ribosomes. Herein, we have demonstrated that energy metabolism plays a crucial role in aminoglycoside tolerance, as knockout strains associated with the tricarboxylic acid cycle (TCA) and the electron transport chain (ETC) exhibited increased tolerance to aminoglycosides in the mid-exponential growth phase of *Escherichia coli* cells. Given that aminoglycoside uptake relies on the energy-driven electrochemical potential across the cytoplasmic membrane, our initial expectation was that these genetic perturbations would decrease the proton motive force (PMF), subsequently affecting the uptake of aminoglycosides. However, our results did not corroborate this assumption. We found no consistent metabolic changes, ATP levels, cytoplasmic pH variations, or membrane potential differences in the mutant strains compared to the wild type. Additionally, intracellular concentrations of fluorophore-labeled gentamicin remained similar across all strains. To uncover the mechanism responsible for the observed tolerance in mutant strains, we employed untargeted mass spectrometry to quantify the proteins within these mutants and subsequently compared them to their wild-type counterparts. Our comprehensive analysis, which encompassed protein-protein association networks and functional enrichment, unveiled a noteworthy upregulation of proteins linked to the TCA cycle in the mutant strains during the mid-exponential growth phase, suggesting that these strains compensate for the perturbation in their energy metabolism by increasing TCA cycle activity to maintain their membrane potential and ATP levels. Furthermore, our pathway enrichment analysis shed light on local network clusters displaying downregulation across all mutant strains, which were associated with both large and small ribosomal binding proteins, ribosome biogenesis, translation factor activity, and the biosynthesis of ribonucleoside monophosphates. These findings offer a plausible explanation for the observed tolerance of aminoglycosides in the mutant strains. Altogether, this research has the potential to uncover mechanisms behind aminoglycoside tolerance, paving the way for novel strategies to combat such cells.

eLife assessment

The findings of this study are **valuable** as they challenge the dogma regarding the link between lowered bacterial metabolism and tolerance to aminoglycosides. The authors propose that the well-known tolerance to AG of mutants such as those of complexes I and II is not due to a decrease in the proton motive force and thus antibiotic uptake. The results presented here are **convincing**.

<https://doi.org/10.7554/eLife.94903.2.sa1>

Introduction

Global public health is currently confronted with a pressing challenge in the form of antimicrobial resistance (Binder et al., 1999). Addressing this issue necessitates a comprehensive understanding of the diverse mechanisms used by bacterial cells to survive antibiotic treatments. These mechanisms can be broadly classified into two groups: reversible and irreversible. Reversible mechanisms, often termed "tolerance" mechanisms, are not genetically inherited and encompass alterations in bacterial growth and behavior. Within this category, antibiotic-tolerant cells, such as persisters, viable but nonculturable cells, and stationary-phase cells, transiently survive high antibiotic concentrations. Their formation is linked to stochastic and/or deterministic processes triggered by various stress factors including the SOS response, stringent response, reactive oxygen species, nutrient depletion, and overpopulation (Amato et al., 2013; Germain et al., 2015; Helaine and Kugelberg, 2014; Molina-Quiroz et al., 2018; Theodore et al., 2013; Van den Bergh et al., 2017; Wood et al., 2013). The reversible nature of these cells allows them to switch between antibiotic-tolerant and sensitive states. On the other hand, irreversible mechanisms, often termed "resistance" mechanisms, are heritable and related to mutagenic processes (Balaban et al., 2019). Antibiotic-resistant mutants, falling into this category, can emerge due to mutations in antibiotic target proteins or proteins that contribute to enhanced repair mechanisms, cell dormancy, drug efflux systems, and alternative mechanisms that circumvent the antibiotic target (C Reygaert, 2018; Darby et al., 2023; Lambert, 2005; Munita and Arias, 2016; Okusu et al., 1996).

Aminoglycosides were among the earliest antibiotics used in clinical practice (Becker and Cooper, 2013; Krause et al., 2016). These antibiotics are a class of naturally occurring or semisynthetic amino-modified sugars known for their broad-spectrum activity against diverse bacterial species (Aggen et al., 2010; Endimiani et al., 2009; Ikäheimo I et al., 2000; Karlowsky et al., 2003; Landman et al., 2010; Ristuccia and Cunha, 1985; Sader et al., 2015). Since the discovery of the first aminoglycoside antibiotic, streptomycin, which was isolated from *Streptomyces griseus* (Kresge et al., 2004; Schatz et al., 2005; Woodruff, 2014), numerous other aminoglycosides have been identified, including gentamicin, kanamycin, tobramycin, amikacin, and neomycin (Becker and Cooper, 2013; Krause et al., 2016). Although the use of aminoglycosides declined for a brief period with the advent of newer antibiotics (such as fluoroquinolones) which were thought to have lower toxicity (Becker and Cooper, 2013; Krause et al., 2016), the rise of resistance to these new drugs has revived interest in aminoglycosides and the development of new ones with improved dosing schemes (Becker and Cooper, 2013; Krause et al., 2016).

Studies on the mechanism of action of aminoglycosides against bacteria have shown that they disrupt protein synthesis by targeting the ribosome. The primary action of aminoglycosides is to bind to the 16S ribosomal RNA (rRNA) of bacteria, a component of the 30S ribosomal subunit (Moazed and Noller, 1987; Recht et al., 1999; Woodcock et al., 1991). The initial entry of

antibiotics into cells causes misreading in protein synthesis as the antibiotics interact with the ribosomes involved in chain elongation. Then, these misread proteins or polypeptides are incorporated into the membrane, causing membrane damage (Davis et al., 1986 [DOI](#)). This increases the amount of antibiotics entering the cells, leading to more misreading, and the formation of more channels, and eventually the complete inhibition of protein synthesis (Davis et al., 1986 [DOI](#)). Moreover, aminoglycosides potentially inhibit ribosome recycling by binding to RNA helix 69 (H69) of the 50S ribosomal subunit, which can also lead to the inhibition of mRNA and tRNA translocation (Borovinskaya et al., 2007 [DOI](#)).

It has been demonstrated that antibiotic-tolerant cells can become susceptible to aminoglycosides by metabolizing certain carbon sources (Allison et al., 2011 [DOI](#)). This susceptibility arises from an enhanced drug uptake as a result of an increase in the electron transport chain activity and membrane potential, facilitated by the breakdown of these specific carbon sources (Allison et al., 2011 [DOI](#)). The process of aminoglycoside uptake is a unique, energy-requiring mechanism where the electrochemical potential across the cytoplasmic membrane and electron flow through membrane-bound respiratory chains are believed to be significant factors (Taber Harry W et al., 1987 [DOI](#)). However, the bactericidal effect of aminoglycosides may not result from the downstream impacts of voltage-dependent drug uptake, but rather from an irregular membrane potential (Bruni and Kralj, 2020 [DOI](#)). Bruni and Kralj suggested that hyperpolarization, stemming from changes in ATP flux due to the reversal of F1Fo-ATPase activity, could potentially intensify aminoglycoside-mediated cell death (Bruni and Kralj, 2020 [DOI](#)).

Previous studies underscore the potential role of metabolic mutations in aminoglycoside tolerance and resistance, a phenomenon that warrants further investigation (Muir et al., 1981 [DOI](#); Shan et al., 2015 [DOI](#)). A recent study, which analyzed genomic alterations in *E. coli* strains, including uropathogenic UTI89 strains, following daily antibiotic exposure (Van den Bergh et al., 2022 [DOI](#)), showed that mutations were predominantly detected in genes of the *nuo* operon, a vital component of bacterial energy metabolism. This highlights a potential link between metabolic adaptations and antibiotic tolerance. Additionally, research led by Collin's group uncovered genes related to central metabolism that contribute to antibiotic resistance in *E. coli* cells exposed to various antibiotics, including aminoglycosides (Lopatkin et al., 2021 [DOI](#)). These findings are corroborated by similar mutations identified in clinical *E. coli* pathogens, as evidenced by the examination of a comprehensive library of 7243 *E. coli* genomes from NCBI Pathogen Detection (Lopatkin et al., 2021 [DOI](#)). In our study, we found that knockout strains with genes related to the tricarboxylic acid cycle (TCA) and the electron transport chain (ETC) displayed increased tolerance to aminoglycosides. Intriguingly, this increased tolerance was not attributed to reduced proton motive force, which affects drug uptake, as evidenced by insignificant alterations in ATP levels or membrane potential in the mutant strains compared to the wild-type strain. We employed untargeted mass spectrometry to quantify proteins in the mutant and wild-type strains, revealing a notable upregulation of proteins associated with the TCA cycle in the mutants. This suggests that these strains compensate for the disruption in their energy metabolism by altering TCA cycle activity to maintain their membrane potential and ATP levels. Moreover, our pathway enrichment analysis underlined local network clusters that were consistently downregulated across all mutant strains. These clusters were related to both large and small ribosomal binding proteins, ribosome biogenesis, translation factor activity, and the biosynthesis of ribonucleoside monophosphates. These findings provide a credible rationale for the observed tolerance to aminoglycosides in the mutant strains.

Results

Deletions of the TCA cycle and ETC genes increased tolerance to aminoglycosides

Given the crucial role of energy metabolism in aminoglycoside tolerance, our initial objective was to assess various knockout strains that involved the deletion of genes associated with the TCA cycle, such as *sucA*, *gltA*, *mdh*, *sdhC*, *icd*, *acnB*, and *fumA*, as well as the NADH dehydrogenase enzyme of ETC, including *nuoM* and *nuoI* from *E. coli* MG1655. To conduct the experiments, both wild-type and mutant strains were cultured overnight and then diluted 100-fold in fresh 2 milliliters (ml) Lysogeny Broth (LB) in test tubes. The cultures were grown until the mid-exponential phase ($t=3.5$ hours) in a shaker at 37°C and 250 rpm. Subsequently, the cultures were exposed to various aminoglycosides (50 µg/ml streptomycin, 50 µg/ml gentamicin, and 50 µg/ml amikacin) for a duration of 5 hours. Samples were collected before and after the treatments and plated on LB agar to quantify the surviving cell fractions. Analysis of the results indicates that most of the knockout strains (Δ *sucA*, Δ *gltA*, Δ *mdh*, Δ *sdhC*, Δ *nuoM*, and Δ *nuoI*) exhibited increased tolerance to streptomycin, gentamicin, and amikacin treatments when compared to the wild-type strain (Figure 1A-C). However, it is worth noting that mutant strains such as Δ *icd*, Δ *acnB*, and Δ *fumA* did not always exhibit increased tolerance (Figure 1A-C), underscoring the complex interplay between energy metabolism and specific antibiotic tolerance, as well as the pleiotropic effects of the gene deletions. Moreover, the surviving cells measured in these assays may not necessarily represent antibiotic-resistant cells, as the antibiotic tolerance assays were conducted at concentrations well above the Minimum Inhibitory Concentration (MIC) levels. The MIC levels of the strains for the tested antibiotics displayed no drastic alterations compared to those of the wild type, despite some minor variations among them (Supplementary Table S1).

To gain a thorough understanding of the observed tolerance, we chose four mutant strains that exhibit either high tolerance (*sucA*, *gltA*, and *nuoI*) or low tolerance (*icd*) to gentamicin for the subsequent assays. Our initial objective was to delineate the time-dependent profiles of antibiotic-tolerant cells within cultures. To achieve this, cells from overnight cultures were diluted in fresh media and cultured for 6 hours in LB in flasks (see Materials and Methods). Hourly samples were collected for cell quantification and antibiotic tolerance assays. We opted for flow cytometry for precise cell quantification, a more reliable method than optical density measurements (Mohiuddin et al., 2020) (Figure 2A). In the antibiotic tolerance assays, samples were treated with gentamicin (50 µg/ml) for 5 hours and plated before and after the treatments (Figure 2B). Notably, the observed tolerance in these mutant strains is not linked to cell growth. For instance, the mutant strains associated with *sucA*, *gltA*, and *nuoI* genes exhibited higher tolerance to aminoglycosides although they did not show significantly altered cell growth compared to the wild type (Figure 2A, B). These mutant strains displayed a substantial increase in the number of tolerant cells (more than 10^4 -fold) between time points $t=3$ and $t=4$ hours during the mid-exponential growth phase when compared to the wild-type strain (Figure 2B). Conversely, while the *icd* mutant exhibited a reduction in cell growth, it was observed to be as sensitive to aminoglycosides as the wild type during the exponential growth phase and the formation of tolerant cells in this mutant strain was only evident after the time point $t=5$ hours (Figure 2A, B). We emphasize that the observed tolerance in the mutant strains is transient or reversible, as they all exhibit high sensitivity to aminoglycosides, similar to the wild type, during the lag phase of growth (Figure 2B). Given that antibiotic concentrations may influence the number of surviving cells, we also treated cells with 5 µg/ml gentamicin at the mid-exponential phase. Although we observed a greater number of surviving cells in all strains when lower gentamicin concentrations were employed compared to higher concentrations, we still observed a similar trend; Δ *sucA*, Δ *gltA*, and Δ *nuoI* exhibited higher levels of surviving cells compared to Δ *icd* and the wild-type strain (Supplementary Figure S1).

Figure 1.

Deletions of the TCA cycle and ETC genes enhanced tolerance to aminoglycosides.

In the mid-exponential phase ($t=3.5$ h), *E. coli* MG1655 wild-type and knockout strains underwent (A) streptomycin, (B) gentamicin, and (C) amikacin treatments at a concentration of 50 $\mu\text{g/ml}$ for a duration of 5 hours in test tubes. Following the treatments, cells were washed to eliminate the antibiotics and then plated on LB agar plates to quantify the surviving cell fractions. CFU: Colony forming units. WT: Wild type. For pairwise comparisons, one-way ANOVA with Dunnett's post hoc test was used where *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$, and ****, $P < 0.0001$. $N=3$. Data points represent mean and standard error.





Figure 2.

The observed tolerance in the mutant strains is not linked to cell growth.

(A) Growth of *E. coli* MG1655 wild type, ΔsucA , ΔgltA , ΔnuoI , and Δicd strains was assessed by measuring the number of cells per ml with flow cytometry. (B) The cells of both the *E. coli* MG1655 WT and mutant strains were collected from culture flasks (see Materials and Methods for details) at indicated time intervals and then subjected to gentamicin treatment. The figure shows the colony-forming unit (CFU) levels of both the treated and untreated cultures, indicating the surviving and total cell populations, respectively. $N=3$. Data points represent mean and standard error.



Energy-dependent aminoglycoside uptake is not a contributing factor

Aminoglycoside uptake is a unique and energy-requiring mechanism that depends on the electrochemical potential across the cytoplasmic membrane (Taber Harry W et al., 1987 [\[4\]](#)). Initially, we hypothesized that the genetic perturbations in these strains may have decreased proton motive force (PMF) and aminoglycoside uptake. The mutant strains exhibiting increased aminoglycoside tolerance demonstrated no consistent pattern in metabolic activities during the mid-exponential growth phase, which was assessed using redox sensor green (RSG) dye for the indicated time points (t=3, 4, 5 and 6 hours) of the cell growth (**Figure 3A** [\[4\]](#)). When RSG molecules are reduced by bacterial reductases, critical components of energy metabolism, these molecules emit green fluorescence. This fluorescence signal should be suppressed by the presence of a metabolic inhibitor, such as carbonyl cyanide 3-chlorophenylhydrazone (CCCP) (**Supplementary Figure S2**). Given that ATP is a pivotal product of PMF, we also quantified intracellular ATP levels in both wildtype and mutant strains, employing the BacTiter-Glo™ Microbial Cell Viability Assay (Catalog# G8230, Promega Corporation, Madison WI). This assay utilizes a single reagent to lyse cells and produce luminescence through the luciferase reaction, with the luminescent signal being directly proportional to the ATP content (**Supplementary Figure S3**). We found no consistent pattern in ATP levels between the antibiotic-sensitive and tolerant strains, particularly during the mid-exponential growth phase (at t=3 and 4 hours) (**Figure 3B** [\[4\]](#)). While the *ΔsucA* mutant generally showed reduced ATP levels, this was not the case for the other strains, which showed increased ATP levels around t=6 hours (**Figure 3B** [\[4\]](#)).

Furthermore, since the proton gradient is a vital element of PMF and a reduced H⁺ ion gradient across the cell membrane is linked to reduced membrane potential, we conducted measurements of intracellular pH using the ratiometric pHluorin, known as a pH-sensitive derivative of green fluorescent protein (GFP) (Miesenböck et al., 1998 [\[4\]](#)). This GFP variant exhibits a bimodal excitation spectrum characterized by peaks at 410 nm and 470 nm, along with an emission maximum at 530 nm (Martinez II et al., 2012 [\[4\]](#)). When subjected to acidification, the excitation at 410 nm diminishes while the excitation at 470 nm concurrently rises, which allows us to construct standard curves for the measurement of intracellular pH (**Supplementary Figure S4**). For these assessments, we introduced pGFPR01 plasmids, where pHluorin is expressed under the arabinose-induced promoter, into the mutant strains utilized. Our results revealed that there was no anticipated acidification of the cytoplasm in the TCA and ETC mutants compared to the wild type, both during the mid-exponential and early stationary phases (**Figure 3C** [\[4\]](#)).

Finally, we utilized fluorophore-labeled aminoglycoside (Gentamicin-Texas Red, or GTTR) to investigate the cellular uptake of the drug. Specifically, cells in the mid-exponential phase of both wild-type and mutant strains (at t=3.5 hours when a significant increase in gentamicin tolerance was observed in the mutant strains) were exposed to GTTR for an hour, followed by the analysis of cells using flow cytometry. A short-term treatment was preferred, in line with a previous study (Bruni and Kralj, 2020 [\[4\]](#)) as cells treated with aminoglycosides become permeabilized at later time points (Bruni and Kralj, 2020 [\[4\]](#)), potentially introducing artificial impacts on drug uptake. Notably, the deletion of genes related to the TCA cycle and the electron transport chain did not induce a significant alteration in GTTR uptake in cells during the exponential phase when compared to the wild type (**Figure 4A, B** [\[4\]](#)). We note that the aminoglycoside concentration utilized in this study exceeds that of previous studies demonstrating the correlation between aminoglycoside uptake and PMF (Taber Harry W et al., 1987 [\[4\]](#)). Given that higher concentrations of aminoglycosides (>30 μg/ml) might obscure energy-dependent aminoglycoside uptake (Taber Harry W et al., 1987 [\[4\]](#)), we investigated a lower concentration of GTTR (5 μg/ml) and observed similar GTTR uptake across all




Figure 3.

Comparable ATP levels, redox activities, and cytoplasmic pH levels were observed between the wild-type and mutant strains.

(A) RSG staining was conducted by suspending wild-type and mutant strains in 0.85% sodium chloride solution during the mid-exponential and early stationary phases, as outlined in the Materials and Methods section. (B) The ATP levels were measured in wild-type and mutant cells during the mid-exponential and early stationary growth phases. (C) A comparison of the cytoplasmic pH of the WT and mutant cells during the mid-exponential and early stationary phases was performed using the ratiometric pHluorin. For pairwise comparisons, one-way ANOVA with Dunnett's post hoc test was used where *, $P < 0.05$, **, $P < 0.01$, ***, and $P < 0.001$. $N \geq 3$. Data points represent mean and standard error.

tested strains (**Supplementary Figure S5**). Collectively, these findings from multiple approaches strongly suggest that energy-dependent drug uptake is not the primary determinant of the observed antibiotic tolerance.

Membrane potential dysregulation is not associated with the observed aminoglycoside tolerance

Fluorescent sensors for voltage and calcium have been utilized to monitor electrophysiology in bacteria at the single-cell level (Bruni and Kralj, 2020), and the findings revealed that the dysregulated membrane voltage was not essential for aminoglycoside uptake or inner membrane pore formation in *E. coli*, but it proved crucial for bactericidal activity (Bruni and Kralj, 2020). To assess the significance of dysregulated PMF, we employed a well-established assay based on 3,3'-Dipropylthiadicarbocyanine [DiSC₃(5)] (Stokes et al., 2020; Wu et al., 1999), a fluorescent dye commonly used for monitoring cell membrane potential. During cell hyperpolarization, DiSC₃(5) infiltrates the cell membrane, undergoing self-quenching and resulting in reduced fluorescence intensity. Conversely, during depolarization, the fluorescence intensity of DiSC₃(5) increases as it exits the cell membrane. Polymyxin B served as a control (**Supplementary Figure S6**), as the polycationic peptide ring of polymyxin B binds to a negatively charged site within the lipopolysaccharide layer of the cellular membrane (Domingues et al., 2012), leading to the dissipation of the electric potential gradient which enhances the fluorescence intensity of DiSC₃(5) (**Supplementary Figure S6**). To assess the effect of gentamicin on membrane potential, metabolically active cells from exponential phase cultures at time points $t=3, 4, 5$, and 6 hours were transferred to a buffer solution containing 1 μ M DiSC₃(5) (**Figure 5**). Upon reaching equilibrium, the cells were treated with gentamicin. At designated time points, samples were collected, and fluorescence levels were measured using a plate reader (**Figure 5**). Similar to Polymyxin B treatments, gentamicin disrupts the electric potential gradient of PMF, leading to an increase in the fluorescence intensity of DiSC₃(5) (**Figure 5**). Despite significant and rapid alterations in membrane potential observed upon gentamicin treatment, there was no significant difference in membrane potential between aminoglycoside-sensitive and tolerant strains (**Figure 5**). Moreover, if any of the gene deletions had an impact on membrane potential, we would have expected to observe altered fluorescence intensity in the specific knockout strain compared to that of the wild type during the equilibrium stage (before the addition of the drug), as previously reported (Mohiuddin et al., 2022). However, no significant difference in DiSC₃(5) fluorescence intensities was observed among the strains tested during the equilibrium phase (the initial 20 minutes, as illustrated in **Figure 5**). Furthermore, we did not observe significant differences in PMF between mutant and wild-type strains when lower concentrations of gentamicin (5 μ g/ml) were used (**Supplementary Figure S7**).

The membrane dysregulation induced by aminoglycosides was previously attributed to the combined activity of NADH dehydrogenase and a reversed F₁F_o-ATPase (Bruni and Kralj, 2020), and the knockout strains targeting the proton-conducting F_o domain ($\Delta atpB$, $\Delta atpE$, $\Delta atpF$), as well as $\Delta atpG$, exhibited increased colony-forming units in response to aminoglycoside treatments in *E. coli* BW25113 (Bruni and Kralj, 2020). However, when we tested ATP synthase knockout strains of *E. coli* BW25113 from the Keio collection, we did not observe a comparable trend as reported earlier (Bruni and Kralj, 2020). In fact, the strains $\Delta atpB$, $\Delta atpE$, $\Delta atpF$, and $\Delta atpG$ exhibited similar sensitivity to gentamicin compared to the wild type under the conditions studied here (**Supplementary Figure S8A**). Interestingly, we observed a drastic increase in gentamicin tolerance in the $\Delta atpC$ mutant strain (**Supplementary Figure S8A**), which was not reported in the previous study (Bruni and Kralj, 2020), possibly due to the different experimental conditions used here. The *atpC* gene encodes an F₁ complex subunit of ATP synthase, promoting motor activity in the direction of ATP production rather than the reversed direction (Bruni and Kralj, 2020; Guo et al., 2019). We acknowledge potential variations between *E. coli* BW25113 and *E. coli* MG1655 strains, due to differences in their genomic DNA. To address this, we deleted two F_o




Figure 4.

Deletion of genes associated with the TCA cycle and ETC did not alter the drug uptake.

(A) Exemplary quantification of GTTR uptake in cells during the exponential growth phase ($t=3.5$ h). (B) GTTR assays were performed on cells in the mid-exponential phase for both the wild-type and mutant strains, followed by fluorescence measurement using flow cytometry after one hour. The concentration of GTTR is 25 $\mu\text{g}/\text{ml}$. For pairwise comparisons, one-way ANOVA with Dunnett's post hoc test was used (no statistical significance was detected). $N=3$. Data points represent mean and standard error.




Figure 5.

The dysregulation of membrane potential is not correlated with the observed aminoglycoside tolerance.

Deletions of the TCA cycle and electron transport chain genes resulted in no significant change in PMF. Mid-exponential and early stationary phase cells of wild-type and mutant strains were stained with DiSC₃(5), and at specified time intervals their fluorescence was measured using a plate reader. Cells were treated with gentamicin (50 μg/ml) after 20 minutes and fluorescence was measured again. N=4. Data points represent mean and standard error.

components ($\Delta atpA$ and $\Delta atpB$) and two F1 components ($\Delta atpC$ and $\Delta atpD$) from *E. coli* MG1655 and assessed their gentamicin survival profiles. While a moderate increase in tolerance was observed for mutant strains of F1 components during the mid-exponential phase (t=3 and 4 hours) compared to the wild type, no clear trend was observed for mutant strains of Fo components (**Supplementary Figure S8B**). The upregulation of tolerance in one of the mutant strains of F1 components ($\Delta atpD$) was also evident during the mid-exponential phase when cells were exposed to a lower concentration of gentamicin (**Supplementary Figure S9**). Overall, despite some variations in tolerance among ATP synthase mutants across different *E. coli* strains (MG1655 vs BW25113) and gentamicin concentrations (**Supplementary Figures S8 and S9**), the observed gentamicin-induced dysregulation in membrane potential may not be the primary factor contributing to the differences in antibiotic tolerance levels between the wild-type and TCA cycle mutant strains studied here.

Proteomic analysis reveals molecular responses in mutant strains, unveiling potential mechanisms underlying aminoglycoside tolerance

In order to gain further insights beyond the conducted experiments and elucidate the mechanism responsible for the observed tolerance in TCA cycle and ETC mutants, we utilized untargeted mass spectrometry to quantify proteins within these mutants and subsequently compared them to their wild-type counterparts. Proteomics data analysis, specifically involving the determination of protein fold-change and calculation of P-value (using *F*-test and *t*-test), was carried out through a process involving data transformation, normalization, and statistical procedures, as detailed elsewhere (Aguilan et al., 2020 [\[link\]](#)). Proteins exhibiting a positive $\log_2$ (fold-change) and a P-value less than 0.05 were categorized as significantly upregulated while those displaying a negative $\log_2$ (fold-change) and a P-value less than 0.05 were categorized as significantly downregulated in the mutant strains compared to the wild type (see **Supplementary Tables S2-S5** for upregulated proteins, and **Supplementary Tables S6-S9** for downregulated proteins). The STRING database was employed to predict both the physical and functional interactions between proteins by inputting the upregulated and downregulated proteins into the STRING interaction network (Szklarczyk et al., 2023 [\[link\]](#)). STRING carries out an automated pathway-enrichment analysis, focusing on the proteins entered and identifying pathways that occur more frequently than expected. The analysis is based on the statistical background of the entire genome and encompasses Gene Ontology annotations (all three domains), KEGG pathways, Uniprot keywords, and the hierarchical clustering of the STRING network itself (see **Supplementary Tables S10-S17** for significantly altered pathways identified). When discussing our findings, we primarily reference the STRING network, as it offers the advantage of broader coverage, including potential novel modules that might not yet be classified as pathways (Szklarczyk et al., 2019 [\[link\]](#)).

In the context of the upregulated protein-protein association networks in the $\Delta sucA$ and $\Delta gltA$ mutant strains, the STRING analysis unveiled significant functional enrichments (**Supplementary Tables S10 and S11**) in the TCA cycle, carbon and pyruvate metabolism, formate c-acetyltransferase activity, and the fatty acid metabolic pathway (**Figure 6A-B** [\[link\]](#), and **Supplementary Tables S10 and S11**). Specifically, key proteins associated with these enrichments encompassed FumA, FumC, AcnA, AcnB, SdhA, SdhB, Dld, TalA, FbaB, FadA, FadB, and Acs (**Figure 6A, B** [\[link\]](#)). Additionally, the $\Delta sucA$ and $\Delta gltA$ mutants exhibited an upregulated cluster of membrane proteins, particularly from the ABC transporter family (MalK, RbsA, RbsB, DppA, OppA, UgpB, LuxS), with some proteins specialized in amino acid transport across the plasma membrane (ArtL, LivJ, GltI, GlnQ, ArgT, HisJ) (**Figure 6A, B** [\[link\]](#)). In the case of the $\Delta gltA$ mutant, a smaller network of upregulated stress-induced proteins was also observed, related to osmotic stress, oxidative stress, and starvation (OsmC, OsmY, YbaY, YgaU, YjbJ, Dps, WrbA) (**Figure 6B** [\[link\]](#)). Regarding the $\Delta nuoI$ mutant, the STRING analysis identified an upregulated functional network associated with carbon and pyruvate metabolism, featuring proteins SucD, FumA, SdhA, Crp, Eda, GalS, Crr, Mdh, PykF, Lpd, GapA, Pfk, PoxB, and TalA, as well as upregulated flagellar proteins

(FlgF, FlgK, FlgM, FliD, FliH, FliJ, FliK, FliN, CheW) (**Figure 6C** [↗](#), and **Supplementary Table S12**). Interestingly, our proteomics data on these three mutant strains, which display higher gentamicin tolerance, reveals upregulation in proteins associated with energy metabolism (e.g., TCA cycle, pyruvate metabolism) (**Figure 6** [↗](#)). This suggests that these mutant strains compensate for metabolic perturbations by enhancing the TCA cycle to preserve their ATP levels, redox activities, and PMF. Indeed, this proteomics data aligns well with measurements of ATP, RSG, and PMF highlighted in previous sections (**Figures 3** [↗](#)-**5** [↗](#)).

In the context of downregulated protein networks, the substantial functional enrichments unveiled through analysis using the STRING database were found to be intricately associated with both the large and small ribosomal subunits, denoted by the specific ribosomal proteins RplX, RplM, RplT, RplY, RplP, RplS, RpsE, RpsG, RpsH, RpsM, RpsT, RpsI, RpsP, RpsK, RpsC, RpsD, RpsL, RpmG, RpmF, and RpmB (**Figure 7A-C** [↗](#), and **Supplementary Tables S14, S15, and S16**). These enrichments extended to domains encompassing translation factor activities, translation processes, and protein export mechanisms, featuring proteins RimM, InfA, InfC, Tsf, YeiP, NusG, Efp, PrfC, Mfd, GlyS, CysS, AspS, ArgS, and Tig (**Figure 7A-C** [↗](#)). Additionally, a noteworthy connection was established with the ribonucleoside monophosphate biosynthetic process, as characterized by proteins CarA, CarB, PyrD, PyrF, and PyrL (**Figure 7A-C** [↗](#)). The consistent downregulation of ribosomal binding proteins, ribosome biogenesis, translation factor activity, and the biosynthesis of ribonucleoside monophosphates is observed in the three mutant strains (Δ sucA, Δ gltA, and Δ nuoI), providing an explanation for the observed aminoglycoside tolerance in these mutants (**Figure 7A-C** [↗](#)). However, genome-level proteomics trends, specifically the drastic upregulation in energy metabolism and the drastic downregulation in ribosomal proteins, were not observed in the Δ icd mutant strain (**Supplementary Figure S10, and Supplementary Tables S13 and S17**). This may be expected, as the Δ icd mutant strain did not consistently exhibit an increased tolerance trend to aminoglycosides compared to Δ sucA, Δ gltA, and Δ nuoI.

Discussion

Our study aimed to explore the impact of gene knockouts associated with the TCA cycle and NADH dehydrogenase enzyme on aminoglycoside tolerance in *E. coli* MG1655. Various knockout strains showed increased tolerance to streptomycin, gentamicin, and amikacin compared to the wild-type strain. However, mutants like Δ icd, Δ acnB, and Δ fumA did not consistently exhibit enhanced tolerance, highlighting the complex relationship between energy metabolism and antibiotic tolerance. The increased tolerance observed in certain TCA cycle gene deletions to gentamicin was also documented in a prior study (Shan et al., 2015 [↗](#)). Our investigation further focused on four selected mutant strains (sucA, gltA, nuoI, and icd), revealing time-dependent profiles of antibiotic-tolerant cells, with a substantial increase observed in the Δ sucA, Δ gltA, and Δ nuoI mutants during the mid-exponential growth phase. The observed tolerance was not linked to altered cell growth, and it appeared to be transient or reversible, as all mutants regained sensitivity to aminoglycosides during the lag phase of growth.

We further investigated the factors influencing aminoglycoside tolerance in *E. coli* MG1655 by examining the role of energy-dependent drug uptake, membrane potential, and genetic perturbations. Studies have shown that cells tolerant to antibiotics can transition to susceptibility to aminoglycosides through the metabolism of specific carbon sources (Allison et al., 2011 [↗](#)). This shift in susceptibility is attributed to increased drug uptake, facilitated by increased activity in the electron transport chain and membrane potential, both triggered by the breakdown of these particular carbon sources (Allison et al., 2011 [↗](#)). Aminoglycoside uptake involves a three-step process: initial ionic binding to cells, followed by two energy-dependent phases (EDP I and EDP II) (Taber Harry W et al., 1987 [↗](#)). The first step, concentration-dependent ionic binding, is not affected by inhibitors of energized uptake (Taber Harry W et al., 1987 [↗](#)). The EDP I phase is gradual, requiring aminoglycoside concentration and a membrane potential for substantial uptake

Figure 6.

Proteomics data on the mutant strains, exhibiting increased gentamicin tolerance, indicates upregulation in proteins linked to energy metabolism.

Cells from both wild-type and mutant strains at the mid-exponential phase (t=3.5 h) were collected after which protein extraction and digestion were carried out for mass spectrometry analysis. The STRING visual network depicts upregulated protein interactions of $\Delta sucA$ (A), $\Delta gltA$ (B) and $\Delta nuoI$ (C) mutants, compared to those of the wild-type strain. The protein clusters and their corresponding gene names are visually distinguished through color-coding on the networks. N= 3.





Figure 7.

Proteomics data on the mutant strains, exhibiting increased gentamicin tolerance, reveals downregulation in proteins associated with ribosomes.

The STRING visual network displays down-regulated protein interactions for $\Delta sucA$ (A), $\Delta gltA$ (B), and $\Delta nuoI$ (C) mutants. N= 3.



(Taber Harry W et al., 1987 [↗](#)). Elevated aminoglycoside concentrations can bypass EDP I. EDP II involves rapid transport across the cytoplasmic membrane, relying on energy from electron transport and potentially ATP hydrolysis. Once inside the cell, studies suggest bactericidal effects through mistranslation and misfolding of membrane proteins, triggering oxidative stress and hydroxyl radical formation, ultimately leading to cell death (Kohanski et al., 2008 [↗](#), 2007 [↗](#)). Despite hypothesizing that reduced PMF and aminoglycoside uptake might be linked to genetic perturbations, the mutant strains with increased tolerance showed no consistent pattern in metabolic activities in our study. Analysis of ATP levels, intracellular pH, and fluorophore-labeled aminoglycoside uptake did not reveal a clear association with antibiotic tolerance. Certainly, our proteomic analysis revealed significant enhancements in upregulated protein networks linked to the TCA cycle, carbon metabolism, pyruvate metabolism, and fatty acid pathways in the mutant strains exhibiting increased gentamicin tolerance. This sheds light on how these mutants manage and maintain their ATP levels, redox activities, PMF, and drug uptake, overcoming genetic disruptions in their metabolism. However, it is crucial to emphasize that our study does not refute the PMF-dependent mechanism of aminoglycoside uptake. The observation of similar drug uptake across all strains tested, including mutants and wild type, can be attributed to the ability of the mutant strains to maintain their PMF under the conditions studied here.

Bruni and Kralj postulated that the bactericidal action triggered by membrane potential might involve mechanisms beyond drug uptake, potentially influenced by the combined effects of pore formation by mistranslated proteins and membrane hyperpolarization (Bruni and Kralj, 2020 [↗](#)). Additionally, their study proposed that the sudden shift in energy demand resulting from ribosome dissociation during aminoglycoside treatment could boost cellular ATP flux, leading to hyperpolarization through the concerted activity of NADH dehydrogenase and a reversed F1F0-ATPase (Bruni and Kralj, 2020 [↗](#)). Interestingly, while our study documented dysregulation in membrane potential after gentamicin treatment, no significant differences were noted between gentamicin-sensitive and tolerant strains. Despite prior research implicating increased survival of specific ATP synthase knockouts in response to aminoglycoside treatments (Bruni and Kralj, 2020 [↗](#)), our study did not report a clear trend regarding the gentamicin tolerance of these knockout strains. We acknowledge that variations in experimental conditions, such as medium composition, bacterial strains, growth phase, treatment time, duration, and antibiotic compositions, may account for the observed differences. Additionally, we recognize that genetic perturbations can have pleiotropic effects, potentially altering cell survival and death mechanisms, which may differ from those observed in the wild-type strain.

We conducted untargeted mass spectrometry to quantify proteins in the TCA-cycle and ETC mutants. Utilizing the STRING database, we predicted functional protein interactions, revealing significant enrichments in downregulated protein networks associated with ribosomal subunits, translation factor activities, protein export mechanisms, and ribonucleoside monophosphate biosynthesis in the mutant strains that displayed higher gentamicin tolerance. The altered levels of ribosomal proteins observed in our mutant strains align well with a prior study that identified respiratory complex I in *E. coli* as a critical mutational target for promoting persister cell formation during the transition to a stationary phase (Van den Bergh et al., 2022 [↗](#)). In that study, point mutations in respiratory complex I, responsible for proton translocation, were found to induce antibiotic tolerance (Van den Bergh et al., 2022 [↗](#)). While they showed that mutations compromised proton translocation, key components such as NADH oxidation, electron transfer, and drug uptake remained largely unchanged (Van den Bergh et al., 2022 [↗](#)). They also demonstrated that the increased persistence correlated with an acidified cytoplasm during the stationary phase, impacting protein translation and contributing to increased antibiotic tolerance, yet no significant differences in pH and persistence were noted between mutant and wild-type strains during the exponential phase (Van den Bergh et al., 2022 [↗](#)). Similarly, our investigation revealed that single gene deletions did not alter cellular PMF, redox activities, membrane potential, and drug uptake. However, the downregulation of ribosomal protein levels reported in our study may not be attributed to a reduction in cytoplasmic pH, as we did not observe

alterations in pH levels in the knockout strains compared to the wild type during the mid-exponential phase. Certain genes linked to the deletion strains in our study, such as *icd*, *sucA*, *mdh*, and *fumA*, do not encode membrane-bound respiratory proteins. Strain-specific differences can significantly influence the outcomes and responses to various experimental conditions. Each strain of *E. coli* or any other microorganism can harbor unique genetic backgrounds, mutation profiles, and physiological characteristics, leading to distinct survival strategies in response to environmental stresses, such as antibiotic exposure. This variability is particularly evident in the *icd* deletion strain, as evidenced by its increased ciprofloxacin persistence in an independent study conducted by Lewis' group (Manuse et al., 2021 [DOI](#)). Furthermore, the same group demonstrated that the *icd* deletion strain exhibited increased sensitivity to gentamicin in another study (Shan et al., 2015 [DOI](#)), consistent with our observation here.

In summary, our study revealed that deletions in TCA cycle and ETC genes, including *sucA*, *gltA*, *mdh*, *sdhC*, *icd*, *acnB*, *fumA*, *nuoM*, and *nuoI*, increased aminoglycoside tolerance. Analyses, including flow cytometry and proteomics, demonstrated enhanced tolerance without significant changes in energy-dependent drug uptake or membrane potential. Our pathway analysis identified a downregulation in large and small ribosomal binding proteins, ribosome biogenesis, translation factor activity, and ribonucleoside monophosphate biosynthesis in all mutant strains, offering a plausible explanation for the observed aminoglycoside tolerance in these mutants. Altogether, our findings underscore the complexity of energy metabolism's role in antibiotic tolerance, providing valuable insights into the molecular mechanisms of aminoglycoside tolerance in specific mutant strains.

Materials and methods

Bacterial strains, chemicals, media, and growth conditions

We employed the in-house *Escherichia coli* K-12 MG1655 strain for our experiments. The gene deletions in *E. coli* K-12 MG1655 were established in our prior studies (Ngo et al., 2024 [DOI](#)) utilizing the Datsenko and Wanner method (Datsenko and Wanner, 2000 [DOI](#)). *E. coli* K-12 BW25113 and its corresponding knockout strains were procured from the Keio collection, purchased from Horizon Discovery (Catalog #OEC4988, Lafayette, CO, USA). The validity of all deletions was confirmed through the use of check primers (Ngo et al., 2024 [DOI](#)). All chemicals utilized in the study were purchased from Fisher Scientific (Atlanta, GA) or VWR International (Pittsburg, PA) unless stated otherwise. The Gentamicin-Texas Red Conjugate (Catalog # 24300) was purchased from AAT Bioquest, Inc (CA, USA). The ATP measurement kit (G8230) was acquired from Promega Corporation (Madison, WI). Standard Lysogeny Broth (LB) broth was prepared by dissolving 5 g yeast extract, 10 g tryptone, and 10 g sodium chloride in 1 liter of deionized (DI) water. LB agar was made by dissolving 40 g of pre-mixed LB agar powder in 1 liter DI water. Sterilization of LB broth and LB agar was done via autoclaving at 121°C and 103.421 kPa. To determine tolerant cells in cultures, gentamicin (50 µg/ml), streptomycin (50 µg/ml), and amikacin (50 µg/ml) were used, and their concentrations were selected to be much higher than the minimum inhibitory concentrations (MICs). Whenever specified, gentamicin at various concentrations (e.g., 5 µg/ml) was also tested. MICs of antibiotics for *E. coli* (Andrews, 2001 [DOI](#)) were determined using MIC Test strips (Fisher Scientific) and the results were presented in **Supplementary Table S1**. Ampicillin (100 µg/ml) was utilized to maintain plasmids in the cells. 0.2% L-arabinose at 13.3 mM was added to the media to induce GFP expression. To decrease the antibiotic concentration below the MIC, a sterile 1X phosphate buffer saline (PBS) solution was utilized to wash the cells. Antibiotics were dissolved in DI water and sterilized using a 0.2 µm PES syringe filter. Overnight cultures were prepared in 14 ml round-bottom tubes (Catalog # 14-959-1B, Fisher Scientific) by incubating cells from a frozen 25% glycerol stock (-80°C) in 2 ml of LB at 37°C and 250 rpm. After 24 hours, cells

from the overnight cultures were diluted 1:100 fold in 2 ml of fresh LB medium in 14-ml round bottom test tubes or 25 ml LB in 250-ml baffled flasks and cultured further to achieve the desired growth phase for the assays.

Cell growth assay

To prepare overnight precultures, cells from frozen stocks were inoculated into 2 ml of LB medium in 14-ml round-bottom tubes and cultured for 24 hours at 37°C with shaking at 250 rpm. The overnight precultures were then diluted 100-fold in 14-ml round-bottom tubes containing 2 ml of LB medium and incubated in the shaker at 250 rpm and 37°C. The growth of the cultures was determined by measuring the number of cells per ml using flow cytometry. To perform this task, cells were transferred to PBS at specified time points and analyzed with a flow cytometer (NovoCyte Flow Cytometer, NovoCyte 3000RYB, ACEA Biosciences Inc., San Diego, CA). The cell populations were delineated on flow diagrams utilizing the forward and side scatter parameters, with controls including PBS with cells and PBS without cells. The inclusion of a PBS solution without cells aids in noise determination. The instrument is capable of quantifying both the number of events and the volume of the solution under analysis. Roughly 30,000-50,000 events were analyzed for each sample.

Clonogenic survival assay

Overnight cultures were prepared in 14-ml round-bottom test tubes by incubating cells from a frozen 25% glycerol stock (-80°C) in 2 ml of LB at 37°C and 250 rpm. After 24 hours, cells from the overnight cultures were diluted 1:100 fold in 2 ml of fresh medium and cultured in the incubator at 37°C with shaking. At the designated growth phase or time points, cells were exposed to antibiotics at the specified concentrations (5 or 50 µg/ml gentamicin, 50 µg/ml streptomycin, and 50 µg/ml amikacin) for a duration of 5 hours. To determine the number of live cells before antibiotic exposure, 10 µl of cell cultures were serially diluted in PBS and plated on an LB agar plate, which was incubated for 16 hours at 37°C. During the antibiotic treatments, 1 ml cultures were collected after 5 hours and washed twice with PBS through centrifugation at 13,300 rpm (17,000 g) for 3 minutes to remove antibiotics. After the final centrifugation, 900 µl of supernatant was removed using a pipette, and the cell pellets were resuspended in the remaining 100 µl of PBS. Next, 10 µl of the cell suspensions were serially diluted in PBS, and 10 µl of the diluted cell suspensions were spotted onto LB agar plates. The plates were then incubated at 37°C for at least 16 hours, and the colony-forming units (CFU) were counted to determine the number of live cells present in the cultures. Survival fractions were calculated by dividing the number of surviving cells (after treatment) by the initial number of cells (before treatment).

To generate survival time profiles, cells from the overnight cultures were diluted 1:100 in 25 ml of fresh LB medium in 250-ml baffled flasks and incubated in the incubator at 37°C with shaking. At the designated time points, 2 ml of cell cultures were transferred to 14-ml round-bottom test tubes and subjected to antibiotic treatment at the specified concentration (50 µg/ml gentamicin) for a duration of 5 hours. Following treatment, cells were collected, washed to remove antibiotics, and plated for CFU enumeration, as previously described.

Redox sensor green dye staining

To measure bacterial reductase and ETC (electron transport chain) activities, we used the *BacLight* Redox Sensor Green Vitality kit (Catalog#B34954, Thermo Fisher) following the manufacturer's instructions. Overnight cultures were prepared in 14 ml Falcon tubes by incubating cells from a frozen 25% glycerol stock (-80°C) in 2 ml of LB at 37°C and 250 rpm. After 24 hours, cells from the overnight cultures were diluted 1:100 fold in 2 ml of fresh medium and cultured in the incubator at 37°C with shaking. For analyzing the cell populations during mid-exponential and early stationary phases ($t = 3\text{h}, 4\text{h}, 5\text{h}, 6\text{h}$), we diluted the cells in 1 ml of 0.85% sodium chloride solution in flow cytometry tubes (5 ml round bottom falcon tubes) by varying amounts (10-, 20-, 20-, 50-fold,

respectively). After that, Redox Sensor Green (RSG) dye was added to the cells at a concentration of 1 μ M, and the samples were incubated at 37°C for 10 minutes before flow cytometry analysis. For the negative controls, cell suspensions were treated with 20 μ M CCCP 5 minutes prior to RSG staining to disrupt membrane electron transport (**Supplementary Figure S2**). Positive controls consisted of mid-exponential-phase cells. The cell populations were gated on flow diagrams using the forward and side scatter parameters of unstained controls. Cells were excited at 488 nm with a solid-state laser, and green fluorescence was collected with a 530/30 bandpass filter.

ATP measurement

The BacTiter-Glo Microbial Cell Viability assay kit (Catalog# G8230, Promega Corporation) was used to measure the intracellular ATP levels of both *E. coli* MG1655 WT and mutant strains during a specified growth phase, following the manufacturer's instructions. To generate a standard curve, ATP solutions of known concentrations were used (**Supplementary Figure S2**). Background luminescence was measured using LB broth.

pH measurement

For pH measurements in *E. coli*, a pGFP01 plasmid was used, in which the GFP derivative ratiometric pHluorin is expressed from the arabinose-induced promoter P_{BAD} . This plasmid was kindly provided by Keith A. Martinez II (Department of Biology, Kenyon College, Gambier, Ohio, USA). A comprehensive pH measurement protocol was obtained from a prior study (Van den Bergh et al., 2022 [DOI](#)). Overnight cultures of WT and mutant strains were prepared in 14 ml tubes by incubating cells carrying the pGFP01 plasmid from a frozen 25% glycerol stock (-80°C) in 2 ml of LB, 0.2% L-arabinose and 100 mg/ml ampicillin, at 37°C and 250 rpm. After 24 hours, cells from the overnight cultures were diluted 1:100 fold in 2 ml of fresh medium, 0.2% L-arabinose and 100 mg/ml ampicillin and cultured in the incubator at 37°C with shaking. Fluorescence was measured at 410 nm and 470 nm using a plate reader at different time points, after measuring and normalizing the optical density of cells in liquid culture. The 410/470 fluorescence ratios were recorded to determine the cytoplasmic pH using the standard curve (**Supplementary Figure S3**). To generate standard pH versus fluorescence ratio curves for *E. coli* MG1655 cells, transmembrane pH was collapsed by adding 40 mM potassium benzoate and 40 mM methylamine hydrochloride to the cells, equalizing the difference between the external and internal pH. The cultures were then buffered to different pH levels ranging from 5 to 10 using a 50 mM concentration of 2-(*N*-morpholino) ethanesulfonic acid (MES) or 3-(*N*-morpholino) propanesulfonic acid (MOPS), and corresponding fluorescence values at 410 nm and 470 nm were obtained using a plate reader. The Boltzmann equation was used to establish the standard curve for each bacterial strain based on the provided data (Martinez II et al., 2012).

DiSC₃(5) assay

The fundamental mechanism of this assay and the assessment of PMF components using DiSC₃(5) have been extensively detailed elsewhere (Farha et al., 2013 [DOI](#); Panta and Doerrler, 2021 [DOI](#); Stokes et al., 2020 [DOI](#)). *E. coli* MG1655 wild-type and mutant cells in both exponential and early stationary phases (at time points $t = 3$ h, 4h, 5h, 6h) were collected and subjected to two washes with an assay buffer containing 5 mM HEPES and 20 mM glucose. The cell density was set to $OD_{600}=0.1$, and the cells were stained with 1 μ M DiSC₃(5). Fluorescence readings were taken at specified intervals using a plate reader, with excitation and emission wavelengths set at 620 nm and 670 nm, respectively. Gentamicin (5 μ g/ml, 50 μ g/ml, or other specified concentrations) was introduced 20 minutes after cells reached equilibrium. At this point, the probe was released into the medium, leading to an upsurge in fluorescence. Polymyxin B (32 μ g/ml) was included as controls and administered 20 minutes after equilibrium to dissipate the electron gradient of PMF. The concentration of polymyxin B was determined based on a prior study (Stokes et al., 2020 [DOI](#)).

Gentamicin uptake assay

Overnight cultures were prepared in 14 ml tubes by incubating cells from a frozen 25% glycerol stock (-80°C) in 2 ml of LB at 37°C and 250 rpm. After 24 hours, cells from the overnight cultures were diluted 1:100 fold in 2 ml of fresh medium and cultured in the incubator at 37°C with shaking. At the mid-exponential phase ($t=3.5$ h), 100 μ l of cell cultures for both wild-type and mutants were exposed to gentamicin-Texas red (GTTR) at a final concentration of 25 μ g/ml or 5 μ g/ml. Untreated mid-exponential phase cells were used as negative controls. The samples containing GTTR were then incubated for 1 hour at 37°C with shaking at 250 rpm. Subsequently, 20 μ l from each sample was washed with 500 μ l of PBS (1X). After the final washing step, cell pellets were resuspended in 500 μ l of PBS. All samples were subjected to analysis using a flow cytometer equipped with lasers emitting light at a wavelength of 561 nm, and the resulting red fluorescence was detected using a 615/20 nm bandpass filter.

Proteomics sample preparation and analysis

Cells from both wild-type and mutant strains at the mid-exponential phase ($t=3.5$ h) were collected after which protein extraction and digestion were carried out for Liquid Chromatography–Mass Spectrometry (LC-MS) analysis by following the Sample Preparation by Easy Extraction and Digestion (SPEED) procedure. The experiments were conducted at the University of Houston Mass Spectrometry Laboratory under a service fee. Comprehensive details regarding the protein isolation and digestion methods can be found elsewhere (Doellinger et al., 2020 [DOI](#)). Briefly, cell pellets were added 20 μ g trifluoroacetic acid (TFA) and incubated at room temperature for 5 min followed by addition of 200 μ l of 2M trisbase. After adding Tris(2-carboxyethyl) phosphine (10 mM) and 2-Chloroacetamide (40 mM), the reaction mixture was heated at 95 °C for 5 min. Digestion was performed by adding trypsin (1/40, w/w) and incubation at 37 °C overnight. The digested peptides were cleaned up using a C18 Ziptip and vacuum dried using a CentriVap (Labconco, Kansas City, MO). Each dried sample was resuspended in 2% acetonitrile (ACN) with 0.1% formic acid (FA) for LC-MS analysis. The method involving LC-MS has been detailed in a separate publication (Qin et al., 2022 [DOI](#)). Specifically, a NanoElute LC system connected to a timsTOF Pro (Bruker Daltonics, Germany) through a CaptiveSpray source was utilized. Samples were loaded onto an in-house packed column (75 μ m x 20 cm, 1.9 μ m ReproSil-Pur C18 particle from Dr. Maisch GmbH, Germany) with a column temperature of 40 °C. Mobile phases included buffer A (0.1% FA in water) and buffer B (0.1% FA in ACN). The short gradient was 0-17.8 minutes, from 2% B to 30% B, followed by 18.3 minutes to 95% B, and 20.7 minutes to 95% B. The parallel accumulation-serial fragmentation (PASEF) mode with 4 PASEF scans per cycle was employed. The electrospray voltage was set at 1.4 kV, and the ion transfer tube temperature was maintained at 180 °C. Full MS scans were conducted across the mass-to-charge (m/z) range of 150–1700. The target intensity value was 2.0×10^5 with a threshold of 2500. A fixed cycle time of 0.53 s was established, and a dynamic exclusion duration of 0.4 minutes with a ± 0.015 amu tolerance was applied. Only peaks with a charge state of ≥ 2 were chosen for fragmentation. The default settings of MSFragger, a database search tool designed for peptide identification in MS-based proteomics (Yu et al., 2020 [DOI](#)), were applied to analyze data obtained from the mentioned instrument. The UniProt-SwissProt *E. coli* K12 database (Taxon ID 83333, downloaded on 6/19/2023, 4518 entries) served as the reference. Fixed modification involved cysteine carbamidomethylation, while variable modifications included methionine oxidation and acetylation. Peptide length was restricted to 7-50, allowing for 2 missed cleavages. Both precursor and product ion masses were set as monoisotopic. The false discovery rate (FDR) was controlled at <1% at the peptide spectrum match, peptide, and protein levels.

Proteomics data analysis

The processing of proteomics data and the calculations of fold change were essentially carried out following the methods described in the paper by Aguilan *et al* (Aguilan *et al.*, 2020 [DOI](#)). In summary, we utilized Excel spreadsheets for key stages of data transformation, normalization, fold change, and P-value calculation. Initially, proteins lacking quantitative values were excluded, and a logarithm transformation was applied to achieve a normal distribution of data. Normalization, using both average and slope methods, was then employed to minimize intragroup variation in technical replicates and log fold change calculations compared to the transformed unnormalized data. Subsequently, missing values were imputed by replacing them with approximated values using the Probabilistic Minimum Imputation method. Following imputation, we determined the relative ratio of each protein in mutant and wild-type strains, along with P-value calculation using the parametric *t*-test. The selection of the *t*-test type involved an *F*-test to evaluate whether the replicates for each protein exhibited homoscedastic (equal variances) or heteroscedastic (unequal variances) characteristics. For the identification of significant networks among input proteins, we utilized the STRING tool V 12.0. This entailed inputting proteins that were significantly upregulated and downregulated based on specified thresholds, as detailed elsewhere (Szkłarczyk *et al.*, 2023 [DOI](#)).

Statistical analysis

All assays were conducted using at least three independent biological replicates. The figures display the mean value and standard error for each data point. Statistical analysis was performed using GraphPad Prism software, with one-way analysis of variance (ANOVA) with Dunnett's post hoc test to determine significance. The P value threshold was set at *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$, ****, and $P < 0.0001$.

Data Availability

The proteomics data will be submitted to one of the NIH-designated repositories. All data generated or analyzed during this study have been incorporated into the manuscript and supplemental files.

Author Details

Mehmet Orman

William A. Brookshire Department of Chemical and Biomolecular Engineering, University of Houston

Contribution: Conceptualization, Data curation, Supervision, Funding acquisition, Investigation, Writing - original draft, Project administration, Writing – review and editing

For correspondence: morman@central.uh.edu

Competing interests: No competing interests declared

Rauf Shiraliyev

William A. Brookshire Department of Chemical and Biomolecular Engineering, University of Houston

Contribution: Data curation, Investigation, Writing - original draft, review and editing

Competing interests: No competing interests declared

Acknowledgment

The authors would like to thank the members of Orman Lab for their help. This study was supported by NSF CAREER 2044375 and NIH/NIAID R01-AI143643.

Proteomics experiments were conducted at the Mass Spectrometry Laboratory of Dr. Chengzhi Cai at the University of Houston, with the associated service fees.

References

- Aggen JB *et al.* (2010) **Synthesis and spectrum of the neoglycoside ACHN-490** *Antimicrob Agents Chemother* **54**:4636–4642 <https://doi.org/10.1128/AAC.00572-10>
- Aguilan JT, Kulej K, Sidoli S (2020) **Guide for protein fold change and: P-value calculation for non-experts in proteomics** *Mol Omics* **16**:573–582 <https://doi.org/10.1039/d0mo00087f>
- Allison KR, Brynildsen MP, Collins JJ (2011) **Metabolite-enabled eradication of bacterial persisters by aminoglycosides** *Nature* **473**:216–220 <https://doi.org/10.1038/nature10069>
- Amato SM, Orman MA, Brynildsen MP (2013) **Metabolic Control of Persister Formation in Escherichia coli** *Mol Cell* **50**:475–487 <https://doi.org/10.1016/j.molcel.2013.04.002>
- Andrews JM (2001) **Determination of minimum inhibitory concentrations** *Journal of Antimicrobial Chemotherapy* **48**:5–16 https://doi.org/10.1093/jac/48.suppl_1.5
- Balaban NQ *et al.* (2019) **Definitions and guidelines for research on antibiotic persistence** *Nat Rev Microbiol* **17**:441–448 <https://doi.org/10.1038/s41579-019-0196-3>
- Becker B, Cooper MA (2013) **Aminoglycoside antibiotics in the 21st century** *ACS Chem Biol* **8**:105–115 <https://doi.org/10.1021/cb3005116>
- Binder S, Levitt AM, Sacks JJ, Hughes JM (1999) **Emerging Infectious Diseases: Public Health Issues for the 21st Century** *Science (1979)* **284**:1311–1313 <https://doi.org/10.1126/science.284.5418.1311>
- Borovinskaya MA, Pai RD, Zhang W, Schuwirth BS, Holton JM, Hirokawa G, Kaji H, Kaji A, Cate JHD (2007) **Structural basis for aminoglycoside inhibition of bacterial ribosome recycling** *Nat Struct Mol Biol* **14**:727–732 <https://doi.org/10.1038/nsmb1271>
- Bruni GN, Kralj JM (2020) **Membrane voltage dysregulation driven by metabolic dysfunction underlies bactericidal activity of aminoglycosides** *Elife* **9**:1–25 <https://doi.org/10.7554/ELIFE.58706>
- C Reygaert W (2018) **An overview of the antimicrobial resistance mechanisms of bacteria** *AIMS Microbiol* **4**:482–501 <https://doi.org/10.3934/microbiol.2018.3.482>

- Darby EM, Trampari E, Siasat P, Gaya MS, Alav I, Webber MA, Blair JMA (2023) **Molecular mechanisms of antibiotic resistance revisited** *Nat Rev Microbiol* **21**:280–295 <https://doi.org/10.1038/s41579-022-00820-y>
- Datsenko KA, Wanner BL (2000) **One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products** *PNAS* **97**:6640–6645 <https://doi.org/10.1073/pnas.120163297>
- Davis BD, Chen L, Tai PC (1986) **Misread protein creates membrane channels: An essential step in the bactericidal action of aminoglycosides** *PNAS* **83**:6164–6168 <https://doi.org/10.1073/pnas.83.16.6164>
- Doellinger J, Schneider A, Hoeller M, Lasch P (2020) **Sample preparation by easy extraction and digestion (SPEED) - A universal, rapid, and detergent-free protocol for proteomics based on acid extraction** *Molecular and Cellular Proteomics* **19**:209–222 <https://doi.org/10.1074/mcp.TIR119.001616>
- Domingues MM, Inácio RG, Raimundo JM, Martins M, Castanho MARB, Santos NC (2012) **Biophysical characterization of polymyxin B interaction with LPS aggregates and membrane model systems** *Biopolymers* **98**:338–344 <https://doi.org/10.1002/bip.22095>
- Endimiani A, Hujer KM, Hujer AM, Armstrong ES, Choudhary Y, Aggen JB, Bonomo RA (2009) **ACHN-490, a neoglycoside with potent in vitro activity against multidrug-resistant Klebsiella pneumoniae isolates** *Antimicrob Agents Chemother* **53**:4504–4507 <https://doi.org/10.1128/AAC.00556-09>
- Farha MA, Verschoor CP, Bowdish D, Brown ED (2013) **Collapsing the proton motive force to identify synergistic combinations against staphylococcus aureus** *Chem Biol* **20**:1168–1178 <https://doi.org/10.1016/j.chembiol.2013.07.006>
- Germain E, Roghanian M, Gerdes K, Maisonneuve E (2015) **Stochastic induction of persister cells by HipA through (p)ppGpp-mediated activation of mRNA endonucleases** *Proc Natl Acad Sci U S A* **112**:5171–5176 <https://doi.org/10.1073/pnas.1423536112>
- Guo H, Suzuki T, Rubinstein JL (2019) **Structure of a bacterial ATP synthase** *Elife* **8** <https://doi.org/10.7554/eLife.43128.001>
- Helaine S, Kugelberg E (2014) **Bacterial persisters: Formation, eradication, and experimental systems** *Trends Microbiol* <https://doi.org/10.1016/j.tim.2014.03.008>
- Ikäheimo I, Syrjälä H, Karhukorpi J, Schildt R, Koskela M (2000) **In vitro antibiotic susceptibility of Francisella tularensis isolated from humans and animals** *Journal of Antimicrobial Chemotherapy* **46**:287–290 <https://doi.org/10.1093/jac/46.2.287>
- Karlowsky JA, Draghi DC, Jones ME, Thornsberry C, Friedland IR, Sahm DF (2003) **Surveillance for antimicrobial susceptibility among clinical isolates of Pseudomonas aeruginosa and Acinetobacter baumannii from hospitalized patients in the United States, 1998 to 2001** *Antimicrob Agents Chemother* **47**:1681–1688 <https://doi.org/10.1128/AAC.47.5.1681-1688.2003>
- Kohanski MA, Dwyer DJ, Hayete B, Lawrence CA, Collins JJ (2007) **A Common Mechanism of Cellular Death Induced by Bactericidal Antibiotics** *Cell* **130**:797–810 <https://doi.org/10.1016/j.cell.2007.06.049>

- Kohanski MA, Dwyer DJ, Wierzbowski J, Cottarel G, Collins JJ (2008) **Mistranslation of Membrane Proteins and Two-Component System Activation Trigger Antibiotic-Mediated Cell Death** *Cell* **135**:679–690 <https://doi.org/10.1016/j.cell.2008.09.038>
- Krause KM, Serio AW, Kane TR, Connolly LE (2016) **Aminoglycosides: An overview** *Cold Spring Harb Perspect Med* **6** <https://doi.org/10.1101/cshperspect.a027029>
- Kresge N, Simoni RD, Hill RL (2004) **Selman Waksman: the Father of Antibiotics** *Journal of Biological Chemistry* **279**:e7–e8 [https://doi.org/10.1016/s0021-9258\(20\)67861-9](https://doi.org/10.1016/s0021-9258(20)67861-9)
- Lambert PA (2005) **Bacterial resistance to antibiotics: Modified target sites** *Adv Drug Deliv Rev* **57**:1471–1485 <https://doi.org/10.1016/j.addr.2005.04.003>
- Landman D, Babu E, Shah N, Kelly P, Bäcker M, Bratu S, Quale J (2010) **Activity of a novel aminoglycoside, ACHN-490, against clinical isolates of Escherichia coli and Klebsiella pneumoniae from New York City** *Journal of Antimicrobial Chemotherapy* **65**:2123–2127 <https://doi.org/10.1093/jac/dkq278>
- Lopatkin AJ, Bening SC, Manson AL, Stokes JM, Kohanski MA, Badran AH, Earl AM, Cheney NJ, Yang JH, Collins JJ (2021) **Clinically relevant mutations in core metabolic genes confer antibiotic resistance** *Science (1979)* **371** <https://doi.org/10.1126/science.aba0862>
- Manuse S, Shan Y, Canas-Duarte SJ, Bakshi S, Sun WS, Mori H, Paulsson J, Lewis K (2021) **Bacterial persisters are a stochastically formed subpopulation of low-energy cells** *PLoS Biol* **19** <https://doi.org/10.1371/journal.pbio.3001194>
- Martinez KA, Kitko RD, Mershon JP, Adcox HE, Malek KA, Berkmen MB, Slonczewski JL (2012) **Cytoplasmic pH Response to Acid Stress in Individual Cells of Escherichia coli and Bacillus subtilis Observed by Fluorescence Ratio Imaging Microscopy** *Appl Environ Microbiol* **78**:3706–3714 <https://doi.org/10.1128/AEM.00354-12>
- Miesenböck G, Angelis DA De, Rothman JE (1998) **Visualizing secretion and synaptic transmission with pH-sensitive green fluorescent proteins** *Nature* **394**:192–195 <https://doi.org/10.1038/28190>
- Moazed D, Noller HF (1987) **Interaction of antibiotics with functional sites in 16S ribosomal RNA** *Nature* **327**:389–394 <https://doi.org/10.1038/327389a0>
- Mohiuddin SG, Kavousi P, Orman MA (2020) **Flow-cytometry analysis reveals persister resuscitation characteristics** *BMC Microbiol* **20**:1–13 <https://doi.org/10.1186/s12866-020-01888-3>
- Mohiuddin SG, Massahi A, Orman MA (2022) **High-Throughput Screening of a Promoter Library Reveals New Persister Mechanisms in Escherichia Coli** *Microbiol Spectr* **10** <https://doi.org/10.1128/spectrum.02253-21>
- Molina-Quiroz RC, Silva-Valenzuela C, Brewster J, Castro-Nallar E, Levy SB, Camilli A (2018) **Cyclic AMP Regulates Bacterial Persistence through Repression of the Oxidative Stress Response and SOS-Dependent DNA Repair in Uropathogenic Escherichia coli** *mBio* **9**:e02144–17 <https://doi.org/10.1128/mBio.02144-17>
- Muir ME, Hanwell DR, Wallace BJ (1981) **Characterization of a respiratory mutant of Escherichia coli with reduced uptake of aminoglycoside antibiotics** *Biochim Biophys Acta* **638**:234–241 [https://doi.org/10.1016/0005-2728\(81\)90232-2](https://doi.org/10.1016/0005-2728(81)90232-2)

- Munita JM, Arias CA (2016) **Mechanisms of Antibiotic Resistance** *Microbiol Spectr* **4** <https://doi.org/10.1128/microbiolspec.VMBF-0016-2015>
- Ngo HG, Mohiuddin SG, Ananda A, Orman MA (2024) **Unraveling Crp/cAMP-mediated metabolic regulation in Escherichia coli persister cells** *Under review*
- Okusu H, Ma D, Nikaido H (1996) **AcrAB Efflux Pump Plays a Major Role in the Antibiotic Resistance Phenotype of Escherichia coli Multiple-Antibiotic-Resistance (Mar) Mutants** *J Bacteriol* **178**:306–308 <https://doi.org/10.1128/jb.178.1.306-308.1996>
- Panta PR, Doerrler WT (2021) **A Burkholderia thailandensis DedA Family Membrane Protein Is Required for Proton Motive Force Dependent Lipid A Modification** *Front Microbiol* **11** <https://doi.org/10.3389/fmicb.2020.618389>
- Qin G, Zhang P, Sun M, Fu W, Cai C (2022) **Comprehensive spectral libraries for various rabbit eye tissue proteomes** *Sci Data* **9** <https://doi.org/10.1038/s41597-022-01241-5>
- Recht MI, Douthwaite S, Dahlquist KD, Puglisi JD (1999) **Effect of Mutations in the A Site of 16 S rRNA on Aminoglycoside Antibiotic-Ribosome Interaction** *J Mol Biol* **286**:33–43 <https://doi.org/10.1006/jmbi.1998.2446>
- Ristuccia AM, Cunha BA (1985) **An overview of amikacin** *Ther Drug Monit* **7**:12–25 <https://doi.org/10.1097/00007691-198503000-00003>
- Sader HS, Rhomberg PR, Farrell DJ, Jones RN (2015) **Arbekacin activity against contemporary clinical bacteria isolated from patients hospitalized with pneumonia** *Antimicrob Agents Chemother* **59**:3263–3270 <https://doi.org/10.1128/AAC.04839-14>
- Schatz A, Bugie E, Waksman SA (2005) **Streptomycin, a substance exhibiting antibiotic activity against gram-positive and gram-negative bacteria** *Clin Orthop Relat Res* **437**:3–6 <https://doi.org/10.1097/01.blo.0000175887.98112.fe>
- Shan Y, Lazinski D, Rowe S, Camilli A, Lewis K (2015) **Genetic basis of persister tolerance to aminoglycosides in Escherichia coli** *mBio* **6** <https://doi.org/10.1128/mBio.00078-15>
- Stokes JM *et al.* (2020) **A Deep Learning Approach to Antibiotic Discovery** *Cell* **180**:688–702 <https://doi.org/10.1016/j.cell.2020.01.021>
- Szklarczyk D *et al.* (2019) **STRING v11: Protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets** *Nucleic Acids Res* **47**:D607–D613 <https://doi.org/10.1093/nar/gky1131>
- Szklarczyk D *et al.* (2023) **The STRING database in 2023: protein-protein association networks and functional enrichment analyses for any sequenced genome of interest** *Nucleic Acids Res* **51**:D638–D646 <https://doi.org/10.1093/nar/gkac1000>
- Harry W Taber, Mueller JP, Miller PF, Arrow AS (1987) **Bacterial Uptake of Aminoglycoside Antibiotics** *Microbiol Rev* **51**:439–457 <https://doi.org/10.1128/mr.51.4.439-457.1987>
- Theodore A, Lewis K, Vulić M (2013) **Tolerance of Escherichia coli to fluoroquinolone antibiotics depends on specific components of the SOS response pathway** *Genetics* **195**:1265–1276 <https://doi.org/10.1534/genetics.113.152306>

Van den Bergh B, Fauvart M, Michiels J (2017) **Formation, physiology, ecology, evolution and clinical importance of bacterial persisters** *FEMS Microbiol Rev* **41**:219–251 <https://doi.org/10.1093/femsre/fux001>

Van den Bergh B *et al.* (2022) **Mutations in respiratory complex I promote antibiotic persistence through alterations in intracellular acidity and protein synthesis** *Nat Commun* **13** <https://doi.org/10.1038/s41467-022-28141-x>

Wood TK, Knabel SJ, Kwan BW (2013) **Bacterial persister cell formation and dormancy** *Appl Environ Microbiol* <https://doi.org/10.1128/AEM.02636-13>

Woodcock J, Moazed D, Cannon M, Davies J, Noller HF (1991) **Interaction of antibiotics with A- and P-site-specific bases in 16S ribosomal RNA** *EMBO J* **1**:3099–3103 <https://doi.org/10.1002/j.1460-2075.1991.tb07863.x>

Woodruff HB (2014) **Selman A. Waksman, winner of the 1952 nobel prize for physiology or medicine** *Appl Environ Microbiol* **80**:2–8 <https://doi.org/10.1128/AEM.01143-13>

Wu M, Maier E, Benz R, Hancock REW (1999) **Mechanism of interaction of different classes of cationic antimicrobial peptides with planar bilayers and with the cytoplasmic membrane of Escherichia coli** *Biochemistry* **38**:7235–7242 <https://doi.org/10.1021/bi9826299>

Yu F, Haynes SE, Teo GC, Avtonomov DM, Polasky DA, Nesvizhskii AI (2020) **Fast Quantitative Analysis of timsTOF PASEF Data with MSFragger and IonQuant** *Molecular and Cellular Proteomics* **19**:1575–1585 <https://doi.org/10.1074/mcp.TIR120.002048>

Editors

Reviewing Editor

Sophie Helaine

Harvard Medical School, Boston, United States of America

Senior Editor

Wendy Garrett

Harvard T.H. Chan School of Public Health, Boston, United States of America

Reviewer #2 (Public Review):

Summary:

This interesting study challenges the dogma regarding the link between bacterial metabolism decrease and tolerance to aminoglycosides (AG). The authors demonstrate that mutants well-known for being tolerant to AG, such as those of complexes I and II, are not so due to a decrease in the proton motive force (PMF) and thus antibiotic uptake, as previously reported in the literature.

Strengths:

This is a complete study that employs several read-outs.

In this revised version, the authors have carefully addressed all the reviewers' comments. I appreciate the effort made in this new version to clarify that this study does not refute the PMF-dependent mechanism of aminoglycoside uptake (in the discussion_ lines 731-734_).

The addition of the requested experiments using lower concentrations of aminoglycosides is a considerable improvement as it allows for comparison with previously published results.

Author response:

The following is the authors' response to the original reviews.

Reviewer #1 (Public Review):

Summary:

In this study, the authors investigate the tolerance of aminoglycosides in E. coli mutants deleted in the Krebs cycle and respiratory chain enzymes. The motivation for this study is unclear. Transport of aminoglycosides is pmf-dependent, as the authors correctly note, and knocking out energy-producing components leads to tolerance of aminoglycosides, this has been well established. In S. aureus, clinically relevant "small colony" strains selected for in the course of therapy with aminoglycosides acquire null mutations in the biosynthesis of heme or ubiquinone, and have been studied in detail. In E. coli, such knockouts have not been reported in clinical isolates, probably due to severe fitness costs.

Response: We sincerely appreciate the time and consideration the reviewer dedicated to evaluating our manuscript. It's important to highlight that while the transport of aminoglycosides is PMF-dependent, recent studies underscore the potential role of metabolic mutations in antibiotic tolerance, a facet that warrants further investigation. For instance, the study by Henimann's and Michiels' groups explored genomic changes in *E. coli* strains (including uropathogenic UTI89 strains) subjected to daily antibiotic exposure (Van den Bergh et al., 2022). Notably, mutations predominantly occurred in genes of the *nuo* operon, a key component of *E. coli* energy metabolism, suggesting a link between metabolic adaptations and antibiotic tolerance. Furthermore, the research by Collin's group revealed previously unrecognized genes related to central metabolism (e.g., *icd*, *gltD*, *sucA*) that contribute to antibiotic resistance in *E. coli* cells exposed to multiple antibiotics, including aminoglycosides (Lopatkin et al., 2021). These findings are corroborated by the presence of similar mutations in clinical *E. coli* pathogens, as evidenced by the analysis of a large library of 7243 *E. coli* genomes from NCBI Pathogen Detection (Lopatkin et al., 2021). The clinical relevance of metabolic mutations in antibiotic tolerance is increasingly recognized, yet their underlying mechanisms remain enigmatic. Therefore, elucidating the role of metabolic pathways in conferring antibiotic tolerance is highly critical. We have updated the introduction to clearly convey our motivation in this study (see page 4).

At the same time, single-cell analysis has shown that individual cells with a decrease in the expression of Krebs cycle enzymes are tolerant of antibiotics and have lower ATP (Manuse et al., PLoS Biol 19: e3001194). The authors of the study under review report that knocking out ICD, isocitrate dehydrogenase that catalyzes the rate-limiting step in the Krebs cycle, has little effect on aminoglycoside tolerance and actually leads to an increase in the level of ATP over time. This observation does not seem to make much sense and contradicts previous reports, specifically that E. coli ICD is tolerant of antibiotics and, not surprisingly, produces less ATP (Kabir and Shimizu, Appl Micro-biol Biotechnol. 2004; 65(1):84-96; Manuse et al., PLoS Biol 19: e3001194). Mutations in other Krebs cycle enzymes, unlike ICD, do lead to a dramatic increase in tolerance of aminoglycosides according to the paper under review. This is all very confusing.

Response: Although our data cannot be directly compared to that of Kabir and Shimizu (Mohiuddin Kabir and Shimizu, 2004), due to the utilization of entirely different experimental procedures and measurement techniques, we can draw some parallels to the


study conducted by Lewis' group (Manuse et al., 2021), despite certain differences in experimental protocols. Furthermore, the reviewer has made strong assertions regarding our manuscript based on the findings of Lewis' group. Thus, we believe it's pertinent to expand our response regarding that study.

In the study of Lewis' group, bacterial cells were inoculated at a ratio of 1:100 into LB medium from an overnight culture (approximately 16 hours). Subsequently, the cultures were incubated at 37°C for approximately 2 hours, and ATP levels were measured using the BacTiter Glo kit (Promega, Madison, WI, USA). ATP levels were then normalized to cell density, determined through optical density measurements, and represented on a linear diagram. As demonstrated in Supplementary Figure S1c of their paper, there was a 10-15% reduction in normalized ATP levels in the *icd* mutant compared to the wild type. In our experiments, cells were grown for 24 hours in overnight cultures, diluted 100-fold in fresh media, and ATP levels were measured at 3, 4, 5, and 6 hours using the same kit. ATP levels were normalized to cell counts quantified by flow cytometry. Upon analyzing our data of the *icd* mutant for around 3 hours (the time point closest to that of the study of Lewis' group), we observed a reduction of approximately 15-20% (without statistical significance) in the *icd* mutant compared to the wild-type (see raw data, linear plot, and logarithmic plot below; Author response image 1), which aligns with the findings of Lewis' group.

We further investigated the gentamicin tolerance of both wild-type and *icd* mutant strains of *E. coli* BW25113 (Author response image 2). Our findings indicate that the increased sensitivity of the *icd* mutant of the MG1655 strain to gentamicin is similar to the observation in the other *E. coli* strain.


Author response image 1.

ATP levels in the *icd* mutant. ATP levels of both the mutant and wild-type strains were measured at t=3 hours of cell growth and normalized to cell counts. The figure presents the raw data (a), linear plot (b), and logarithmic plot (c) of the same dataset. This data corresponds to the first panel of Figure 3B in the manuscript.



Author response image 2.

Gentamicin tolerance of wild-type and *icd* mutant strains of *E. coli* BW25113. Both wild type and mutant strains were treated with gentamicin (50 µg/ml) for 5 hours at the mid-exponential phase. Cells were plated before and after treatment for CFU/ml counts. The dashed line represents the limit of detection. CFU: Colony forming units.



We think that there are two primary reasons why our study cannot contradict the findings of the Lewis group:

Firstly, our study cannot be directly compared to theirs, as they did not comprehensively explore the impact of gene deletions on cell metabolism beyond the measurement of ATP levels at a single time point (Manuse et al., 2021). Our study encompasses various metabolic parameters such as cellular ATP, redox status, proton motive force (PMF), intracellular pH, and drug uptake throughout the exponential and/or early stationary phase. Additionally, we conducted proteomic analysis for five different strains including mutants and wild type. Moreover, we performed pathway enrichment analysis grounded in the statistical background of the entire genome, encompassing various functional pathway classification frameworks such as Gene Ontology annotations, KEGG pathways, and Uniprot keywords. The results of these pathway enrichment analyses are now available in the Supplementary File (see Supplementary Tables 11-17 in the current manuscript). Thus, we believe it is unjust to deem our study contradictory compared to the Lewis group's study, which does not have a comprehensive analysis of the metabolism of the mutant strains they investigated.

Secondly, our study cannot be compared to that specific study (Manuse et al., 2021) due to the utilization of a distinct antibiotic (ciprofloxacin). Cell tolerance is heavily reliant on the mechanism of action of the antibiotic used. Therefore, the reviewer should have focused on studies closely related to aminoglycoside tolerance. Our study is not confusing or contradictory, as Lewis' group also demonstrated that the tolerance of the *icd* mutant to gentamicin was significantly reduced while the tolerance of other TCA cycle mutant strains was increased in a different study (Shan et al., 2015). However, they did not delve into the metabolism of these mutant strains, as we did. We now mention this point in our manuscript (see pages 14-15).

Apart from the confusing data, it is not clear what useful information may be obtained from the choice of the experimental system. The authors examine exponentially growing cells of E. coli for tolerance of aminoglycosides. The population at this stage of growth is highly susceptible to aminoglycosides, and only some rare persister cells can survive. However, the authors do not study persisters. A stationary population of E. coli is tolerant of aminoglycosides, and this is clinically relevant, but this is not the subject of the study.

Response: Respectfully, we must express our disagreement with the reviewer's comments. Our experimental system is meticulously organized and logically structured. Mutant strains such as *gltA*, *sucA*, and *nuoI* deletions exhibit increased tolerance to all aminoglycosides tested, with their fractions clearly increasing around the mid-exponential phase between 3-4 hours (refer to Figure 2B in our manuscript). This surge in tolerance is evident at the population level as well (as depicted in Figure 1A in our manuscript, where certain mutant strains demonstrate complete survival to streptomycin, with survival fractions nearing 1). Given the pronounced increase observed around the mid-exponential phase, we primarily characterize the metabolism of these cells during this growth phase.

It's essential to note that any investigation into antibiotic tolerance and/or resistance holds immense significance, regardless of the growth phase under scrutiny, as antibiotic tolerance/resistance poses a substantial healthcare challenge. Additionally, metabolic mutant strains do not necessarily entail severe fitness costs, as evidenced by Figure S2A published by the Lewis group (Manuse et al., 2021), a finding consistent with our study (see Figure 2B in our manuscript). This phenomenon could confer a survival advantage to bacterial cells, as they may acquire metabolic mutations to bolster their tolerance without incurring significant fitness costs. Furthermore, numerous studies suggest that bacterial cells may opt for the evolutionary pathway leading to increased tolerance before acquiring resistance mechanisms (Levin-Reisman et al., 2017; Santi et al., 2021). The presence of metabolic mutations in clinical

E. coli pathogens has also been confirmed through the analysis of a large library of 7243 *E. coli* genomes from NCBI Pathogen Detection by Collin's group (Lopatkin et al., 2021). Consequently, comprehending the tolerance mechanisms of metabolic mutations holds paramount importance.

References

- Levin-Reisman I, Ronin I, Gefen O, Braniss I, Shores N, Balaban NQ. 2017. Antibiotic tolerance facilitates the evolution of resistance. *Science* (1979) 355:826–830. doi:10.1126/science.aaj2191
- Lopatkin AJ, Bening SC, Manson AL, Stokes JM, Kohanski MA, Badran AH, Earl AM, Cheney NJ, Yang JH, Collins JJ. 2021. Clinically relevant mutations in core metabolic genes confer antibiotic resistance. *Science* (1979) 371. doi:10.1126/science.aba0862
- Manuse S, Shan Y, Canas-Duarte SJ, Bakshi S, Sun WS, Mori H, Paulsson J, Lewis K. 2021. Bacterial persisters are a stochastically formed subpopulation of low-energy cells. *PLoS Biol* 19. doi:10.1371/journal.pbio.3001194
- Mohiuddin Kabir M, Shimizu K. 2004. Metabolic regulation analysis of *icd*-gene knockout *Escherichia coli* based on 2D electrophoresis with MALDI-TOF mass spectrometry and enzyme activity measurements. *Appl Microbiol Biotechnol* 65:84–96. doi:10.1007/s00253-004-1627-1
- Santi I, Manfredi P, Maffei E, Egli A, Jenal U. 2021. Evolution of Antibiotic Tolerance Shapes Resistance Development in Chronic *Pseudomonas aeruginosa* Infections. doi:10.1128/mBio.03482-20
- Shan Y, Lazinski D, Rowe S, Camilli A, Lewis K. 2015. Genetic basis of persister tolerance to aminoglycosides in *Escherichia coli*. *mBio* 6. doi:10.1128/mBio.00078-15
- Van den Bergh B, Schramke H, Michiels JE, Kimkes TEP, Radzikowski JL, Schimpf J, Vedelaar SR, Burschel S, Dewachter L, Lončar N, Schmidt A, Meijer T, Fauvart M, Friedrich T, Michiels J, Heinemann M. 2022. Mutations in respiratory complex I promote antibiotic persistence through alterations in intracellular acidity and protein synthesis. *Nat Commun* 13:546. doi:10.1038/s41467-022-28141-x

Reviewer #2 (Public Review):

Summary:

This interesting study challenges a dogma regarding the link between bacterial metabolism decrease and tolerance to aminoglycosides (AG). The authors demonstrate that mutants well-known for being tolerant to AG, such as those of complexes I and II, are not so due to a decrease in the proton motive force (PMF) and thus antibiotic uptake, as previously reported in the literature.

Strengths:

This is a complete study. These results are surprising and are based on various read-outs, such as ATP levels, pH measurement, membrane potential, and the uptake of fluorophore-labeled gentamicin. Utilizing a proteomic approach, the authors show instead that in tolerant mutants, there is a decrease in the levels of proteins associated with ribosomes (targets of AG), causing tolerance.

Response: We sincerely appreciate the reviewer for taking the time to read our manuscript and offer valuable suggestions.

Weaknesses:

The use of a single high concentration of aminoglycoside: my main comment on this study concerns the use of an AG concentration well above the MIC (50 µg/ml or 25 µg/ml for uptake experiments), which is 10 times higher than previously used concentrations (Kohanski, Taber) in study showing a link with PMF. This significant difference may explain the discrepancies in results. Indeed, a high concentration of AG can mask the effects of a metabolic disruption and lead to less specific uptake. However, this concentration highlights a second molecular level of tolerance. Adding experiments using lower concentrations (we propose 5 µg/ml to compare with the literature) would provide a more comprehensive understanding of AG tolerance mechanisms during a decrease in metabolism.

Another suggestion would be to test iron limitation (using an iron chelator as DIP), which has been shown to induce AG tolerance. Can the authors demonstrate if this iron limitation leads to a decrease in ribosomal proteins? This experiment would validate their hypothesis in the case of a positive result. Otherwise, it would help distinguish two types of molecular mechanisms for AG tolerance during a metabolic disruption: (i) PMF and uptake at low concentrations, (ii) ribosomal proteins at high concentrations.

Response: While we acknowledge the intriguing possibility of exploring whether iron limitation results in a reduction of ribosomal proteins, we believe that this topic falls slightly outside the scope of our current study. This area warrants independent investigation since our current research did not specifically focus on iron-limited environments (LB medium is iron-rich, as referenced (Abdul-tehrani et al., 1999; Rodríguez-Rojas et al., 2015)). However, we fully concur with the notion that experimental outcomes may be contingent upon the concentration of aminoglycosides (AG). Hence, we repeated the critical experiments using a lower concentration of gentamicin (5 µg/mL), as suggested by the reviewer. Before delving into a discussion of these results, we wish to emphasize two key points. Firstly, the majority of our metabolic measurements, including ATP levels, redox activities, intracellular pH, and metabolomics, were conducted in mutant and wild-type cells in the absence of drugs. Our objective was to elucidate the impact of genetic perturbations of the TCA cycle on cell metabolism. Secondly, it's important to emphasize that our study does not invalidate the hypothesis that AG uptake is proton motive force (PMF)-dependent. We observed similar drug uptake across the strains tested, which is reasonable considering that their energy metabolism and PMF are not significantly altered compared to the wild type (at least we did not observe a consistent trend in their metabolic levels). Consequently, our study does not necessarily contradict with previous claims (Taber Harry W et al., 1987). We have now clarified this point in the manuscript (see pages 1 and 13).

When we employed a lower gentamicin concentration, we still noted a significant elevation in tolerance among the *gltA*, *sucA*, and *nuoI* mutant strains compared to the wild type. Also, it remained evident that the observed tolerance in the mutant strains cannot be ascribed to differences in drug uptake or impaired PMF, as the levels of drug uptake and the disruption of PMF by gentamicin (at lower concentrations) in the mutant strains were comparable to those of the wild type. Moreover, since our metabolic measurements and proteomics analyses failed to reveal any notable alterations in energy metabolism in these strains, the consistency in drug uptake levels across both mutant and wild-type strains, even at lower concentrations, further bolsters the validity of our findings obtained at higher gentamicin concentrations. The new results have been incorporated into the Supplementary file (see Supplementary Figures S1, S5, S7, and S9) and discussed throughout the manuscript.

Recommendations for the authors:

Reviewer #2 (Recommendations For The Authors):

Line 120: Luria-Bertani (LB), used Lysogeny Broth.

Line 180: "RSG dye can be reduced by bacterial reductases of PMF" to be reformulated.

Response: The suggested corrections have been incorporated into the manuscript.

References

Abdul-tehrani H, Hudson AJ, Chang Y, Timms AR, Hawkins C, Williams JM, Harrison PM, Guest JR, Andrews SC. 1999. Ferritin Mutants of *Escherichia coli* Are Iron Deficient and Growth Impaired, and fur Mutants are Iron Deficient, *Journal of Bacteriology*.

Rodríguez-Rojas A, Makarova O, Müller U, Rolff J. 2015. Cationic Peptides Facilitate Iron-induced Mutagenesis in Bacteria. *PLoS Genet* 11. doi:10.1371/journal.pgen.1005546

Taber Harry W, Mueller JP, Miller PF, Arrow AS. 1987. Bacterial Uptake of Aminoglycoside Antibiotics. *Microbiol Rev* 51:439–457. doi:10.1128/mr.51.4.439-457.1987

<https://doi.org/10.7554/eLife.94903.2.sa2>