

Bridging the 3D geometrical organisation of white matter pathways across anatomical length scales and species

Reviewed Preprint

v2 • December 20, 2024

Revised by authors

Reviewed Preprint

v1 • April 10, 2024

Hans Martin Kjer , Mariam Andersson, Yi He, Alexandra Pacureanu, Alessandro Daducci, Marco Pizzolato, Tim Salditt, Anna-Lena Robisch, Marina Eckermann, Mareike Toepperwien, Anders Bjorholm Dahl, Maria Louise Elkjær, Zsolt Illes, Maurice Ptito, Vedrana Andersen Dahl, Tim B Dyrby 

Danish Research Centre for Magnetic Resonance, Center for Functional and Diagnostic Imaging and Research, Copenhagen University Hospital Amager and Hvidovre, Hvidovre, Denmark • Department of Applied Mathematics and Computer Science, Technical University of Denmark, Kongens Lyngby, Denmark • Guangdong Provincial Engineering Research Center of Molecular Imaging, The Fifth Affiliated Hospital, Sun Yat-sen University, Zhuhai, China • ESRF - The European Synchrotron, Grenoble, France • Department of Computer Science, University of Verona, Verona, Italy • Institut für Röntgenphysik, Universität Göttingen, Göttingen, Germany • Department of Neurology, Odense University Hospital, Odense, Denmark • Institute of Molecular Medicine, University of Southern Denmark, Odense, Denmark • BRIDGE—Brain Research—Inter-Disciplinary Guided Excellence, Department of Clinical Research, University of Southern Denmark, Odense, Denmark • Rheumatology Research Unit, Odense University Hospital, Odense, Denmark • School of Optometry, University of Montreal, Montreal, Canada • Department of Neuroscience, Faculty of Health Science, University of Copenhagen, Copenhagen, Denmark

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **valuable** study presents new observations on white matter organisation at the micron scale, using a combination of synchrotron imaging and diffusion MRI across two species. Notably, the authors provide **solid** evidence for the fasciculation of axons within major fibre bundles into laminar structures, though these structures are not consistently observed across modalities or species. The study will be of general interest to neuroanatomists and those interested in white matter imaging.

<https://doi.org/10.7554/eLife.94917.2.sa3>

Abstract

We used diffusion MRI and x-ray synchrotron imaging on monkey and mice brains to examine the organisation of fibre pathways in white matter across anatomical scales. We compared the structure in the corpus callosum and crossing fibre regions and investigated the differences in cuprizone-induced demyelination mouse brains versus healthy controls. Our findings revealed common principles of fibre organisation that apply despite the varying

patterns observed across species; small axonal fasciculi and major bundles formed laminar structures with varying angles, according to the characteristics of major pathways. Fasciculi exhibited non-straight paths around obstacles like blood vessels, comparable across the samples of varying fibre complexity and demyelination. Quantifications of fibre orientation distributions were consistent across anatomical length scales and modalities, whereas tissue anisotropy had a more complex relationship, both dependent on the field-of-view. Our study emphasises the need to balance field-of-view and voxel size when characterising white matter features across length scales.

1. Introduction

The connectome is the map of axonal connections between different brain regions. Mapping of the connectome across several anatomical length scales can unravel its hierarchical organisation, which bears a close association with normal brain function and with disruptions in disease states. The major white matter pathways can be outlined at millimetre image resolution, e.g. through photographic microdissection or tractography on diffusion magnetic resonance imaging (dMRI) (K. G. Schilling et al. 2021 [\[1\]](#)). They include the association, projection, and interhemispheric pathways, which are further categorised into functionally related tracts such as the pyramidal tract and the inferior fronto occipital fasciculus. At millimetre resolution, the white matter tracts appear to form a structural backbone, but examination at higher resolution shows a particular topological and spatial organisation of the axonal fasciculi within (Sarubbo et al. 2019 [\[2\]](#)). A fasciculus is a bundle of axons that travel together over short or long distances. Its size and shape can vary depending on its internal organisation and its relationship to neighbouring fasciculi. The pyramidal tract, for example, has a topological organisation into parallel fasciculi determined by the homunculus organisation of the primary motor cortex. Hence, diseases such as amyotrophic lateral sclerosis that over time results in the loss of different types of motor control, correlate in the pyramidal tract with a topological dependent axonal degeneration (Sach et al. 2004 [\[3\]](#)).

The conventional model of white matter (WM) architecture has assumed a parallel within-tract topological organisation of fibres. Based on MRI tractography, Van Weeden proposed a sheet-like axonal organisation between tracts, with frequent right angle crossings (Wedeen et al. 2012 [\[4\]](#)), an interpretation that has been debated (Catani, Bodi, and Dell'Acqua 2012 [\[5\]](#)). Indeed, the limited spatial resolution afforded by diffusion MRI calls into question the modelling accuracy of sub-voxel fibre crossings and parallel fibres (Maier-Hein et al. 2017 [\[6\]](#)). Subsequently, Schilling et al. explored a potential resolution limit of crossing fibres by mapping the distribution of fibre orientations from histological images using the structure tensor model (K. Schilling et al. 2017 [\[7\]](#)). They found that even at the scale of isotropic 32 μm , voxels still contain many crossing fibres. Despite the limited resolution of dMRI, the water diffusion process can reveal microstructural geometrical features, such as axons and cell bodies, though these features are compounded at the voxel level. Consequently, estimating microstructural characteristics depends on biophysical modelling assumptions, which can often be simplistic due to limited knowledge of the 3D morphology of cells and axons and their intermediate-level topological organisation within a voxel. Thus, complementary high-resolution imaging techniques that directly capture axon morphology and fasciculi organisation in 3D across different length scales within an MRI voxel are essential for understanding anatomy and improving the accuracy of dMRI-based models (Alexander et al. 2019 [\[8\]](#)).

MicroCT from lab sources or large-scale synchrotron facilities has been demonstrated as useful for validation of low-resolution MRI in small samples. Combining structure tensor analysis and streamline tractography applied to the microCT, Trinkel et al. demonstrated virtual mapping of the murine brain connectome of major pathways in comparison to tractography from diffusion MRI

(Trinkle et al. 2021 [↗](#)). In their work, they did not provide insights into the organisation of interfacing fasciculus bundles and were not able to resolve finer structures such as axons when using the resolution of MRI due to limited resolution.

X-ray holographic nanotomography (XNH) is another complementary imaging technique that enables nanoscopic image resolutions of intact axonal white matter in 3D using phase-contrast synchrotron imaging. By stacking image XNH volumes, Andersson et al. obtained an anisotropic FOV of 600 micrometres, which is of comparable scale to a typical MRI voxel (M. Andersson et al. 2020 [↗](#)). Through segmentation of the larger diameter axons, they characterised the microdispersion of their trajectories using a measure of tortuosity, thereby revealing that the cross-sectional morphological changes along individual axons are imposed by structures in the local environment, i.e., cells, vacuoles, and blood vessels (M. Andersson et al. 2020 [↗](#)). Such 3D characteristics have provided new insights that improved microstructure MRI diffusion models (Lee et al. 2020 [↗](#); M. Andersson et al. 2022 [↗](#)). However, the Andersson study provided no analysis of the organisational features at the scale of axon fasciculus level due to the technical difficulty of segmenting full volumes. We thus see a need for a sufficiently high-resolution imaging modality with a larger FOV for exploring and quantifying axon fasciculus organisation (especially their interfaces) in simple and complex white matter regions. In this regard, Mortazavi et al., combined a neuronal tracer with structure tensor analysis applied to histological images to show that labelled axons of the major white matter pathways in the *centrum semiovale* cross each other in a grid-pattern (Mortazavi et al. 2018 [↗](#)). Moreover, Zhou et al. combined two different fluorescence neuronal tracers to show vertical stacking of parallel interhemispheric projections in the corpus callosum of mice (Zhou et al. 2013 [↗](#)). However, such approaches capture only a small number of labelled axons.

Recent advances in synchrotron imaging introduced hierarchical phase-contrast CT of the intact human brain, with selection of different FOVs and image resolution (down to 2.5 μm) within a single imaging session (Walsh et al. 2021 [↗](#)). Ideally, sub-micrometre imaging techniques are needed to be comparable to the microstructural features that diffusion MRI is sensitive to.

In this work, we applied a multimodal imaging approach, aiming to bridge the gap between diffusion MRI and the scale of the axonal fasciculi, aiming to reveal the 3D white matter organisation across different anatomical length scales throughout intact monkey and mouse brains, and in a murine demyelination model. We study two white matter regions of differing organisational complexity: The relatively simple and homogenous *corpus callosum* (CC), and secondly a complex crossing fibre region in the *centrum semiovale* (CS). We use the term “anatomical length scales” throughout to designate anatomical features of different magnitudes – axons, bundles of axons (fasciculi), and tracts – that may have differing scales between mouse and monkey brains. To perform this multi-scale investigation, we apply a multi-modal imaging approach combining conventional diffusion MRI with sub-micron x-ray synchrotron phase-contrast tomography at two different synchrotron facilities: the Deutsches Elektronen-Synchrotron (DESY), beamline P10; and the European Synchrotron (ESRF), beamline ID16A. After diffusion MRI, we imaged volumes of interest at DESY (550 nm voxel size) and then at ESRF (75-100 nm voxel sizes). Each step produced images with increasing resolution, but in incrementally smaller FOVs. From the diffusion MRI data, we estimated tissue anisotropy using the diffusion tensor model (Basser, Mattiello, and LeBihan 1994 [↗](#)) and micro-tensor model (Kaden et al. 2016 [↗](#)). Additionally, we modelled the fibre orientation distribution (FOD) with the multi-fibre constrained spherical deconvolution (CSD) model (Tournier, Calamante, and Connelly 2007 [↗](#)). In the x-ray synchrotron data, we applied a scale-space structure tensor analysis, which allowed for the quantification of structure tensor-derived tissue anisotropy and FOD in the same anatomical regime indirectly detected by dMRI. Additionally, we performed tractography on the main (or dominant) direction obtained from the structure tensor, and modelled axonal fasciculi based on the clustering of the tractography streamlines, with quantification of their deviations from linearity, i.e., tortuosity. Through these methods, we demonstrated organisational principles of

white matter that persists across anatomical length scales and species. These principles govern the organisation of axonal fasciculi into sheet-like laminar shapes (structures with a predominant planar arrangement). Interestingly, while these principles remain consistent, they result in varied structural organisations in different species. We found that these laminae have differing inclination angles with respect to each other, depending on their presence within parallel or crossing major pathways. Furthermore, using a mouse model of focal demyelination induced by cuprizone (CPZ) treatment, we investigate the inflammation-related influence on axonal organisation. This is achieved through the same structure tensor-derived micro-anisotropy and tractography streamline metrics.

2. Results

2.1. Multi-scale, multi-modal imaging data in the monkey CC

We first acquired an ex vivo diffusion MRI image of the whole monkey brain, which was reconstructed with isotropic $0.5 \times 0.5 \times 0.5 \text{ mm}^3$ voxel sizes (**Fig. 1A**) (see Methods section for details). We then obtained a thin needle biopsy sample of 1 mm diameter from the mid-body of the corpus callosum (CC) (**Fig. 1B**), which we imaged twice with phase contrast tomography; at DESY with 550 nm voxel size, providing fascicular resolution, and at ESRF with 75 nm voxel size, providing axonal resolution. At ESRF, we stacked four consecutive image volumes to create a large effective FOV (**Fig. 1C**). The relative sizes of an MRI voxel and the various synchrotron volumes are shown in **Figure 1D**. The dark image intensities in the synchrotron x-ray images (**Fig. 1B, C**) are due to osmium staining of the myelin.

As seen in the image acquired at DESY in **Figure 1B**, the staining only penetrated the outer rim of the biopsy sample. At this image resolution, resolvable structures include the largest axons (with diameters roughly $>2.5 \mu\text{m}$), blood vessels, clusters of cell bodies, and vacuoles. Cell bodies and vacuoles appear bright in contrast to their outlining myelinated axons, which appeared to be oriented mainly in the left-right (L-R) axis, as expected for a CC sample.

The high-resolution ESRF data has a field-of-view (FOV) of $0.21 \times 0.21 \times 0.21 \text{ mm}$. **Fig. 1B** depicts the placement of the FOV in the osmium-stained rim of the CC sample. The four stacked scans provide an extended FOV of $0.21 \times 0.21 \times 0.78 \text{ mm}$ to match better that of the DESY volume and an MRI voxel (**Fig. 1C**). In this data, it is possible to observe and quantify fine microstructural details, i.e., axon morphology, myelin thickness, nodes of Ranvier, clusters of cell bodies, vacuoles, and blood vessels (for a detailed description see (M. Andersson et al. 2020)).

2.2. Corpus Callosum: A “straight fibre” region in the monkey brain

The CC, the largest white matter tract in the brain, has high tissue anisotropy, being dominated by densely packed axon fasciculi that all run in the right-left direction between the two hemispheres (M. Andersson et al. 2020). The directional colour-coding of the three eigenvectors from the diffusion MRI tensor model suggests that the axonal directions are locally consistent and symmetric around the midsagittal plane (**Fig. 2A**). The MRI voxel corresponding to the location of the synchrotron samples is located in the mid-body, approximately 2 mm lateral to the midsagittal plane (the white box in **Fig. 2A**). In this voxel, the estimated orientations of the diffusion tensor and CSD models (**Figs. 2B** and **2C**) both have a main R-L component (red) with a strong I-S inclination (blue), and very little response in the A-P direction (green). The multi-fibre (CSD) model shows a single fibre peak, with a relatively narrow and isotropic spread of the fibre orientation distribution (FOD) when mapped onto a spherical polar histogram (**Fig. 2B**). This is consistent with a homogeneous microstructural environment dominated by densely packed parallel axons. Indeed, the diffusion tensor model predicted a high fractional anisotropy (FA) value of 0.861 (Basser and Pierpaoli 1996) and even higher FA anisotropy in the micro-tensor domain of 0.997, i.e., the micro (μ)FA calculated as (Kaden et al. 2016).




Figure 1

Sample comparison.

A) Rendering of the vervet monkey brain structural MRI close to the mid-sagittal plane. **B)** Large field-of-view (FOV) scanning of the biopsy, DESY. The small orange box near the red arrow in A) indicates the location and size of the FOV within the MRI data. **C)** The stack of 4 small FOVs scans obtained at ESRF. The cylinder in B) indicates the relative size of the FOV within the DESY scan. **D)** Relative size comparison of one MRI voxel, the biopsy sample prior to staining and fixation, and various synchrotron FOVs. For cylindrical FOVs, the first number indicates the diameter, and the second number is the height. Sample orientations are related to the whole brain in A): R: Right, L: Left, I: Inferior, S: superior, A: Anterior, P: posterior.




Figure 2

dMRI-based orientations and tensor shapes.

A) The diffusion tensor model showing principal directions in a sagittal slice of the monkey corpus callosum (CC). The voxel corresponding to the biopsy sampling location (the synchrotron FOVs) is outlined in a white box. **B)** The fibre orientation distribution (FOD) of the CSD model in the selected diffusion MRI voxel is represented as a spherical polar histogram. **C)** The corresponding glyph representations of the diffusion tensor and CSD.

Our analysis of the high-resolution DESY data of the CC sample is shown in **Figure 3** (A, C, and E). The structure tensor was estimated based on a single set of scale-space parameters, corresponding to an integration patch size of 12 microns (see methods for further details), which is sufficient to detect all tissue-relevant orientation features. Interestingly, the structure tensor directional colour map in **Fig. 3A** (yellow-green bands) indicates axonal fasciculi disposed with a mainly planar organisation, thus forming laminae with a thickness of up to 40–45 μm projecting throughout the whole cross-section of the CC sample. The laminar thickness was determined by manual measurements on laminae visually identified in the 3D volume. These laminae have inclination angles up to ~ 35 degrees from the right-left (R-L) axis. **Figure 3C** shows a subset of the structure tensors visualised as 3D glyphs overlaid onto an image from the synchrotron volume. At this image resolution, the structure tensor directions follow either blood vessels or axonal structures.

Upon mapping the structure tensor principal directions within the image volume onto a spherical polar histogram (**Fig. 3E**), we observe a FOD having a single fibre peak. The I-S inclination angle is smaller than that of CSD from diffusion MRI (**Figs. 2B, C**). **Figure 3F** illustrates the elongated shape of the structure tensor FOD towards the I-S axis, reflecting a degree of fibre dispersion originating mainly from the axonal laminar organisation visible in **Fig. 3A** (regions marked in white square).

From the ultra-high image resolution ESRF data shown in **Figure 3** (B, D), we estimated the structure tensor at different scales, corresponding to patch sizes ranging from 9.2 - 2 μm . This allowed for the detection of tissue anisotropy on different anatomical length scales, i.e., corresponding to blood vessels, large axons, and small axons. The structure tensor directional colouring (**Fig. 3B**) does not clearly capture the layered laminar organisation seen in the DESY data. This may be explained by the smaller FOV (210 μm) not covering a volume in which the laminar organisation appears.

The direction vectors of the structure tensor analysis from all four stacked ESRF volumes are converted to spherical polar coordinates in **Figure 3H** (see **supplementary Fig. S3** for each volume individually). The histogram resembles that representing the DESY data (**Fig. 3E**), except that the main direction of the ESRF data has a lower inclination angle in the I-S axis. An exact co-registration of the two synchrotron volumes was not feasible, and some discrepancies between the FODs are to be expected. The spread and shape of the FODs are quantified using the Orientation Dispersion Index (ODI) and Dispersion Anisotropy (DA) (see Methods section). For the DESY and ESRF samples, the ODI values are 0.038 and 0.069, respectively, while the DA values are 0.55 and 0.59. Despite the different FOVs, these values are similar. The low ODI indicates a high degree of axon alignment, as expected in the CC. The mid-level DA values suggest some anisotropic spread of the directions, reflecting the angled laminar organisation observed in the DESY sample. Interestingly, the DA value for the ESRF sample is almost identical, despite the laminar bands being less visually apparent.

In **Figure 3G**, we present the calculated distribution of FA values of the structure tensor for the four stacked ESRF volumes, intended to match approximately a single MRI voxel. The mean FA value of the ESRF data was higher than in the DESY data (0.67 ± 0.05 versus 0.54 ± 0.09 , respectively) and the ESRF data distribution was narrower than in the DESY data, i.e., (ESRF median/Inter quartile range (IQR): 0.68/0.06, DESY median/IQR: 0.55/0.12). The high anisotropy at the micrometre-scale image resolutions is in accord with the FA value of 0.86 and μFA of 0.99 in the dMRI voxel, as shown in **Fig. 2C**. Although the two synchrotron volumes originated from the same biopsy, the FA distributions are not identical. Visual inspection reveals that smaller axons and cell bodies are hardly distinguishable in the DESY data (**Fig. 3A**) but are clearly evident in the higher resolution ESRF data (**Fig. 3B**).




Figure 3

Structure tensor shape for corpus callosum (CC) sample.

A-B) 3D renderings from respectively the DESY and ESRF CC biopsy sample with structure tensor main direction colour coding (in accordance with the colour sphere). The white markings illustrate bands where axons are oriented at an angle compared to their surroundings, indicating a laminar organisation. The red cylinder on (A) shows the scale difference between the DESY and ESRF data. The position of the sample within the CC is marked on the MR image in **Figures 1A** and **2A**. **C-D)** Selected coronal slice overlaid with a regularly spaced subset of structure tensor glyphs, coloured according to their predominant direction. **E-F)** Spherical polar histograms of the DESY structure tensor main directions (FOD) and the corresponding glyph. **G)** Kernel density estimates of structure tensor FA values. **H-I)** Spherical histogram of the ESRF structure tensor main directions (FOD) and the corresponding glyph. NB: We exclude contributions from the voxel of blood vessels in FA histograms and FODs.

2.3. Deep white matter: A “complex tissue” region in the monkey brain

We next investigated the axonal organisation in the *centrum semiovale* (CS) in deep white matter. The region contains up to three major pathways that are potentially crossing, i.e., the corticospinal tract, the CC, and the superior-longitudinal fasciculus (Schmahmann and Pandya 2009 [\[1\]](#)). We imaged this complex tissue sample with dMRI and ultra-high resolution synchrotron imaging at the ESRF, and then applied a structure tensor analysis using the same range of scale-space parameters as for the CC sample, i.e., focusing on contrasted tissue features of size in the range of 9.2 - 2 μm .

Figure 4A [\[2\]](#) shows the CSD FOD multi-fibre reconstructions from dMRI of a region encompassing the sample puncture. Due to the image resolution of dMRI and the known regional complexity of fibre organisation, most voxels contained more than one fibre peak, which indicates the expected presence of crossing fibres. **Fig. 4A** [\[2\]](#) (white circle) highlights the dMRI voxel best matching the FOD in the synchrotron imaging volume; it contains three fibre components, as shown by the spherical polar histogram (**Fig. 4B** [\[2\]](#)) and the corresponding glyph (**Fig. 4C** [\[2\]](#)). The two largest peaks (“parallel fibres”) are S-I directed (blue) and the smaller peaks point mostly in the R-L direction (red).

The synchrotron data from *centrum semiovale* shows axons that visually tend to organise as laminae similarly to the CC. The laminar thickness ranges from as little as 5-10 μm to as much as 35-40 μm . Some laminae are predominantly populated by axons of small diameter, too small to segment individually at this image resolution. Other laminae include a mixture of populations of different diameters. Some axons intermingle with a neighbouring lamina as shown in **Figure 4E** [\[2\]](#). The laminar findings are consistent across the four stacked sub-volumes (see the supplementary animation, Video 1).

The structure tensor FOD within the four stacked ESRF volumes (**Fig. 4F, G** [\[2\]](#)) (see **supplementary Fig. S3** [\[2\]](#) for each volume individually) has one strong S-I and one weak R-L directed peak, much as seen in the diffusion MRI (**Fig. 4B, C** [\[2\]](#)). The largest discrepancy between the FODs of the two modalities lies in the presence of a secondary S-I directed peak in the dMRI data (**Fig. 4B, C** [\[2\]](#)). Interestingly, visual inspection of the colour-coded structure tensor directions in **Fig. 4E** [\[2\]](#) shows the existence of voxels whose primary direction is along the A-P axis. However, this represents a small enough portion of the volume that it does not appear as a distinct peak on the FOD.

Estimation of ODI and DA was not performed in this case, as these indices are uninterpretable when the FOD has contributions from multiple WM pathways. It was not possible to robustly isolate the different pathways at the voxel level in this sample.

Finally, the estimated FA of the diffusion tensor within the single dMRI voxel (0.35) was much lower than that in the micro-tensor (0.99). The low FA value of the diffusion tensor model is expected as it cannot handle the complex fibre crossings (Kaden et al. 2016 [\[3\]](#); Jespersen et al. 2013 [\[4\]](#); Lasič et al. 2014 [\[5\]](#)), whereas the micro-tensor is modelled to be independent of the axonal organisation.

The structure tensor FA histogram from the ESRF data (**Figure 4H** [\[2\]](#)) had a median value of 0.66 and IQR 0.07. Interestingly, the structure tensor FA histograms of the “complex region” and the “straight fibre” regions (**Figure 3G** [\[2\]](#) vs. **4H** [\[2\]](#), respectively) are almost identical, suggesting that the given scale-space structure tensor is independent of fibre organisation, similar to the diffusion micro-tensor model (μFA).




Figure 4

Results in complex monkey deep white matter (WM) region.

A) Anatomy and CSD glyphs seen in the volume surrounding the local neighbourhood of the sampled biopsy location (white circle). **B and C)** Results in the matching MRI voxel, showing FOD (CSD) and the corresponding glyph. **D)** The directional colour map used throughout the figure. **E)** Example of rendering from an ESRF volume with colouring corresponding to the structure tensor main direction. Marking with dashed lines indicates clear fasciculi. **F and G)** Structure tensor directional statistics from the stacked FOV, showing the structure tensor FOD and corresponding glyph. **H)** Structure tensor shape statistics from the stacked FOV, showing the kernel density estimate of FA values. The red curve is a copy of the FA distribution from the ESRF CC region ([Fig. 3G](#)).

2.4. Axon fasciculi trajectories in straight and complex tissue samples

To explore the macroscopic axonal organisation and trajectory variations of fasciculi at the MRI sub-voxel level, we applied streamline tractography (Tournier et al. 2019) to the main direction of the structure tensor analysis of the synchrotron data. We manually drew seeding regions to delineate where we expected axons to project towards within the sampled volume (see Methods section). To ease visualisation and quantification, we used QuickBundle clustering (Garyfallidis et al. 2012) to group neighbouring streamlines with similar trajectories into a centroid streamline. This centroid streamline serves as an approximation of the actual trajectory of a fasciculus. For simplicity, we designate such a centroid streamline as streamline.

Tractography in the DESY corpus callosum data revealed evenly distributed streamlines throughout the sample (Figure 5A). The streamlines all follow a primary L-R direction, and a large portion of the trajectories follow a small S-I bending (coloured purple), which is in good agreement with the local FODs both from structure tensor and CSD dMRI data (Figure 2 vs. 3). Depicting the portion of streamlines without a strong S-I direction component with a separate colour (Fig. 5A, cyan streamlines, and the supplementary animation, Video 2) conveys the laminar organisation also observed in the structure tensor analysis (Fig. 3A). The inclination angle between the trajectory of the two laminae is about 35 degrees.

For the ESRF data, the streamlines typically represent the trajectory of a small number of axons or, in some cases, of single large diameter axons. As such, the streamlines in ESRF data (Fig. 5C) better reflect detailed axonal trajectory variations as compared to the DESY data. For example, streamlines in the CC show non-straight trajectories that circumvent neighbourhood obstacles such as clusters of cell bodies, other axons, and blood vessels (orange structures). Interestingly, we observed a few axon fasciculi crossing the FOV orthogonally to the otherwise dominant R-L organisation. This phenomenon was particularly evident along the blood vessels, as illustrated in Figure 5C (red streamlines).

As expected, streamlines in the ESRF *centrum semiovale* data, followed the main direction of the laminae identified in the structure tensor directional map seen in Fig. 4F. Figure 5D shows the trajectory of streamlines coloured according to the local direction, i.e., R-L (red), A-P (green) and S-I (blue). Independent of the main direction, we observe streamlines skirting around local obstacles and other axons, resulting in more complex and less straight trajectories than in the CC region.

To quantify the shape of fasciculi trajectories through the image volume, we characterised each streamline with two scalar values: (1) the streamline tortuosity index, which is a unitless score in $[1, \infty]$ describing the deviation from linearity, and (2) the calculated maximum physical deviation from a straight line of each streamline, which depicts the maximal axonal “amplitude” or dispersion. Both scalar metrics are independent of the general fasciculus orientation, but reflect micro-dispersion that can impact dMRI estimations (Nilsson et al. 2012; M. Andersson et al. 2020). We estimated the distributions of tortuosity and maximum deviation for all streamlines in both the DESY and ESRF samples (Fig. 5B).

In the CC, the median tortuosity index of the distribution was almost identical in the DESY (1.01, IQR: 0.01) and ESRF sample (1.01, IQR: 0.01). A tortuosity index close to unity is equivalent to highly straight fibres, consistent with both the visual inspection of the streamlines (Fig. 5A and 5C), the estimated ODI, and likewise with our expectation for the CC.




Figure 5

Structure tensor-based tractography for the monkey samples.

A) Streamlines in the CC sample from DESY. Purple streamlines have a strong upwards directional component, unlike the cyan streamlines (assuming that streamlines travel from right to left). **B)** The statistics of streamlines quantification using tortuosity index and maximum deviation. **C)** A selection of streamlines in the CC within a single ESRF scan. Cyan streamlines have R-L as the strongest directional component, whereas red streamlines do not. The orange structure is a segmented blood vessel. **D)** Streamlines in the complex *centrum semiovale* region (CS) within a single ESRF scan. Streamlines are coloured according to their local main direction. Orange structures represent segmented blood vessels.

In contrast, the median observed maximum deviation within the samples differed distinctly between DESY (17.1 μm , IQR: 10.7 μm) and ESRF (2.8 μm , IQR: 3.1 μm). In ESRF data, the small FOV limits the analysis to capture only microscopic effects. Therefore, the maximum deviation is driven by the interaction of the fasciculus with neighbourhood obstacles, i.e., micro-dispersion. We observed maximum deviations as high as 12 μm , which were due to obstacles like blood vessels, thus in good agreement with observations reported in (M. Andersson et al. 2020). In DESY data, deviations were as high as 40 μm , which we attribute to more macroscopic effects such as the bending of pathways, as observed in **Figure 5A**. We suppose that this may arise from the U-shaped macroscopic trajectories of CC fibres towards the cortex.

In the ESRF *centrum semiovale* sample, the median tortuosity index was significantly higher and more variable (1.04, IQR: 0.07) than in the CC. This agrees with the more complex crossing fibre trajectories observed in **Fig. 5D** (and the supplementary animation, Video 3). Interestingly, the distribution of maximum deviations had a peak around 5.3 μm and a median of 7.1 μm (IQR: 9.6 μm). While it is an increase compared to the CC, the magnitude of streamline dispersion is on a comparable scale.

We conducted three non-parametric statistical tests of significance ($\alpha = 0.05$) between the two CC samples (DESY vs. ESRF) and between the two ESRF samples (CC vs. CS). These tests included: a Kolmogorov-Smirnov (two-sample) test for equality of distributions, a Wilcoxon rank sum test for equality of medians, and a Brown-Forsythe test for equality of variance (Hollander, Wolfe, and Chicken 2015; Brown and Forsythe 1974; Massey 1951). For streamline tortuosity (**Fig 5B**, top), the null hypothesis was accepted for all three tests between CC samples ($p = 0.2$, $p = 0.7$, $p = 0.2$), but rejected for all tests comparing CC and CS ($p < 0.001$). For streamline maximum deviation (**Fig 5B**, bottom), all six null hypotheses were rejected ($p < 0.001$).

2.5. Multi-scale, multi-modal imaging data in healthy and demyelination mouse brains

Finally, we investigated the organisation of fasciculi in both healthy mouse brains and a murine model of focal demyelination induced by five weeks of cuprizone (CPZ) treatment. This allowed for the exploration of the disease-related influence on axonal organisation, particularly under inflammation-like conditions with high glial cell density at the demyelination site (He et al. 2021). The experimental setup for DESY and ESRF is similar to that described for the monkey, with the exception that we did not perform dMRI and synchrotron imaging on the same brains, and only collected MRI data for healthy mouse brains. This approach allowed us to apply the same structure tensor and tractography streamline analysis used previously, but in a healthy versus disease comparison, demonstrating the methodology's ability to provide insights into pathological conditions.

The dMRI is a whole-brain FOV, whereas the FOVs of DESY and ESRF are similar to those for the monkey (see **Table 1**). The image voxels of dMRI, DESY and ESRF have isotropic side lengths of 125 μm , 550 nm and 75-100 nm respectively. Although the mouse brain image resolution of the DESY and ESRF data is similar to that acquired on the monkey, the white matter structures in the mouse brain are considerably smaller in proportion, and the resolvable anatomical length scale differs. In the mouse, the FOV of the DESY data included both the splenium region of corpus callosum and parts of the neighbouring Cingulum pathway (**Fig. 6A**). Further, in the ESRF data, the image resolution did not suffice to outline the diameters of even the largest axons (**Fig. 6B**), in contrast to the monkey CC. However, blood vessels and glial cells are outlined as the stained myelinated axons skirt around them. Unlike in the monkey samples, we did not observe vacuoles in the mouse samples.




Figure 6

Overview of the mouse datasets.

A) Location of the sample within the mouse brain and a coronal slice from the DESY dataset, with indications of anatomic regions: cc = corpus callosum, cg = cingulum, ctx = cortex, and ccp = cortical projections. The blue box indicates the size of a DWI voxel. **B)** The co-registered slice from the ESRF volume (position indicated by the red frame). Labels indicating blood vessels (bv) and cells (c). **C)** Coronal slice from the DESY dataset of a CPZ-treated mouse, with indication of anatomic regions: cc = corpus callosum, cg = cingulum, alv = alveus, hps = hippocampal striatum, hpp = hippocampal pyramidal layer, ccp = cortical projections. The dashed purple line shows a demyelinated region of the CC. **D)** The co-registered slice from the ESRF volume (indicated by the red frame). Labels indicate blood vessels (bv), cells (c), and myelin 'debris'/macrophages (m). **E)** 3D rendering of a local segmentation in the corresponding region in D). The blood vessel is coloured in red, cells in blue, and 'myelin debris' in green. This segmentation was done manually using ITK-SNAP (RRID: SCR_002010).

ID	Specimen	Synchrotron	Sample	FOV [mm]	Voxel size [nm]
1	Monkey	DESY	Corpus callosum (CC), midbody	0.54 x 0.99 x 0.99	550
2		ESRF		0.78 x 0.21 x 0.21*	100
3		ESRF	Centrum semiovale (CS)	0.78 x 0.21 x 0.21*	100
4	Mouse	DESY	Corpus callosum, splenium & cingulum	0.63 x 0.35 x 0.35	550
5		ESRF		0.24 x 0.24 x 0.24	75
6	CPZ Mouse	DESY		0.64 x 0.38 x 0.50	550
7		ESRF		0.30 x 0.30 x 0.30	100

Table 1

Data samples included in this study.

All voxel sizes are isotropic. The FOVs of ID-2 and ID-3 are given as four stitched scans (marked with *). Individual FOVs were 0.21 x 0.21 x 0.21 mm.

DESY and ESRF data both revealed demyelinated regions in the CC of the cuprizone-treated mouse. The lesion area is demarcated by an intensity gradient increasing from dark (intact myelination) toward a brighter signal (demyelination) (**Fig. 6C, D**). In the demyelinated area, the ESRF data revealed a higher density of cell bodies and small accumulations of heavily stained material. Cellular structures were manually segmented in a small representative region to reveal their 3D shapes (**Fig. 6E**).

Preservation of the organisation of major pathways in healthy and demyelination mouse brains

In dMRI of the healthy mouse brain, the colour-coded diffusion tensor directions of the first (major) principal direction have the same R-L direction (red) as seen in the monkey (**Fig. 7A** vs. **2A**). However, **Figure 7A** shows a shift of the second and third principal directions in the mouse CC, as compared to the monkey.

In the healthy and demyelination mouse brains, we use only a single scale-space level for the structure tensor patch size (see method section) both in the DESY and ESRF data sets. The integration patch sizes correspond to 18.7 μm for the DESY and 9.4 μm for the ESRF samples (see **Table 2**).

In the DESY data from healthy mice, the directional colour coding of the structure tensor was in accord with the diffusion tensor MRI (**Figure 7A** vs. **B**). We found sporadic directional deviations in the coronal view, represented as local regions with different directional colours in **Fig. 7B**, which indicate local crossing axon fasciculi and/or blood vessels. Nevertheless, visualisation of orientations did not reveal any axonal organisation in the mouse CC due to the lack of local angular contrast, unlike the clear laminar structures seen in the monkey sample (**Fig. 3A**). Any parallel organisation in tissue remains undetectable because our visual contrast relies on angular differences.

The two major pathways, the CC and cingulum, appear in the mouse as two sharply separated structures, facilitating the generation of image masks for regionally independent quantifications of the structure tensor analysis. These structures cross almost orthogonally, as shown in the 3D glyphs of the structure tensor FODs from both the ESRF and DESY datasets (**Fig. 7E**). In all cases, the ODI were estimated to be below 0.06, indicating highly directed pathways, similar to those values measured in the monkey CC. The DA indices show more variation: In the CC, DA was 0.49 and 0.32 for the DESY and ESRF samples, respectively, and in the cingulum, 0.78 and 0.29. These values are on the same scale as those for the monkey and are non-zero, suggesting some axonal organisation is present, even though it was not visually confirmed.

The structure tensor FA histograms of the mouse CC and cingulum, are both shifted toward lower values in the DESY data as compared to the ESRF data (**Fig. 7G**). The two histograms overlap in the DESY data, whereas the ESRF data shows a shift towards higher FA values in the cingulum as compared to CC, indicating a dependency of the estimated FA on image resolution relative to the size of structures. From the ESRF data we were able to roughly segment cell clusters and blood vessels. Since the volume consists of about 13.5% cells/blood vessels in the cingulum, versus only 8.5% in CC, the higher density of extra-axonal structures may factor into the differences in estimated FA values.

In the CPZ-treated mouse brain, the directional colour-coded structure tensors in the DESY data match that in the healthy mouse, even in the demyelination region, which shows reduced image contrast (**Fig. 7C** vs. **D**). The fact that there even is directional contrast in the demyelinated region, depicts the advantage of phase-contrast imaging, where the obtained contrast is not solely dependent on absorption. Note that the demyelination region is confined to axons within the CC, with sparing of the cingulum. The lower image contrast (lack of strong edges) in the demyelinated




Figure 7

Mouse diffusion tensor and structure tensor results

A) and B): The directional components of the diffusion tensor and the structure tensor in a coronal slice from a healthy mouse brain. The white frame in (A) indicates the approximate size and location of the DESY FOV, i.e., the black frame in (B). **C) and D)** The main structure tensor direction from the DESY data overlaid on a slice from a healthy mouse brain (C) and a CPZ-treated mouse (D). **E)** Structure tensor FODs from a healthy mouse along with corresponding glyphs. The glyph colouring indicates whether the FOD contribution is from the CC (red/blue) or cingulum (yellow/cyan). **F)** structure tensor FODs from the CPZ-treated mouse CC (DESY), split into contributions from a normal appearing region and a demyelinated area. **G) and H)** structure tensor FA distributions from various regions of healthy and lesioned mouse samples.

ID	Voxel size [nm]	Scaling factor	ST-parameters (ρ, σ) [voxels]		ST conversion (γ)	Kernel size [μm]
1	550	2	2.5	0.5	0.30	12
2	100	4	8 scales*	8 scales*	0.30	[9.2 - 2]
3	100	4	8 scales*	8 scales*	0.30	[9.2 - 2]
4	550	2	4	1	0.35	18.7
5	75	5	6	1	0.25	9.4
6	550	2	4	1	0.35	18.7

* Scale space structure tensor parameters:

$\rho = [5.50, 4.50, 3.50, 3.50, 2.50, 2.50, 1.50, 1.00]$

$\sigma = [3.00, 2.75, 2.50, 1.50, 1.50, 1.00, 1.00, 0.50]$

Table 2

Structure tensor parameter values used for the samples in this study (see [Table 1](#) for sample ID). The kernel size represents the width of the ρ -kernel converted to physical distance in accordance with the voxel size of the specific dataset.

region contributes to the blurrier tensor shapes and lower FA values as compared to the normal-appearing white matter (mean FA: 0.44 vs. 0.52), as shown in **Fig. 7H**. The FODs are similar (**Fig. 7F**) with ODI = 0.03 and 0.04 for the normal-appearing and lesioned regions respectively as expected for the highly aligned axons in CC. In the same order DA = 0.18 and 0.40.

Axon fasciculi trajectories in healthy and demyelination mouse brains

For better comparisons, we registered the ESRF data of mouse CC onto the DESY data. In healthy and cuprizone-lesioned mice, **Figure 8A, B** shows that the CC and cingulum are both mappable using streamlines. Irrespective of image resolution, the cingulum did have a lower tortuosity and maximal deviation than the CC (**Fig. 8C, E**), which could be due to differences in densities of extra-axonal structures. This is suggested by the streamlines projecting through the demyelinated region (**Fig. 8B**), which contains more cells (**Fig. 6D, E**). Indeed, both indices broadened with lower image resolution in combination with a larger FOV that covers macroscopic shape changes. An example is the bending of corpus callosum detected by the longer streamlines.

Like before, we conducted three non-parametric statistical tests of significance ($\alpha = 0.05$): a Kolmogorov-Smirnov test, a Wilcoxon rank-sum test, and a Brown-Forsythe test. These tests compared measurements between the two pathways from each dataset (CC vs. Cg) and within the same pathway across different modalities (DESY vs. ESRF). For tortuosity (**Fig. 8C**), the null hypotheses for distribution and median equality were rejected in all cases ($p < 0.001$). The null hypotheses for variance equality were accepted when comparing the CC across samples ($p = 0.35$, red vs. blue) and comparing the CC and cingulum in the DESY dataset ($p = 0.66$, red vs. yellow), but were otherwise rejected ($p < 0.001$). For maximum deviation (**Fig. 8E**), all 12 null hypotheses were rejected ($p < 0.001$).

Interestingly, both in healthy and demyelination samples, some streamlines from the CC corpus deviate to project into the cortex, as illustrated in **Fig. 6** and **Fig. 8A, B** (pink extensions of streamlines). Those that project into the cortex also tend to originate from a more superiorly placed seed point. As such, the streamlines visually present a vertically layered organisation within the CC.

3. Discussion

By combining diffusion MRI and high-resolution phase-contrast synchrotron imaging, we quantified the 3D organisation of fibre pathways in white matter across a range of anatomical length scales. Importantly, our findings revealed common principles of fibre organisation in both monkeys and mice despite differences between species; small axonal fasciculi and major bundles formed sheet-like laminar structures, with relative angle that depended on the characteristics of the major pathway to which they belonged. By applying a scale-space structure tensor analysis and streamline tractography to the synchrotron datasets, we quantified the micro-dispersion of individual axons and fasciculi. Interestingly, the dispersion magnitude is indicative of fasciculi that skirt around obstacles in the white matter such as cells and blood vessels, and the results are observed across both white matter complexity (straight vs crossing fibre region) and pathology. Our dMRI and scale-space structure tensor analysis allowed us to compare and quantify tissue anisotropies and fibre orientations. We found that FODs were comparable across image resolutions and modalities. The observed discrepancies can be attributed to the fact that the FOVs are not exactly matched. Estimates of structure tensor derived microscopic FA show a clear pattern across modalities. However, to achieve good specificity in the anatomical feature correlation the diffusion tensor and structure tensor models must be sensitive to the same anatomical length scale.




Figure 8

Mouse tractography results.

A) Tractography visualisation from a healthy mouse brain sample. Red and Blue: Streamlines in CC. Yellow and Cyan: Streamlines in cingulum. Pink streamlines are extrapolations of CC streamlines that represent plausible cortical projections through the cingulum. **B)** Tractography visualisation from DESY data of a CPZ-treated mouse brain. **C) and E)** Tractography streamline centroid statistics from the healthy mouse brain synchrotron volumes. The curves are kernel density estimates of the tortuosity and maximum deviation, respectively. **D) and F)** Statistics for the streamlines in the corpus callosum of the DESY healthy and CPZ mouse brain samples.

Laminar organisation across image resolutions and modalities

Our white matter samples from monkey and mouse brains were geometrically organised as stacked laminae across anatomical length scales extending from individual axons to axon fasciculi and the major WM pathways.

In the monkey CC DESY data, which has a field of view (FOV) comparable to a dMRI voxel, a columnar laminar organisation at a macroscopic level was visually revealed from the structure tensor (ST) direction colouring. However, this laminar organisation was not visible in the higher-resolution ESRF data for the same tissue sample. Although the two samples were not co-registered, the size of a single ESRF FOV within the DESY sample is illustrated in [Fig. 3A](#). This demonstrates the possibility of placing the ESRF sample where the observed laminar structure is absent. Consequently, knowledge of the tissue structural organisation and its orientation is important to fully benefit from the stacked FOV of the ESRF sample and when choosing appropriate minimal FOV sizes in future experiments.

Interestingly, when characterising FODs with measures like ODI and DA as indicators of fibre organisation, rather than relying on visualisation, results from large- and small-FOV data show no discrepancies. This statistical approach discards the spatial context (visually perceived as laminae), highlighting the need to combine both methods.

Invasive tracer studies and histology in monkeys ([Caminiti et al. 2009](#); [Howard et al. 2023](#)) and histology of humans ([Mollink et al. 2017](#)) previously hinted at a columnar organisation in the ventral part of midsagittal CC. However, the earlier approaches failed to show the finer 3D columnar organisation of 10 µm thick laminae as revealed by the present structure tensor analysis of the synchrotron data. The occurrence of these thin layers with slightly different angles may be confirmed by visual inspection of polarised light imaging (PLI) data collected in a human brain coronal CC slice (in [Fig. 6](#) of ([Mollink et al. 2017](#))). In a coronal view of the mid-sagittal region, PLI showed fibres arising from the right and left hemispheres reaching the midsagittal CC as a layered structure, based on its orientational colour coding. The greater slice thickness (in PLI 100 µm) in a coronal view may have resulted in a partial volume effect arising from the thinner non-parallel columnar laminar sheets that we observed in the sagittal view of our synchrotron data ([Fig. 3A](#)).

Structure tensor analysis of the present ESRF data in the complex fibre region clearly shows the crossing of three pathways composed of laminar sheets of fasciculi ([Fig. 4E](#)). Compared to the parallel within-tract axonal fasciculi in the monkey CC, the boundaries between fasciculi in the complex fibre region are thin (10s of micrometres), and therefore likely to be less well defined when visualised as streamlines. However, we did not see such an intermingling of streamlines at the interface between the crossing of the larger well-defined tracts such as the CC and cingulum in the mouse brain ([Fig. 7B-D](#)).

The inclination angle between fasciculi

The structure tensor method performed very well in analysing the fasciculi without segmentation, through its detection of tracts and fasciculi based on their angular differences and highlighting of their sheet-like and laminar organisation. The observed tendencies regarding fasciculi organisation can be grouped into two: Within-WM pathways where inclination angles are low, including 0-degree parallel alignment, and between-WM pathways where the angles are high.

Within major WM pathways, neighbouring axonal fasciculi appeared to follow the same principal direction but could have inclination angles up to the order of 35 degrees in both species.

In the monkey CC (mid-body), we observed laminar organisation indicated by clear spatial angular differences in the ST directions in the sample (**Fig. 3A**). Quantifications of the FOD shape showed DA indices of 0.55 and 0.59 for the DESY and ESRF samples, respectively. In contrast, the mouse CC (splenium) did not visually reveal a similar angled laminar organisation (**Fig. 7C**), and the DA indices were lower, at 0.49 and 0.32, respectively. Two possible explanations exist. First, the within-pathway laminar organisation may not be identical across the entire CC. Consequently, more scans from other CC regions would be required to confirm. Second, the different species might account for the differences. Larger brains like the monkey might foster a different level of within-pathway axon organisation compared to the smaller mouse. Although we could not visually detect laminar organisation from the colour coding of the ST direction in the mouse, the non-zero DA values suggest some level of organisation. This is supported by our streamline tractography, which indicates a vertical layered organisation (**Fig. 8A, B**). It further aligns with studies using histological tracer mapping that shows a stacked parallel organisation of callosal projections in mice, between cortex regions M1 and S1 (Zhou et al. 2013). Nevertheless, we cannot rely solely on voxel-wise ST directions to fully describe axonal organisation, as this method does not contrast almost parallel fasciculi (inclination angles approaching 0 degrees). Analysing patterns in tractography streamlines would be an interesting future direction for this purpose.

Interestingly, the default multi-fibre dMRI CSD model for the monkey CC showed only a single fibre direction with an isotropic FOD, indicating a coherent fibre pathway (**Fig. 2C**). This approach did not reveal any of the layering and organisation evident from structure tensor analysis of the synchrotron data (**Fig. 3**). Nevertheless, when investigating the directions of the diffusion tensor model we see signs that the diffusion signal is indeed sensitive to the complexity of fasciculi organisation; the principal direction agrees with the CSD and is locally consistent across the entire CC. However, the 2nd and 3rd eigenvectors are also informative, in that they behave symmetrically across the midline, and are locally consistent, although their order can shift at specific locations. This is likely an indication of the local organisation, similar to PLI-based observations (Mollink et al. 2017). Additionally, Tax et al. have demonstrated the calculation of a grid-crossing sheet probability index from diffusion MRI data, which suggested the presence of sheet-like features in a crossing fibre region (Tax et al. 2016), which is in line with our findings in the synchrotron data. Note that the method by Tax et al. only detects sheet-like structures crossing on a grid and does not reveal laminar structures with lower inclination angles, as we observed in the monkey CC.

By further exploring the diffusion tensor modelling of the CC in the Human Connectome Data set (N=4, Fig. S1) we found an order of the 2nd and 3rd eigenvectors resembling that in the dMRI of the monkey. This observation supports the future exploration of the diffusion-weighted signal to reveal insights about axonal organisation within a WM pathway. For the present, we have demonstrated axonal organisation only in selected regions of the WM, but further research may establish more generalised rules governing fasciculi organisation in WM pathways, and potential species-related differences.

Between-WM pathways, the fasciculi cross with high inclination angles approaching 90 degrees. This is readily detectable in diffusion MRI multi-fibre CSD modelling (Tournier et al. 2004) and in the structure tensor analysis. We observed this right angle crossing at the *centrum semiovale*, a complex fibre region with the crossing of three different pathways (**Figure 4**). Using tracer labelling of axonal projections, Mortazav et al. revealed grid-like right-angle crossing of axons in a similar region in the monkey brain (Mortazavi et al. 2018). Tractography and micro-dissection findings of Wedeen et al. of the same region similarly showed grid-like crossing of laminar from different pathways (Wedeen et al. 2012).

Interestingly, we observed in the ESRF data a single axonal fasciculus composed of a few axons that crossed the main callosum pathway at an almost 90-degree angle running parallel to a blood vessel (**Fig. 5C**). Such phenomena may be overlooked in lower resolution data and have such a

small contribution to the diffusion MRI signal as to be dismissed as noise/artefact. Without prior knowledge of their existence, such crossings are apt to be disregarded by modelling choices, such as orders of spherical harmonics, model regularisation, and number of fibre populations.

Our findings of laminar organisation and the distinction between high and low inclination angles seem to support a simple topological rule across anatomical length scales. This holds especially for the low inclination cases, which retain their laminar organisation with minimal intermingling in the neighbourhood, meaning that we expect that topological organisation should be kept over a distance. If so, the (inclination angle) information might serve to form rules for low-resolution diffusion MRI based tractography about how best to project through crossing fibre and bottleneck regions, which is currently a source of false-positives trajectories (Maier-Hein et al. 2017 [\[1\]](#)). The reason is that standard tractography methods do not “remember” or follow anatomical organisation rules as they trace through complex regions. Our findings on pathway lamination and inclination angles—low for parallel-like trajectories and high for crossing-like trajectories—can help incorporate trajectory memory into these methods, reducing the risk of false trajectories.

We believe our observed topological rule of white matter laminar organisation can be explained by a biological principle known from studies of nervous tissue development. The first axons to reach their destination, guided by their growth cones, are known as “pioneering” axons. “Follower” axons use the shaft of the pioneering axon for guidance to efficiently reach the target region (Breau and Trembleau 2023 [\[2\]](#)). Axons can form a fasciculus by fasciculating or defasciculating along their trajectory through a zippering or unzipping mechanism, controlled by chemical, mechanical, and geometrical parameters. Zippering “glues” the axons together, while unzipping allows them to defasciculate at a low angle (Šmít et al. 2017 [\[3\]](#)). Although speculative, the zippering mechanism may be responsible for forming the laminar topology observed across length scales. The defasciculation effect can explain our results in the corpus callosum (CC) of monkeys, with laminar structures at low angles (~35 degrees) also observed by (Innocenti et al. 2019 [\[4\]](#); Caminiti et al. 2009 [\[5\]](#)), as well as in other major pathways (Sarubbo et al. 2019 [\[6\]](#)). In contrast, a fasciculation mechanism may be observed in the mouse CC (0 degrees). If the geometrical angle between two axons is high, i.e., toward 90 degrees, the zippering mechanism will not occur, and the two axons (fasciculi) will cross (Šmít et al. 2017 [\[3\]](#)). This supports our and other findings that crossing fasciculi or pathways occur at high angles toward 90 degrees in the fully matured brain (Wedeen et al. 2012 [\[7\]](#)). Once myelination begins, the zippering mechanism is lost (Šmít et al. 2017 [\[3\]](#)), suggesting that laminar topology is established at the earliest stages of brain maturation.

Sources to the non-straight trajectories of axon fasciculi

Applying streamline tractography to the main fibre orientations in the structure tensor analysis enabled a geometric quantification of the non-linearity of axon fasciculi trajectories as a tortuosity metric (Pingel et al. 2022 [\[8\]](#)). Surprisingly, we observed that the maximal amplitude of the axon fasciculi was comparable when measured in the crossing and straight fibre regions. This suggests that the geometrical variation in an axon fasciculus trajectory may hold similarly throughout the white matter.

We also detected differences in minor pathways: In the mouse brain ESRF data, axon fasciculi were straighter (i.e., having lower tortuosity) in the cingulum than those in the corpus callosum. Similarly in monkey ESRF data, fasciculi were straighter in the CC than in the crossing fibre region.

For the monkey and mouse brain samples, the distributions of maximal deviation in the ESRF and DESY data showed main peaks around 5-7 μm . This range of maximal deviation aligns with the typical radius of the cell bodies. Indeed, axons and fasciculi have been observed to skirt around

oligodendrocytes in the ESRF monkey data set (M. Andersson et al. 2020 [↗](#)). In addition, the maximal deviation is in accord with the size of a single axonal fasciculi (Fig. 5C [↗](#)) or blood vessels.

Even in the case of regional demyelination in the cuprizone-treated mice, we saw minor changes in tortuosity compared to normal axon fasciculi (Fig. 8D [↗](#)). We suppose that microglia and macrophages invading the site of demyelination are sufficiently similar in size to oligodendrocytes that they have little net effect on the distribution of tortuosity. The measured max deviation was higher in the cuprizone-treated mouse (Fig. 8F [↗](#)), but given the magnitude of approx. 30 μm , this effect is more likely a description of the macroscopic bending of the CC, rather than a result of the demyelination. As our streamlines went through both demyelinated and normal appearing regions of the CC, we experienced a similar dilemma found in MRI-based tractography applied in humans. The streamlines are based on only the direction of the tensor and are not guaranteed to be sensitive to local pathology. Additionally, demyelination is a dynamic process. We only sampled a single time point, and more must be included to explore if tractography-based analysis has the sensitivity to correlate with the temporal dimension of demyelination.

Diffusion MRI is sensitive to the micro-dispersion effects (Lee et al. 2019 [↗](#); M. Andersson et al. 2020 [↗](#)), but our findings suggest that the amplitude of micro-dispersion is primarily related to the sizes of extra-axonal structures independent of the complexity of the major white matter pathway. We also observe that the tortuosity can change between different white matter pathways. Assuming that the local distributions of cell bodies, vessels, and axon fasciculi can generalise within a pathway, then we can expect the micro-dispersion effects on the dMRI signal to be homogeneous.

Notably, high-resolution light-microscopy-techniques such as Polarised Light imaging with in-plane 5 μm and 30-100 μm thick slices (Ayer et al. 2011 [↗](#)) as well as light-sheet imaging are expected to be too low in resolution to detect the micro-dispersion effects.

Fibre orientation distributions across image resolutions depends on FOV

We found a general agreement in the FODs between the different image modalities and across image resolutions. This is in line with other studies comparing dMRI with structure tensor analysis using both single and multi-fibre models (Khan et al. 2015 [↗](#); Budde and Frank 2012 [↗](#); Leuze et al. 2021 [↗](#); K. Schilling et al. 2016 [↗](#)).

The FOD discrepancies that we did observe could be attributed to the differences in the FOVs across the imaging modalities. As shown in Fig. 1 [↗](#), we prepared biopsy samples for synchrotron imaging that was on a sufficiently large scale to cover several MRI voxels of isotropic 500 μm size. Indeed, the DESY sample covered multiple MRI voxels, except that a relatively large unstained region in the middle of the sample was not included in the generated FOD. Similarly, the four stacked image volumes of the ESRF measuring 780 μm in length by 210 μm in diameter only covered a fraction of the DESY volume and MRI image voxel. Since the FOD is dependent upon fibre organisation, the observed FOD differences reflect actual anatomical differences, for example with more volumetric weighing in one fibre direction. Therefore, although FODs are independent of image resolution, care must be taken when comparing two modalities without covering the same 3D volume, as observed in the CC (Fig. 3E, F [↗](#) vs. Fig. 3H, I [↗](#)). Examples are validation studies comparing the FOD from diffusion MRI fibre models with that derived from 2D or 3D histology which only rarely cover the same identical volume (Khan et al. 2015 [↗](#); Budde and Frank 2012 [↗](#)). In cases of differing FOV, there may be a risk of misattributing an actual difference in anatomical information across length scales as a methodological difference.

Tissue micro anisotropy across modalities, image resolutions, and pathology

We show that tissue anisotropy metrics are comparable across modalities but vary with the anatomical length scale. The simple FA metric in the diffusion tensor MRI model was, as expected, sensitive to the average tissue organisation/anisotropy in the voxel. Thus, we found higher anisotropy in the CC (**Fig. 2C**) as compared to the *centrum semiovale*, a more complex fibre region, much as observed by others (Andersen et al. 2020). In contrast, the μ FA dMRI metric (Kaden et al. 2016; Jespersen et al. 2014; Jespersen et al. 2013; Lasič et al. 2014) is sensitive to tissue anisotropy only in micro-domains of 10 μ m length scale (molecular displacement). The μ FA generates a mean anisotropy value within a voxel, returning similarly high FA values in both the CC and *centrum semiovale*, thereby confirming the expected independence of FA on the structural organisation (Andersen et al. 2020).

In the DESY and ESRF data, anisotropy measured by the structure tensor model is also determined within a micro-domain controlled by the patch-size and produces a distribution of values from the full volume. We introduced a scale-space parameter that automatically adjusts the patch-size to ensure optimal sensitivity to anisotropic features. Patch-sizes of 18 μ m for DESY and 5 μ m for ESRF data depict similarly sized micro-domains as in μ FA diffusion MR model and were shown to be largely independent of axonal organisation (**Fig. 4H** and **7G**).

The same anatomical features, namely cell membranes and myelin, dominate the contrast of diffusion vs. structure tensor techniques. In dMRI, these structures are the main sources for restricted and hindered water diffusion, and their osmium staining for the synchrotron scans provides strong image gradients modelled by the structure tensor. However, tissue preparation differs substantially for the two image modalities. In the case of ex vivo MRI, Sun et al. showed that anisotropy of perfusion fixated, hydrated tissue should closely match that in vivo (Sun et al. 2005). However, synchrotron imaging calls for an extra dehydration step before tissue embedding in EPON, which changes the intra- and extracellular volume fractions due to tissue shrinkage (Korogod, Petersen, and Knott 2015). Similarly, there are shrinkage effects between dehydrated and hydrated synchrotron image samples (Töpperwien et al. 2019). Given the large differences in tissue processing the obtained anisotropy measures naturally differ across the modalities. Nevertheless, as both the diffusion- and structure tensor models were sensitive to the same anatomical features, the observed strong correlation in FA values was expected.

Across image resolutions, the distributions of structure tensor anisotropy also changed, being generally lower and broader in DESY data compared to ESRF data. This could reflect greater partial volume effects in the lower resolution DESY data, thus decreasing the apparent separation of cells and axons. Such image blurring can change the image gradient information used by the structure tensor model, thereby making the anisotropy metric dependent on the image resolution. Therefore, structure tensor quantification of anisotropy metrics calls for invariant image resolution.

In our comparison of structure tensor anisotropy in healthy mice versus cuprizone-induced demyelination, the data clearly shows a lowered anisotropy distribution in focally demyelinated regions compared with healthy WM regions (**Fig. 7H**). We observed an increased extra-axonal content in the lesion area, thus in agreement with the observed higher density of cells in demyelinated tissue (He et al. 2021). By outlining the various cellular structures in 3D, we could visualise what we believe to be cell bodies, macrophages, and the myelin debris engulfed inside macrophages (**Fig. 6E**).

Interestingly, despite reduced anisotropy in demyelinated regions, the structure tensor still detects a clear directionally dependent anisotropy as in the healthy brain, consistent with the persistence of axons in this animal model despite demyelination (He et al. 2021 [↗](#)). We did not collect diffusion MRI for the cuprizone-lesioned brains. However, we recently reported relatively preserved μ FA values in demyelinated regions of the rat CC (He et al. 2021 [↗](#)). Similar findings were made in a group of multiple sclerosis patients compared to normal (Andersen et al. 2020 [↗](#)) supporting our synchrotron anisotropy results.

Limitations

Sample size: Increasing the number of samples across both species and examining laminar organisation at various length scales in more regions would strengthen our findings. However, securing beamtime at two different synchrotron facilities to scan the same sample with varying image resolutions is a limiting factor. Beamline development for multi-resolution experimental setups, along with faster acquisition methods, is a rapidly advancing field. For instance, the Hierarchical Phase-Contrast Tomography (HiP-CT) imaging beamline at ID-18 at the ESRF, enables multi-resolution imaging within a single session to address this challenge, though it is currently limited to a resolution of 2.5 μ m (Walsh et al. 2021 [↗](#)).

Registration: It was not possible to realise perfect correlational cell-to-cell imaging between the three very different modalities and experimental set-ups. This resulted in minor cross-modal differences between FODs, for example in **Fig. 3** [↗](#), where we see a FOD tilt difference due to either misalignment or an anatomical difference, albeit without affecting the interpretation of quantitative measures.

MRI-to-synchrotron registration is challenging due both to the large resolution difference and different contrast mechanisms. Our best option was, therefore, to rely on prior knowledge of fibre orientations to match approximately the samples. Given that we know the site of biopsy sampling, we could confine our search to a small neighbourhood of possible/potential MRI voxels and identify the one giving the best visual match of fibre orientation.

The matching between the different synchrotron volumes was easier and achievable manually for the mouse samples, as they contained large global features in the form of both the CC and cingulum and even a little beyond. Despite our efforts, we could not obtain similar matching for the monkey samples. Due to the imaging at different beamlines, the samples had been physically moved and repositioned in a new setup, thus losing alignment. This process gradually becomes easier and more streamlined as the various synchrotron beamlines develop. Walsh et al. recently succeeded in applying several zoom-ins on a low resolution overview scan of an intact brain within the same scanning session at the ESRF BM05 beamline (Walsh et al. 2021 [↗](#)). This approach removes the need to collect small selective biopsy samples for the different experiments. However, the finest pixel size in Walsh et al. was isotropic 2.5 μ m, which might be too coarse to perform the structure tensor analysis that we undertook to map fibre organisation across anatomical length scales to be compared with diffusion MRI.

Resolution: The different imaging setups each have characteristic image resolutions and native voxel sizes, but these are not the only relevant factors in this study. The targeted anatomical features are a primary consideration. Our examination of the monkey and mouse samples at the same beamlines gave approximately the same image resolution, voxel size, and FOV size. Nevertheless, the anatomical scales are quite different simply because the mouse brain is smaller, with a brain volume ratio of approximately 1:190. Thus, the same synchrotron FOV covered only a small part of the CC in the monkey, versus a large portion of the major callosal pathway and part of the cingulum in the mouse. When defining terminologies such as “high image resolution,” it is essential to relate it to the size of the anatomical structures of interest (Dyrby et al. 2014 [↗](#)).

A limiting consequence of having samples imaged at differing anatomical scales is that certain measures become inherently hard to compare in a normalised way. The tractography-based metrics—tortuosity and maximum deviation—serve as good examples of this resolution and FOV dependence. In the ESRF samples, the anatomical scale was at the level of individual axons, and the streamline metrics primarily reflect micro-scale effects from the extra-axonal environment, such as the influence of cells and blood vessels. In comparison, the larger anatomical scale in the DESY samples represents the level of fasciculi and above, with metrics influenced by macroscopic effects, such as the bending of the CC pathway. Both scales are interesting and can provide valuable insights, but caution is required when comparing the numbers, especially for cross-species studies where there is a significant difference in brain volume ratios.

Within the same species, assuming perfect co-registration of samples, it would be possible to perform correlative imaging and analysis. This would allow validation of whether tractography streamlines could be reproduced at different image resolutions within the same normalised FOV. Although this was not possible with the current data and experimental setup, it would be an interesting point to pursue in future work.

Additionally, the image resolution does not determine voxel size, since acquired data of fixed resolution can be up- or down-sampled by interpolation. Indeed, interpolation does not change the image information (Dyrby et al. 2014 [DOI](#)), while the image resolution is fundamentally defined by the “quality” of the imaging setup. As summarised in **Table 2** [DOI](#), we interpolated by down sampling for ease of processing and visualisation. Nonetheless, we continue referring to the datasets according to the original reconstructed voxel size (e.g., 75 nm of the ESRF data), as this reflects the approximate inherent image resolution. We minimised the effects of interpolation on the results by specifying the parameters and converting quantities to physical distances whenever possible. Further, we did not target anatomical features such as axon geometries, which cannot be robustly disentangled/quantified after downsampling, although other approaches may serve that purpose (M. Andersson et al. 2020 [DOI](#), 2022 [DOI](#)). Therefore, if the results are to be reproduced by others, then comparable imaging set-ups should be used.

5. Materials and methods

Monkey

The tissue came from a 32-month-old female perfusion-fixated vervet (*Chlorocebus aethiops*) monkey brain, obtained from the Montreal Monkey Brain Bank. The monkey, cared for on the island of St. Kitts, had been treated in line with a protocol approved by The Caribbean Primate Center of St. Kitts.

Mice

C57BL/6 female mice were obtained from Taconic Ltd. (Ry, Denmark). Mice were bred at the Biomedical Laboratory, University of Southern Denmark according to protocols and guidelines approved by the Danish Animal Health Care Committee (2014-15-00369). All animal experiments complied with the EU Directive 2010/63/EU for animal experiments.

Repeated oral administration of the copper chelator bis-cyclohexanone-oxalyldihydrazone (cuprizone) leads to demyelination and oligodendrocyte loss notably in the CC, and thus serves as a model for the demyelination lesions in patients with multiple sclerosis (MS) (Torkildsen et al. 2008 [DOI](#)). Cuprizone (Sigma Aldrich, MO, USA) was administered as 0.4% cuprizone in powdered standard chow to female mice aged 8-9 weeks for five weeks. Control mice were kept on a normal diet. During experiments, mice were weighed every second day to monitor the characteristic weight loss due to cuprizone exposure, with euthanasia of mice losing more than 20% of their

baseline body weight. After 5 weeks on cuprizone diet, the mice were euthanized with an overdose of pentobarbital (Glostrup Apotek, Glostrup, Denmark) followed by perfusion with DPB and 4% paraformaldehyde (PFA). The brains were stored in 4% PFA at 4°C.

Diffusion MRI

Two acquisition setups were used to collect the ex vivo diffusion MRI data sets on whole brains from the monkey (N=1) and normal mouse (N=1). The protocol for the monkey brain was from (M. Andersson et al. 2020 [\[1\]](#)) and includes three shells: b-values of [2011, 2957, 9259] s/mm², (gradient strength (G) = [300, 219, 300] mT/m, gradient duration (δ) = [5.6, 7.0, 10.5] ms, gradient separation (Δ) = [12.1, 20.4, 16.9] ms), [84, 87, 68] non-collinear diffusion encoding directions and the number of b=0 s/mm² was [15, 16, 13]. The higher b-value was adjusted to the lowered ex vivo diffusivity compared to in vivo (Dyrby et al. 2011 [\[2\]](#)). Correspondingly, the mouse brain protocol is from (Perens et al. 2023 [\[3\]](#)) and includes a single shell: b-value of 4000 s/mm² (G = 456 mT/m, δ = 5 ms, Δ = 13 ms), 60 non-collinear diffusion encoding directions and five b=0 s/mm². Indeed, whole-brain MRI scanning was performed on both the normal and cuprizone mice prior to synchrotron sample preparation. However, the MRI image quality was poor and could not be improved, as this was only realised after the tissue had been processed for synchrotron imaging. Therefore, the collected MRI mice data from (Perens et al. 2023 [\[3\]](#)) was included in the study.

Tissue preparation in both species followed a standard pipeline for diffusion MRI ex vivo (Dyrby et al. 2011 [\[2\]](#)). To reduce susceptibility artefacts and avoid air bubbles, we scanned the monkey brain in a double sealed plastic bag filled with PBS, whereas the mouse brain was kept in the skull and placed in a sealed plastic bag with PBS. The monkey brain data was collected on an experimental 4.7 Tesla Agilent MRI scanner, whereas the mouse brain data were collected on an experimental 7 Tesla Bruker Biospec MRI scanner. We used a quadrature radio frequency volume coil for the monkey, and a 2-parallel cryo-coil probe for the mouse brains. The isotropic image voxels were 0.55 mm for the monkey brain and 0.125 mm for the mouse. The monkey brain acquisition used an optimised three-shell ActiveAx MRI protocol based on a maximal gradient strength of 300 mT/m for ex vivo tissue as in (Dyrby et al. 2013 [\[4\]](#)). The mouse acquisition used a single-shell diffusion MRI protocol for diffusion tensor imaging (Perens et al. 2023 [\[3\]](#)). All whole-brain diffusion MRI data sets are available at www.drcmr.map/ [\[5\]](#).

Before local fibre modelling, the diffusion MRI datasets were denoised (Veraart, Fieremans, and Novikov 2016 [\[6\]](#)) and processed in the MRTrix3 software toolbox (RRID: SCR_006971) (M. Andersson et al. 2020 [\[1\]](#); Perens et al. 2023 [\[3\]](#)) to remove Gibbs ringing artefacts (Kellner et al. 2016 [\[7\]](#)). Then, we fitted the single-fibre diffusion tensor and the multi-fibre constrained-spherical deconvolution models in the MRTrix3 software toolbox (Tournier et al. 2019 [\[8\]](#)). For the monkey data, we fitted the constrained-spherical deconvolution using only a single-shell b-value i.e., 9686 s/mm², from which we estimated the fibre orientation distribution (Tournier et al. 2004 [\[9\]](#)). The diffusion tensors were fitted with a single-shell b-value, i.e., 2957 s/mm² for the monkey (M. Andersson et al. 2020 [\[1\]](#)) and 4000 s/mm² for the mouse (Perens et al. 2023 [\[3\]](#)). From the diffusion tensor, we estimated the three Eigenvectors representing fibre orientations as well as the fractional anisotropy (FA) metric (Basser, Mattiello, and LeBihan 1994 [\[10\]](#)). Since the FA value of microstructure anisotropy is biased by fibre architecture (M. Andersson et al. 2020 [\[1\]](#)), we also fitted a micro-tensor model to estimate μFA from the monkey three-shell diffusion MRI data set of monkey (Kaden et al. 2016 [\[11\]](#)). The diffusion tensor model assumes a single tensor per voxel, whereas the micro-tensor model assumes a micro-tensor regime with many micro-tensors on the diffusion length scale. Hence, the micro-tensor model is not sensitive to fibre organisation (Kaden et al. 2016 [\[11\]](#)) (Lasič et al. 2014 [\[12\]](#); Jespersen et al. 2013 [\[13\]](#)).

Diffusion MRI human

We used data from four subjects of the Human Connectome Project (HCP) Adult Diffusion database to compute the eigenvalues and eigenvectors of the diffusion tensor (Basser, Mattiello, and LeBihan 1994) via a two-step weighted and iterated least-squares method (Veraart et al. 2013) as implemented in MRtrix3 (Tournier et al. 2019). Data were denoised as indicated in (Pizzolato et al. 2023) using a Rician variance stabilisation transform (Foi 2011) in combination with PCA optimal shrinkage (Gavish and Donoho 2017), with subsequent application of Gibbs ringing removal (Kellner et al. 2016) and eddy current distortion correction (J. L. R. Andersson and Sotiropoulos 2016). For the estimation of the tensor, we selected only the $b=0$ and the 64 volume-directions corresponding to $b=1000\text{s/mm}^2$. The resulting eigenvector colour-coded maps (Pajevic and Pierpaoli 2000) are shown in Figure S1, which is organised similarly to **Figure 2A**.

Tissue preparation for synchrotron imaging

To perform the SRI experiments, small tissue samples extracted from the monkey and normal (N=1) and cuprizone (N=1) mouse brains were processed and embedded in EPON.

The monkey brain was sliced in the sagittal plane with a monkey brain matrix. Cylindrical samples of 1 mm diameter extracted from the mid sagittal CC and the centrum semiovale (CS) with a biopsy punch. After post-fixation in 2.5% glutaraldehyde for 24 hours, they were stained by immersion in 0.5% osmium tetroxide (OsO_4), followed by an embedding in EPON resin and shaped into blocks measuring 1 x 1 x 4 mm. For details of sample preparation, see (M. Andersson et al. 2020).

The brains of the normal and cuprizone mice were cut into 1 mm thick coronal slices using a mouse brain matrix. For the cuprizone mouse, we selected a slice where a demyelination lesion in the white matter was visible. The same slice position was selected in the normal mouse brain for comparison. After selecting the slice containing the splenium of the CC, we carefully excised the part of the splenium traversing the mid-sagittal plane and extending approximately 2 mm into the left hemisphere, using a scalpel under a microscope. The mouse brain samples were then processed as described above. Once the EPON resin had polymerized, we used a metallographic grinder to polish the blocks to have a smooth surface, a thickness of approximately 700 μm , and a length of a few millimetres.

Synchrotron imaging

The specimens were imaged at beamline ID16A of the European Synchrotron Radiation Facility (ESRF) with x-ray nano-holotomography, as described in (M. Andersson et al. 2020). In short, the samples were illuminated with a nano-focused cone (Da Silva et al. 2017) 17 keV x-ray beam. The samples were rotated over 180 degrees, and tomographic scans were acquired at four different propagation distances (Hubert et al. 2018). Each scan consisted of 1800 projections with exposure times of 0.22 s, and a pixel size of 100 nm or 75 nm, taking approximately 4 hours to acquire. Upon performing the phase retrieval and tomographic reconstruction, the resulting volumes had dimensions 2048 x 2048 x 2048 voxels. In the case of the healthy mouse sample, the reconstruction was performed in an extended FOV providing a volume of 3200^3 voxels as presented in **Table 1**. For the monkey CC and CS samples, we collected four consecutive image volumes with a small overlap to extend the total FOV.

The specimens were also imaged at Deutsches Elektronen-Synchrotron (DESY), at the synchrotron radiation nano-CT end station (GINIX) of the P10/PETRA III beamline (Salditt et al. 2015). Here, a 13.8 keV x-ray beam illuminated the EPON samples, which were rotated through 180 degrees to acquire tomographic scans at the lens-coupled detector (XSight Micron, Rigaku). Each tomographic scan consisted of 1000 projections with 20 ms of exposure. The detector was placed in the direct-contrast regime of the sample, and a phase retrieval was performed with an in-house Bronnikov-

aided correction-based algorithm (Lohse et al. 2020 [DOI](#); De Witte et al. 2009 [DOI](#)), prior to tomographic reconstruction, which produced volumes of voxel size 550 nm and variable dimensions, as presented in **Table 1** [DOI](#). Only one volume was collected per sample.

Note that several separate beamline experiments were conducted to collect the volumes listed in **Table 1** [DOI](#). In the first two experiments, samples from the monkey brain were scanned at ESRF and DESY, respectively. The samples from the mouse brain were imaged in two subsequent experiments. Consequently, the location of the identified demyelinating lesion in the cuprizone mice, which cannot be precisely controlled, did not match the location of the CC biopsies in the monkey.

Due to the size of the data selected, processed volumes, masks and results are available at <https://zenodo.org/records/10458911> [DOI](#). Other datasets can be shared on request.

Image registration

Finding spatial correspondences within and across image modalities was largely a manual process. For aligning DESY/ESRF images to MRI, photo documentation of tissue extraction sites was visually compared with the dMRI scan, and the best matching dMRI voxel or region was selected manually. For registering ESRF images to DESY scans, manual initialization in ITK-SNAP was followed by rigid registration refinement, employing rigid transformation and mutual information-based image similarity. Stitching individual ESRF fields of view involved translation-only registration, where the approximate translation shift - known from the scan setup - was manually refined using the ITK-SNAP (RRID: SCR_002010).

Structure tensor analysis

For each synchrotron volume we computed the 3D structure tensor for each voxel (Jeppesen et al. 2021 [DOI](#); Khan et al. 2015 [DOI](#)). The processing involves three steps: (1) Computation of the image gradients, by filtering with the derivative of a Gaussian kernel, where the parameter σ is the standard deviation of the Gaussian; (2) Calculating the outer product of the gradient with itself, yielding a tensor in each image voxel; (3) Aggregation of tensor information within a local neighbourhood, by filtering with a Gaussian kernel i.e., also known as the patch size, where the parameter ρ is the standard deviation of the Gaussian.

Similar to Diffusion Tensor in MRI (Basser, Mattiello, and LeBihan 1994 [DOI](#)), the eigen-decomposition of the structure tensor defines a 3D ellipsoid, whose axes are scaled according to the eigenvalues λ_1^* , λ_2^* and λ_3^* (where $\lambda_1^* > \lambda_2^* > \lambda_3^*$ and may be normalised to ensure that $\sum \lambda_i^* = 1$), and their orientations are defined by the three orthogonal eigenvectors $\mathbf{v}_1$, $\mathbf{v}_2$, and $\mathbf{v}_3$. Throughout this paper, the estimated structure tensor decomposition is by default converted to a diffusion-like tensor, as the structure tensor and the diffusion tensor are ‘inverted’ to one another (Khan et al. 2015 [DOI](#)). We use the following single parameter (γ) model for converting the eigenvalues of the structure tensor:

$$\lambda_i = e^{-\lambda_i^*/\gamma}$$

Afterwards, the values are normalised to have $\sum \lambda_i = 1$.

From the eigenvalues, the anisotropy of the structure tensor i.e., its shape is characterised by the same Fractional Anisotropy (FA) metrics as defined for the diffusion tensor (Basser, Mattiello, and LeBihan 1994 [DOI](#)) i.e.,

$$FA = \sqrt{\frac{3}{2} \frac{(\lambda_1 - \hat{\lambda})^2 + (\lambda_2 - \hat{\lambda})^2 + (\lambda_3 - \hat{\lambda})^2}{\lambda_1^2 + \lambda_2^2 + \lambda_3^2}}$$

where $\hat{\lambda} = (\lambda_1 + \lambda_2 + \lambda_3)/3$ is the mean of the three eigenvalues.

We introduce a **scale-space structure tensor** approach to ease parameter tuning and provide a scale invariant analysis. The variation of axon diameters within some of the samples makes it challenging to capture all relevant structural orientation information with a single set of (σ, ρ) -parameters, i.e., for a single patch size. Therefore, we employed a ‘scale space structure tensor’ approach (Lindeberg 1998 [DOI](#)). Here, the structure tensor is computed multiple times using a suite of varying (σ, ρ) -parameters called *scales*. Finally, for each voxel, we search for the dominating scale from which we retain the tensor. The criterion for selecting the dominant scale is based on the scale-wise relative maximum fractional anisotropy (FA) value. Let FA_j^i be the calculated FA-value in voxel j at scale i using the parameter set $(\sigma, \rho)^i$. The dominant scale index, d_j , for voxel j is then selected as

$$d_j = \operatorname{argmax}_i \left(\frac{FA_j^i}{\max_i(FA_j^i)} \right), \quad j = 1, 2, \dots, N$$

In other words, at a given scale, the maximum observed FA value across all voxels, $\max_i(FA_j^i)$, is used as a normalisation factor. With this approach, we select the structure tensor that provides the most anisotropic response across the set of manually defined scales. This ensures that the tensor-derived quantities—eigenvectors, eigenvalues, and FA—become independent of the axon diameter. Using the FA value for scale selection is reasonable, as we focus on the fibre-like axons. Figure S2 illustrates the scale-space structure tensor concept applied to the monkey CC ESRF data, where the FA of three selected scales of gradually decreasing patch sizes (scales 2, 6, and 8), are shown to emphasise distinctly different microstructural features. The dominant scale illustration shows how scales with large patch sizes are selected where the axons are large.

The scale-space approach was applied only to the monkey ESRF samples, with the range of scales listed in **Table 2** [DOI](#). In all other cases, the image resolution was too low compared to the anatomical scale for structures to visually present with distinguishably different sizes. Therefore, there was no benefit in the scale-space approach. Instead, the standard ST analysis with a fixed kernel size was used.

Structure tensor based quantifications

The ST analysis, whether using the scale-space or standard variant with a single scale, provides one tensor per voxel. Each sample is then quantified by the statistical distribution of FA values and the FOD based on the principal directions. The distributions are estimated from the combination of the following inclusion and exclusion regions of interest (ROIs).

- - Inclusion ROIs: In the mouse samples, ROIs defining the CC and the cingulum were generated, allowing for FA and FOD quantifications for each pathway separately.
- - Exclusion ROIs: In all samples, an ROI representing non-stained or non-tissue regions within the FOV was used to exclude those voxels. In the monkey samples, an ROI representing blood vessels was further applied for additional voxel exclusion. Notably, including blood vessels in the results had minimal influence, as they constituted a small percentage of the volume and it would not alter any conclusions.

The ROI segmentation masks were generated either manually using ITK-SNAP RRID: SCR_002010 or through intensity thresholding followed by morphological operations.

The statistical FA distributions were generated based on the selected voxels by making a kernel density estimation of the probability density function over the FA values.

The fibre Orientation Distribution (FOD) was generated by binning the principal (unit) vectors of all selected voxels into a spherical histogram parametrized by azimuth and elevation angles. The poles for this representation are poorly defined and hard to visualise. Therefore, the polar direction is aligned along the anatomical axis where we expect the least directional contribution for the particular sample. Additionally, since the bins in the parameterization do not have equal areas, the actual area is estimated and used to normalise the histogram, obtaining a fibre orientation based probability density function.

To characterise the FOD, we fitted a spherical Bingham distribution to the FOD using the *mtex* toolbox (<https://mtex-toolbox.github.io/BinghamODFs.html>). From the Bingham parameters, we derive two indices: the Orientation Dispersion Index (ODI), which describes the dispersion of the fibres on the surface of the unit sphere, and the Dispersion Anisotropy Index (DA), which expresses the anisotropic shape of the fibre dispersion (Tariq et al. 2016). Both metrics range from 0 to 1. A low ODI indicates a population of fibres with a narrow spread (a focused main direction), while a low DA suggests that the shape of the fibre dispersion is close to isotropic. It should be noted that the use of ODI and DA is only meaningful for a single pathway FOD.

Structure tensor based tractography

The principle and the process of structure tensor based tractography are much the same as in diffusion MRI-based tractography, but applied to the synchrotron volumes using the direction vectors of the structure tensor and the deterministic *FACT* algorithm (Tournier et al. 2019). The process is controlled by defining **seeding point regions**, **masks** for rejection, inclusion / termination, and various **streamline filtering** parameters, such as minimum length. We utilise an *a priori* anatomical understanding of the sample and its axonal organisation to manually generate the seeding region(s). For example, in the CC samples, the axons should primarily run in the L-R direction. We therefore create a mask at the left and right ends of the sample (with some margin from the sample edge). When seeding points in the left region of interest (ROI), streamlines must reach the right ROI to be included, and vice versa.

In several instances, we employed inclusion or rejection masks, which defined the permitted boundaries of travel for the streamlines. In essence, these masks should roughly represent axonal tissue segmentations. Depending on the sample, such a mask can be generated by thresholding either FA - or image intensity values, followed by morphological operations (opening, closing) to close holes and remove small spurious regions.

Streamline Clustering: The output of tractography is unstructured, thus often resulting in an overabundance of streamlines, which may be hard to interpret. It is then beneficial to apply a streamline clustering method, which can collect multiple streamlines into meaningful axonal bundles/fasciculi, while filtering away lone and spurious streamlines. To this end, we used the QuickBundles method (Garyfallidis et al. 2012), known for its simplicity and scalability, with manual selection of the distance threshold parameter for each sample individually.

Streamline Analysis: We use two different metrics to quantify the clustered tractography streamlines: Tortuosity and maximum deviation.

The tortuosity index is a single number representing the non-straightness of the trajectory of each streamline. It is calculated as the ratio of the length of a straight line, d , (between the streamline endpoints) and the piecewise length of the actual streamline trajectory, L .

$$\tau = L/d$$

The tortuosity index is bound between $[1, \infty]$. Following this definition, perfectly straight streamlines have a score of 1, and erratic streamlines have higher values. The various tortuosity indices are finally summarised in a histogram for each sample.

The maximum deviation is a supplementary index that describes for each streamline the largest observed physical deviation from a straight line between the given endpoints, which might be described as the maximum amplitude. This index may be easier to interpret in anatomical terms, compared to the unit-less tortuosity.

Similar to the tortuosity measure, we define the direct vector, $\mathbf{v}_d$, as going from the streamline starting point to the end point, i.e., $|\mathbf{v}_d| = d$. Additionally, we let $\mathbf{v}_i$ be the vector extending from the streamline starting point to either of the N streamline sampling points. We then measure the orthogonal distance from each sample point of the streamline to this direct vector by finding the projection of $\mathbf{v}_i$ onto $\mathbf{v}_d$. The maximum deviation, d_{max} , is then defined as the largest observed distance between all sample points and $\mathbf{v}_d$.

$$d_{max} = \operatorname{argmax}_i \left(\left\| \mathbf{v}_i - \frac{\mathbf{v}_i \cdot \mathbf{v}_d}{\mathbf{v}_d \cdot \mathbf{v}_d} \mathbf{v}_d \right\| \right), \quad i = 2:N-1$$

Statistical tests of significance were applied to both described streamline measures where relevant. Specifically, we conducted:

- Two-sample Kolmogorov-Smirnov tests to assess whether two selected distributions are identical. The null hypothesis is that the two selected distributions are equal (Massey 1951 [↗](#)).
- Two-sided Wilcoxon rank sum tests to compare the medians of two selected distributions. The null hypothesis is that the median values are equal (Hollander, Wolfe, and Chicken 2015 [↗](#)).
- Brown-Forsythe tests to examine the variance of the two selected distributions. The null hypothesis is that the distributions have equal variance (Brown and Forsythe 1974 [↗](#)).

All tests were performed at significance level 0.05, and we used the standard implementations in MATLAB 2021b (MathWorks, Massachusetts, USA)

Acknowledgements

We thank Susanne Sørensen for her assistance with the tissue preparation, and Johanna Perens from Gubra A/S for preparing the mouse MRI data. The authors acknowledge Professor Paul Cumming for critical reading of the manuscript.

We acknowledge DESY (Hamburg, Germany), a member of the Helmholtz Association HGF, for the provision of experimental facilities. Parts of this research were carried out at PETRA III and we would like to thank Dr. Michael Sprung for assistance in using the GINIX setup at P10. Beamtime was allocated for proposal(s) I-20170269 EC and I-20180267 EC.

We acknowledge the European Synchrotron Radiation Facility (ESRF) for provision of synchrotron radiation facilities under proposal numbers LS-2702 and LS-2840 and we would like to thank Peter Cloetens for assistance and support in using beamline ID16A.

M.A. and H.M.K. were supported by Capital Region Research Foundation Grant A5657 (principal investigator: T.B.D.). M.L.E. is grateful for the financial support from Lundbeckfonden (R347-2020-2454). Z.I. is grateful for the financial support from Lundbeckfonden R118-A11472, Scleroseforeningen A41354, and Independent Research Fund Denmark (DFF 9039-00370B). The project has received funding from the European Research Council (ERC) under the European Union's Horizon Europe research and innovation programme (grant agreement No. 101044180) (Principal Investigator: T.B.D.).

The human MRI Data were in part provided by the Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at Washington University.

Author contributions

Conceptualization: T.B.D.; Methodology: H.M.K, M.A, Y.H., A.P, A.D, M.P, T.S., A.R, M.E, M.T., A.D, M.L, Z.I, M.P, V.A., T.B.D. Investigation: H.M.K, M.A, T.B.D, Visualization: H.M.K., M.A. Supervision: T.B.D. Writing—original draft: H.M.K., M.A., T.B.D. Writing— review and editing: H.M.K, M.A, Y.H., A.P, A.D, M.P, T.S., A.R., M.E, M.T., A.D, M.L, Z.I, M.P, V.A, T.B.D.

Supplementary




Figure S1

HCP dataset diffusion tensor modelling.

The RGB colour-coded first, second, and third eigenvectors (ordered by decreasing eigenvalues) for the four subjects in both sagittal and coronal crop-out views.




Figure S2

Example of applying the scale-space structure tensor. (Top row)

A region of a slice from the ESRF monkey CC sample, where the FA is estimated from 8 scales (see [Table 2](#)), here showing only the scales 2, 6, and 8. The red transparent overlay shows the thresholded FA response ($FA > 0.7$). Notice how large, medium, and small cross-sectional axons respond differently at the different scales (represented by the blue, light blue, and orange circle, respectively). **(Bottom row, left):** The final thresholded dominant FA response ($dFA > 0.7$), shows high FA values for all axons regardless of the diameter. **(Bottom row, mid and right):** The same subset of voxels coloured according to the dominant scale index for the same 2D slice region and in a 3D rendering respectively. Notice how the low scales (large kernels) are selected for large axons, and similarly high scales (small kernels) selected for the small axons.




Figure S3

ST-derived statistics from the individual stacked FOVs in the monkey ESRF CC and CS sample. (Rows 1-4)

Individual FODs showing minor variation as the FOV placement changes. Across the CC, the most notable change is the anisotropic shape of the peak. Across the CS, the prominence of the L-R directed peak (near the red points) represents the largest variation. **(Row 5):** FA distributions of the four individual FOVs plotted together, showing almost no difference in the statistics.

References

- Alexander Daniel C., Dyrby Tim B., Nilsson Markus, Zhang Hui (2019) **Imaging Brain Microstructure with Diffusion MRI: Practicality and Applications** *NMR in Biomedicine* **32**
- Andersen Kasper Winther, Lasič Samo, Lundell Henrik, Nilsson Markus, Topgaard Daniel, Sellebjerg Finn, Szczepankiewicz Filip, Siebner Hartwig Roman, Blinkenberg Morten, Dyrby Tim B. (2020) **Disentangling White-Matter Damage from Physiological Fibre Orientation Dispersion in Multiple Sclerosis** *Brain Communications* **2**
- Andersson Jesper L. R., Sotiropoulos Stamatios N. (2016) **An Integrated Approach to Correction for off-Resonance Effects and Subject Movement in Diffusion MR Imaging** *NeuroImage* **125**:1063–78
- Andersson Mariam, Kjer Hans Martin, Rafael-Patino Jonathan, Pacureanu Alexandra, Pakkenberg Bente, Thiran Jean-Philippe, Ptito Maurice, et al. (2020) **Axon Morphology Is Modulated by the Local Environment and Impacts the Noninvasive Investigation of Its Structure-Function Relationship** *Proceedings of the National Academy of Sciences of the United States of America* **117**:33649–59
- Andersson Mariam, Pizzolato Marco, Kjer Hans Martin, Skodborg Katrine Forum, Lundell Henrik, Dyrby Tim B. (2022) **Does Powder Averaging Remove Dispersion Bias in Diffusion MRI Diameter Estimates within Real 3D Axonal Architectures?** *NeuroImage* **248**
- Axer Markus, Amunts Katrin, Grässel David, Palm Christoph, Dammers Jürgen, Axer Hubertus, Pietrzyk Uwe, Zilles Karl (2011) **A Novel Approach to the Human Connectome: Ultra-High Resolution Mapping of Fiber Tracts in the Brain** *NeuroImage* **54**:1091–1101
- Basser P. J., Mattiello J., LeBihan D. (1994) **MR Diffusion Tensor Spectroscopy and Imaging** *Biophysical Journal* **66**:259–67
- Basser P. J., Pierpaoli C. (1996) **Microstructural and Physiological Features of Tissues Elucidated by Quantitative-Diffusion-Tensor MRI** *Journal of Magnetic Resonance. Series B* **111**:209–19
- Breau Marie Anne, Trembleau Alain (2023) **Chemical and Mechanical Control of Axon Fasciculation and Defasciculation** *Seminars in Cell & Developmental Biology* **140**:72–81
- Brown Morton B., Forsythe Alan B. (1974) **Robust Tests for the Equality of Variances** *Journal of the American Statistical Association* **69**:364–67
- Budde Matthew D., Frank Joseph A. (2012) **Examining Brain Microstructure Using Structure Tensor Analysis of Histological Sections** *NeuroImage* **63**:1–10
- Caminiti Roberto, Ghaziri Hassan, Galuske Ralf, Hof Patrick R., Innocenti Giorgio M. (2009) **Evolution Amplified Processing with Temporally Dispersed Slow Neuronal Connectivity in Primates** *Proceedings of the National Academy of Sciences of the United States of America* **106**:19551–56

Catani Marco, Bodi Istvan, Dell'Acqua Flavio (2012) **Comment on 'The Geometric Structure of the Brain Fiber Pathways.'** *Science* **337**

Da Silva Julio Cesar, Pacureanu Alexandra, Yang Yang, Bohic Sylvain, Morawe Christian, Barrett Raymond, Cloetens Peter (2017) **Efficient Concentration of High-Energy X-Rays for Diffraction-Limited Imaging Resolution** *Optica* **4**:492–95

De Witte Yoni, Boone Matthieu, Vlassenbroeck Jelle, Dierick Manuel, Van Hoorebeke Luc (2009) **Bronnikov-Aided Correction for X-Ray Computed Tomography** *Journal of the Optical Society of America. A, Optics, Image Science, and Vision* **26**:890–94

Dyrby Tim B., Baaré William F. C., Alexander Daniel C., Jelsing Jacob, Garde Ellen, Søgaard Lise V. (2011) **An Ex Vivo Imaging Pipeline for Producing High-Quality and High-Resolution Diffusion-Weighted Imaging Datasets** *Human Brain Mapping* **32**:544–63

Dyrby Tim B., Lundell Henrik, Burke Mark W., Reislev Nina L., Paulson Olaf B., Ptito Maurice, Siebner Hartwig R. (2014) **Interpolation of Diffusion Weighted Imaging Datasets** *NeuroImage* **103**:202–13

Dyrby Tim B., Søgaard Lise V., Hall Matt G., Ptito Maurice, Alexander Daniel C. (2013) **Contrast and Stability of the Axon Diameter Index from Microstructure Imaging with Diffusion MRI** *Magnetic Resonance in Medicine: Official Journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine* **70**:711–21

Foi Alessandro (2011) **Noise Estimation and Removal in MR Imaging: The Variance-Stabilization Approach** *In* :1809–14

Garyfallidis Eleftherios, Brett Matthew, Correia Marta Morgado, Williams Guy B., Nimmo-Smith Ian (2012) **QuickBundles, a Method for Tractography Simplification** *Frontiers in Neuroscience* **6**

Gavish Matan, Donoho David L. (2017) **Optimal Shrinkage of Singular Values** *IEEE Transactions on Information Theory / Professional Technical Group on Information Theory* **63**:2137–52

He Yi, Aznar Susana, Siebner Hartwig R., Dyrby Tim B. (2021) **In Vivo Tensor-Valued Diffusion MRI of Focal Demyelination in White and Deep Grey Matter of Rodents** *NeuroImage. Clinical* **30**

Hollander Myles, Wolfe Douglas A., Chicken Eric (2015) **Nonparametric Statistical Methods. Wiley Series in Probability and Statistics** Hoboken, NJ: Wiley-Blackwell

Howard Amy F. D., Huszar Istvan N., Smart Adele, Cottaar Michiel, Daubney Greg, Hanayik Taylor, Khrapitchev Alexandre A., et al. (2023) **An Open Resource Combining Multi-Contrast MRI and Microscopy in the Macaque Brain** *Nature Communications* **14**

Hubert Maxime, Pacureanu Alexandra, Guilloud Cyril, Yang Yang, da Silva Julio C., Laurencin Jerome, Lefebvre-Joud Florence, Cloetens Peter (2018) **Efficient Correction of Wavefront Inhomogeneities in X-Ray Holographic Nanotomography by Random Sample Displacement** *Applied Physics Letters* **112**

Innocenti Giorgio M., Dyrby Tim B., Girard Gabriel, St-Onge Etienne, Thiran Jean-Philippe, Daducci Alessandro, Descoteaux Maxime (2019) **Topological Principles and Developmental Algorithms Might Refine Diffusion Tractography** *Brain Structure & Function* **224**:1–8

- Jeppesen N., Mikkelsen L. P., Dahl A. B., Christensen A. N., Dahl V. A. (2021) **Quantifying Effects of Manufacturing Methods on Fiber Orientation in Unidirectional Composites Using Structure Tensor Analysis** *Composites. Part A, Applied Science and Manufacturing* **149**
- Jespersen Sune Nørhøj, Lundell Henrik, Sønderby Casper Kaae, Dyrby Tim B. (2013) **Orientationally Invariant Metrics of Apparent Compartment Eccentricity from Double Pulsed Field Gradient Diffusion Experiments** *NMR in Biomedicine* **26**:1647–62
- Kaden Enrico, Kelm Nathaniel D., Carson Robert P., Does Mark D., Alexander Daniel C. (2016) **Multi-Compartment Microscopic Diffusion Imaging** *NeuroImage* **139**:346–59
- Kellner Elias, Dhital Bibek, Kiselev Valerij G., Reisert Marco (2016) **Gibbs-Ringing Artifact Removal Based on Local Subvoxel-Shifts** *Magnetic Resonance in Medicine* <https://doi.org/10.1002/mrm.26054>
- Khan Ahmad Raza, Cornea Anda, Leigland Lindsey A., Kohama Steven G., Jespersen Sune Nørhøj, Kroenke Christopher D. (2015) **3D Structure Tensor Analysis of Light Microscopy Data for Validating Diffusion MRI** *NeuroImage* **111**:192–203
- Korogod Natalya, Petersen Carl C. H., Knott Graham W. (2015) **Ultrastructural Analysis of Adult Mouse Neocortex Comparing Aldehyde Perfusion with Cryo Fixation** *eLife* **4** <https://doi.org/10.7554/eLife.05793>
- Lasič Samo, Szczepankiewicz Filip, Eriksson Stefanie, Nilsson Markus, Topgaard Daniel (2014) **Microanisotropy Imaging: Quantification of Microscopic Diffusion Anisotropy and Orientational Order Parameter by Diffusion MRI with Magic-Angle Spinning of the Q-Vector** *Frontiers of Physics* **2** <https://doi.org/10.3389/fphy.2014.00011>
- Lee Hong-Hsi, Jespersen Sune N., Fieremans Els, Novikov Dmitry S. (2020) **The Impact of Realistic Axonal Shape on Axon Diameter Estimation Using Diffusion MRI** *NeuroImage* **223**
- Lee Hong-Hsi, Yaros Katarina, Veraart Jelle, Pathan Jasmine L., Liang Feng-Xia, Kim Sungheon G., Novikov Dmitry S., Fieremans Els (2019) **Along-Axon Diameter Variation and Axonal Orientation Dispersion Revealed with 3D Electron Microscopy: Implications for Quantifying Brain White Matter Microstructure with Histology and Diffusion MRI** *Brain Structure & Function* **224**:1469–88
- Leuze C., Goubran M., Barakovic M., Aswendt M., Tian Q., Hsueh B., Crow A., et al. (2021) **Comparison of Diffusion MRI and CLARITY Fiber Orientation Estimates in Both Gray and White Matter Regions of Human and Primate Brain** *NeuroImage* **228**
- Lindeberg Tony (1998) **Feature Detection with Automatic Scale Selection** *International Journal of Computer Vision* **30**:79–116
- Lohse Leon M., Robisch Anna Lena, Töpperwien Mareike, Maretzke Simon, Krenkel Martin, Hagemann Johannes, Salditt Tim (2020) **A Phase-Retrieval Toolbox for X-Ray Holography and Tomography** *Journal of Synchrotron Radiation* **27**:852–59
- Maier-Hein Klaus H., Neher Peter F., Houde Jean-Christophe, Côté Marc-Alexandre, Garyfallidis Eleftherios, Zhong Jidan, Chamberland Maxime, et al. (2017) **The Challenge of Mapping the Human Connectome Based on Diffusion Tractography** *Nature Communications* **8**
- Massey Frank J. (1951) **The Kolmogorov-Smirnov Test for Goodness of Fit** *Journal of the American Statistical Association* **46**:68–78

- Mollink Jeroen, Kleinnijenhuis Michiel, van Cappellen van Walsum Anne-Marie, Sotiropoulos Stamatis N., Cottaar Michiel, Mirfin Christopher, Heinrich Mattias P., et al. (2017) **Evaluating Fibre Orientation Dispersion in White Matter: Comparison of Diffusion MRI, Histology and Polarized Light Imaging** *NeuroImage* **157**:561–74
- Mortazavi Farzad, Oblak Adrian L., Morrison Will Z., Schmahmann Jeremy D., Eugene Stanley H., Wedeen Van J., Rosene Douglas L. (2018) **Geometric Navigation of Axons in a Cerebral Pathway: Comparing dMRI with Tract Tracing and Immunohistochemistry** *Cerebral Cortex* **28**:1219–32
- Nilsson Markus, Lätt Jimmy, Ståhlberg Freddy, van Westen Danielle, Hagglätt Håkan (2012) **The Importance of Axonal Undulation in Diffusion MR Measurements: A Monte Carlo Simulation Study** *NMR in Biomedicine* **25**:795–805
- Pajevic S., Pierpaoli C. (2000) **Color Schemes to Represent the Orientation of Anisotropic Tissues from Diffusion Tensor Data: Application to White Matter Fiber Tract Mapping in the Human Brain** *Magnetic Resonance in Medicine: Official Journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine* **43**
- Perens Johanna, Salinas Casper Gravesen, Roostalu Urmas, Skytte Jacob Lercke, Gundlach Carsten, Hecksher-Sørensen Jacob, Dahl Anders Bjørholm, Dyrby Tim B. (2023) **Multimodal 3D Mouse Brain Atlas Framework with the Skull-Derived Coordinate System** *Neuroinformatics* **21**:269–86
- Pingel Jessica, Kjer Hans Martin, Biering-Sørensen Fin, Feidenhans'l Robert, Dyrby Tim B. (2022) **3D Synchrotron Imaging of Muscle Tissues at Different Atrophic Stages in Stroke and Spinal Cord Injury: A Proof-of-Concept Study** *Scientific Reports* **12**
- Pizzolato Marco, Canales-Rodríguez Erick Jorge, Andersson Mariam, Dyrby Tim B. (2023) **Axial and Radial Axonal Diffusivities and Radii from Single Encoding Strongly Diffusion-Weighted MRI** *Medical Image Analysis* **86**
- Sach Miriam, Winkler Gerhard, Glauche Volkmar, Liepert Joachim, Heimbach Bernhard, Koch Martin A., Büchel Christian, Weiller Cornelius (2004) **Diffusion Tensor MRI of Early Upper Motor Neuron Involvement in Amyotrophic Lateral Sclerosis** *Brain: A Journal of Neurology* **127**:340–50
- Salditt Tim, Osterhoff Markus, Krenkel Martin, Wilke Robin N., Priebe Marius, Bartels Matthias, Kalbfleisch Sebastian, Sprung Michael (2015) **Compound Focusing Mirror and X-Ray Waveguide Optics for Coherent Imaging and Nano-Diffraction** *Journal of Synchrotron Radiation* **22**:867–78
- Sarubbo Silvio, Petit Laurent, De Benedictis Alessandro, Chioffi Franco, Pito Maurice, Dyrby Tim B. (2019) **Uncovering the Inferior Fronto-Occipital Fascicle and Its Topological Organization in Non-Human Primates: The Missing Connection for Language Evolution** *Brain Structure & Function* **224**:1553–67
- Schilling Kurt, Gao Yurui, Janve Vaibhav, Stepniewska Iwona, Landman Bennett A., Anderson Adam W. (2017) **Can Increased Spatial Resolution Solve the Crossing Fiber Problem for Diffusion MRI?** *NMR in Biomedicine* **30** <https://doi.org/10.1002/nbm.3787>
- Schilling Kurt G., Rheault François, Petit Laurent, Hansen Colin B., Nath Vishwesh, Yeh Fang-Cheng, Girard Gabriel, et al. (2021) **Tractography Dissection Variability: What Happens When 42 Groups Dissect 14 White Matter Bundles on the Same Dataset?** *NeuroImage* **243**

Schilling Kurt, Janve Vaibhav, Gao Yurui, Stepniewska Iwona, Landman Bennett A., Anderson Adam W. (2016) **Comparison of 3D Orientation Distribution Functions Measured with Confocal Microscopy and Diffusion MRI** *NeuroImage* **129**:185–97

Schmahmann Jeremy, Pandya Deepak (2009) **Fiber Pathways of the Brain** USA: Oxford University Press

Šmít Daniel, Fouquet Coralie, Pincet Frédéric, Zapotocky Martin, Trembleau Alain (2017) **Axon Tension Regulates Fasciculation/defasciculation through the Control of Axon Shaft Zippering** *eLife* **6**

Sun Shu-Wei, Neil Jeffrey J., Liang Hsiao-Fang, He Yong Y., Schmidt Robert E., Hsu Chung Y., Song Sheng-Kwei (2005) **Formalin Fixation Alters Water Diffusion Coefficient Magnitude but Not Anisotropy in Infarcted Brain** *Magnetic Resonance in Medicine: Official Journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine* **53**:1447–51

Tariq Maira, Schneider Torben, Alexander Daniel C., Wheeler-Kingshott Claudia A. Gandini, Zhang Hui (2016) **Bingham-NODDI: Mapping Anisotropic Orientation Dispersion of Neurites Using Diffusion MRI** *NeuroImage* **133**:207–23

Tax Chantal M. W., Haije Tom Dela, Fuster Andrea, Westin Carl-Fredrik, Viergever Max A., Florack Luc, Leemans Alexander (2016) **Sheet Probability Index (SPI): Characterizing the Geometrical Organization of the White Matter with Diffusion MRI** *NeuroImage* **142**:260–79

Töpperwien Mareike, Markus Andrea, Alves Frauke, Salditt Tim (2019) **Contrast Enhancement for Visualizing Neuronal Cytoarchitecture by Propagation-Based X-Ray Phase-Contrast Tomography** *NeuroImage* **199**:70–80

Torkildsen O., Brunborg L. A., Myhr K-M, Bø L. (2008) **The Cuprizone Model for Demyelination** *Acta Neurologica Scandinavica. Supplementum* **188**:72–76

Tournier J-Donald, Calamante Fernando, Connelly Alan (2007) **Robust Determination of the Fibre Orientation Distribution in Diffusion MRI: Non-Negativity Constrained Super-Resolved Spherical Deconvolution** *NeuroImage* **35**:1459–72

Tournier J-Donald, Calamante Fernando, Gadian David G., Connelly Alan (2004) **Direct Estimation of the Fiber Orientation Density Function from Diffusion-Weighted MRI Data Using Spherical Deconvolution** *NeuroImage* **23**:1176–85

Tournier J-Donald, Smith Robert, Raffelt David, Tabbara Rami, Dhollander Thijs, Pietsch Maximilian, Christiaens Daan, Jeurissen Ben, Yeh Chun-Hung, Connelly Alan (2019) **MRtrix3: A Fast, Flexible and Open Software Framework for Medical Image Processing and Visualisation** *NeuroImage* **202**

Trinkle Scott, Foxley Sean, Kasthuri Narayanan, La Rivière Patrick (2021) **Synchrotron X-Ray Micro-CT as a Validation Dataset for Diffusion MRI in Whole Mouse Brain** *Magnetic Resonance in Medicine: Official Journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine* **86**:1067–76

Veraart Jelle, Fieremans Els, Novikov Dmitry S. (2016) **Diffusion MRI Noise Mapping Using Random Matrix Theory** *Magnetic Resonance in Medicine: Official Journal of the Society of Magnetic Resonance in Medicine / Society of Magnetic Resonance in Medicine* **76**:1582–93

Veraart Jelle, Sijbers Jan, Sunaert Stefan, Leemans Alexander, Jeurissen Ben (2013) **Weighted Linear Least Squares Estimation of Diffusion MRI Parameters: Strengths, Limitations, and Pitfalls** *NeuroImage* **81**:335–46

Walsh C. L., Tafforeau P., Wagner W. L., Jafree D. J., Bellier A., Werlein C., Kühnel M. P., et al. (2021) **Imaging Intact Human Organs with Local Resolution of Cellular Structures Using Hierarchical Phase-Contrast Tomography** *Nature Methods* **18**:1532–41

Wedeen Van J., Rosene Douglas L., Wang Ruopeng, Dai Guangping, Mortazavi Farzad, Hagmann Patric, Kaas Jon H., Tseng Wen-Yih I. (2012) **The Geometric Structure of the Brain Fiber Pathways** *Science* **335**:1628–34

Zhou Jing, Wen Yunqing, She Liang, Sui Ya-Nan, Liu Lu, Richards Linda J., Poo Mu-Ming (2013) **Axon Position within the Corpus Callosum Determines Contralateral Cortical Projection** *Proceedings of the National Academy of Sciences of the United States of America* **110**:E2714–23

Author information

Hans Martin Kjer

Danish Research Centre for Magnetic Resonance, Center for Functional and Diagnostic Imaging and Research, Copenhagen University Hospital Amager and Hvidovre, Hvidovre, Denmark, Department of Applied Mathematics and Computer Science, Technical University of Denmark, Kongens Lyngby, Denmark
ORCID iD: [0000-0001-7900-5733](https://orcid.org/0000-0001-7900-5733)

For correspondence: hmkj@dtu.dk

Mariam Andersson

Danish Research Centre for Magnetic Resonance, Center for Functional and Diagnostic Imaging and Research, Copenhagen University Hospital Amager and Hvidovre, Hvidovre, Denmark, Department of Applied Mathematics and Computer Science, Technical University of Denmark, Kongens Lyngby, Denmark
ORCID iD: [0000-0001-7181-3007](https://orcid.org/0000-0001-7181-3007)

Yi He

Danish Research Centre for Magnetic Resonance, Center for Functional and Diagnostic Imaging and Research, Copenhagen University Hospital Amager and Hvidovre, Hvidovre, Denmark, Guangdong Provincial Engineering Research Center of Molecular Imaging, The Fifth Affiliated Hospital, Sun Yat-sen University, Zhuhai, China

Alexandra Pacureanu

ESRF - The European Synchrotron, Grenoble, France
ORCID iD: [0000-0003-2306-7040](https://orcid.org/0000-0003-2306-7040)

Alessandro Daducci

Department of Computer Science, University of Verona, Verona, Italy
ORCID iD: [0000-0002-4677-6678](https://orcid.org/0000-0002-4677-6678)

Marco Pizzolato

Department of Applied Mathematics and Computer Science, Technical University of Denmark, Kongens Lyngby, Denmark
ORCID iD: [0000-0003-2455-4596](https://orcid.org/0000-0003-2455-4596)

Tim Salditt

Institut für Röntgenphysik, Universität Göttingen, Göttingen, Germany
ORCID iD: [0000-0003-4636-0813](https://orcid.org/0000-0003-4636-0813)

Anna-Lena Robisch

Institut für Röntgenphysik, Universität Göttingen, Göttingen, Germany

Marina Eckermann

Institut für Röntgenphysik, Universität Göttingen, Göttingen, Germany, ESRF - The European Synchrotron, Grenoble, France
ORCID iD: [0000-0001-6418-4310](https://orcid.org/0000-0001-6418-4310)

Mareike Toepperwien

Institut für Röntgenphysik, Universität Göttingen, Göttingen, Germany

Anders Bjorholm Dahl

Department of Applied Mathematics and Computer Science, Technical University of Denmark, Kongens Lyngby, Denmark
ORCID iD: [0000-0002-0068-8170](https://orcid.org/0000-0002-0068-8170)

Maria Louise Elkjær

Department of Neurology, Odense University Hospital, Odense, Denmark, Institute of Molecular Medicine, University of Southern Denmark, Odense, Denmark
ORCID iD: [0000-0002-8596-142X](https://orcid.org/0000-0002-8596-142X)

Zsolt Illes

Department of Neurology, Odense University Hospital, Odense, Denmark, Institute of Molecular Medicine, University of Southern Denmark, Odense, Denmark, BRIDGE—Brain Research—Inter-Disciplinary Guided Excellence, Department of Clinical Research, University of Southern Denmark, Odense, Denmark, Rheumatology Research Unit, Odense University Hospital, Odense, Denmark
ORCID iD: [0000-0001-9655-0450](https://orcid.org/0000-0001-9655-0450)

Maurice Ptito

School of Optometry, University of Montreal, Montreal, Canada, Department of Neuroscience, Faculty of Health Science, University of Copenhagen, Copenhagen, Denmark
ORCID iD: [0000-0002-5173-5862](https://orcid.org/0000-0002-5173-5862)

Vedrana Andersen Dahl

Department of Applied Mathematics and Computer Science, Technical University of Denmark, Kongens Lyngby, Denmark
ORCID iD: [0000-0001-6734-5570](https://orcid.org/0000-0001-6734-5570)

Tim B Dyrby

Danish Research Centre for Magnetic Resonance, Center for Functional and Diagnostic Imaging and Research, Copenhagen University Hospital Amager and Hvidovre, Hvidovre,

Denmark, Department of Applied Mathematics and Computer Science, Technical University of Denmark, Kongens Lyngby, Denmark
ORCID iD: [0000-0003-3361-9734](https://orcid.org/0000-0003-3361-9734)

For correspondence: timd@drcmr.dk

Editors

Reviewing Editor

Saad Jbabdi

University of Oxford, Oxford, United Kingdom

Senior Editor

Yanchao Bi

Beijing Normal University, Beijing, China

Reviewer #1 (Public Review):

This study presents valuable observations of white matter organisation from diffusion MRI and two types of synchrotron imaging in both monkeys and mice. Cross-modality comparisons are interesting as the different methods are able to probe anatomical structures at different length scales, from single axons in high-resolution synchrotron (ESRF) imaging, to clusters of axons in lower-resolution synchrotron (DEXY) data, to axon populations at the mm-scale in diffusion MRI. By acquiring all modalities in monkey and mouse ex vivo samples, the authors can observe principles of fibre organisation, and characterise how fibre characteristics, such as tortuosity and micro-dispersion, vary across select brain regions and in healthy tissue versus a demyelination model.

One very interesting result is the observation of apparent laminar organisation of fibres in ex vivo monkey white matter samples. DESY data from the corpus callosum shows fibres with two dominant orientations (one L-R, one slightly inclined), clustered in laminar structures within this major fibre bundle. Thanks to the authors providing open data, I was able to look through the raw DESY volume and observe regions with different "textures" (different orientations) in the described laminar arrangement. That this organisation can be observed by eye, as well as by structure tensor, is fairly convincing.

<https://doi.org/10.7554/eLife.94917.2.sa2>

Reviewer #2 (Public Review):

Summary:

In this work, the authors combine diffusion MRI and high-resolution x-ray synchrotron phase-contrast imaging in monkey and mouse brains to investigate the 3D organization of brain white matter across different scales and species. The work is at the forefront of the anatomical investigation of the human connectome and aligns with several current efforts to bridge the resolution gap between what we can see in vivo at the millimeter scale and the complexity of the human brain at the sub-micron scale. The authors compare the 3D white matter organization across modalities within 2 small regions in one monkey brain (body of the corpus callosum, centrum semiovale) and within one region (splenium of the corpus callosum) in healthy mice and in one murine model of focal demyelination. The study compares measures of tissue anisotropy and fiber orientations across modalities, performs a qualitative comparison of fasciculi trajectories across brain regions and tissue conditions using streamlined tractography based on the structure tensor, and attempts to quantify the

shape of fasciculi trajectories by measuring the tortuosity index and the maximum deviation for each reconstructed streamline. Results show measures of anisotropy and fiber orientations largely agree across modalities, especially for larger FOV data. The high-resolution data allows us to explore the fiber trajectories in relation to tissue complexity and pathology. The authors claim the study reveals new common organization principles of white matter fibers across species and scales, for which axonal fasciculi arrange into sheet-like laminar structures.

Strengths:

The aim of the study is of central importance within present efforts to bridge the gap between macroscopic structures observable in vivo in humans using conventional diffusion MRI and the microscopic organization of white matter tissue. Results obtained from this type of study are important to interpret data obtained in vivo, inform the development of novel methodologies, and expand our knowledge of the structural and thus functional organization of brain circuits.

Multi-scale data acquired across modalities within the same sample constitute extremely valuable data that is often hard to acquire and represent a precious resource for validation of both diffusion MRI tractography and microstructure methods.

The inclusion of multi-species data adds value to the study, allowing the exploration of common organization principles across species.

The addition of data from a murine cuprizone model of focal demyelination adds interesting opportunities to study the underlying biological changes that follow demyelination and how these impact tissue anisotropy and fiber trajectories. These data can inform the interpretation and development of diffusion MRI microstructure models.

[Editors' note: The Reviewing Editor considers that the authors addressed the reviewers' questions adequately. The original reviews are here: <https://elifesciences.org/reviewed-preprints/94917/reviews>]

<https://doi.org/10.7554/eLife.94917.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Reviewer #1 (Public Review):

This study presents valuable observations of white matter organisation from diffusion MRI and two types of synchrotron imaging in both monkeys and mice. Cross-modality comparisons are interesting as the different methods are able to probe anatomical structures at different length scales, from single axons in high-resolution synchrotron (ESRF) imaging, to clusters of axons in lower-resolution synchrotron (DEXY) data, to axon populations at the mm-scale in diffusion MRI. By acquiring all modalities in monkey and mouse ex vivo samples, the authors can observe principles of fibre organisation, and characterise how fibre characteristics, such as tortuosity and micro-dispersion, vary across select brain regions and in healthy tissue versus a demyelination model. The results are solid, though some statements (in the abstract/discussion) do not appear to be fully supported, and statistical tests would help confirm whether tissue characteristics are similar/different between different conditions.

R1.1: Thank you for the kind feedback. We have included statistical tests in the paper for tissue characteristics where appropriate.

Due to the very high number of sample points (one per voxel) within the 3D synchrotron volumes, testing for statistical significance is challenging for the structure tensor-based tissue fractional anisotropy (FA) metric. This causes any standard statistical test to have sufficient power to evaluate even minute differences between the volumes as statistically significant with high confidence. In other words, the null hypothesis (H0) will always be rejected with $p = 0$, regardless of the practical significance of the difference. Therefore, we have not added statistical analysis for FA results.

For the tractography based metrics, the number of sample points (one per streamline) is not as high as that for the structure tensor FA, thus making it more reasonable to test for statistical significance. The statistical analyses performed included tests for equality of distributions (Two-sample Kolmogorov-Smirnov tests), equality of medians (Two-sided Wilcoxon rank sum tests), and equality of variance (Brown-Forsythe tests). The results are described in relation to Figure 5(B, D), Figure 8(CF), and detailed in the Methods section.

One very interesting result is the observation of apparent laminar organisation of fibres in ex vivo monkey white matter samples. DESY data from the corpus callosum shows fibres with two dominant orientations (one L-R, one slightly inclined), clustered in laminar structures within this major fibre bundle. Thanks to the authors providing open data, I was able to look through the raw DESY volume and observe regions with different "textures" (different orientations) in the described laminar arrangement. That this organisation can be observed by eye, as well as by structure tensor, is fairly convincing. As not all readers will download the data themselves, the manuscript could benefit from additional figures/videos to demonstrate (1) the quality of the DESY data and (2) a more 3D visualisation of the laminar structures (where the coronal plane shows convincing columnar structure or stripes). Similarly in Figure 5A, though this nicely depicts two populations with different orientations, it is somewhat difficult to see the laminar structure in the current image.

ESRF data of the centrum semiovale (CS) contributes evidence for similar laminar structures in a crossing fibre region, where primarily AP fibres are shown to cluster in 3 laminar structures. As above, further visualisations of the ESRF volume in the CS (as shown in Figure 4E) would be of value (e.g. showing consistency across the 4 volumes, 2D images showing stripey/columnar patterns along different axes, etc).

R1.2: Conveying complex 3D geometry through 2D still images is indeed challenging, and we greatly appreciate the reviewer's comments and suggestions. To better communicate the understanding of the 3D anatomical environments, we have taken the following actions:

- (1) To enhance insights into the tractography results in Figures 5A and 5D, we have rendered and added animations of the tractography scenes as supplemental material.
- (2) To visually support 3D insights concerning the consistency of the laminar organisation of the callosal fibres, we have replaced the 2D slice views in Figures 3A and 3B with 3D renderings similar to the one in Figure 4E.
- (3) An animation of Figure 4E was created to display the colour-coded structure tensor directions of all four stacked scans. This animation visually supports the complexity of the fibre orientation and the layered structural laminar organisation of the CS sample.

A key limitation of this result is that, though the DESY data from the CC seems convincing, the same structures were not observed in high-resolution synchrotron (ESRF) data of the same tissue sample in the corpus callosum. This seems surprising and the manuscript does not provide a convincing explanation for this inconsistency. The authors argue that this is due to the limited FOV of the ESRF data (~200x200x800 microns). However, the

observed laminar structures in DESY are ~40 microns thick, and ERSF data from the CST suggests laminar thicknesses in the range of 5-40 microns with a similar FOV. This suggests the ERSF FOV would be sufficient to capture at least a partial description of the laminar organisation. Further, the DESY data from the CC shows columnar variations along the LR axis, which we might expect to be observed along the long axis of the ESRF volume of the same sample. Additional analyses or explanations to reconcile these apparently conflicting observations would be of value. For example, the authors could consider down-sampling the ESRF data in an appropriate manner to make it more similar to the DESY data, and running the same analysis, to see if the observed differences are related to resolution (i.e. the thinner laminar structures cluster in ways that they look like a thicker laminar structure at lower resolution), or crop the DESY data to the size of the ESRF volume, to test whether the observed differences can be explained by differences in FOV. Laminar structures were not observed in mouse data, though it is unclear if this is due to anatomical differences or somewhat related to differences in data quality across species.

R1.3: We have clarified and expanded upon the results regarding the laminar organisation observed in the monkey CC DESY data. As noted in R1.2, we replaced the 2D images in Figures 3A (DESY) and 3B (ESRF) with 3D renderings to better display the spatial outline of the laminar organisation in the volumes. The reviewer is correct that, although the smaller field of view (FOV) of the ESRF data should allow us to at least partially capture parts of the laminar organisation observed in the larger FOV of the DESY data, this is not guaranteed. It depends on how the smaller FOV is positioned relative to the structural organisation, and since we lack co-registration, we do not know this. It should now be visually evident that the ESRF FOV can be placed such that it does not cover the observed laminae, a point which is now also emphasised in the Discussion.

Secondly, it is important to emphasise that the voxel colouring using the primary structure tensor direction is just a visualisation technique, which has limitations when it comes to assessing laminar organisation. Mapping 3D directions to RGB colours is inherently difficult and will always have ambiguities. If we had used the standard R-G-B to LR-AP-IS colouring in Figure 3, the laminar organisation would not be evident. Additionally, the laminae will only be visible when there are clear angular differences. There can still be a layered organisation even if we don't observe it, which is the case for the mouse. The primary direction differences of these layers could be very low (i.e., parallel layers), and consequently not visually evident. This point has been clarified in both the Results and Discussion sections.

Finally, in response to R1.6, we have added analyses regarding the shape of the FOD, specifically estimating the Orientation Dispersion Index (ODI) and Dispersion Anisotropy (DA). This provides further context to the reviewer's comments about the discrepancies in laminar organisation. We have reflected on the relationship between DA and the visually observed laminar organisation, and this has been integrated into the relevant parts of the Results and Discussion sections.

The changes to manuscript reflecting the statements above are listed here:

The Discussion section (page 21): "In the monkey CC DESY data, which has a field of view (FOV) comparable to a dMRI voxel, a columnar laminar organisation at a macroscopic level was visually revealed from the structure tensor (ST) direction colouring. However, this laminar organisation was not visible in the higher-resolution ESRF data for the same tissue sample. Although the two samples were not co-registered, the size of a single ESRF FOV within the DESY sample is illustrated in Fig. 3A. This demonstrates the possibility of placing the ESRF sample where the observed laminar structure is absent. Consequently, knowledge of the tissue structural organisation and its orientation is important to fully benefit from the

stacked FOV of the ESRF sample and when choosing appropriate minimal FOV sizes in future experiments.

Interestingly, when characterising FODs with measures like ODI and DA as indicators of fibre organisation, rather than relying on visualisation, results from large- and small-FOV data show no discrepancies. This statistical approach discards the spatial context (visually perceived as laminae), highlighting the need to combine both methods.”

The Results section (page 8): “The mid-level DA values suggest some anisotropic spread of the directions, reflecting the angled laminar organisation observed in the DESY sample. Interestingly, the DA value for the ESRF sample is almost identical, despite the laminar bands being less visually apparent.”

The Results section (page 17): “Nevertheless, visualisation of orientations did not reveal any axonal organisation in the mouse CC due to the lack of local angular contrast, unlike the clear laminar structures seen in the monkey sample (Fig. 3A). Any parallel organisation in tissue remains undetectable because our visual contrast relies on angular differences.”

The Discussion section (page 22): “In the monkey CC (mid-body), we observed laminar organisation indicated by clear spatial angular differences in the ST directions in the sample (Fig. 3A). Quantifications of the FOD shape showed DA indices of 0.55 and 0.59 for the DESY and ESRF samples, respectively. In contrast, the mouse CC (splenium) did not visually reveal a similar angled laminar organisation (Fig. 7C), and the DA indices were lower, at 0.49 and 0.32, respectively. Two possible explanations exist. First, the within-pathway laminar organisation may not be identical across the entire CC. Consequently, more scans from other CC regions would be required to confirm. Second, the different species might account for the differences. Larger brains like the monkey might foster a different level of within-pathway axon organisation compared to the smaller mouse. Although we could not visually detect laminar organisation from the colour coding of the ST direction in the mouse, the non-zero DA values suggest some level of organisation. This is supported by our streamline tractography, which indicates a vertical layered organisation (Fig. 8A, B). It further aligns with studies using histological tracer mapping that shows a stacked parallel organisation of callosal projections in mice, between cortex regions M1 and S1 (Zhou et al. 2013). Nevertheless, we cannot rely solely on voxel-wise ST directions to fully describe axonal organisation, as this method does not contrast almost parallel fasciculi (inclination angles approaching 0 degrees). Analysing patterns in tractography streamlines would be an interesting future direction for this purpose.”

The authors further quantify various other characteristics of the white matter, such as micro-dispersion, tortuosity, and maximum displacement. Notably, the microscopic FA calculated via structure tensor is fairly consistent across regions, though not modalities. When fibre orientations are combined across the sample, they are shown to produce similar FODs to dMRI acquired in the same tissue, which is reassuring. As noted in the text, the estimates of tortuosity and max displacement are dependent on the FOV over which they are calculated. Calculating these metrics over the same FOV, or making them otherwise invariant to FOV, could facilitate more meaningful comparisons across samples and/or modalities.

R1.4: This raises an interesting point about the necessity of normalising the FOV to obtain invariant, tractography-based metrics of tortuosity and maximum deviation across different samples and modalities. In general, achieving this is challenging, and in this study, it is practically not possible. Between species, we encounter significant differences in brain volume ratios, which complicates the establishment of a common reference FOV due to the distinct anatomical organisation of monkey and mouse brains (see our response to R1.8).

Within species, we would encounter challenges due to missing contrast—such as issues with staining—and the lack of perfect co-registration.

The Discussion section (page 28) has been extended to reflect this: "Within the same species, assuming perfect co-registration of samples, it would be possible to perform correlative imaging and analysis. This would allow validation of whether tractography streamlines could be reproduced at different image resolutions within the same normalised FOV. Although this was not possible with the current data and experimental setup, it would be an interesting point to pursue in future work."

Though the results seem solid, some statements, particularly in the abstract and discussion, do not seem to be fully supported by the data. For example, the abstract states "Our findings revealed common principles of fibre organisation in the two species; small axonal fasciculi and major bundles formed laminar structures with varying angles, according to the characteristics of major pathways.", though the results show "no strong indication within the mouse CC of the axonal laminar organisation observed in the monkey". Similarly, the introduction states: "By these means, we demonstrated a new organisational principle of white matter that persists across anatomical length scales and species, which governs the arrangement of axons and axonal fasciculi into sheet-like laminar structures." Further comments on the text are provided below.

R1.5: We understand that it can be misunderstood that the laminar organisation is identical in monkeys and mice, which is not the case. For example, we show that in the corpus callosum, pathways are parallel in the mouse but not in the monkey. We have clarified that while the principle of layered laminar organisation of pathways is shared between monkeys and mice, species-specific differences do exist.

We have made the following clarifying changes to the manuscript:

The Abstract (page 2): "Our findings revealed common principles of fibre organisation that apply despite the varying patterns observed across species"

The Introduction (page 4-5): "Through these methods, we demonstrated organisational principles of white matter that persists across anatomical length scales and species. These principles govern the organisation of axonal fasciculi into sheet-like laminar shapes (structures with a predominant planar arrangement). Interestingly, while these principles remain consistent, they result in varied structural organisations in different species."

The Discussion (page 21): "despite species differences".

One observation not notably discussed in the paper is that the spherical histograms of Figure 3E/H appear to have an anisotropic spread of the white points about 0,0. It would be interesting if the authors could comment on whether this could be interpreted as the FOD having asymmetric dispersion and if so, whether the axis of dispersion relates to the fibre orientations of the laminar structures.

R1.6: That is a good point, and to address it, we have fitted spherical Bingham distributions to the FODs, allowing us to quantify their shapes. From each Bingham distribution, we derived two wellknown indices from the diffusion MRI community: the Orientation Dispersion Index (ODI) and Dispersion Anisotropy (DA) index. The ODI explains the dispersion of fibres for a single bundle FOD, whereas DA expresses the shape of the FOD on the unit sphere surface, i.e., the degree of anisotropy. We have integrated the Bingham-based analysis into the Methods, Discussion, and Results sections concerning Figures 3 and 7, but not Figure 4, which contains multiple fibre bundles that we cannot separate on a voxel level. The analysis does not impact the overall message and conclusion but adds interesting context to the discussion around laminar organisation.

A limitation of the study is that it considers only small ex vivo tissue samples from two locations in a single postmortem monkey brain and slightly larger regions of mouse brain tissue. Consequently, further evidence from additional brain regions and subjects would be required to support more generalised statements about white matter organisation across the brain.

R1.7: Collecting more samples from various locations in the brain would provide valuable insights into the consistency of white matter organisation across anatomical length scales, as well as the structure tensor based anisotropy and tortuosity metrics. However, being awarded beamtime at two different synchrotron facilities to scan the same sample with different imaging setups is practically challenging. At the ESRF, we have gathered additional image volumes from other white matter regions of the monkey brain that support all our findings, which will be published separately. X-ray synchrotron imaging technology is advancing rapidly, with faster acquisition times enabling more image volumes to be stitched together. This extends the FOV and allows for a more robust statistical description of the anatomy. Consequently, future studies with an extended FOV and varying image resolutions could utilise a single synchrotron facility to collect additional samples, further supporting our findings.

The Discussion section (page 27) has been extended to reflect this: “Increasing the number of samples across both species and examining laminar organisation at various length scales in more regions would strengthen our findings. However, securing beamtime at two different synchrotron facilities to scan the same sample with varying image resolutions is a limiting factor. Beamline development for multiresolution experimental setups, along with faster acquisition methods, is a rapidly advancing field. For instance, the Hierarchical Phase-Contrast Tomography (HiP-CT) imaging beamline at ID-18 at the ESRF, enables multi-resolution imaging within a single session to address this challenge, though it is currently limited to a resolution of 2.5 μm (Walsh et al. 2021).”

Given the monkey results, the mouse study (section 2.5 onwards) lacks some motivation. In particular, it is unclear why a demyelination model was studied and if/how this would link to the laminar structure observed in the monkey data. Further, it is unclear how comparable tortuosity/max deviation values are across species, considering the differences in data quality and relative resolution, given that the presented results show these values are very modality-dependent.

R1.8: We have clarified the motivation for including the mouse part of the study in both the Introduction and the Results sections.

The Introduction section (page 5): “Furthermore, using a mouse model of focal demyelination induced by cuprizone (CPZ) treatment, we investigate the inflammation-related influence on axonal organisation. This is achieved through the same structure tensor-derived micro-anisotropy and tractography streamline metrics.”

The Results section (page 15): “Finally, we investigated the organisation of fasciculi in both healthy mouse brains and a murine model of focal demyelination induced by five weeks of cuprizone (CPZ) treatment. This allowed for the exploration of the disease-related influence on axonal organisation, particularly under inflammation-like conditions with high glial cell density at the demyelination site (He et al. 2021). The experimental setup for DESY and ESRF is similar to that described for the monkey, with the exception that we did not perform dMRI and synchrotron imaging on the same brains, and only collected MRI data for healthy mouse brains. This approach allowed us to apply the same structure tensor and tractography streamline analysis used previously, but in a healthy versus disease comparison, demonstrating the methodology’s ability to provide insights into pathological conditions.”

Across species, the comparison of tortuosity and maximum deviation must be approached with caution. On one hand, we observe a comparable influence of the extra-axonal environment in both the monkey and mice, as discussed in the section “Sources to the non-straight trajectories of axon fasciculi.” On the other hand, the anatomical scale and relative image resolution are significant factors, as correctly pointed out. In the mouse, for instance, the measures are influenced by white matter pathway macroscopic effects, making cross-species comparison challenging to perform in a normalised way.

The limitations section of the Discussion (page 28) has been updated to reflect this: “A limiting consequence of having samples imaged at differing anatomical scales is that certain measures become inherently hard to compare in a normalised way. The tractography-based metrics—tortuosity and maximum deviation—serve as good examples of this resolution and FOV dependence. In the ESRF samples, the anatomical scale was at the level of individual axons, and the streamline metrics primarily reflect micro-scale effects from the extra-axonal environment, such as the influence of cells and blood vessels. In comparison, the larger anatomical scale in the DESY samples represents the level of fasciculi and above, with metrics influenced by macroscopic effects, such as the bending of the CC pathway. Both scales are interesting and can provide valuable insights in their own right, but caution is required when comparing the numbers, especially for cross-species studies where there is a significant difference in brain volume ratios.”

The paper introduces a new method of "scale-space" parameters for structure tensors. Since, to my understanding, this is the first description of the method, some simple validation of the method would be welcomed. Further, the same scale parameters are not used across monkeys and mice, with a larger kernel used in mice (Table 2) which is surprising given their smaller brain size. Some explanation would be helpful.

R1.9: We have expanded the description of the scale-space structure tensor approach in the Methods section. Specifically, we have elaborated on the empirical process used to select the scale-space parameters shown in Table 2 and explained why multiple scales were applied only to the monkey samples scanned at ESRF (see Table 2, sample IDs 2 and 3) but not to the other datasets. Additionally, we have added a supplementary figure to assist in illustrating the concept.

Reviewer #2 (Public Review):

Summary:

In this work, the authors combine diffusion MRI and high-resolution x-ray synchrotron phase-contrast imaging in monkey and mouse brains to investigate the 3D organization of brain white matter across different scales and species. The work is at the forefront of the anatomical investigation of the human connectome and aligns with several current efforts to bridge the resolution gap between what we can see in vivo at the millimeter scale and the complexity of the human brain at the sub-micron scale. The authors compare the 3D white matter organization across modalities within 2 small regions in one monkey brain (body of the corpus callosum, centrum semiovale) and within one region (splenium of the corpus callosum) in healthy mice and in one murine model of focal demyelination. The study compares measures of tissue anisotropy and fiber orientations across modalities, performs a qualitative comparison of fasciculi trajectories across brain regions and tissue conditions using streamlined tractography based on the structure tensor, and attempts to quantify the shape of fasciculi trajectories by measuring the tortuosity index and the maximum deviation for each reconstructed streamline. Results show measures of anisotropy and fiber orientations largely agree across modalities, especially for larger FOV data. The high-resolution data allows us to

explore the fiber trajectories in relation to tissue complexity and pathology. The authors claim the study reveals new common organization principles of white matter fibers across species and scales, for which axonal fasciculi arrange into sheet-like laminar structures.

Strengths:

The aim of the study is of central importance within present efforts to bridge the gap between macroscopic structures observable in vivo in humans using conventional diffusion MRI and the microscopic organization of white matter tissue. Results obtained from this type of study are important to interpret data obtained in vivo, inform the development of novel methodologies, and expand our knowledge of the structural and thus functional organization of brain circuits.

Multi-scale data acquired across modalities within the same sample constitute extremely valuable data that is often hard to acquire and represent a precious resource for validation of both diffusion MRI tractography and microstructure methods.

The inclusion of multi-species data adds value to the study, allowing the exploration of common organization principles across species.

The addition of data from a murine cuprizone model of focal demyelination adds interesting opportunities to study the underlying biological changes that follow demyelination and how these impact tissue anisotropy and fiber trajectories. These data can inform the interpretation and development of diffusion MRI microstructure models.

Weaknesses:

The main claim of a newly discovered laminar organization principle that is consistent across scales and species is not supported strongly enough by the data. The main evidence in support of the claim comes from the larger FOV data obtained from the body of the corpus callosum in the monkey brain. A laminar organization principle is partially shown in the centrum semiovale in the monkey brain and it is not shown in mice data. Additionally, the methods lack details to help the correct interpretation of these findings (e.g., how were these fasciculi defined?; how well do they represent different axonal populations?; what is the effect of blood vessels on the structure tensor reconstruction?; how was laminar separation quantified?) and the discussion does not provide a biological background for this organization. The corpus callosum sample suggests axons within a bundle of fibers are organized in a sheet-like fashion, while data from the centrum semiovale suggest fibers belonging to different fiber bundles are organized in a sheet-like arrangement. While I acknowledge the challenges in acquiring such high-resolution data, additional samples from different regions in the same animals and from different animals would help strengthen this claim.

R2.1

- how were these fasciculi defined?

In the introduction (page 3), we have clarified our definition of an axon fasciculus: “A fasciculus is a bundle of axons that travel together over short or long distances. Its size and shape can vary depending on its internal organisation and its relationship to neighbouring fasciculi.”

Additionally, we emphasise in the Results section (page 12) that the centroid streamlines are not guaranteed to be actual fasciculi, but rather representations of them. The paragraph now states: “To ease visualisation and quantification, we used QuickBundle clustering (Garyfallidis et al. 2012) to group neighbouring streamlines with similar trajectories into a centroid

streamline. This centroid streamline serves as an approximation of the actual trajectory of a fasciculus.”

- what is the effect of blood vessels on the structure tensor reconstruction?

Fair point, that was not clear from our description. The clarification contains two parts. First, the estimation of the structure tensor occurs in all voxels, and in that sense, the blood vessels respond very similarly to axons. Second, when it comes to sample statistics derived from the structure tensor analysis (FA histograms and the FODs), they will have an influence, albeit a small one, given the low volume percentage of the blood vessels within the FOVs. In the monkey samples, segmenting the blood vessels was achievable with little effort, allowing us to exclude their contribution from FA statistics and FODs. To make this clear, we have added a paragraph to the Methods section (page 34) titled “Structure tensor-based quantifications,” reflecting this clarification. Additionally, we have restructured the entire structure tensor methods description (starting on page 32) as part of the reviewer comments in R1.6 and R1.9.

- how was laminar separation quantified?

We have added a clarification in Results section (page 7): “The laminar thickness was determined by manual measurements on laminae visually identified in the 3D volume”.

- discussion does not provide a biological background for this organization.

A good point. Including the biological background is relevant as it supports the laminar organisation of white matter pathways observed in our findings and those of others.

We have added a section on this background in the Discussion (page 24): “We believe our observed topological rule of white matter laminar organisation can be explained by a biological principle known from studies of nervous tissue development. The first axons to reach their destination, guided by their growth cones, are known as “pioneering” axons. “Follower” axons use the shaft of the pioneering axon for guidance to efficiently reach the target region (Breau and Trembleau 2023). Axons can form a fasciculus by fasciculating or defasciculating along their trajectory through a zippering or unzipping mechanism, controlled by chemical, mechanical, and geometrical parameters. Zippering “glues” the axons together, while unzipping allows them to defasciculate at a low angle (Šmít et al. 2017). Although speculative, the zippering mechanism may be responsible for forming the laminar topology observed across length scales. The defasciculation effect can explain our results in the corpus callosum (CC) of monkeys, with laminar structures at low angles (~35 degrees) also observed by (Innocenti et al. 2019; Caminiti et al. 2009), as well as in other major pathways (Sarubbo et al. 2019). In contrast, a fasciculation mechanism may be observed in the mouse CC (0 degrees). If the geometrical angle between two axons is high, i.e., toward 90 degrees, the zippering mechanism will not occur, and the two axons (fasciculi) will cross (Šmít et al. 2017). This supports our and other findings that crossing fasciculi or pathways occur at high angles toward 90 degrees in the fully matured brain (Wedeen et al. 2012). Once myelination begins, the zippering mechanism is lost (Šmít et al. 2017), suggesting that laminar topology is established at the earliest stages of brain maturation.”

- additional samples from different regions in the same animals and from different animals would help strengthen this claim

Reviewer #1 also pointed to the inclusion of additional samples, and this is now discussed as part of the study limitations on page 27 (see also R1.7).

The main goal of the study is to bridge the organization of white matter across anatomical length scales and species. However, given the substantial difference in FOVs between the two imaging modalities used, and the absence of intermediate-resolution data, it remains difficult to effectively understand how these results can be used to

inform conventional diffusion MRI. In this sense, the introduction does not do a good enough job of building a strong motivation for the scientific questions the authors are trying to answer with these experiments and for the specific methodology used.

R2.2: Indeed, this is an essential point now emphasised in the introduction, page 3, which now states: "Despite the limited resolution of dMRI, the water diffusion process can reveal microstructural geometrical features, such as axons and cell bodies, though these features are compounded at the voxel level. Consequently, estimating microstructural characteristics depends on biophysical modelling assumptions, which can often be simplistic due to limited knowledge of the 3D morphology of cells and axons and their intermediate-level topological organisation within a voxel. Thus, complementary highresolution imaging techniques that directly capture axon morphology and fasciculi organisation in 3D across different length scales within an MRI voxel are essential for understanding anatomy and improving the accuracy of dMRI-based models(Alexander et al. 2019)."

Additionally, in the introduction, page 4, we have made the following changes to strengthen the link across modalities, such that it now states: "In the x-ray synchrotron data, we applied a scale-space structure tensor analysis, which allowed for the quantification of structure tensor-derived tissue anisotropy and FOD in the same anatomical regime indirectly detected by dMRI."

The cuprizone data represent a unique opportunity to explore the effect of demyelination on white matter tissue. However, this specific part of the study is not well motivated in the introduction and seems to represent a missed opportunity for further exploration of the qualitative and quantitative relationship between diffusion MRI and sub-micron tissue information (although unfortunately not within the same brain sample). This is especially true considering the diffusion MRI protocol for mice would allow extrapolation of advanced measures from different tissue compartments.

R2.3: A similar point was raised by Reviewer 1 (R1.8), and we have clarified the motivation for including the healthy mice and the demyelination samples.

Recommendations for the authors:

Reviewer #1 (Recommendations For The Authors):

Many thanks to the authors for providing open data. This was very helpful when reviewing the manuscript and is a valuable resource for the community.

R1.10: We are happy to share our data with the community. Understanding anatomy in 3D is hard to achieve through still images and animations, so the ability to explore it on your own is quite important. The link to the data repository has been added in the Methods section in the following paragraph: "Due to the size of the data selected, processed image volumes, masks and results are available at <https://zenodo.org/records/10458911>. Other datasets can be shared on request."

One confusing element of the paper is that orientations (or axes) do not seem to be consistent across samples/modalities. For example, the green tensors in Figures 3 C and D are tilted up/down in opposite directions and the streamlines in Figure 5A seem opposite (SL) from what we would expect from Figure 2A (SR). Having consistent orientations across modalities and images would help the reader. When colouring tensors (e.g. in Figure 3), the authors could consider a 3D colour scheme (similar to that used by diffusion MRI) rather than colouring by only inclination, as this would provide useful information on whether different laminae have similar orientations, as implied by the tractography in Figure 4.

R1.11: Thank you for spotting the suboptimal consistency between Figures 2, 3, and 5. Figure 2 has been corrected and updated. The left-right direction in the coronal views was not correctly displayed. Additionally, the glyph directions have been updated in Figures 2 and 3.

By default, we use the “standard” RGB colour scheme used in dMRI. However, for the monkey CC—essentially Figure 3—this did not effectively illustrate our findings. We decided to use a different directional colour encoding scheme, which captures the angular deviation from the L-R axis. This was to assist in the visualisation of the inclination angle between the laminars. We have used the same colour scheme for the tensors in Figure 3 to avoid confusion.

On a general note, the standard colour scheme has uniform “colour contrast” in all directions, but when there is only a single dominant direction in the sample, it can make sense to concentrate the colour contrast in that axis.

Results: "even higher FA anisotropy in the micro-tensor domain of 0.997, i.e., the micro (μ)FA (20, 21)." I understand these references lead to a definition of μ FA that is based on multiple diffusion tensor encodings which is quite different from that suggested by Kaden. It may be preferable to reference Kaden directly (since I understand this is the method used) to avoid confusion.

R1.12: Correctly spotted, and we now reference the method from Kaden et al. and use the other references elsewhere when relevant.

"and scanned the mouse brain in a whole." - typo?

R1.13: Thank you for spotting the typo. The mouse brain was kept in the skull during MRI scanning, which has been clarified in the Methods section.

The crossing fibre region appears to be sometimes referred to as the centrum semiovale, and other times as the CST. CS seems the better description and keeping this naming consistent would avoid confusion to the reader.

R1.14: Well spotted, thank you. We have replaced the usage of Corticospinal Tract (CST) with centrum semiovale (CS) where relevant.

Direct comments on the text:

Abstract: "Individual axon fasciculi exhibited tortuous paths in a manner independent of fibre complexity and demyelination"

Do statistical comparisons of the various distributions support this? The data shows somewhat increased tortuosity in the CST compared to the CC, and somewhat lower tortuosity in CPZ tissue.

R1.15: The intention of the text was not to point to the comparison of tortuosity, but rather to highlight the maximum deviation. We observe a high probability density of maximum deviations at approximately 5-10 microns in all samples, which corresponds to the size of structures in the extraaxonal environment, such as blood vessels and cells.

Additionally, we understand that the original statement might imply an expectation of a statistical analysis demonstrating independence, which is not the case. To clarify, we have reformulated the sentence in the Abstract (page 2) to address these points: “Fasciculi exhibited non-straight paths around obstacles like blood vessels, comparable across the samples of varying fibre complexity and demyelination.”

Abstract: "A quantitative analysis of tissue anisotropies and fibre orientation distributions gave consistent results for different anatomical length scales and modalities, while being dependent on the field-of-view."

To my understanding, the FODs here from different modalities are calculated over different FOVs (in monkeys at least), and FODs are only presented for a single FOV for each modality, meaning it is difficult to separate the effects of modality from FOV. The microscopic anisotropy is also noticeably different across modalities (DESY < ESRF < dMRI).

R1.16: That is a fair point. Our statement was trying to capture too much condensed content to be correctly interpretable. We have reformulated the sentence to state: "Quantifications of fibre orientation distributions were consistent across anatomical length scales and modalities, whereas tissue anisotropy had a more complex relationship, both dependent on the field-of-view".

While it is true that we only present the ST-derived quantifications – FOD and FA statistics – for a single FOV per modality and sample, the results shown for the ESRF monkey samples (Figures 3 and 4) are a merge of four individually processed volumes. The quantifications of each individual subFOV have now been added as a supplementary figure (Figure S3) to highlight the consistency of the methodology and the effect of shifting the FOV position. In the case of the mouse, we have two volumes from different mice, which also display similar FOD and FA statistics.

Abstract: "Our study emphasises the need to balance field-of-view and voxel size when characterising white matter features across anatomical length scales."

This point does not seem very well explored in the paper, rather it is an observation of the limitations of the different imaging modalities. For example, there aren't analyses to compare metrics from highresolution data at different FOVs (i.e. by taking neighbourhoods of different sizes), nor are metrics compared from data at different resolutions and the same FOV.

R1.17: The question is related to R1.16, R1.4, and R1.8, and we have addressed this point in our responses to those comments.

Figure 7 - Taking into account the eigenvalues can be helpful when interpreting the secondary and tertiary eigenvectors of tensors (V2 and V3). It would be interesting to know whether the eigenvalues $L2 \approx L3$ are approximately equal (suggesting isotropic diffusion about V1, where the definition of V2 versus V3 isn't very meaningful), or if L2 is noticeably larger than L3 (suggesting anisotropic diffusion about V1, potentially similar to the anisotropic dispersion discussed above).

R1.18: It would be interesting to explore the eigenvalues of the structure tensor in more detail, as has been done for the diffusion tensor. However, we believe this belongs to future work, as such additional detailed methodological analysis would complicate the already complex story. As mentioned in response to R1.10, most processed data has been made publicly available, and the rest can be requested (due to the storage size of the data sets) to perform such additional analysis.

Discussion: "Importantly, our findings revealed common principles of fibre organisation in both monkeys and mice; small axonal fasciculi and major bundles formed sheet-like laminar structures," See above regarding the lack of evidence for laminar structures in mouse data.

R1.19: We have reformulated the text for clarification as part of R1.3. Additionally, we added FOD quantifications to support why we do not observe an apparent laminar organisation in the mouse CC— please see our response to R1.6.

Discussion: "Interestingly, the dispersion magnitude is indicative of fasciculi that skirt around obstacles in the white matter such as cells and blood vessels, and the results are largely independent of both white matter complexity (straight vs crossing fibre region) and pathology." Again, do statistical tests of the various distributions support this?

R1.20: As part of R1.1, we have added statistical tests of significance for the quantifications of how max deviation changes when bending around objects. Indeed, the distributions are not statistically the same, and we do not wish to convey that sentiment, but they are comparable in the object sizes that they detect. As done in the abstract, we have reformulated the sentence to avoid misunderstanding and have replaced “largely independent” with “observed across.”

Discussion: "Tax et al. have demonstrated the calculation of a sheet probability index from diffusion MRI data, which suggested the presence of sheet-like features in the CC"

My understanding was that this was observed in crossing fibre regions, such as where fibres projecting with the CC cross the CST, but not the main body of the CC itself. Tax defines sheet structure as "composed of two tracts that cross each other on the same surface in certain regions along their trajectories." Is this a different phenomenon to the laminar structures observed here (where we observe fibres within a single tract being locally organised into laminar structures)?

R1.21: Thank you for pointing our attention to this. We have corrected the section in the Discussion (page 23), so it now states: “Additionally, Tax et al. have demonstrated the calculation of a gridcrossing sheet probability index from diffusion MRI data, which suggested the presence of sheet-like features in a crossing fibre region (Tax et al. 2016), which is in line with our findings in the synchrotron data. Note that the method by Tax et al. only detects sheet-like structures crossing on a grid and does not reveal laminar structures with lower inclination angles, as we observed in the monkey CC.”

Discussion: "We found that FODs were consistent across image resolutions and modalities, but only given that the FOV is the same." See above.

R1.22: As part of our response to R1.6, we quantified the FODs using the ODI and DA indices, which should help support our statement. Nevertheless, we have toned down the statement and reformulated the text as follows: “We found that FODs were comparable across image resolutions and modalities. The observed discrepancies can be attributed to the fact that the FOVs are not exactly matched.”

Discussion: "microscopic FA were highly correlated across modalities."

The data shows FA is considerably lower in DESY to ESRF; within modality FA is quite consistent irrespective of tissue region; and differences between the CC and CG shown in ESRF data in mice are not repeated in DESY. It is unclear from the current data if this would lead to a high correlation across modalities. Some evidence would be helpful.

R1.23: This is a fair point; we have not performed a correlation analysis. However, the pattern we observe for the synchrotron samples is as follows: When the anatomical length scale increases (becomes more macroscopic), the FA distribution shifts to lower values. This reflects the scale of information captured with the ST analysis (see also R1.9). Therefore, the

most interesting comparison of FA statistics occurs when the resolution and anatomical length scale are approximately the same. The sentence in question has been reformulated to the following: "Estimates of structure tensor derived microscopic FA show a clear pattern across modalities."

Discussion: "If so, the (inclination angle) information might serve to form rules for low-resolution diffusion MRI based tractography about how best to project through bottleneck regions, which is currently a source of false-positives trajectories (6)."

This is an interesting idea but it is unclear to me how this inclination information would help track through bottlenecks where, by definition, fibres are passing through with the same orientation. Some further explanation would be helpful.

R1.24: We have elaborated on the section in the Discussion (page 23), explaining how this can be used to improve tractography tracing through complex regions: "The reason is that standard tractography methods do not "remember" or follow anatomical organisation rules as they trace through complex regions. Our findings on pathway lamination and inclination angles—low for parallel-like trajectories and high for crossing-like trajectories—can help incorporate trajectory memory into these methods, reducing the risk of false trajectories".

Reviewer #2 (Recommendations For The Authors):

Below I report comments that if addressed I believe would improve the clarity and readability of the manuscript.

- Figures 1 and 2 would be more meaningful if combined into one figure. This would allow for a direct visual comparison of the two modalities. If space is needed, I believe the second row of Figure 1 (coronal views of CC) does not add much information. It is often hard to navigate the different orientations of the tissue in the images; thus any effort in trying to help the reader visually clarify would improve readability.

R2.4: We considered the reviewer's suggestion to merge Figures 2 and 3. However, this made both the figures and the main text additionally complex, so we chose to retain the original figure layout. Secondly, Figure 3 utilises a non-standard directional colormap. Keeping the colormap consistent within each figure is a feature we wish to preserve. In response to R1.11, the figures have been updated to have more consistent orientations for the monkey samples.

In Figure 2, the second row, showing a coronal view of the CC, is essential for comparison with human data in Figure S1. It highlights where we observed the columnar laminar organisation and their inclination angle, as also detected by DTI.

- Figure 4 shows synchrotron data revealing an anterior-posterior component within the centrum semiovale that is not necessarily seen in the dMRI data. Could the authors comment on this?

R2.5: Thank you for pointing this out. We have now addressed this in the Results section (page 10), where we describe the observation in detail: "Interestingly, visual inspection of the colour-coded structure tensor directions in Fig. 4E shows the existence of voxels whose primary direction is along the A-P axis. However, this represents a small enough portion of the volume that it does not appear as a distinct peak on the FOD."

- The authors claim they observed several purple axons crossing orthogonally in Figure 5c. However, that is not necessarily clear in the figure.

R2.6: We appreciate the feedback. We have now coloured the streamlines of the crossing fasciculi in Figure 5C in red.

- Figure 5 would benefit from adding the color encoding scheme for Figure 5d, as sometimes this is not necessarily consistent.

R2.7: We appreciate the feedback. We have added an indication of the standard directional colour coding to Figure 5D.

- Figure 5d shows interesting data from the complex region. However, it is hard to visualize and it looks like there are not many streamlines traveling entirely I-S? Maybe a different orientation of the sample would help visualization.

R2.8: A similar point was raised by Reviewer 1 (see R1.2). We have added an animation of the scene to assist in the interpretation of the 3D organisation within this complex sample.

- The concept of axon fasciculi is not necessarily immediately clear. Adding an explanation for what the authors refer to when using this term would improve clarity.

R2.9: In the introduction, we now state our conceptual definition of an axon fasciculus as a number of axons that follow each other (see also R2.1).

- The methods do not provide details on how structure tensor FA is measured.

R2.10: Thank you for pointing this out. We have restructured and expanded the structure tensor description in the Methods section (see also R1.9 and R2.1), which now includes the definition of FA.

- Why didn't the authors select the same cc region for both mice and monkeys? It seems this would have increased the strength of the comparison.

R2.11: We agree. The reason lies in the chronology of experiments and the fact that we cannot control where demyelination takes place. We have added a clarifying description in the Methods section (page 31): “Note that several separate beamline experiments were conducted to collect the volumes listed in Table 1. In the first two experiments, samples from the monkey brain were scanned at ESRF and DESY, respectively. The samples from the mouse brain were imaged in two subsequent experiments. Consequently, the location of the identified demyelinating lesion in the cuprizone mice, which cannot be precisely controlled, did not match the location of the CC biopsies in the monkey.”

- While it is mentioned in the results, the methods do not explain how vessel segmentations or cell segmentation in mice was performed and for which datasets it was performed.

R2.12: For the small ROI shown in Figure 6, the labelling was a manual process using the software ITK-SNAP, which has now been clarified in the corresponding figure caption. The generation of ROI masks and blood vessel segmentations involved a combination of intensity thresholding, morphological operations, and manual labelling in ITK-SNAP. This has been clarified in the restructured and expanded description of structure tensor analysis in the Methods section (starting on page 32).

- From the methods it is hard to understand (1) how many mice were used; (2) why dMRI was done on a different sample; (3) whether the same selenium region was selected for

| *both healthy and CPZ animals; (4) how the registration across samples was performed.*

R2.13: We appreciate the feedback and have inserted clarifying statements in the relevant parts of the Methods section. (1) The total number of mice included was three: one normal, one cuprizone, and one normal for MRI scanning. (2) The quality of the collected dMRI on the mouse was too poor to use, and it could not be redone as the brain had already been sliced and prepared for synchrotron experiments. (3) The same splenium section was selected for both healthy and cuprizone mice. (4) A paragraph on image registration has been added.

| *- Diffusion MRI method sections would benefit from additional details on the protocols used.*

R2.14: Thank you for pointing this out. We have added more details about the diffusion MRI protocols, including the b-value, gradient strength, and other relevant parameters.

<https://doi.org/10.7554/eLife.94917.2.sa0>