

Distinct catecholaminergic pathways projecting to hippocampal CA1 transmit contrasting signals during navigation in familiar and novel environments

Reviewed Preprint

v3 • August 23, 2024

Revised by authors

Reviewed Preprint

v2 • July 30, 2024

Reviewed Preprint

v1 • February 26, 2024

Chad M Heer, Mark EJ Sheffield 

The Department of Neurobiology, The University of Chicago, Chicago, USA

 https://en.wikipedia.org/wiki/Open_access Copyright information

Abstract

Neuromodulatory inputs to the hippocampus play pivotal roles in modulating synaptic plasticity, shaping neuronal activity, and influencing learning and memory. Recently it has been shown that the main sources of catecholamines to the hippocampus, ventral tegmental area (VTA) and locus coeruleus (LC), may have overlapping release of neurotransmitters and effects on the hippocampus. Therefore, to dissect the impacts of both VTA and LC circuits on hippocampal function, a thorough examination of how these pathways might differentially operate during behavior and learning is necessary. We therefore utilized 2-photon microscopy to functionally image the activity of VTA and LC axons within the CA1 region of the dorsal hippocampus in head-fixed male mice navigating linear paths within virtual reality (VR) environments. We found that within familiar environments some VTA axons and the vast majority of LC axons showed a correlation with the animals' running speed. However, as mice approached previously learned rewarded locations, a large majority of VTA axons exhibited a gradual ramping-up of activity, peaking at the reward location. In contrast, LC axons displayed a pre-movement signal predictive of the animal's transition from immobility to movement. Interestingly, a marked divergence emerged following a switch from the familiar to novel VR environments. Many LC axons showed large increases in activity that remained elevated for over a minute, while the previously observed VTA axon ramping-to-reward dynamics disappeared during the same period. In conclusion, these findings highlight distinct roles of VTA and LC catecholaminergic inputs in the dorsal CA1 hippocampal region. These inputs encode unique information, with reward information in VTA inputs and novelty and kinematic information in LC inputs, likely contributing to differential modulation of hippocampal activity during behavior and learning.

eLife assessment

This study presents **important** findings on the differential activity of noradrenergic and dopaminergic input to dorsal hippocampus CA1 in head-fixed mice traversing a runway in a virtual environment that is familiar or novel. The data are rigorously analysed, and the observed divergence in the dynamics of activity in the dopaminergic and noradrenergic axons is **solid**. Future studies, using specific manipulations of the two distinct midbrain inputs combined with behavioral testing, are required to strengthen the claim that distinct signals to the hippocampus cause distinct behavioral effects.

<https://doi.org/10.7554/eLife.95213.3.sa4>

Introduction

Catecholamines have a well established role in hippocampal function. Both dopamine and norepinephrine have been shown to impact hippocampal dependent learning and memory (Silva et al., 2012 [DOI](#); Retailleau and Morris, 2018 [DOI](#); Tsetsenis et al., 2019 [DOI](#); Hansen and Manahan-Vaughan, 2014 [DOI](#); Gibbs and Summers, 2002 [DOI](#); Thomas, 2015 [DOI](#); André et al., 2015 [DOI](#)), alter synaptic plasticity (Hagena and Manahan-Vaughan, 2013 [DOI](#); Hansen and Manahan-Vaughan, 2014 [DOI](#); Chu et al., 2011 [DOI](#); Edison and Harley, 2012 [DOI](#); Hagena and Manahan-Vaughan, 2012 [DOI](#); Goh and Manahan-Vaughan, 2013 [DOI](#)), modulate cell excitability (Edelmann and Lessmann, 2018 [DOI](#); Segal et al., 1991 [DOI](#)), and influence the formation and stability of place cells (Kentros et al., 2004 [DOI](#); Retailleau and Morris, 2018 [DOI](#)), hippocampal neurons that selectively fire at specific locations in an environment (O'Keefe and Dostrovsky, 1971 [DOI](#)). Traditionally, the main source of dopamine to the dorsal hippocampus was thought to be sparse inputs from the ventral tegmental area (VTA), while locus coeruleus (LC) inputs provided the main source of norepinephrine.

VTA DA inputs to dorsal CA1 of the hippocampus mainly innervate stratum oriens (Takeuchi et al., 2016 [DOI](#); Adeniyi et al., 2020 [DOI](#); Adeyelu and Ogundele, 2023 [DOI](#)), and their activity bidirectionally modulates Schaffer Collateral (CA3-CA1) synapses (Rosen et al., 2015 [DOI](#)), enhances persistence of reward-location associations (McNamara et al., 2014 [DOI](#)), and drives place preference (Mamad et al., 2017 [DOI](#)). VTA-hippocampus input activity can also bias place field location (Mamad et al., 2017 [DOI](#)), improve place field stability across days (McNamara et al., 2014 [DOI](#)), and drive reward expectation dependent enhancement of place field quality (Krishnan et al., 2022 [DOI](#)). However, many of the effects of DA modulation of the hippocampus have now been attributed to LC inputs as their activity enhances the strength of Schaffer Collateral synapses (Takeuchi et al., 2016 [DOI](#)), improves memory retention (Kempadoo et al., 2016 [DOI](#)), improves place field stability across days (Wagatsuma et al., 2018 [DOI](#)), and can bias place fields to a location when paired with a reward (Kaufman et al., 2020 [DOI](#)) through DA mechanisms. Although many of the effects of LC and VTA are overlapping, potentially indicating shared mechanisms of action, they are believed to play different roles in spatial learning and memory (Duszkiewicz et al., 2019 [DOI](#)). LC inputs influence the encoding of novel environments (Kempadoo et al., 2016 [DOI](#); Wagatsuma et al., 2018 [DOI](#)), while VTA DA inputs increase persistence of reward context associations (McNamara et al., 2014 [DOI](#)), alter hippocampal neurons firing rate (Adeniyi et al., 2020 [DOI](#)), and their suppression can evoke place avoidance (Mamad et al., 2017 [DOI](#)). It is possible that these differences arise because of the differences in activity observed between LC and VTA DA neurons. Therefore, characterizing the encoding properties of LC and VTA inputs directly in the hippocampus during navigation and spatial learning would provide important insights into the specific roles of these distinct neuromodulatory pathways.

Additionally, recent findings indicate considerable heterogeneity in the activity of VTA (Engelhard et al., 2019 [↗](#)) and LC (Uematsu et al., 2017 [↗](#); Noei et al., 2022 [↗](#); Chandler et al., 2014 [↗](#)) neurons, highlighting the need for projection specific recordings. Therefore, we functionally imaged VTA DA and LC axons with single axon resolution in dCA1 of mice as they navigated familiar and novel virtual reality (VR) environments for rewards. We observed distinct encoding properties between these sets of inputs during navigation and in response to environmental novelty. During the approach to a previously rewarded location, the activity of most VTA DA axons ramped up. In contrast, most LC axons did not show this ramping activity and instead predicted the start of motion. Additionally, a majority of LC axons and some VTA axons showed activity associated with the animal's velocity. Following exposure to a novel environment, VTA axon ramping-to-reward signals greatly reduced but LC axon activity sharply increased. These findings support distinct roles for VTA and LC inputs to the hippocampus in spatial navigation of rewarded and novel environments.

Results

To record the activity of dopaminergic inputs to the dorsal hippocampus, we expressed axon-GCaMP6s or axon-GCaMP7b in LC or VTA neurons of different mice. We utilized the NET-cre mouse line (Wagatsuma et al., 2018 [↗](#)) to restrict expression to noradrenergic LC neurons, and the DAT-Cre line (Zhuang et al., 2005 [↗](#)) to restrict expression to dopaminergic VTA neurons (**Fig. 1B** [↗](#)). Mice were then head-fixed and trained to run a linear virtual reality (VR) track for water rewards delivered through a stationary waterspout when they reached the end of the virtual track (**Fig. 1A** [↗](#)). Following reward delivery, mice were teleported to the beginning of the track and allowed to complete another lap. On experiment day mice navigated the familiar, rewarded VR environment for 10 min while 2-photon microscopy was used to image the calcium activity of LC (87 ROIs from 22 imaging sites in 16 mice) or VTA (9 ROIs from 8 imaging sites in 8 mice) axons in the dorsal CA1 (**Fig. 1C** [↗](#)).




Figure 1.

Distinct activity dynamics in VTA and LC axons during navigation of familiar environments

a, Experimental setup (top). Example virtual reality environment. **b**, Schematic representation of injection procedure (left). Representative coronal brain sections immunostained for Tyrosine Hydroxylase (TH) from a DAT-Cre mouse showing overlapping expression of axon-GCaMP (green) and TH (red) in VTA neurons (top) and from a NET-Cre mouse showing overlapping expression of axon-GCaMP (green) and TH (red) in LC neurons (bottom). **c**, Example CA1 field of view of VTA axons (top) and LC axons (bottom). Extracted regions of interest used for example VTA and LC activity throughout the figure. **d**, Example DAT-Cre mouse (left) and NET-Cre mouse (right) with aligned reward delivery (top, green), mouse track position (black), $\Delta F/F$ from example ROI (VTA-orange, LC-blue), and mouse velocity (bottom, gray). **e, i**, Position binned $\Delta F/F \pm$ s.e.m. in example VTA (orange) and LC (blue) ROIs during navigation of the familiar rewarded environment. **ii**, Population average position binned $\Delta F/F \pm$ s.e.m. in VTA ROIs (orange, $n = 9$ ROIs from 8 mice) and LC ROIs (blue, $n = 87$ ROIs from 27 sessions in 17 mice). Linear regression analysis (on all data points, not means) shows that the population of VTA ROIs increase activity during approach of the end of the track while the population of LC ROIs have consistent activity throughout all positions. Linear regression, F test, VTA, $P < 1e - 21$, LC, $P < 0.01$. **iii**, The LC data set was resampled 1000x using $n = 9$ ROIs to match the number of VTA ROIs and the slope and intercept of the regression line were measured each time (blue dots). The VTA slope is steeper than all LC slopes indicating a stronger positive relationship between position and activity for VTA axons. **iv**, Linear regression of position binned activity of individual VTA (orange diamonds), and LC (blue, circles) ROIs. The majority (8/9) of VTA ROIs show a significant positive relationship with position while LC ROIs show both a positive (25/87) and negative (37/87) relationship. **f, i**, Same example ROIs as (d) binned by velocity. **ii**, Same data as (d, ii,) binned by velocity. Linear regression shows that the population of VTA and LC ROIs have a significant relationship with velocity. Linear regression, F test, VTA, $P < 0.05$, LC, $P < 1e - 68$. **iii**, Resampling shows the VTA slope and intercept is within the resampled LC slopes and intercepts indicating similar relationships with velocity. **iv**, Linear regression of individual VTA and LC axons shows the majority (63/87) of LC ROIs have a significant positive relationship with velocity while only 2 VTA ROIs show this relationship. **g**, Same example ROIs as (d) aligned to motion onset. **ii**, Same data as (d, ii,) aligned to motion onset. Linear regression shows that the population of VTA axons have a negative slope prior to motion onset while LC axons have positive slope. Linear regression, F test, VTA, $P < 0.01$, LC, $P < 1e - 65$. **iii**, Resampling shows the VTA slope is negative while all resampled LC slopes are positive. **iv**, Linear regression of individual VTA and LC ROIs shows the majority (56/87) of LC ROIs have a significant positive slope prior to motion onset while the majority (6/9) of VTA ROIs have a negative slope.

© 2024, BioRender Inc. The graphics in panels 1A and B were created with [BioRender](#) are not made available under the same license as the Reviewed Preprint, and are © 2024, BioRender Inc.

Based on the z-axis depth of the recording planes, and the presence of increased autofluorescence in stratum pyramidal, we determined all 9 VTA axons were in *Stratum Oriens*, while for LC recordings, 18 sessions (78 axons in 11 mice) occurred in *Stratum Oriens* and 5 sessions (9 axons in 5 mice) in *Stratum Pyramidalis*. Example VTA (left, orange) and LC (right, blue) axon calcium activity aligned to the animal's behavior are shown in [Fig. 1D](#). Axons from both brain regions showed periodic activity linked to the animals' exploration of the VR environment.

Distinct activity dynamics in VTA and LC inputs during rewarded navigation of a familiar environment

To examine axon activity further, we first looked at the mean activity across all axons as a function of normalized track position ([Fig. 1E](#)). As previously reported ([Krishnan et al., 2022](#)), VTA DA axons increase activity along the track, peaking at the reward location at the end of the track. In contrast, LC input activity remains relatively constant across all positions along the track ([Fig. 1Eii](#)). To examine if this difference could be due to the lower sample size of VTA axons

compared to LC axons, the LC axons were down-sampled to match the VTA sample size ($n = 9$ ROIs in 8 mice) and the slope and intercepts of the down-sampled data was found. This was repeated 1000 times and did not generate any LC data points that overlap with VTA data demonstrating the difference in relationship between position and activity was not due to the different sample sizes (**Fig. 1Eiii**). We also examined the position related activity of individual VTA and LC axons and observed a positive relationship between position and activity in 88.9% of VTA ROIs (8/9 in 7 mice) but only 28.7% of LC ROIs (25/87 from 12 sessions in 12 mice) while 42.5% of LC ROIs (37/87 from 16 sessions in 9 mice) had a negative relationship between position and activity (**Fig. 1Eiv**). To account for the different track lengths between VTA and LC recordings, we also looked at the virtual distance from the rewarded end of the track as well as time from reward (**Supp. Fig. 1**). The same trends were seen for VTA and LC axons, with 8 VTA ROIs showing positive relationships with both distance and time from reward, and 21 LC ROIs showing a positive relationship with distance from reward and 31 LC ROIs showing a positive relationship with time to reward. This suggests that track length does not influence the encoding properties of VTA and LC axons.

For a subset of VTA ($n = 6$) and LC ROIs ($n = 26$), the reward at the end of the track was removed, and the activity of these axons in the unrewarded condition was examined (**Fig. 2A**). While the slope of the VTA population activity across positions significantly decreased (**Fig. 2Bi**) as expected and previously reported (Krishnan et al., 2022), the LC population activity across positions did not significantly change (**Fig. 2Cii**). This confirmed that the ramping activity in VTA axons was due to the animal's proximity to an expected reward and this signal was not present in the average activity of LC axons.

Next, we investigated the mean activity of these axons as a function of velocity. The population mean of both VTA and LC axons increased as velocity increased (**Fig. 1Fi**). This is consistent with the finding that LC inputs to dCA1 encode velocity (Kaufman et al., 2020) and the finding that some DA VTA neurons encode kinematics (Engelhard et al., 2019). Again, to account for differences in sample size, we down sampled the LC axons 1000 times and found the slope and y-intercept of each sampling. The overlap of the VTA and LC slopes and intercepts confirms we cannot conclude any differences in velocity related activity in the VTA and LC axon populations (**Fig. 1Fiii**). However, analyzing individual ROIs, we observed a statistically significant positive relationship between velocity and activity in the majority, 72.4%, of LC axons (63/87 ROIs from 21 sessions in 13 mice), while only 28.6% (2/9 ROIs in 2 mice) of VTA axons showed a positive relationship with velocity. (**Fig. 1Fiv**). The strong velocity correlated activity in a small subset of VTA DA axons indicates heterogeneity in the activity of these inputs similar to what is observed in VTA DA somas (Engelhard et al., 2019). To determine if the velocity correlated activity in VTA DA axons is confounded by the ramping to reward activity, we examined this activity in the unrewarded condition where the ramping to reward activity is absent. Here, the VTA population activity across all velocities is decreased but the slope, or the relationship between velocity and activity, remains unchanged (**Fig. 2Bii**). This is consistent with a subset of VTA axons encoding velocity information, while the overall decrease in activity is explained by the decrease in reward related activity demonstrated in **Fig. 2Bi**. However, the relationship between velocity and LC activity is unchanged in the unrewarded environment (**Fig. 2Cii**), indicating condition invariant velocity encoding in LC axons.

Finally, we examined the activity of LC and VTA axons during rest and the transition to movement. The population of LC axons ramped up in activity during the 2 s leading up to motion onset (**Fig. 1Gi**). This is consistent with reports of activity of LC axons in cortical areas (Reimer et al., 2016) showing LC activity prior to motion onset. In contrast, VTA axons show decreasing activity during the 2 s leading up to motion onset (**Fig. 1Gi**). This ramping down in VTA axon activity is likely due to most periods of immobility occurring between reward delivery and the start of the next lap, during which we previously demonstrated reward related activity decays in VTA axons (Krishnan et al., 2022). Indeed, in the unrewarded condition the negative slope of the VTA population activity leading up to motion onset disappears (**Fig. 2Biii**), indicating this




Figure 2.

Removal of reward restructures VTA but not LC input activity during spatial navigation:

a. Experimental Paradigm. **b.i.** Population average position binned $\Delta F/F \pm$ s.e.m. of VTA ROIs ($n = 6$ ROIs in 6 mice) in the rewarded (VTA rew, orange) and unrewarded conditions (VTA unrew, gray). Linear regression, F test, Rewarded, $P < 1e - 22$, Unrewarded, $P < 0.001$. **ii.** Same data as (b,i) binned by velocity. Linear regression, F test, Rewarded, $P < 0.05$, Unrewarded, $P < 0.05$. **iii.** Same data as (b,i) aligned to motion onset. Linear regression, F test, Rewarded, $P < 1e - 4$, Unrewarded, $P < 0.05$. **c. i.** Population average position binned $\Delta F/F \pm$ s.e.m. of LC ROIs ($n = 26$ ROIs in 7 sessions in 4 mice) in the rewarded (LC rew, blue) and unrewarded conditions (LC unrew, gray). Linear regression, F test, Rewarded, $P < 0.01$, Unrewarded, $P < 0.01$. **ii.** Same data as (c,i) binned by velocity. Linear regression, F test, Rewarded, $P < 1e - 12$, Unrewarded, $P < 1e - 15$. **iii.** Same data as (c,i) aligned to motion onset. Linear regression, F test, Rewarded, $P < 1e - 21$, Unrewarded, $P < 1e - 26$. The slope of each unrewarded measure was compared to the familiar laps using a one-way ANCOVA with Tukey HSD post hoc test. * $P < 0.05$, ** $P < 0.001$, *** $P < 1e - 4$.

© 2024, BioRender Inc. The graphics in panel 2A were created with [BioRender](#) are not made available under the same license as the Reviewed Preprint, and are © 2024, BioRender Inc.

relationship is largely driven by the presence of reward rather than motion onset. However, the LC population activity remains unchanged in the unrewarded environment (**Fig 2C** [.iii.](#)), suggesting this activity is related to the lead up to motion initiation. These differences in activity are not an artifact of lower sample size of VTA ROIs as shown by down-sampling the LC ROIs activity 1000 times and measuring the slopes and intercepts of the down-sampled data did not generate any data points that overlapped with the VTA slope and intercept (**Fig 1G** [.iii.](#)). In further support of distinct activity profiles leading up to motion onset, we found that the majority (6/9 ROIs in 6 mice) of VTA ROIs decreased in activity leading up to motion onset but only 2/87 LC ROIs in one session decreased in activity, while the majority, (56/87 ROIs in 17 sessions in 9 mice), of LC ROIs increased in activity leading up to motion onset (**Fig. 1G** [.iv.](#)). Together, this analysis demonstrates overlapping but distinct activity in VTA and LC axons during spatial navigation with VTA axons showing strong activity correlated with distance to reward and some velocity correlated activity, while LC axons demonstrate activity correlated to velocity and time to motion onset.

Because the DAT-Cre mice were injected with a virus for expression of either axon-GCaMP6s (4 mice) or axon-GCaMP7b (4 mice), we asked whether these two GCaMP variants led to different activity dynamics (**Supp. Fig. 2** [.i.](#)). The axon-GCaMP6s (5 ROIs) and axon-GCaMP7b (4 ROIs) both had a positive relationship with position, negative relationship with time to motion initiation, and a neutral relationship with velocity, although the GCaMP6s relationships were stronger (**Supp. Fig. 2 A, C** [.i.](#)). Importantly, when we compared only axon-GCaMP6s expressing VTA ROIs with the axon-GCaMP6s expressing LC ROIs (**Supp. Fig. 3** [.i.](#)), we saw similar relationships as the comparisons using all VTA ROIs (**Fig. 1** [.i.](#)).

Environmental Novelty induces activity in LC but not VTA inputs

Both LC and VTA neurons have been shown to respond to novel environments ([Takeuchi et al., 2016](#) [.i.](#)). Therefore, we aimed to examine the activity of VTA DA and LC inputs to the hippocampus during exposure to novel VR environments. Following 10 minutes in the familiar environment, mice were teleported to a novel VR environment of the same track length, with a reward at the same position at the end of the track. Following teleportation, we found the running speed of both DAT-Cre and NET-Cre mice transiently decreased (**Fig. 3A** [.i.](#)) and they spend less time immobile (**Fig. 3A** [.i.](#)), demonstrating mice recognize they are navigating a novel environment. While the velocity quickly recovered for both groups of mice, the freezing ratio, or amount of time spent immobile, never recovered in the first ten laps in the novel environment for the NET-Cre mice indicating some novelty induced changes in behavior persist.

We aligned VTA and LC axon activity to the switch to the novel environment and investigated changes in activity due to exposure to novelty. To test whether the mean axon activity is significantly elevated or lowered, we defined a baseline by generating 1000 shuffles of the axon traces across the entire recording sessions, downsampling the shuffled data 1000 times to match the VTA ($n = 7$) and LC ($n = 87$) sample sizes, and calculating the mean and 95% CI of the shuffled data. After teleportation, the periodic activity observed in the mean of VTA axons, likely reflecting ramping-to-reward signals in each individual axon, disappeared (**Fig. 3B** [.i.](#)). This is evident in the traces of most of the individual VTA ROIs showing a loss of the ramping-to-reward signal (**Fig. 3C** [.i.](#)), and in the VTA population position binned activity showing a significant reduction in ramping activity (**Supp. Fig. 4** [.i.](#)). However, 1 VTA ROI showed an increase in activity immediately following exposure to novelty (**Fig. 3C** [.i.](#)), indicating heterogeneity across VTA axons in CA1 and the lack of a novelty signal on average may be due to a small sample size.

Strikingly, LC axons show a dramatic increase in mean activity that remained elevated for > 1 minute following exposure to the novel environment (**Fig. 3B** [.i.](#)) similar with findings that LC cell body activity is elevated for minutes following exposure to environmental novelty ([Takeuchi et al., 2016](#) [.i.](#)). A significant increase in activity above baseline activity following the switch to a novel environment can be seen in 36 LC axon ROIs (from 15/22 sessions in 10/16 mice) (**Fig 3C** [.i.](#)). To




Figure 3.

Exposure to a novel environment causes an abrupt and sustained increase in activity in LC but not VTA inputs to dCA1

a, i, Experimental Paradigm. **ii**, Behavior from example mouse during the transition from the familiar VR environment to a novel VR environment showing the animals track position (top, black) and velocity (bottom, gray). **iii**, The average running velocity $\pm$ s.e.m. of all mice during the transition to a novel VR environment (top). The average freezing ratio of all mice $\pm$ s.e.m., calculated as the time spent immobile (velocity < 5cm/s) divided by the total lap time. Each lap was compared to the final lap in the familiar environment using a one-way ANOVA with Tukey HSD post hoc test (Blue *, NET-Cre mice $P < 0.05$; Orange *, DAT-Cre mice $P < 0.05$). **b**, Mean normalized fluorescence of all VTA ROIs (top, n = 7) and LC ROIs (bottom, n = 87) aligned to the switch to the novel environment. To define a baseline and 95% CI (gray shaded region), 1000 shuffles were created from the calcium traces and down sampled to match the sample size and averaged. This was repeated 1000 times and the mean and 95% CI of this shuffled data was determined for each frame. Red lines indicate periods where two or more consecutive frames passed above the % CI of the shuffled baseline. **c**, Normalized $\Delta F/F$ activity of all VTA ROIs (top) and LC ROIs (bottom) aligned to the switch to the novel VR environment. **d**, The normalized fluorescence of all LC ROIs binned by lap (left) or into 50 frame bins (right). The baseline and 95% CI (gray shaded region) was defined using the same method as in (b). Red lines indicate bins above the baseline 95% CI.

further characterize this activity, we found the mean population activity for each lap and separately for 10 s time bins leading up to and following exposure to the novel environment. This analysis shows that LC activity is significantly elevated above baseline for 6 consecutive laps and approximately 90 seconds following exposure to the novel environment (**Fig. 3D**). These findings demonstrate that LC inputs signal environmental novelty, supporting a role for these inputs in novelty encoding in the hippocampus.

Novelty-induced changes in behavior explains the late but not early increases in LC activity

It is possible that the change in the amount of time the animals spend running versus immobile in the novel environment could explain the increase in LC activity in the novel environment, as LC activity is related to behavior (**Fig. 1**). For instance, LC axons show elevated activity during motion versus rest (**Fig. 1G**). Therefore, an increase in the time spent in motion upon exposure to the novel environment could lead to an increase in LC activity. To account for the differences in time spent running vs immobile between the two environments, we removed any periods where the mice were immobile to isolate the effects of novelty from changes in behavior (**Fig. 4B**). When only looking at activity during running in both environments, we found that LC axon activity is elevated for 2 laps, or 40 s, in the novel environment (**Fig. 4B-E**). Additionally, when we look only at activity when the mice were immobile, we see that activity is elevated for 30s in the novel environment (**Supp. Fig. 5B**). Together, this indicates that there are two separate components that drive LC axon activity during the initial exposure to the novel environment. One, a shorter purely novelty-induced increase in activity which occurs during the first 2 laps, or about 40 s, in the novel environment. Two, a behavior-induced increase in LC activity caused by an increase in the percentage of time spent running that extends beyond the increase in the novelty-induced activity for 6 laps or 90 s.

If the short, novelty-induced signal in LC axons is an additional signal riding on top of the behavior correlated signals - position, velocity, and motion onset-we would expect a disruption of these behavioral correlations during the initial lap in a novel environment. To test this we examined these behavioral correlations lap-by-lap following exposure to the novel environment. Indeed, the slopes of the position binned, velocity binned, and motion onset aligned data are all significantly more negative in the first lap in the novel environment than the final laps of the familiar environment (**Fig. 4F-H**). This is consistent with a decaying novelty signal that peaks at the start of the first lap and rides on top of these behaviorally-correlated signals. This produces an elevation in activity at positions near the start of the first lap that is lower at positions near the end of the first lap, causing a negative slope relationship between position and LC axon activity on the first lap (**Fig. 4F**; light green line). Further, low velocities occur at the start of each lap compared to the end of the lap. Because the novelty signal is highest when animals are running slowest, the novelty signal flattens the velocity-LC activity relationship (**Fig. 4G**; light green line). Lastly, rest periods typically occur at the start of the track. Therefore, motion onset encoding on the first lap in the novel environment occurs when the novelty signal is highest, again, flattening the relationship (**Fig. 4H**; light green line). By the third lap in the novel environment, where the novelty-induced signal is no longer observable, the relationships between LC activity and position is no longer different than the relationship in the familiar environment (**Fig. 4F**, -green). The relationship between LC activity and motion onset is also no longer different by the third lap in the novel environment (**Fig. 4H**). Although the relationship between velocity and LC activity is different in the third novel lap than that of the final familiar laps, by the final lap in the novel environment it is no longer different than the familiar relationship(**Fig. 4G**). Together this demonstrates that the relationships between LC activity and behavior in the novel environment quickly return to those in the familiar environment. Interestingly, the slope is significantly increased in the final lap of the novel environment (**Fig. 4F**), potentially indicating the development of activity at the novel reward location as has been previously reported (Kaufman et al., 2020). Altogether, examining the lap-by-lap dynamics of the position, velocity, and motion




Figure 4.

Novelty-induced changes in behavior explain the late but not early increases in LC activity

a, Good behavior from example mouse following removal of freezing periods (velocity < 0.2 cm/s) during the transition from the familiar to a novel VR environment showing the animals track position (top, black) and velocity (bottom, gray). **b**, Normalized $\Delta F/F$ activity of all LC ROIs aligned to the switch to the novel VR environment following removal of freezing periods. **c**, Mean normalized F of all LC ROIs (bottom, $n = 87$) aligned to the switch to the novel environment. To define a baseline and 95% CI (gray shaded region), 1000 shuffles were created from the calcium traces and down sampled to match the sample size and averaged. This was repeated 1000 times and the mean and 95% CI of this shuffled data was determined for each frame. Red lines indicate periods where two or more consecutive frames passed above the 95% CI of the shuffled baseline. **d,e**, The normalized F of all LC ROIs binned by lap (d) or into 50 frame bins (e). The baseline and 95% CI (gray shaded region) was defined using the same method as in (c). Red lines indicate bins above the baseline 95% CI. **f-h**, Population average position binned (f), velocity binned (g), and motion onset aligned (h) $\Delta F/F \pm$ s.e.m. in LC ROIs ($n = 90$) in the final laps of the familiar environment (light blue), and the first (orange), third (dark blue), and final laps (green) of the novel environment. Linear regression, F test, position binned (f) fam last, $P < 1e - 4$, nov 1, $P < 1e - 68$, nov 3, $P = 0.48$, nov last, $P < 0.001$; velocity binned (g) fam last, $P < 1e - 31$, nov 1, $P < 0.05$, nov 3, $P < 1e - 5$, nov last, $P < 1e - 13$; motion onset aligned (h) fam last, $P < 1e - 29$, nov 1, $P < 0.001$, nov 3, $P < 1e - 9$, nov last, $P < 1e - 18$. The slope of each lap was compared to the final familiar laps using a one-way ANCOVA with Tukey HSD post hoc test. * $P < 0.01$, ** $P < 0.001$, *** $P < 1e - 4$.

onset activity indicates that environmental novelty induces a sharp increase in LC input activity during the first two laps in the novel environment while also inducing a change in behavior that leads to increased LC input activity in subsequent laps.

We also examined the activity of a subset of LC axons (50 ROIs from 11 sessions in 9 mice) during the transition from darkness to the familiar VR environment. These axons showed elevated activity for 20s following initial exposure to the familiar environment (**Supp. Fig. 5 A**). While this activity did not persist as long as the activity following exposure to the novel environment, it indicates these axons may be generally responsive to abrupt and unpredicted changes to the animal's environment. This fits with the proposed role of the LC in arousal, with novelty driving greater arousal than abrupt exposure to a familiar environment.

Discussion

During spatial navigation in a familiar environment, activity of VTA DA inputs to dCA1 were strongly modulated by position relative to reward, ramping up as mice approached the end of the track where reward was located. This activity could be related to the animal's distance or time from the rewarded location. Further experiments should be conducted to distinguish between time and distance such as those conducted in VTA DA soma recordings (Kim et al., 2020). We have previously shown that this activity is dependent on the history of reward delivery and reflects the animals' reward expectation (Krishnan et al., 2022). VTA axons in dCA1 also showed decreasing activity during rest prior to motion onset. However, after removing reward, VTA axons showed no activity prior to motion onset indicating this relationship was driven by reward related activity. This ramping activity cannot be explained by changes in the animals' sensory experience, as the VR environment and waterspout position remained unchanged in the unrewarded condition. Additionally, on the initial laps with no reward, the ramping activity is still present (Krishnan et al., 2022) indicating this activity is not directly related to the delivery of water but is instead caused by the animal's internal state of reward expectation.

On average, VTA axons showed activity modulated by velocity in a familiar rewarded environment. This relationship was largely due to the activity of two VTA axons that were strongly modulated by velocity, suggesting that there is heterogeneity in the population of VTA axons in dCA1. A positive relationship between average VTA axon activity and velocity persisted in the unrewarded condition and in the novel environment, indicating velocity encoding in these inputs that is not a result of reward related activity. This heterogeneity in the encoding across individual VTA axons is consistent with studies demonstrating heterogeneous encoding of behavioral variables in VTA DA cell bodies, including activity related to rewards and kinematics (Engelhard et al., 2019).

Our findings that VTA DA axons show no novelty induced activity and instead show reduced activity following exposure to a novel environment, is in contrast with several studies showing novelty induced activity in VTA DA cell bodies (Takeuchi et al., 2016; Duszkievicz et al., 2019; Lisman and Grace, 2005), indicating potential heterogeneity in VTA neurons in response to novelty. It is possible that some VTA DA inputs to dCA1 respond to novel environments, and the small number of axons recorded here are not representative of the whole population. Another possibility is that the lack of a novelty response we observe is due to differences in experimental design. Here, mice learned to approach a location for reward which has been shown to lead to ramping activity in dopaminergic VTA neurons (Howe et al., 2013; Krishnan et al., 2022; Kim et al., 2020; London et al., 2018; Jeong et al., 2022). Following exposure to novelty, the disappearance of reward related activity could obscure any novelty induced increases in activity. In addition, as noted above, on average we did observe a velocity associated signal in VTA axons. When mice were exposed to the novel environment their velocity initially decreased. This would be expected to reduce the average signal across the VTA axon population relative to the higher

velocity in the familiar environment. It is possible that this decrease could somewhat mask a subtle novelty induced signal in VTA axons. Therefore, additional experiments should be conducted to investigate the heterogeneity of these axons and their activity under different experimental conditions during tightly controlled behavior.

LC axons showed no position encoding. Instead, they were modulated by velocity and ramped up in activity prior to motion initiation, consistent with recordings of LC axons by others in dCA1 (Kaufman et al., 2020 [DOI](#)) and in the cortex (Reimer et al., 2016 [DOI](#)), respectively. An important question is how is this LC axon activity impacting hippocampal neurons during navigation in a familiar environment? It is possible that LC axons during navigation provide increases in excitability that promotes place cell activity as both dopamine and norepinephrine in the hippocampus can impact cell excitability (Segal et al., 1991 [DOI](#); Edelmann and Lessmann, 2011 [DOI](#), 2018 [DOI](#)). Additionally, place cells are flexible during spatial navigation with new place fields forming in familiar environments (Sheffield et al., 2017 [DOI](#); Dong et al., 2021 [DOI](#)) and shifting position with time/experience (Dong et al., 2021 [DOI](#)). Place fields can also shift to follow changing reward (Gauthier and Tank, 2018 [DOI](#)) and object locations (Bourboulou et al., 2019 [DOI](#)). Dopamine and norepinephrine have also been shown to impact hippocampal synaptic plasticity (Zhang et al., 2009 [DOI](#); Edelmann and Lessmann, 2011 [DOI](#); Hagena and Manahan-Vaughan, 2012 [DOI](#); Goh and Manahan-Vaughan, 2013 [DOI](#)). Therefore, LC inputs may promote the plasticity necessary for place cells to flexibly adapt to changes in a familiar environment (Kaufman et al., 2020 [DOI](#); Redondo and Morris, 2011 [DOI](#)). In other words, LC inputs could allow the hippocampus to be flexible during navigation through their impacts on synaptic plasticity (Takeuchi et al., 2016 [DOI](#); Yamasaki and Takeuchi, 2017 [DOI](#); Duzskiewicz et al., 2019 [DOI](#)).

Exposure to environmental novelty leads to an increase in dopamine in the dorsal hippocampus (Ihalainen et al., 1999 [DOI](#)) and promotes synaptic plasticity (Li et al., 2003 [DOI](#); Hagena and Manahan-Vaughan, 2012 [DOI](#)), hippocampal replay (McNamara et al., 2014 [DOI](#); Dupret et al., 2010 [DOI](#)) and memory persistence (Li et al., 2003 [DOI](#); Cohen et al., 2017 [DOI](#)). In our experiment, exposure to a novel environment caused an increase in LC axon activity but not in VTA DA axon activity, supporting findings that novel experiences induce activity of LC neurons (Takeuchi et al., 2016 [DOI](#)). As discussed above, the slowing down of animal behavior in the novel environment could have decreased LC axon activity and reduced the magnitude of the novelty signal we detected during running. The novelty signal we report here may therefore be an under estimate of its magnitude under matched behavioral settings. The increased activity of LC neurons in the novel environment could increase hippocampal neuron activity (Wagatsuma et al., 2018 [DOI](#)), increase efficacy of Schaffer Colateral synapses (Takeuchi et al., 2016 [DOI](#)), stabilize place cells across days (Wagatsuma et al., 2018 [DOI](#)) and enhance memory persistence (Wagatsuma et al., 2018 [DOI](#); Takeuchi et al., 2016 [DOI](#); Chowdhury et al., 2022 [DOI](#)) through dopamine receptor dependent mechanisms. Importantly, while LC inputs to CA1 have been shown to cause an increase in activity (Wagatsuma et al., 2018 [DOI](#); Chowdhury et al., 2022 [DOI](#)), shape over-representation of novel reward locations (Kaufman et al., 2020 [DOI](#)), and modulate memory linking (Chowdhury et al., 2022 [DOI](#)), they have not been shown to play a role in the formation of contextual memories (Chowdhury et al., 2022 [DOI](#)) or stabilization of place cell maps across days (Wagatsuma et al., 2018 [DOI](#)) in dCA1. However, it is possible this novelty induced activity in LC inputs to dCA1 impacts the formation of instant place fields observed in novel environments (Sheffield et al., 2017 [DOI](#); Dong et al., 2021 [DOI](#)). Instant place fields form on the first lap of a novel environment, right when the LC novelty signal is highest in dCA1, suggesting LC inputs may play a role in their formation or stabilization. Therefore, further experiments should investigate the role of LC axons on dCA1 place fields on a trial-by-trial basis.

While LC neurons have been shown to impact novelty encoding through dopaminergic mechanisms (Wagatsuma et al., 2018 [DOI](#); Takeuchi et al., 2016 [DOI](#); Chowdhury et al., 2022 [DOI](#)), this does not exclude the possibility that they also release norepinephrine during exposure to novelty and exploration of a familiar environment. Indeed, hippocampal levels of norepinephrine also increase during exposure to environmental novelty (Lima et al., 2019 [DOI](#); Moreno-Castilla et al.,

2017 [↗](#)), but how this norepinephrine release effects hippocampal function is not well understood. Additionally, it is not known whether norepinephrine and dopamine are released from the same LC inputs or from distinct sets of LC inputs. Dopamine is in the synthesis pathway of norepinephrine and in LC neurons it is loaded into vesicles where it is then converted to norepinephrine by dopamine *p*-hydroxylase (Cimarusti et al., 1979 [↗](#)). It is possible that high levels of activity of LC inputs, like those occurring during exposure to novelty, lead to release of vesicles before dopamine can be converted to norepinephrine thus leading to the release of dopamine under these conditions. However, low levels of LC activation, like those observed during familiar environment exploration, may provide time for dopamine to be converted to norepinephrine and thus lead to the release of norepinephrine from LC terminals under these conditions. Further experiments investigating the dynamics of dopamine conversion and release from LC terminals in the hippocampus should be conducted to test this hypothesis.

Here we show that LC input activity is modulated by velocity, time to motion onset, abrupt exposure to a familiar environment and abrupt exposure to novel environments. Each of these conditions is associated with an increase in arousal, and LC activity has been strongly linked to arousal levels (Aston-Jones and Bloom, 1981 [↗](#); Berridge and Waterhouse, 2003 [↗](#); Carter et al., 2010 [↗](#)). Therefore, rather than encoding each of these variables independently, LC inputs are likely encoding the animals arousal level during spatial navigation. It has been shown that attention and arousal levels impact tuning properties in many cortical areas and this is thought to be mediated through LC activity (Shulman et al., 1979 [↗](#); Bouret and Sara, 2002 [↗](#); Martins and Froemke, 2015 [↗](#); Waterhouse and Navarra, 2019 [↗](#)). Similarly, changes in the animals' brain state, including changes in attention (Kentros et al., 2004 [↗](#)) and engagement (Pettit et al., 2022 [↗](#)), alter the tuning properties of place cells. This indicates arousal could impact the function of hippocampal neurons through these LC inputs, either directly or through astrocytes (Peter Rupprecht et al., 2023 [↗](#)).

The distinct activity dynamics exhibited by LC and VTA DA axons during spatial navigation of familiar and novel environments underscore their distinct contributions to hippocampal dependent learning and memory processes. Notably, these findings reinforce the notion that VTA DA inputs play a pivotal role in the ongoing maintenance and updating of associations between expected rewards and the locations that lead to them, while LC axons appear to be integral to the process of encoding memories of entirely new environments and stimuli.

Methods

Subjects

All experimental and surgical procedures were in accordance with the University of Chicago Animal Care and Use Committee guidelines. For this study, we used 10–20 week-old male Slc6a3Cre^{+/−}(DAT-Cre^{+/−}) mice obtained from JAX labs and Slc6a2 Cre^{+/−}(NET-Cre^{+/−})(23–33 g) mice obtained from the Tonegawa lab (Wagatsuma et al., 2018 [↗](#)). Male mice were used over female mice due to the size and weight of the headplates (9.1 mm × 31.7 mm, 2 g) which were difficult to firmly attach on smaller female skulls. Mice were individually housed in a reverse 12 h light/dark cycle at 72 °F and 47% humidity, and behavioral experiments were conducted during the animal's dark cycle.

Mouse surgery and viral injections

Mice were anesthetized (1–2% isoflurane) and injected with 0.5 ml of saline (intraperitoneal injection) and 0.05 ml of Meloxicam (1–2 mg/kg, subcutaneous injection) before being weighed and mounted onto a stereotaxic surgical station (David Kopf Instruments). A small craniotomy (1–1.5 mm diameter) was made over the ventral tegmental area (VTA) (± 0.5 mm lateral, 3.1mm caudal of Bregma) of DAT-Cre^{+/−} mice or over the locus coeruleus (LC) (± 0.875 mm lateral, -5.45 mm caudal

of Bregma). The genetically-encoded calcium indicator, pAAV-hsyn-Flex-Axon-GCaMP6s (pAAV-hSynapsin1-FLEEx-axon-GCaMP6s was a gift from Lin Tian (Addgene viral prep # 112010-AAV5 ; <http://n2t.net/addgene:112010> ; RRID:Addgene 112010) was injected into the VTA of DAT-Cre⁺/- mice (200 nL at a depth of 4.4 mm below the surface of the dura) or the LC of NET-Cre⁺/- mice (200 nL at a depth of 3.65 mm below Bregma). For a subset (4/8) of VTA recordings, a different GCaMP variant, pAAV-Ef1A-Flex-Axon-GCaMP7b, was injected due to the difficulty finding and recording VTA axons in dCA1 (pAAV-Ef1a-Flex-Axon-GCaMP7b (pAAV-Ef1a-Flex-Axon-GCaMP7b was a gift from Rylan Larsen - Addgene plasmid # 135419; <http://n2t.net/addgene:135419> ; RRID: Addgene 135419). Following injections, the site was covered up using dental cement (Metabond, Parkell Corporation) and a metal head-plate (9.1 mm $\times$ 31.7 mm, Atlas Tool and Die Works) was also attached to the skull with the cement. To reduce bleeding and swelling during the hippocampal window implantation, mice were separated into individual cages and water restricted for 3 weeks (0.8–1.0 ml per day). Mice then underwent surgery to implant a hippocampal window as previously described (Dombeck et al., 2010). Following implantation, the headplate was attached with the addition of a head-ring cemented on top of the head-plate which was used to house the microscope objective and block out ambient light. Post-surgery mice were given 2–3 ml of water/day for 3 days to enhance recovery before returning to the reduced water schedule (0.8–1.0 ml/day).

Behavior and virtual reality

Our virtual reality (VR) and treadmill setup was designed similar to previously described setups (Sheffield et al., 2017 ; Heys et al., 2014). The virtual environments that the mice navigated through were created using VIRMEn (Aronov and Tank, 2014). 2 m (DAT-Cre mice) or 3 m (NET-Cre mice) linear tracks rich in visual cues were created that evoked numerous place fields in mice as they moved along the track at all locations (Fig. 1) (Bourboulou et al., 2019). Mice were head restrained with their limbs comfortably resting on a freely rotating styrofoam wheel ('treadmill'). Movement of the wheel caused movement in VR by using a rotatory encoder to detect treadmill rotations and feed this information into our VR computer, as in refs. (Sheffield et al., 2017 ; Heys et al., 2014). Mice received a water reward (4 μ L) through a waterspout positioned directly in front of the animal's mouth upon completing each traversal of the track (a lap), which was associated with a clicking sound from the solenoid. Licking was monitored by a capacitive sensor attached to the waterspout. Upon receiving the water reward, a short VR pause of 1.5 s was implemented to allow for water consumption and to help distinguish laps from one another rather than them being continuous. Mice were then virtually teleported back to the beginning of the track and could begin a new traversal. Mouse behavior (running velocity, track position, reward delivery, and licking) was collected using a PicoScope Oscilloscope (PICO4824, Pico Technology, v6.13.2). Behavioral training to navigate the virtual environment began 4–7 days after window implantation (30 min per day) and continued until mice reached >2 laps per minute, which took 10–14 days (although some mice never reached this threshold). Mice that reached this behavioral threshold and had adequate GCaMP expression were imaged the following day. Out of the 36 NET-Cre mice injected, 15 were never recorded for either failing to reach behavioral criteria, or a lack of visible expression in axons. Out of the 54 DAT-Cre mice injected, images were never conducted in 36 for lack of expression or failing to reach behavioral criteria.

Two-photon imaging

Imaging was done using a laser scanning two-photon microscope (Neurolabware). Using a 8 kHz resonant scanner, images were collected at a frame rate of 30 Hz with bidirectional scanning through a 16x/0.8 NA/3 mm WD water immersion objective (MRP07220, Nikon). GCaMP6s and GCaMP7b were excited at 920 nm with a femtosecond pulsed two photon laser (Insight DS + Dual, Spectra-Physics) and emitted fluorescence was collected using a GaAsP PMT (H11706, Hamamatsu). The average power of the laser measured at the objective ranged between 50–80 mW. A single imaging field of view (FOV) between 400–700 μ m equally in the x/y direction was positioned to collect data from as many VTA or LC axons as possible. Time-series images were

collected from 3–5 planes spaced 2 μm apart using an electric lens to ensure axons remained in a field of view and reduce power going to an individual plane. Images were collected using Scanbox (v4.1, Neurolab-ware) and the PicoScope Oscilloscope (PICO4824, Pico Technology, v6.13.2) was used to synchronize frame acquisition timing with behavior.

Imaging sessions

The familiar environment was the same environment that the animals trained in. The experiment protocol for single day imaging sessions is shown in [Fig. 1](#). Each trial lasted 8–12 min and was always presented in the same order. Mice were exposed to the familiar rewarded environment for 10 minutes, then were immediately teleported to the start of a novel rewarded VR environment and allowed to navigate for 10 minutes. Mice on average ran 19 ± 3.8 (mean $\pm$ 95% CI) laps in the familiar environment, at which point they were teleported to the novel environment and imaging continued for 30 ± 5.4 laps. The Novel-rewarded environment (N) had distinct visual cues, colors and visual textures, but the same dimensions (2 or 3 m linear track) and reward location (end of the track) as the familiar environment. Imaging sessions with large amounts of drift or bleaching were excluded from analysis (8 sessions for NET mice, 6 sessions for LC Mice). Sample size for NET-Cre mice was based off previous experiments examining calcium activity of individual somas and axons (Reimer et al., 2016; Engelhard et al., 2019). For DAT-Cre mice sample size was largely limited by the difficulty of recordings and fields of view only containing one axon.

Histology and brain slices imaging

We checked the expression axon-GCaMP to confirm expression was restricted to the VTA of DAT-Cre mice and the LC of NET-Cre. Mice were deeply anesthetized with isoflurane and perfused with 10 ml phosphate-buffered saline (PBS) followed by 20 mL 4% paraformaldehyde in PBS. The brains were removed and immersed in 30% sucrose solution overnight before being sectioned at 30 μm -thickness on a cryostat. Brain slices were collected into well plates containing PBS. Slices were washed 5 times with PBS for 5 min then were blocked in 1% Bovine Serum Albumin, 10% Normal goat serum, 0.1% Triton X-100 for 2 h. Brain slices were then incubated with 1:500 rabbit- α -TH (MAB318, Sigma Aldrich) and 1:500 mouse- α -GFP (SAB2702197, Sigma Aldrich) in blocking solution at 4 $^{\circ}\text{C}$. After 48 h, the slices were incubated with 1:1000 goat- α -rabbit Alexa Fluor 647nm secondary antibody (A32731, ThermoFisher) and 1:1000 goat- α -mouse Alexa Fluor 488nm (A32723, ThermoFisher) for 2 h. Brain slices were then collected on glass slides and mounted with a mounting media with DAPI (SouthernBiotech DAPI-Fluoromount-G Clear Mounting Media, 010020). The whole-brain slices were imaged under $\times 10$ and $\times 40$ with a Caliber I.D. RS-G4 Large Format Laser Scanning Confocal microscope from the Integrated Light Microscopy Core at the University of Chicago.

Image processing and ROI selection

Time-series images were preprocessed using Suite2p (v0.10.1)79. Movement artifacts were removed using rigid and non-rigid transformations and assessed to ensure absence of drifts in the z-direction. Datasets with z-drifts were discarded, as determined by visually assessing imaging sessions, followed by using the registration metrics output by suite2p (8 sessions). For axon imaging, ROIs were first defined using Suite2p and manually inspected for accuracy. ROIs were then hand drawn over all segments of Suite2p defined active axons using ImageJ to ensure all axon segments were included for analysis. Fluorescent activity for each ROI was extracted and highly correlated ROIs (Pearson correlation coefficient ≥ 0.7) were combined and their fluorescent activity was extracted. To be included Baseline corrected $\Delta F/F$ traces across time were then generated for each ROI using both a small window of 300 frames for lap by lap analysis, and a larger sliding window of 2000 frames to avoid flattening slow signals for novelty response analysis. Additional ROIs were drawn over autofluorescent structures that were not identified by suite2p. These “blebs” were processed in the same way as axon ROIs and used as controls to check for imaging and motion artifacts.

To remove low signal to noise axons, we defined the SNR of each ROI using the power spectrum of their fluorescent activity similar to (Reimer et al., 2016 [DOI](#)). For frequencies above 1Hz, the power was defined as noise because this sits outside of the range of frequencies possible for GCaMP6s fluorescence. The SNR ratio was then defined as the ratio of the peak power between 0.5 Hz and 1Hz over the average power between 1 Hz and 3 Hz. The SNR of “blebs” was also determined and any axon with a SNR greater than 1.5 std from the mean of the “blebs” SNR was used for analysis (110/231 LC ROIs, 9/9 VTA ROIs).

Additionally, it was observed that a subset of axon ROIs would greatly increase fluorescence at seemingly random time points and remain elevated for the rest of the trial. This activity could be due to the axons being unhealthy and filling with calcium. Therefore, we identified these axons using the *cusum* function in matlab to detect changes in mean activity that remained elevated for at least 2000 frames or at least 500 frames if they were still elevated at the end of the recording session and removed them from analysis (23/110 LC ROIs, 0/7 VTA ROIs).

Behavioral Analysis

Mouse velocity was calculated as the change in VR position divided by the sampling rate and smoothed using a Savitzky-Golay filter with a 7 frame window and 5 degree polynomial. To find the lap mean velocity, periods where the mice were immobile (velocity < 5 cm/s) were removed and the average velocity during the remaining frames was calculated. The lap mean freezing ratio was calculated as the number of frames spent immobile (velocity < 5 cm/s) divided by the total number of frames for each lap.

Axon Imaging Analysis

For the three measures below, to avoid weighting axons with a high SNR more than others each ROI was normalized by $(\Delta F/F * \Delta F/F_{min}) / (\Delta F/F_{max} * \Delta F/F_{min})$ where $\Delta F/F_{min}$ is the 1st quantile and $\Delta F/F_{max}$ was the 99 quantile for each ROI. The 1st and 99th quantiles were used in order to avoid normalize to noisy outlier data points.

Position binned fluorescence

To find the position binned fluorescent activity of each ROI, the track was divided into 5 cm bins. For each lap, the average fluorescence in each bin was calculated for each ROI. The position binned fluorescence was then averaged across all laps in each environment to find the mean position binned activity in the familiar and novel environments.

Velocity binned fluorescent activity

To find the velocity binned fluorescent activity for each ROI, the velocity was divided into 1 cm/s bins from 1 to 30 cm/s. Velocities above this 14 cm/s were excluded from figures because not all mice ran faster than 14 cm/s. For each lap, the ROIs average fluorescence in each velocity bin was calculated and then averaged across all laps in each environment to find the velocity binned activity in the familiar and novel environments.

Motion initiation aligned fluorescence

Periods where mice were immobile (velocity < 5 cm/s) for at least 1.5s then proceed to run (velocity > 5cm/s) for at least 3s were identified. The fluorescent activity for ROIs for these periods was aligned to the frame mice began running (velocity crossed above 5cm/s). The average aligned fluorescent activity of each ROI was then determined for each environment.

Linear regression analysis

To assess dynamics between each of the above measures and calcium activity of LC and VTA axons, we performed linear regression on the population's familiar environment data and significance was assessed with an F test. To compare the dynamics between LC and VTA axons, we performed exact testing based on Monte-Carlo resampling (1000 resamples with sample size matching the lower sample size condition) as detailed in legends (**Fig. 1E** [↗](#)).

To assess the changing position and velocity encoding of LC axons following exposure to a novel environment, we performed linear regression on the population fluorescence data of the average of the last 4 laps in the familiar environment, and each of the first three laps in the novel environment for each measure. The significance for the fit of each line was assessed with an F test, and an ANCOVA was conducted to test for differences in slope between the four laps. The same process was conducted for the motion initiation dynamics, but only using ROIs in mice who paused within the first 2 laps and 30s following exposure to the novel environment.

Novel response analysis

To examine the response of LC and VTA axons to the novel VR environment, the fluorescence data was normalized by the mean for each ROIs and aligned to the frame where the mice were switched to the novel environment and the mean normalized F for LC and VTA ROIs at each time point was calculated. Baseline fluorescent activity was then calculated for LC and VTA ROIs separately by generating 1000 shuffled traces of the ROIs calcium activity and subsampling down to the sample size (90 for LC; 7 for VTA) 1000 times and finding the mean of the subsampled shuffles. The mean and 95% CI of all 1000 subsamples was found and the mean activity of LC and VTA ROIs was considered significantly elevated when it passed above the 95% CI of the shuffled data. The same process was repeated to define a baseline for the time binned data (fluorescent activity divided into 50 frame bins) and the lap binned data (mean activity for each lap).

Additionally, to account for changes in behavior between the familiar and novel environments, periods where the animals were immobile (velocity < 0.2 cm/s) were removed and running periods were concatenated together and aligned to the switch to the novel environment. Here, we again defined a baseline for the time mean traces, time binned activity, and the lap binned activity using the above bootstrapping approach.

Code availability

Scripts used for data analysis are available on Github https://github.com/chadheer/LC_VTA_paper [↗](#)

Acknowledgements

We thank Dr. Seetha Krishnan and Dr. Douglas Goodsmith for feedback on the manuscript. We also thank Dr. Antoine Madar for extensive and helpful discussions on using appropriate statistical analysis used in this paper. This work was supported by The Whitehall Foundation, The Searle Scholars Program, The Sloan Foundation, The University of Chicago Institute for Neuroscience start-up funds, the NIH 1DP2NS111657-01, and the NIH BRAIN Initiative RF1NS127123 awarded to M.S. and a T32 training grant T32DA043469 from National Institute on Drug Abuse awarded to C.H.

Author contributions


C.H. and M.S. conceived and designed the experiments. C.H. collected and analyzed the data. C.H. and M.S. interpreted the data and wrote the manuscript.

Declaration of interests

The authors declare no competing interests.


Supplementary Figure 1.


a, i, Population activity $\Delta F/F \pm$ s.e.m. binned by the virtual distance to reward for VTA ROIs (orange, 200m track $n = 9$ ROIs in 8 mice) and LC ROIs (blue, 300m track, $n = 87$ ROIs from 27 sessions in 17 mice) in the familiar environment. Linear regression, F test, VTA, $P = 1.83e - 28$, LC, $P = 8.77e - 06$. **ii,** The LC data set was resampled 1000x using $n = 9$ axons to match the number of VTA ROIs and the slope and intercept of the regression line were measured each time (blue dots). The VTA slope is steeper than all LC slopes indicating a stronger positive relationship between position and activity for VTA inputs. **iii,** Linear regression of position binned activity of individual VTA (orange diamonds), and LC (blue, circles) axons. The majority (8/9) of VTA axons show a significant positive relationship with position while LC axons show both a positive (21/87 axons from 9 sessions in 8 mice) and negative (32/87 axons from 15 sessions in 9 mice) relationship. **b, i,** Same data as (a, i), averaged by time to reward. Linear regression shows that the population of VTA axons has a significant positive relationship with time to reward. Linear regression, F test, VTA, $P = 4.44e - 25$, LC, $P = 0.119$. **ii,** Resampling shows the VTA slope is above the resampled LC slopes indicating VTA ROIs have a stronger positive relationship with time to reward. **iii,** Linear regression of individual VTA and LC axons shows the majority (8/9) of VTA axons have a significant positive relationship with time to reward while LC axons show both a significant positive (31/87 axons from 14 sessions in 11 mice) and negative (17/87 axons from 7 sessions in 4 mice) relationship



Supplementary Figure 2.

a, Population average position binned $\Delta F/F \pm$ s.e.m. of VTA GCaMP6s ROIs (orange, $n = 5$ ROIs in 4 mice) and VTA GCaMP7b ROIs (dark orange, $n = 4$ ROIs, in 4 mice) in the familiar environment. Linear regression, F test, GCaMP6s, $P = 2.4264e - 32$, GCaMP7b, $P = 1.9576e - 14$. **b,** Same data as (a) binned by velocity. Linear regression, F test, GCaMP6s, $P = 0.21076$, GCaMP7b, $P = 0.14113$. **c,** Same data as (a) aligned to motion onset. Linear regression, F test, GCaMP6s, $P = 2.0974e - 8$, GCaMP7b, $P = 0.026893$. The slopes for the two GCaMP variants were compared using a one-way ANCOVA with Tukey HSD post hoc test * $P < 0.05$, ** $P < 0.001$, *** $P < 1e - 4$.






Supplementary Figure 3.

a, i, Population average position binned $\Delta F/F \pm$ s.e.m. in axon-GCaMP6s expressing VTA ROIs (orange, $n = 5$ ROIs in 4 mice) and LC ROIs (blue, $n = 87$ ROIs from 27 sessions in 17 mice). Linear regression analysis (on all data points, not means) shows that the population of VTA ROIs increase activity during approach of the end of the track while the population of LC ROIs have consistent activity throughout all positions. Linear regression, F test, VTA, $P = 2.42e - 32$, LC, $P = 0.00396$. **ii**, The LC data set was resampled 1000x using $n = 5$ axons to match the number of VTA ROIs and the slope and intercept of the regression line were measured each time (blue dots). The VTA slope is steeper than all LC slopes indicating a stronger positive relationship between position and activity for VTA inputs. **iii**, Linear regression of position binned activity of individual VTA (orange diamonds), and LC (blue, circles) axons. The majority (4/5) of VTA axons show a significant positive relationship with position while LC axons show both a positive (25/87) and negative (37/87) relationship. **b, i**, Same data as (a, i,) binned by velocity. Linear regression shows that the population of LC ROIs have a significant relationship with velocity. Linear regression, F test, VTA, $P = 0.211$, LC, $P < 1e - 68$. **ii**, Resampling shows the VTA slope is within the resampled LC slopes indicating similar relationships with velocity. **iii**, Linear regression of individual VTA and LC axons shows the majority (63/87) of LC axons have a significant positive relationship with velocity while only 1 VTA axon shows this relationship. **c, i** Same data as (a, i,) aligned to motion onset. Linear regression shows that the population of VTA axons have a negative slope prior to motion onset while LC axons have positive slope. Linear regression, F test, VTA, $P = 2.10e - 081$, LC, $P = 5.51e - 66$. **ii**, Resampling shows the VTA slope is negative while all resampled LC slopes are positive. **iii**, Linear regression of individual VTA and LC axons shows the majority (56/87) of LC axons have a significant positive slope prior to motion onset while the majority (3/5) of VTA axons have a negative slope.


Supplementary Figure 4.

a,i, Population average position binned $\Delta F/F \pm$ s.e.m. of VTA ROIs ($n = 7$ ROIs in 7 mice) in the familiar (orange) and novel (green) rewarded environments. Linear regression, F test, Rewarded, $P < 1e - 21$, Unrewarded, $P < 0.01$. **ii**, Same data as (a,i) binned by velocity. Linear regression, F test, Rewarded, $P < 0.05$, Unrewarded, $P = 0.57$. **iii**, Same data as (a,i) aligned to motion onset. Linear regression, F test, Rewarded, $P < 0.01$, Unrewarded, $P < 0.001$. The slope of each novel measure was compared to the familiar laps using a one-way ANCOVA with Tukey HSD post hoc test. * $P < 0.05$, ** $P < 0.001$, *** $P < 1e - 4$.



Supplementary Figure 5.

a, i, Mean normalized fluorescence of LC ROIs ($n = 50$, 11 sessions in 9 mice) aligned to the switch from dark to the familiar environment. To define a baseline and 95% CI (gray shaded region), 1000 shuffles were created from the calcium traces and down sampled to match the sample size and averaged. This was repeated 1000 times and the mean and 95% CI of this shuffled data was determined for each frame. Red lines indicate periods where two or more consecutive frames passed above the % CI of the shuffled baseline. **a,ii**, The normalized fluorescence of all LC binned into 50 frame bins. The baseline and 95% CI (gray shaded region) was defined using the same method as in (a). Red lines indicate 2 or more consecutive bins above the baseline 95% CI. **b, i**, Mean normalized fluorescence of LC ROIs ($n = 87$) during immobile periods aligned to the switch from the familiar to the novel environment. The baseline and 95% CI (gray shaded region) was defined using the same method as in (a). Red lines indicate 2 or more consecutive bins above the baseline 95% CI. **b, ii** The normalized fluorescence of LC axons during immobile periods binned into 50 frame bins. The baseline and 95% CI (gray shaded region) was defined using the same method as in (a) Red lines indicate 2 or more consecutive bins above the baseline 95% CI.



References

- Adeniyi P. A., Shrestha A., Ogundele O. M (2020) **Distribution of VTA Glutamate and Dopamine Terminals, and their Significance in CA1 Neural Network Activity** *Neuroscience* **446**:171–198
- Adeyelu T., Ogundele O. M (2023) **VTA multifaceted modulation of CA1 local circuits** *Neurobiology of Learning and Memory* **202**
- André M. A. E., Wolf O. T., Manahan-Vaughan D (2015) **Beta-adrenergic receptors support attention to extinction learning that occurs in the absence, but not the presence, of a context change** *Frontiers in Behavioral Neuroscience* **9**
- Aronov D., Tank D. W (2014) **Engagement of neural circuits underlying 2D spatial navigation in a rodent virtual reality system** *Neuron* **84**:442–456
- Aston-Jones G., Bloom F. E (1981) **Norepinephrine-containing locus coeruleus neurons in behaving rats exhibit pronounced responses to non-noxious environmental stimuli** *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* **1**:887–900
- Berridge C. W., Waterhouse B. D (2003) **The locus coeruleus–noradrenergic system: modulation of behavioral state and state-dependent cognitive processes** *Brain Research Reviews* **42**:33–84
- Bourboulou R., Marti G., Michon F.-X., El Feghaly E., Nouguié M., Robbe D., Koenig J., Epsztein J (2019) **Dynamic control of hippocampal spatial coding resolution by local visual cues** *eLife* **8**
- Bouret S., Sara S. J (2002) **Locus coeruleus activation modulates firing rate and temporal organization of odour-induced single-cell responses in rat piriform cortex** *The European Journal of Neuroscience* **16**:2371–2382
- Carter M. E., Yizhar O., Chikahisa S., Nguyen H., Adamantidis A., Nishino S., Deisseroth K., de Lecea L. (2010) **Tuning arousal with optogenetic modulation of locus coeruleus neurons** *Nature Neuroscience* **13**:1526–1533
- Chandler D. J., Gao W.-J., Waterhouse B. D (2014) **Heterogeneous organization of the locus coeruleus projections to prefrontal and motor cortices** *Proceedings of the National Academy of Sciences of the United States of America* **111**:6816–6821
- Chowdhury A., Luchetti A., Fernandes G., Filho D. A., Kastellakis G., Tzivilaki A., Ramirez E. M., Tran M. Y., Poirazi P., Silva A. J (2022) **A locus coeruleus-dorsal CA1 dopaminergic circuit modulates memory linking** *Neuron*, page S **896627322007073**
- Chu H.-Y., Wu Q., Zhou S., Cao X., Zhang A., Jin G.-Z., Hu G.-Y., Zhen X (2011) **SKF83959 suppresses excitatory synaptic transmission in rat hippocampus via a dopamine receptor-independent mechanism** *Journal of Neuroscience Research* **89**:1259–1266

- Cimarusti D., Saito K., Vaughn J., Barber R., Roberts E., Thomas P (1979) **Immunocytochemical localization of dopamine--hydroxylase in rat locus coeruleus and hypothalamus** *Brain Research* **162**:55–67
- Cohen J. D., Bolstad M., Lee A. K (2017) **Experience-dependent shaping of hippocampal CA1 intracellular activity in novel and familiar environments** *eLife* **6**
- Dombeck D. A., Harvey C. D., Tian L., Looger L. L., Tank D. W (2010) **Functional imaging of hippocampal place cells at cellular resolution during virtual navigation** *Nature Neuroscience* **13**:1433–1440
- Dong C., Madar A. D., Sheffield M. E. J (2021) **Distinct place cell dynamics in CA1 and CA3 encode experience in new environments** *Nature Communications* **12**
- Dupret D., O'Neill J., Pleydell-Bouverie B., Csicsvari J (2010) **The reorganization and reactivation of hippocampal maps predict spatial memory performance** *Nature Neuroscience* **13**:995–1002
- Duszkiewicz A. J., McNamara C. G., Takeuchi T., Genzel L (2019) **Novelty and Dopaminergic Modulation of Memory Persistence: A Tale of Two Systems** *Trends in Neurosciences* **42**:102–114
- Edelmann E., Lessmann V (2011) **Dopamine Modulates Spike Timing Dependent Plasticity and Action Potential Properties in CA1 Pyramidal Neurons of Acute Rat Hippocampal Slices** *Frontiers in Synaptic Neuroscience* **3**
- Edelmann E., Lessmann V (2018) **Dopaminergic innervation and modulation of hippocampal networks** *Cell and Tissue Research* **373**:711–727
- Edison H. T., Harley C. W (2012) **Medial and lateral perforant path evoked potentials are selectively modulated by pairing with glutamatergic activation of locus coeruleus in the dentate gyrus of the anesthetized rat** *Hippocampus* **22**:501–509
- Engelhard B. *et al.* (2019) **Specialized coding of sensory, motor and cognitive variables in VTA dopamine neurons** *Nature* **570**:509–513
- Gauthier J. L., Tank D. W (2018) **A Dedicated Population for Reward Coding in the Hippocampus** *Neuron* **99**:179–193
- Gibbs M. E., Summers R. J (2002) **Role of adrenoceptor subtypes in memory consolidation** *Progress in Neurobiology* **67**:345–391
- Goh J. J., Manahan-Vaughan D (2013) **Hippocampal long-term depression in freely behaving mice requires the activation of beta-adrenergic receptors** *Hippocampus* **23**:1299–1308
- Hagena H., Manahan-Vaughan D (2012) **Learning-facilitated long-term depression and long-term potentiation at mossy fiber-CA3 synapses requires activation of -adrenergic receptors** *Frontiers in Integrative Neuroscience* **6**
- Hagena H., Manahan-Vaughan D (2013) **Differentiation in the protein synthesis-dependency of persistent synaptic plasticity in mossy fiber and associational/commissural CA3 synapses in vivo** *Frontiers in Integrative Neuroscience* **7**

- Hansen N., Manahan-Vaughan D (2014) **Dopamine D1/D5 Receptors Mediate Informational Saliency that Promotes Persistent Hippocampal Long-Term Plasticity** *Cerebral Cortex* **24**:845–858
- Heys J. G., Rangarajan K. V., Dombeck D. A (2014) **The functional micro-organization of grid cells revealed by cellular-resolution imaging** *Neuron* **84**:1079–1090
- Howe M. W., Tierney P. L., Sandberg S. G., Phillips P. E. M., Graybiel A. M (2013) **Prolonged dopamine signalling in striatum signals proximity and value of distant rewards** *Nature* **500**:575–579
- Ihalainen J., Riekkinen Jr P., Feenstra M (1999) **Comparison of dopamine and noradrenaline release in mouse prefrontal cortex, striatum and hippocampus using microdialysis** *Neuroscience Letters* **277**:71–74
- Jeong H., Taylor A., Floeder J. R., Lohmann M., Mihalas S., Wu B., Zhou M., Burke D. A., Namboodiri V. M. K (2022) **Mesolimbic dopamine release conveys causal associations** *Science* **378**
- Kaufman A. M., Geiller T., Losonczy A (2020) **A Role for the Locus Coeruleus in Hippocampal CA1 Place Cell Reorganization during Spatial Reward Learning** *Neuron* **105**:1018–1026
- Kempadoo K. A., Mosharov E. V., Choi S. J., Sulzer D., Kandel E. R (2016) **Dopamine release from the locus coeruleus to the dorsal hippocampus promotes spatial learning and memory** *Proceedings of the National Academy of Sciences* **113**:14835–14840
- Kentros C. G., Agnihotri N. T., Streater S., Hawkins R. D., Kandel E. R (2004) **Increased Attention to Spatial Context Increases Both Place Field Stability and Spatial Memory** *Neuron* **42**:283–295
- Kim H. R. *et al.* (2020) **A Unified Framework for Dopamine Signals across Timescales** *Cell* **183**:1600–1616
- Krishnan S., Heer C., Cherian C., Sheffield M. E. J (2022) **Reward expectation extinction restructures and degrades CA1 spatial maps through loss of a dopaminergic reward proximity signal** *Nature Communications* **13**
- Li S., Cullen W. K., Anwyl R., Rowan M. J (2003) **Dopamine-dependent facilitation of LTP induction in hippocampal CA1 by exposure to spatial novelty** *Nature Neuroscience* **6**:526–531
- Lima K. R., da Silva de Vargas L., Ramborger B., Roehrs R., Sevenster D., Izquierdo I., D’Hooge R., Mello-Carpes P. B. (2019) **Noradrenergic and dopaminergic involvement in novelty modulation of aversive memory generalization of adult rats** *Behavioural Brain Research* **371**
- Lisman J. E., Grace A. A (2005) **The Hippocampal-VTA Loop: Controlling the Entry of Information into Long-Term Memory** *Neuron* **46**:703–713
- London T. D., Licholai J. A., Szczot I., Ali M. A., LeBlanc K. H., Fobbs W. C., Kravitz A. V (2018) **Coordinated Ramping of Dorsal Striatal Pathways preceding Food Approach and Consumption** *The Journal of Neuroscience* **38**:3547–3558

- Mamad O., Stumpp L., McNamara H. M., Ramakrishnan C., Deisseroth K., Reilly R. B., Tsanov M (2017) **Place field assembly distribution encodes preferred locations** *PLOS Biology* **15**
- Martins A. R. O., Froemke R. C (2015) **Coordinated forms of noradrenergic plasticity in the locus coeruleus and primary auditory cortex** *Nature Neuroscience* **18**:1483–1492
- McNamara C. G., Tejero-Cantero, Trouche S., Campo-Urriza N., Dupret D. (2014) **Dopaminergic neurons promote hippocampal reactivation and spatial memory persistence** *Nature Neuroscience* **17**:1658–1660
- Moreno-Castilla P., Pérez-Ortega R., Violante-Soria V., Balderas I., Bermúdez-Rattoni F (2017) **Hippocampal release of dopamine and norepinephrine encodes novel contextual information** *Hippocampus* **27**:547–557
- Noei S., Zouridis I. S., Logothetis N. K., Panzeri S., Totah N. K (2022) **Distinct ensembles in the noradrenergic locus coeruleus are associated with diverse cortical states** *Proceedings of the National Academy of Sciences* **119**
- O’Keefe J., Dostrovsky J (1971) **The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely-moving rat** *Brain Research* **34**:171–175
- Peter Rupprecht, Duss Sian N, Becker Denise, Lewis Christopher M, Bohacek Johannes, Helmchen Fritjof (2023) **Centripetal integration of past events by hippocampal astrocytes and its regulation by the locus coeruleus** *bioRxiv*, page 2022
- Pettit N. L., Yuan X. C., Harvey C. D (2022) **Hippocampal place codes are gated by behavioral engagement** *Nature Neuroscience* **25**:561–566
- Redondo R. L., Morris R. G. M (2011) **Making memories last: the synaptic tagging and capture hypothesis** *Nature Reviews. Neuroscience* **12**:17–30
- Reimer J., McGinley M. J., Liu Y., Rodenkirch C., Wang Q., McCormick D. A., Tolias A. S (2016) **Pupil fluctuations track rapid changes in adrenergic and cholinergic activity in cortex** *Nature Communications* **7**
- Retailleau A., Morris G (2018) **Spatial Rule Learning and Corresponding CA1 Place Cell Reorientation Depend on Local Dopamine Release** *Current Biology* **28**:836–846
- Rosen Z. B., Cheung S., Siegelbaum S. A (2015) **Midbrain dopamine neurons bidirectionally regulate CA3-CA1 synaptic drive** *Nature Neuroscience* **18**:1763–1771
- Segal M., Markram H., Richter-Levin G (1991) **Actions of norepinephrine in the rat hippocampus** *Progress in Brain Research* **88**:323–330
- Sheffield M. E., Adoff M. D., Dombeck D. A (2017) **Increased Prevalence of Calcium Transients across the Dendritic Arbor during Place Field Formation** *Neuron* **96**:490–504
- Shulman G. L., Remington R. W., McLean J. P (1979) **Moving attention through visual space** *Journal of Experimental Psychology. Human Perception and Performance* **5**:522–526
- Silva W. C. d., Köhler C. C., Radiske A., Cammarota M (2012) **D1/D5 dopamine receptors modulate spatial memory formation** *Neurobiology of Learning and Memory* **97**:271–275

- Takeuchi T. *et al.* (2016) **Locus coeruleus and dopaminergic consolidation of everyday memory** *Nature* **537**:357–362
- Thomas S. A (2015) **Neuromodulatory Signaling in Hippocampus-Dependent Memory Retrieval** *Hippocampus* **25**:415–431
- Tsetsenis T., Badyna J. K., Subramaniam M., Li R., Yang K., Dani J. A. (2019) **Dopamine and Norepinephrine Modulate the Formation of Aversive Memories in the Hippocampus** *SSRN Electronic Journal*
- Uematsu A., Tan B. Z., Ycu E. A., Cuevas J. S., Koivumaa J., Junyent F., Kremer E. J., Witten I. B., Deisseroth K., Johansen J. P (2017) **Modular organization of the brainstem noradrenergic system coordinates opposing learning states** *Nature Neuroscience* **20**:1602–1611
- Wagatsuma A., Okuyama T., Sun C., Smith L. M., Abe K., Tonegawa S (2018) **Locus coeruleus input to hippocampal CA3 drives single-trial learning of a novel context** *Proceedings of the National Academy of Sciences* **115**:E310–E316
- Waterhouse B. D., Navarra R. L (2019) **The locus coeruleus norepinephrine system and sensory signal processing: A historical review and current perspectives** *Behavioral Consequences of Noradrenergic Actions in Sensory Networks* **1709**:1–15
- Yamasaki M., Takeuchi T (2017) **Locus Coeruleus and Dopamine-Dependent Memory Consolidation** *Neural Plasticity* **2017**:1–15
- Zhang J.-C., Lau P.-M., Bi G.-Q (2009) **Gain in sensitivity and loss in temporal contrast of STDP by dopaminergic modulation at hippocampal synapses** *Proceedings of the National Academy of Sciences* **106**:13028–13033
- Zhuang X., Masson J., Gingrich J. A., Rayport S., Hen R (2005) **Targeted gene expression in dopamine and serotonin neurons of the mouse brain** *Journal of Neuroscience Methods* **143**:27–32

Editors

Reviewing Editor
Mihaela Iordanova
 Concordia University, Montreal, Canada

Senior Editor
Lu Chen
 Stanford University, Stanford, United States of America

Reviewer #1 (Public Review):

Summary:

Heer and Sheffield used 2 photon imaging to dissect the functional contributions of convergent dopamine and noradrenaline inputs to the dorsal hippocampus CA1 in head restrained mice running down a virtual linear path. Mice were trained to collect water reward at the end of the track and on test days, calcium activity was recorded from dopamine (DA) axons originating in ventral tegmental area (VTA, n=7) and noradrenaline axons from

the locus coeruleus (LC, $n=87$) under several conditions. When mice ran laps in a familiar environment, VTA DA axons exhibited ramping activity along the track that correlated with distance to reward and velocity to some extent, while LC input activity remained constant across the track, but correlated invariantly with velocity and time to motion onset. A subset of recordings taken when the reward was removed showed diminished ramping activity in VTA DA axons, but no changes in the LC axons, confirming that DA axon activity is locked to reward availability. When mice were subsequently introduced to a new environment, the ramping to reward activity in the DA axons disappeared, while LC axons showed a dramatic increase in activity lasting 90s (6 laps) following the environment switch. In the final analysis, the authors sought to disentangle LC axon activity induced by novelty vs. behavioral changes induced by novelty by removing periods in which animals were immobile, and established that the activity observed in the first 2 laps reflected novelty-induced signal in LC axons.

The revised manuscript included additional evidence of increased (but transient) signal in LC axons after a transition to a novel environment during periods of immobility, and also that a change from dark to familiar environment induces a peak in LC axon activity, showing that LC input to dCA1 may not solely signal novelty.

Strengths:

The results presented in this manuscript provide insights into the specific contributions of catecholaminergic input to the dorsal hippocampus CA1 during spatial navigation in a rewarded virtual environment, offering a detailed analysis at the resolution of single axons. The data analysis is thorough and possible confounding variables and data interpretation are carefully considered.

Weaknesses:

Aspects of the methodology, data analysis, and interpretation diminish the overall significance of the findings, as detailed below.

The LC axonal recordings are well powered, but the DA axonal recordings are severely underpowered, with recordings taken from a mere 7 axons (compare to 87 LC axons). Additionally, 2 different calcium indicators with differential kinetics and sensitivity to calcium changes (GCaMP6S and GCaMP7b) were used ($n=3$, $n=4$ respectively) and the data pooled. This makes it very challenging to draw any valid conclusions from the data, particularly in the novelty experiment. The surprising lack of novelty-induced DA axon activity may be a false negative. Indeed, at least 1 axon (axon 2) appears to be showing novelty-induced rise in activity in Figure 3C. Changes in activity in 4/7 axons are also referred to as a 'majority' occurrence in the manuscript, which again is not an accurate representation of the observed data.

The authors conducted analysis on recording data exclusively from periods of running in the novelty experiment to isolate the effects of novelty from novelty-induced changes in behavior. However, if the goal is to distinguish between changes in locus coeruleus (LC) axon activity induced by novelty and those induced by motion, analyzing LC axon activity during periods of immobility would enhance the robustness of the results.

The authors attribute the ramping activity of the DA axons to the encoding of the animals' position relative to reward. However, given the extensive data implicating the dorsal CA1 in timing, and the remarkable periodicity of the behavior, the fact that DA axons could be signalling temporal information should be considered.

The authors should explain and justify the use of a longer linear track (3m, as opposed to 2m in the DAT-cre mice) in the LC axon recording experiments.

AFTER REVISIONS:

The authors have addressed my concerns in a thorough manner. The reviewer also appreciates the increased transparency of reporting in the revised manuscript.

Listed below are some remaining comments.

The increase in LC activity with any change in environment (from familiar to novel or from dark to familiar) suggests that LC input acts not solely as a novelty signal, but as a general arousal or salience signal in response to environmental changes. Based on this, I have a couple of questions:

- Is the overall claim that LC input to the dHC signals novelty still valid based on observed findings - as claimed throughout the manuscript?
- Would the omission of a reward be considered a salient change in the environment that activates LC signals, or is the LC not involved with processing reward-related information? Has the activity of LC and VTA axons been analysed in the seconds following reward presentation and/or omission?

<https://doi.org/10.7554/eLife.95213.3.sa3>

Reviewer #2 (Public Review):

Summary:

The authors used 2-photon Ca^{2+} -imaging to study the activity of ventral tegmental area (VTA) and locus coeruleus (LC) axons in the CA1 region of the dorsal hippocampus in head-fixed male mice moving on linear paths in virtual reality (VR) environments.

The main findings were as follows:

- In a familiar environment, activity of both VTA axons and LC axons increased with the mice's running speed on the Styrofoam wheel, with which they could move along a linear track through a VR environment.
- VTA, but not LC, axons showed marked reward position-related activity, showing a ramping-up of activity when mice approached a learned reward position.
- In contrast, activity of LC axons ramped up before initiation of movement on the Styrofoam wheel.
- In addition, exposure to a novel VR environment increased LC axon activity, but not VTA axon activity.

Overall, the study shows that the activity of catecholaminergic axons from VTA and LC to dorsal hippocampal CA1 can partly reflect distinct environmental, behavioral and cognitive factors. Whereas both VTA and LC activity reflected running speed, VTA, but not LC axon activity reflected approach of a learned reward and LC, but not VTA, axon activity reflected initiation of running and novelty of the VR environment.

I have no specific expertise with respect to 2-photon imaging, so cannot evaluate the validity of the specific methods used to collect and analyse 2-photon calcium imaging data of axonal activity.

Strengths:

- (1) Using a state-of-the-art approach to record separately the activity of VTA and LC axons with high temporal resolution in awake mice moving through virtual environments, the authors provide convincing evidence that activity of VTA and LC axons projecting to dorsal CA1 reflect partly distinct environmental, behavioral and cognitive factors.
- (2) The study will help a) to interpret previous findings on how hippocampal dopamine and norepinephrine or selective manipulations of hippocampal LC or VTA inputs modulate

behavior and b) to generate specific hypotheses on the impact of selective manipulations of hippocampal LC or VTA inputs on behavior.

Comments on revised version:

I thank the authors for including a sample size justification.

The justification is based on previous studies using similar sample sizes to characterize behavioral correlates of LC and VTA activity and on practical reasons. I note that to improve reproducibility, it would be preferable to have predefined target sample sizes based on predefined plans for statistical analysis.

<https://doi.org/10.7554/eLife.95213.3.sa2>

Reviewer #3 (Public Review):

Summary:

Heer and Sheffield provide a well-written manuscript that clearly articulates the theoretical motivation to investigate specific catecholaminergic projections to dorsal CA1 of the hippocampus during a reward-based behavior. Using 2-photon calcium imaging in two groups of cre transgenic mice, the authors examine activity of VTA-CA1 dopamine and LC-CA1 noradrenergic axons during reward seeking in a linear track virtual reality (VR) task. The authors provide a descriptive account of VTA and LC activities during walking, approach to reward, and environment change. Their results demonstrate LC-CA1 axons are activated by walking onset, modulated by walking velocity, and heighten their activity during environment change. In contrast, VTA-CA1 axons were most activated during approach to reward locations. Together the authors provide a functional dissociation between these catecholamine projections to CA1. A major strength to their approach is the methodological rigor of 2-photon recording, data processing, and analysis approaches to accommodate their unequal LC-CA1 and VTA-CA1 sample sizes. These important systems neuroscience studies provide solid evidence that will contribute to the broader field of navigation and memory.

Weaknesses:

The conclusions of this manuscript are mostly well supported by the data. However, increasing the sample size of the VTA-CA1 group and using experimental methods that are identical among LC-CA1 and VTA-CA1 groups would help to fully support the author's conclusions.

<https://doi.org/10.7554/eLife.95213.3.sa1>

Author response:

The following is the authors' response to the previous reviews.

Recommendations for the authors:

Reviewer #1 (Recommendations For The Authors):

Please reorder the supplementary figures in the order they are referred to in the Results section for ease of reading. Supp Fig 5 b - should read 'Mean normalized fluorescence of LC ROIs (n = 87) during immobile periods aligned to the switch from familiar to novel environment.'

We thank the reviewer for highlighting these issues and have reordered the supplementary figures and edited the figure legends appropriately.

Reviewer #2 (Recommendations For The Authors):

The authors should include sample size justifications (e.g. based on previous studies, considerations of statistical power, practical considerations, or a combination of these factors).

In response to this concern, we have added a statement to the “Imaging Sessions” section of the methods. Here we highlight sample sizes were largely based on previous studies and/or limited by the difficulty of recordings and the limited number of visible axons per imaging session.

Reviewer #3 (Recommendations For The Authors):

The addition of Supp. Fig 5 partially addresses my previous point 3. However, the claim of dissociation between VTA-CA1 and LC-CA1 would be strengthened by showing that VTA-CA1 axons do not respond to the darkness -> familiar environment in Supp Fig 5. This is particularly important given that (1) the additional 2 VTA-CA1 axons in the revision were not recorded during transitions to novel environments and (2) the overall concern of the reviewers that the low n and heterogeneity of the VTA-CA1 dataset may lead to a false negative. Providing VTA-CA1 data for the darkness -> familiar environment would provide a within-manuscript replication that these axons are not responding to environment changes; a major claim of this manuscript.

While we agree that data of VTA-CA1 axons during the switch from darkness to the familiar environment would provide additional evidence that these axons are not responding to environment changes, unfortunately, VTA axons were not recorded during the switch from familiar to novel.

<https://doi.org/10.7554/eLife.95213.3.sa0>