

The relationship between gut and nasopharyngeal microbiome composition can predict the severity of COVID-19

Reviewed Preprint

v2 • December 4, 2024

Revised by authors

Reviewed Preprint

v1 • March 26, 2024

Benita Martin-Castaño, Patricia Diez-Echave, Jorge García-García , Laura Hidalgo-García, Antonio Jesús Ruiz-Malagon, José Alberto Molina-Tijeras, María Jesús Rodríguez-Sojo, Anaïs Redruello, Margarita Martínez-Zaldívar, Emilio Mota, Fernando Cobo, Marta Alvarez-Estevéz, Federico García, Concepción Morales-García, Silvia Merlos, Paula García-Flores, Manuel Colmenero-Ruiz, José Hernández-Quero, María Nuñez, María Elena Rodríguez-Cabezas, Ángel Carazo, Javier Martín, Rocío Morón , Alba Rodríguez-Nogales, Julio Gálvez

Centro de Salud Las Gabias, Distrito Granada-Metropolitano, Granada, Spain • Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain • Department of Pharmacology, Center for Biomedical Research (CIBM), University of Granada, Granada, Spain • Servicio Microbiología, Hospital Universitario Clínico San Cecilio, Granada, Spain • Centro de Salud “Salvador Caballero”, Distrito Granada-Metropolitano, Granada, Spain • Servicio Microbiología, Hospital Universitario Virgen de las Nieves, Granada, Spain • CIBER de Enfermedades Infecciosas (CIBER-Infecc), Instituto de Salud Carlos III, Madrid, Spain • Respiratory Medicine Department, Hospital Universitario Virgen de las Nieves, Granada, Spain • Servicio de Medicina Intensiva, Hospital Universitario Clínico San Cecilio, Granada, Spain • Servicio de Enfermedades Infecciosas, Hospital Universitario Clínico San Cecilio, Granada, Spain • Servicio Farmacia Hospitalaria, Hospital Universitario Clínico San Cecilio, Granada, Spain • CIBER de Epidemiología y Salud Pública (CIBER-ESP), Instituto de Salud Carlos III, Madrid, Spain • Department of Cell Biology and Immunology, Institute of Parasitology and Biomedicine López-Neyra, CSIC, Granada, Spain • CIBER de Enfermedades Hepáticas y Digestivas (CIBER-EHD), Instituto de Salud Carlos III, Madrid, Spain

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This potentially **valuable** work characterizes the changes in the microbial composition of the nasal and fecal microbiomes in COVID-19 patients based on disease severity. This study enhances the understanding of COVID-19 severity predictors by identifying changes in bacterial species abundance in nasopharyngeal and fecal samples as a biomarker for predicting disease severity. The methods and statistics used appear to be **solid** and in line with the standards of the field.

<https://doi.org/10.7554/eLife.95292.2.sa3>

Abstract

Coronavirus disease 2019 (COVID-19) is a respiratory illness caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) that displays great variability in clinical

phenotype. Many factors have been described to be correlated with its severity, and microbiota could play a key role in the infection, progression, and outcome of the disease. SARS-CoV-2 infection has been associated with nasopharyngeal and gut dysbiosis and higher abundance of opportunistic pathogens. To identify new prognostic markers for the disease, a multicenter prospective observational cohort study was carried out in COVID-19 patients divided into three cohorts based on symptomatology: mild (n=24), moderate (n=51), and severe/critical (n=31). Faecal and nasopharyngeal samples were taken, and the microbiota was analyzed. Linear discriminant analysis identified *M. salivarium*, *P. dentalis*, and *H. parainfluenzae* as biomarkers of severe COVID-19 in nasopharyngeal microbiota, while *P. bivia* and *P. timonensis* were defined in faecal microbiota. Additionally, a connection between faecal and nasopharyngeal microbiota was identified, with a significant ratio between *P. timonensis* (faeces) and *P. dentalis* and *M. salivarium* (nasopharyngeal) abundances found in critically ill patients. This ratio could serve as a novel prognostic tool for identifying severe COVID-19 cases.

Introduction

Coronavirus disease 2019 (COVID-19) is a respiratory illness caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The data reported in November 2023 revealed that over 700 million people have been infected with the virus [1]. The appearance of mutations and variants of concern has caused several additional waves of infection and compromise the effectiveness of existing vaccines and antiviral drugs [2]. Moreover, SARS-CoV-2 infection has shown to cause long-term effects on human health, although the mechanisms are still poorly described [3]. Even though most COVID-19 cases are mild, disease can be severe, resulting in hospitalisation, respiratory failure, or even death [1]. Therefore, a remarkable feature of SARS-CoV-2 infection is the great variability in clinical phenotype among infected people. Many factors can correlate with COVID-19 disease severity, including age, gender, body mass index, previous comorbidities, immune responses, and genetics [4–6]. Of note, and unfortunately, the determinants of infection outcome and the pathogenic mechanisms are not completely understood yet.

SARS-CoV-2 primarily infects the respiratory tract by binding to angiotensin-converting enzyme 2 (ACE2) receptor [7]. However, a growing body of evidence suggests that SARS-CoV-2 can also infect other organs since viral particles and nucleic acids have been found in different biological samples, like sputum, bronchoalveolar lavage fluid, faeces, blood, and urine [8, 9]. Thus, studies employing single-cell RNA sequencing have demonstrated that ACE2 is present in various organs and tissues, including the gastrointestinal tract, where ACE2 receptors have been reported to be highly expressed [10]. This supports several lines of evidence suggesting a substantial involvement of the gastrointestinal tract in the pathogenesis of the disease, including the ability of SARS-CoV-2 to infect and replicate in intestinal enterocytes [11], or to increase expression of the viral entry receptor (ACE2 receptor) and several membrane-bound serine proteases (such as transmembrane protease serine 2 (TMPRSS2) and TMPRSS4) in intestinal epithelial cells [12].

In addition, reports have shown that SARS-CoV-2 infection can disrupt nasopharyngeal and intestinal microbiota composition and promote dysbiosis, reducing diversity and increasing the presence of opportunistic pathogens, including *Staphylococcus*, *Corynebacterium* and *Acinetobacter* bacteria [13, 14], which can make patients more susceptible to secondary infections, rising morbidity and mortality [15]. Besides local changes in the respiratory tract, alterations in the distal gut microbiota have also been observed in SARS-CoV-2 infection [16–19]. Previous studies have evidenced a relevant connection between the microbiome in the nasopharynx and the gut, and preliminary data have suggested a bidirectional interaction that could play a role in the development of the immune response both in healthy and pathological conditions, including SARS-CoV-2 infection [20]. In this sense, dysbiosis has been associated not

only with the severity of the disease but also with the recovery processes [13, 19]; however, little is understood of the link with the establishment of different symptomatic profiles in this condition, and to date, few studies have focused on the association between COVID-19 severity and the nasopharyngeal and faecal microbiota, being examined simultaneously. Considering that the efficacy of the COVID-19 vaccines and antiviral drugs against SARS-CoV-2 is compromised with the emergence of mutations and new variants of the virus [2], new therapeutic approaches and prognostic tools are still necessary. Therefore, the characterization of the nasopharyngeal and intestinal microbiome will allow to identify predictive biomarkers for the diagnosis and prognosis of the disease, as well as possible therapeutic targets in the management of SARS-CoV-2.

Results

Study patients characteristics

A total of 106 patients who had laboratory confirmation of SARS-CoV-2 infection were included in the present study. Of note, none of these patients reported another viral infection at the time of enrolment. Based on the clinical spectrum criteria reported in the COVID-19 treatment guidelines, patients were categorised into 3 groups: mild (24 patients), moderate illness that needed hospitalisation in Respiratory Unit (51 patients), and severe symptomatology and admitted in the ICU (31 patients) (Table 1). The age of the patients significantly increased with the severity of the symptoms. Patients included in the severe group were significantly older than those with mild or moderate symptoms ($p < 0.05$; ANOVA). Patient inclusion was carried out evenly in terms of gender as 52 women and 54 men were recruited; nevertheless, a gender-related impact on the clinical course of these patients was observed since males were predominantly in the group of patients with severe symptoms when compared with the mild illness group ($p < 0.05$; Fisher's test). Regarding the symptomatology, mild patients showed symptoms of a mild respiratory infection, including fever, cough and headache. 33% of mild patients displayed dyspnoea and low oxygen saturation, 26% reported gastrointestinal complaints (including stomach ache, digestive discomfort and diarrhoea); and 4% reported high respiratory and heart rates. In contrast, moderate and severe patients showed higher frequencies of dyspnoea, low oxygen saturation and increased respiratory or heart rates ($p < 0.05$; Fisher's test). When the different comorbidities were considered, patients in the group with severe symptoms showed a higher percentage of cardiomyopathy compared to those from the mild or moderate cohorts (42% vs 3% and 6%, respectively; $p < 0.05$; Fisher's test). However, no significant differences were found in the prevalence of the other pathologies among groups. Regarding blood-based biomarkers, platelets ($p < 0.05$; ANOVA), D-dimer, ferritin and C reactive protein ($p < 0.05$; Kruskal-Wallis) correlated with the severity of the symptoms, being the severe group significantly different.

Bacterial composition differs between sample type and severity index in SARS-CoV-2 infected patients

Nasopharyngeal swabs and faeces were obtained from all the patients included in the study within seven days of symptom onset and used for characterization of the microbiota composition. In the nasopharyngeal microbiota, α -diversity was reduced in the moderate and severe groups compared to the mild group ($p < 0.05$ and $p = 0.07$ respectively; ANOVA) (Figure 1A). Conversely, when α -diversity was examined in stool samples, no significant modifications were observed among groups (Figure 1B). On the other hand, β -diversity analysis through the Bray-Curtis index revealed statistical differences between groups for both nasopharyngeal swabs and stools samples ($p < 0.001$; PERMANOVA) (Figure 1C and 1D). As a result, microbiota is grouped in three separate clusters depending on the severity of the symptoms, which indicates that microbiota differences are based on changes in bacterial composition.

	Mild (n=24)	Moderate (n=51)	Severe (n=31)			
Clinical variables				a p-value	b p-value	c p-value
Mean age (SD), years	43 ± 12	54 ± 14 ^b	62 ± 11 ^{a,c}	0.000001	0.0041	0.017
Male, n (%)	8 (33)	24 (47) ^b	22 (71) ^{a,c}	0.0071	0.032	0.041
Symptoms, n (%)						
Presence of dyspnoea	6 (33)	38 (75) ^b	26 (84) ^a	0.00018	0.001	-
Presence of gastrointestinal alteration	5 (26)	17 (33)	10 (33)		-	
High respiratory rate	1 (4)	11 (22)	19 (63) ^{a,c}	0.00018	-	0.004
Low sPO2	6 (33)	22 (43)	22 (71) ^{a,c}	0.001	-	0.021
High heart rate	1 (4)	14 (27) ^b	17 (55) ^{a,c}	0.0008	0.02	0.01
Comorbidities, n (%)						
Obesity	5 (26)	14 (27)	10 (32)		-	
Diabetes	4 (20)	9 (18)	8 (26)		-	
Asthma	3 (3)	3 (6)	2 (7)		-	
Cardiomyopathy	3 (3)	3 (6)	13 (42) ^{a,c}	0.02	0.001	-
Plasma determinations						
Mean lymphocytes (SD), 103/μL	1.1 ± 0.6	1.4 ± 0.6	1.2 ± 2.7		-	
Median neutrophils (IQR), 103/μL	6 [5.5;6.6]	6.4 [4.2;8.6]	7.9 [5.4;10.9]		-	
Mean platelets (SD), 103/μL	329.6 ± 8.5	257.4 ± 115 ^b	276.4 ± 93 ^a	0.031	0.018	-
Median D dimer (IQR), mg/L	0.39 [0.2;0.8]	0.6 [0.3;1]	1.6 [0.9;4.3] ^{a,c}	0.0001	-	0.0001
Median ferritin (IQR), ng/L	157 [126;179]	487 [274;1027] ^b	829 [488;1376] ^a	0.0001	0.0001	-
Median C reactive protein (IQR), mg/L	3.4 [2.6;4]	18.2 [7.8;41.9] ^b	162 [65;210] ^{a,c}	0.0001	0.0002	0.0001

Table 1.

Clinical data description of enrolled patients.

Differences were displayed as: ^a p<0.05 Severe sv. Mild; ^b p<0.05 Moderate vs. Mild; ^c p<0.05 Severe vs. Moderate. ANOVA or Kruskal were employed for numerical variables and Fisher's test for categorical variables.




Figure 1.

Nasopharyngeal and gut microbiota composition is modified depending on the severity of COVID-19 symptoms.

(A) Alpha diversity index for nasopharyngeal swab samples microbiota. (B) Alpha diversity index for stool samples microbiota. (C) PCoA for Bray-Curtis index of nasopharyngeal swab microbiota. (D) PCoA for Bray-Curtis index of stool samples microbiota. Values are represented as mean $\pm$ SD. Differences were displayed as: ^a $p < 0.05$ Severe vs. Mild; ^b $p < 0.05$ Moderate vs. Mild; ^c $p < 0.05$ Severe vs. Moderate. PERMANOVA test was employed to determine Bray-Curtis significance differences.

Similarly, the characterization of microbiota abundance composition revealed heterogeneity in the microbiota profile associated with severity and disease progression in these patients. Relative abundance levels for both phylum and genera were evaluated using the median abundance as a threshold. Specifically, genera with a median relative abundance greater than 0.5% were considered for detailed analysis.

At phylum level, in nasopharyngeal microbiota, the abundance of *Bacillota* was increased while the abundance of *Bacteroidota* and *Actinobacteroidota* was reduced in patients with severe symptomatology (**Figure 2A** [↗](#)). Conversely, the faecal microbiota of the three groups presented a more homogeneous distribution than the nasopharyngeal one, being the most abundant phyla *Bacillota* and *Bacteroidota* (**Figure 2B** [↗](#)). However, patients with severe symptoms, thus with a worse prognosis, showed a lower abundance of *Bacteroidota* (**Figure 2B** [↗](#)). Exact values of relative abundance for different phyla are shown in supplementary file 1.

At genus level, the nasopharyngeal microbiome composition revealed that symptom severity was associated with a higher number of detected genera (**Figure 2C** [↗](#)). Of note, significant differences in genus abundance for each group of patients were found. Thus, mild patients presented a significantly higher abundance of *Alistipes*, *Muribaculaceae*, *Lactobacillus* and *Lachnospiraceae* ($p < 0.05$; Kruskal-Wallis), whereas the group with moderate symptoms showed a significant increase in *Alcaligenes*, *Pseudorobacter* and *Pseudomonas* ($p < 0.05$; Kruskal-Wallis). Additionally, patients with more severe disease had significantly higher relative abundance of *Acinetobacter*, *Actinomyces*, *Anaerococcus*, *Atopobium*, *Campylobacter*, *Dolosigranulum*, *Enterobacter*, *Enterococcus*, *Fingoldia*, *Fusobacterium*, *Gemella*, *Haemophilus*, *Lawsonella*, *Leptotrichia*, *Megasphaera*, *Neisseria*, *Serratia*, *Rotia* and *Veillonella* ($p < 0.05$; Kruskal-Wallis). Unlike nasopharyngeal swabs, stool samples showed a reduction in the number of detected genera as symptom severity increased (**Figure 2D** [↗](#)). Precisely, mild patients showed more presence of *Blautia*, *Muribaculaceae* and different members of the *Clostridia* class (*Clostridia*, *Coprococcus* and *Ruminococcus*) than the other groups ($p < 0.05$; Kruskal-Wallis). In moderate ill patients *Bacteroides*, *Barnesiella*, *Faecalibacterium*, *Parabacteroides* and *Streptococcus* were significantly increased ($p < 0.05$; Kruskal-Wallis). Remarkably, *Alistipes*, *Anaerococcus*, *Dialister*, *Fingoldia*, *Lachnocostidium*, *Prevotella* or *Peptoniphilus* were more abundant in patients with severe symptoms ($p < 0.05$; Kruskal-Wallis).

Relative abundance levels for genera identified bacteria and exact p-values can be found in supplementary files 2 and 3.

Differences in bacteria composition could be used as biomarkers to predict disease severity and outcome in SARS-CoV-2 infection

We investigated whether some specific taxa could be associated with the severity of the symptoms. ASVs were evaluated to determine core taxa along with the specific bacteria of each group of patients and samples (**Figure 3A,B** [↗](#)). Core taxa were identified using a detection threshold of 0.1 and a prevalence threshold of 0.1. In nasopharyngeal swabs, Venn diagram revealed that the three cohorts shared 51 core taxa. 60 specific bacteria were identified in patients with mild symptoms while 32 were observed in patients with moderate symptoms and 8 in patients with severe symptoms. When stools were considered, the three cohorts shared 159 core taxa. 27 were specific of mild patients, another 33 of patients with moderate symptoms and 27 of the severe patients (For more details see supplementary file 4).

Besides, the linear discriminant analysis (LEfSe) was performed to identify differential microorganisms for each group of patients (**Figure 3C,D** [↗](#)). In nasopharyngeal samples from mild patients, *Burkholderia* sp., *Paraburkholderia* sp. and *Massilia* sp. were determined; in moderate patients, *Pseudomonas veronii*, *Stenotrophomonas rhizophila* and *Azotobacter chroococcum*; and in severe patients, *Mycoplasma salivarium*, *Prevotella dentalis*, *Leptotrichia* and *Haemophilus*




Figure 2.

Microbiota composition of nasopharyngeal and stool samples at phylum level is slightly modified by COVID-19 symptoms severity. In contrast, at genus level, severity increases the total amount of detected bacteria in nasopharyngeal swabs while in stool samples it is promoting a reduction.

(A) Representation of the relative abundance of the main phyla in nasopharyngeal swab samples. (B) Representation of the relative abundance of the main phyla in stool samples. (C) Relative abundance of the principal genera detected in nasopharyngeal swab samples. (D) Relative abundance of the principal genera detected in nasopharyngeal swab samples. For C and D only those ASVs with a median higher to 0.5 were chosen.




Figure 3.

Differential analysis expression of microbiota composition from nasopharyngeal and stool samples revealed the presence of specific bacteria related to COVID-19 severity index.

(A) Venn diagram showing ASVs distribution in nasopharyngeal swab samples. (B) Venn diagram showing ASVs distribution in stool samples. (C) LEfSe plot of taxonomic biomarkers present in nasopharyngeal swab samples (p value = 0.01 and LDA value = 4). (D) LEfSe plot of taxonomic biomarkers present in stool samples (p value = 0.01 and LDA value = 4). Venn diagrams were acquired with the following parameters: detection level = 0.01 and prevalence level = 0.01.

parainfluenzae. On the other hand, in stool samples, *Bacteroides coprocola*, *Veillonella sp.*, *Ruminococcus bicirculans* and *Sutterella stercoricanis* were identified as predictors of mild condition; *Prevotella stercorea*, *Bacteroides cellulosilyticus*, *Streptococcus salivarius*, *Bacteroides stercoris* and *Prevotella copri* as predictors of moderate status; and *Escherichia*, *Enterococcus durans*, *Alistipes onderdonkii*, *Prevotella timonensis* and *Prevotella bivia* as markers of severe condition. Unlike the Venn diagram, the presence of a species in the LEfSe analysis does not imply its absence in the other groups, but rather indicates significant differences in abundance. Hence, detailed abundance of these bacteria can be found in supplementary files 5 and 6.

To further assess the role of these potential biomarkers in the prediction of COVID-19 severity, a correlation analysis was performed between these species and symptomatology. In summary, biomarkers identified in nasopharyngeal swabs in patients with severe condition, *M. salivarium* and *Leptotrichia*, showed a positive correlation with D dimer and cardiomyopathy, respectively ($R > 0.3$). In addition, the other two biomarkers related to the highest severity, *H. parainfluenzae* and *P. dentalis*, also showed a positive association with CRP, D dimer and cardiomyopathy, even though it was not statistically significant ($R = 0.2$). More precisely, *M. salivarium* and *Leptotrichia* along with *P. dentalis* showed a significantly negative correlation towards lymphocytes count ($p < 0.05$; $R < -0.25$) (Figure 4A). In the case of stool samples, biomarkers for mild symptomatology (*B. coprocola*, *R. bicirculans* and *S. stercoricanis*) presented a negative correlation with CRP levels and respiratory rate ($p < 0.05$; $R < -0.25$). In contrast, the severe biomarkers identified, presented important correlations towards dyspnoea, cardiomyopathy or respiratory rate. Concretely, *P. bivia* and *P. timonensis*, revealed a positive correlation with D dimer, ferritin levels and respiratory rate, as well as a negative correlation with lymphocyte count ($R > 0.6$) (Figure 4B). Additionally, these two species showed a significant positive correlation against CRP levels and a negative correlation towards lymphocyte count ($p < 0.05$). Exact correlation and p-values are shown in supplementary file 7.

Identification of a novel microbiome-based COVID-19 prognosis approach

Considering the possible identified biomarkers and their relationship towards clinical variables, we proposed a new approach to predict disease severity in patients suffering SARS-CoV-2 infection by evaluating correlation among specific bacteria abundance from nasopharyngeal and stool samples. The analysis revealed no important associations between nasopharyngeal and faecal microbiota in mild and moderate patients (Figure 5A,B). Nonetheless, in patients with severe symptoms the Spearman's rho coefficient showed a significant positive correlation between *P. timonensis* towards *P. dentalis* and *M. salivarium* (Figure 5C). Consequently, the ratios *P. timonensis* / *M. salivarium* and *P. timonensis* / *P. dentalis* were calculated to determine whether they could be used as a predictor of COVID-19 severity. Both seemed to be significantly increased in patients with severe symptoms compared to those with mild or moderate symptomatology ($p < 0.05$; Kruskal-Wallis) (Figure 5D,E). As a result, the ratio between the abundance of these bacteria could serve as reliable predictors of severity of COVID-19.

Discussion

Recent findings have evidenced the prominent role of the microbiome in several diseases, including those caused by viruses, proving that commensal microbiota could either enhance or hinder viral infections [30, 31]. A large body of evidence indicates that respiratory virus infections can induce a microbial imbalance in the airways, increasing the risk of secondary bacterial infections and worsening the prognosis [32]. Besides, alterations in the gut microbiota have also been noted during respiratory virus infections, likely influenced by the "gut-lung axis" [33]. In this regard, different studies have tried to establish relationships between microbiota




Figure 4.

Whereas mild biomarkers showed negative correlations towards clinical variables, severe biomarkers presented positive correlations.

(A) Correlation plot of nasopharyngeal swab biomarkers and clinical variables. **(B)** Correlation plot of stool samples biomarkers and clinical variables. RR: respiratory rate; HR: heart rate; GI: gastrointestinal alterations. Correlation was calculated by taking into account abundance levels of each bacteria against the clinical variable of severe ill patients. Red asterisks stand for p-values < 0.05 and correlations greater than or equal to 0.3 or less than or equal to -0.25.




Figure 5.

The existence of a relationship between the abundance of nasopharyngeal severe biomarkers and stool severe biomarkers allows the employment of an abundance ratio between them as a new tool for predicting COVID-19 severity.

(A) Correlation plot among biomarkers found in nasopharyngeal swab and stool samples in mild condition. (B) Correlation plot among biomarkers found in nasopharyngeal swab and stool samples in moderate condition. (C) Correlation plot among biomarkers found in nasopharyngeal swab and stool samples in severe condition. (D) Ratio of the abundance between *P. timonensis* (stool) and *M. salivarium* biomarkers. (E) Ratio of the abundance between *P. timonensis* (stool) and *P. dentalis* (nasopharyngeal swab) biomarkers. ^a $p < 0.05$ Severe vs. Mild; ^b $p < 0.05$ Moderate vs. Mild.

composition and SARS-CoV-2 infection [34, 35] but just a few have explored microbiota compositional variations among patients with different symptomatology [36]. However, only a few studies have explored the nasopharyngeal-faecal axis as a potential biomarker of severity in patients infected with SARS-CoV-2. Hence, the present study provides some biomarkers located in nasopharyngeal and gut microbiota that could help to predict COVID-19 severity.

In this context, 106 patients with SARS-CoV-2 infection were included in this study and classified into 3 severity groups according to their symptoms. To ensure results were derived only from SARS-CoV-2 infection, included patients were not positive for other viral infections. Considering clinical parameters, the current study confirmed that age and gender are positively associated with more severe symptoms like dyspnoea, increased heart and respiratory rates together with lower oxygen saturation. Results that are in line with other COVID-19 studies conducted globally, which have shown that men and women are disproportionately affected. Data revealed that males suffered from more severe disease than females, including higher ICU admission rates, dyspnoea, increased heart rate, being all clinical signs for higher severity of the COVID-19 disease [37, 38]. Regarding some biochemical parameters those groups of patients with moderate or severe symptoms showed significantly increased levels of D-dimer, ferritin and CRP, which are typically described as biomarkers for COVID-19 severity [39–41]. Interestingly, previous studies have reported that alterations in the gut microbiota composition in COVID-19 patients could be also associated with disease severity, measured with the elevated concentration of the blood markers described above. [42–44]. In this sense, we have confirmed these results in our study as specific bacteria from severe condition positively correlates with the levels of ferritin, CRP, D-dimer or lymphocytes. Consequently, the microbiota could be key in the clinical phenotype of COVID-19 patients, although its specific contribution to the progression of the infection and a poor prognosis needs to be better understood.

The results of alpha diversity revealed only substantial changes in richness in the nasopharyngeal microbiome associated with moderate and severe symptoms. Although most of the studies have proposed that SARS-CoV-2 infection is associated with lower microbial diversity in nasopharyngeal samples [13, 45, 46], others did not find differences in alpha diversity composition among groups with different symptomatology [47, 48]. This could be explained by the diverse methods of sample collection, and also the SARS-CoV-2 variant and the treatment with antibiotics [49].

Furthermore, in the present study, beta diversity analysis showed that in both types of samples, every group of patients presented distinctly differentiated clusters. As previously reported, COVID-19 disease severity is more dependent on the presence or absence of certain bacteria rather than alterations in bacterial diversity and richness [50]. This argument was supported by the characterization of bacterial microbiota composition at phyla and genera levels. In nasopharyngeal swabs, the abundance of *Bacteroidota* and *Actinobacteriota* was reduced compared to mild patients, which goes in accordance with previous reports where these bacteria have been previously linked to a better prognosis of SARS-CoV-2 infection, since can exert beneficial effects by preventing respiratory diseases, including COVID-19 [51–55]. Moreover, the abundance of *Bacillota* and *Pseudomonadota* was increased in patients with severe symptomatology, thus supporting previous studies in which higher counts of *Bacillota* (*Staphylococcus* sp. and *Streptococcus* sp.) and *Pseudomonadota* (*Pseudomonas* sp.) were associated with moderate and severe symptoms of COVID-19 [56].

Regarding genera composition, *Alistipes* and *Muribaculaceae* were highly abundant in patients with mild symptoms. These bacteria have been well characterised in gut microbiota, but little is known about their presence in nasopharyngeal microbiota. Different experimental studies in mice have suggested their role in viral infections. Thus, *Muribaculaceae* was found in the lung microbiota in SARS-CoV-2 infected mice that were treated with a selective inhibitor of the main protease (M^{pro}) [57]. In the case of *Alistipes*, in a study conducted in children infected with

respiratory syncytial virus (RVS), its abundance was reduced in the nasopharyngeal microbiota in comparison with non infected subjects [58]. Overall, these genera could be associated with a protective role against viral infection, and their higher presence in nasopharyngeal samples from mild COVID-19 patients could prevent the progression to severe disease. In contrast, the increased presence of other genera, such as *Corynebacterium*, *Acinetobacter*, *Staphylococcus* and *Veillonella*, was associated with the severity of SARS-CoV-2 infection. This is supported by previous studies in which these genera were associated with both disease severity and systemic inflammation [59]. In addition, higher abundance of *Enterococcus* was observed in severe patients, thus confirming other studies in critically ill patients [60].

On the contrary, findings at phylum level in stool samples did not reveal notable modifications among patient groups. However, in mild ill patients, different genera from *Clostridia* class (*Clostridia*, *Coproccoccus*, *Dorea*, *Lachnospiraceae*, *Roseburia* and *Ruminococcus*), and *Barnesiella* were identified as highly abundant. While the class *Clostridia* was associated with a reduced production of proinflammatory cytokines in COVID-19 patients and in those who recovered from the infection [61, 62], *Barnesiella* prevents colonisation by antibiotic-resistant bacteria such as *Enterococcus*, which has been reported as a frequent cause of systemic infection in critically ill COVID-19 patients [63, 64]. Thus, the reduction of gut abundance of *Barnesiella* and *Clostridia* members could be associated with more severe symptoms in COVID-19 patients. Moreover, the specific genera detected in the severe illness group, *Lachnocostridium*, *Anaerococcus* and *Peptoniphilus*, have been previously recognised as opportunistic pathogens, which could induce gut inflammation and contribute to a poor prognosis [65].

Considering the variations observed in both nasopharyngeal and gut microbiota composition and their miscellaneous effects, the identification of specific bacteria could be used as a biomarker to predict COVID-19 severity. Accordingly, the present study has identified unique ASVs for the different severities of COVID-19. For nasopharyngeal samples, species belonging to the genus *Lactobacillus* (*L. fermentum* or *L. reuteri*) or *Prevotella* (*P. pallens*, *P. ori* and *P. shahii*) have been pinpointed. Interestingly, mild patients have also shown a higher content of *Lactobacillus*, which could also contribute to the host defence against viral infection at early stages, as reported in asymptomatic COVID-19 patients [66]. In the case of *Prevotella* sp., its role in COVID-19 infection has not been clearly elucidated, but previously published microbiome analysis has revealed that its abundance was higher in mild patients [67]. Nevertheless others have suggested that it could be a biomarker of critical phenotype in COVID-19 patients [68, 69]. It is interesting to highlight that *Anaerococcus prevotii* was one of the species exclusively found in stools in mild patients. This microorganism has already been linked to lower inflammation in COVID-19 patients [70]. Conversely, *Coprobacillus cateniformis* was only found in severe patients, and could be involved in the development of a worse condition in these patients through ACE2 upregulation [17]. Even though these bacteria were found to be unique for each group, LEfSe was performed to obtain specific severity prognostic biomarkers depending on the abundance threshold. Therefore, for mild patients, *Burkholderia* and *Paraburkholderia* were identified in nasopharyngeal swabs and *B. coprocola* and *R. bicirculans* in stool samples. Although the information regarding the first two species in humans is limited, a few studies have reported their presence in the commensal human microbiota [71, 72]. Contrarily, *B. coprocola* and *R. bicirculans* have been found in both healthy and COVID-19 patients, being the abundance of *R. bicirculans* reduced in infected subjects [73].

Correspondingly, in patients with moderate symptoms, *P. veronii* was detected in nasopharyngeal samples whereas *P. stercorea*, *B. cellulosilyficus*, *B. stercoris* and *P. copri* were identified in stool samples. In general, these findings agree with previous studies [74]. Xu *et al.* found that infected patients showed higher abundance of *B. cellulosilyficus* [75], whereas *B. stercoris* and *P. copri* were associated with ACE2 upregulation and increased proinflammatory cytokine production, respectively, in COVID-19 patients [73, 76].

Likewise, in critically ill patients, the biomarkers detected for nasopharyngeal microbiota were *M. salivarium*, *P. dentalis*, *Leptotrichia* and *H. parainfluenzae*, while, in stool samples, *Escherichia* sp., *E. durans*, *A. Onderdonkii*, *P. timonensis* and *P. bivia* were the species recognised as biomarkers. Both *P. bivia* and *P. timonensis* have been defined as unique microorganisms in COVID-19 patients microbiota [73, 77], whilst a higher abundance of *M. salivarium*, *H. parainfluenzae* and *E. durans* were related to poor prognosis [78–80]. When considering *A. onderdonkii*, it is a short-chain fatty acid-producing bacteria that has been reported to be inversely associated with COVID-19 severity [17, 50]; however, there are conflicting evidences regarding its pathogenicity that indicate that *A. onderdonkii* may have protective effects against some diseases, including COVID-19 [17]. Nonetheless, considering that Zuo *T et al.* employed a different methodology to analyse microbiota composition from stool samples, *A. onderdonkii* could be considered as a biomarker of severe condition in SARS-CoV-2 infected subjects.

Of note, the use of these bacteria as biomarkers of severity in SARS-CoV-2 infection is further supported by the fact that these species exhibited positive correlations with critical clinical variables mentioned above. Specifically, possible severe biomarkers from nasopharyngeal samples such as *M. salivarium*, *H. parainfluenzae* and *P. dentalis* showed a negative correlation towards lymphocytes count. This also happens for stool severe biomarkers *P. bivia* and *P. timonensis*. As it is well established that microbiota can modify blood cells count [81], it would not be a surprise that these bacteria could promote COVID-19 severity by lowering lymphocyte count. In addition, *P. bivia* and *P. timonensis* showed positive correlation with ferritin, CRP and D-dimer levels, as well as cardiomyopathy and respiratory rates. Several studies have revealed both the relationship between CRP levels, gut microbiota and COVID-19 severity [82], as well as the positive correlation of specific bacteria with D-dimer, CRP and the levels of pro-inflammatory mediators in plasma [83].

Finally, although the composition of the nasal and faecal microbiota in COVID-19 patients has been previously studied [32], the interaction between these microbiotas has not been extensively evaluated. In this study, we analyzed the relationship between specific bacteria from nasopharyngeal and stool samples, confirming a correlation between the abundances of species in these different sample types. Concretely, a strong positive correlation between *P. timonensis* (in stool samples) towards *P. dentalis* and *M. salivarium* (in nasopharyngeal samples) was found in severe patients. To maximise the potential use of these biomarkers, the ratio of the abundance of these species was also significantly increased within the highest severity of this condition. As a result, the ratio proposed in this study could be used as a novel predictor to identify critically ill COVID-19 patients as the ratio *Bacillota* and *Bacteroidetes* has been used as a marker of dysbiosis [84]. In this case, the ratio *P. timonensis*/*P. dentalis* and *P. timonensis* /*M. Salivarium* could be a prognostic tool for severe SARS-CoV-2, and its increase could be associated with a higher risk COVID-19 severity development.

Materials and methods

Ethics approval

The study was conducted in accordance with the declaration of Helsinki and the protocol approved by the Clinical Research Ethics Committee of Granada (CEIC) (ID of the approval omicovid-19 1133-N-20). All patients provided written informed consent before being included in the study. The samples were managed by the ibs.GRANADA Biobank following the protocols approved by the Andalusian Biomedical Research Ethics Coordinating Committee.

Subject recruitment and sample collection

A multicenter prospective observational cohort study was carried out between September 2020 and July 2021. Patients with SARS-CoV-2 infection were recruited from the University Hospital San Cecilio, the University Hospital Virgen de las Nieves, and the Primary Care centres, Salvador Caballero and Las Gabias in Granada (Spain). These patients were Caucasian and laboratory-confirmed SARS-CoV-2 positive by quantitative reverse transcription polymerase chain reaction (RT-qPCR) performed on nasopharyngeal swabs collected by healthcare practitioners. No confounding factors such as other viral infections were present. Patients were classified in three groups based on severity profile following the described guidelines [21]; mild (n=24), moderate (n=51) and severe/critical (n=31). Mild illness includes individuals who have any of the various signs and symptoms of COVID-19 (e.g., fever, cough, sore throat, malaise, headache, muscle pain, nausea, vomiting, diarrhoea, loss of taste and smell) but do not have shortness of breath, dyspnoea, or abnormal chest imaging. Moderate illness includes patients that show fever and respiratory symptoms with radiological findings of pneumonia. Severe/critical illness includes patients with any of the following criteria: respiratory distress (≥ 30 breaths/min), oxygen saturation $\leq 93\%$ at rest, arterial partial pressure of oxygen (PaO_2)/fraction of inspired oxygen (FiO_2) ≤ 300 mmHg, respiratory failure and requiring mechanical ventilation, shock, and with other organ failures that required intensive care unit (ICU) care.

Nasopharyngeal swabs and stools samples from patients were collected by healthcare staff in the moderate and severe/critical cohorts, while the patients of the mild cohort provided stools self-sampled at home. Stools and nasopharyngeal swabs were collected in collection tubes containing preservative media (OMNIGene®•GUT, DNAGENOTEK®, Ottawa, Ontario, Canada) and stored at -80°C until processing.

Microbial DNA extraction, library preparation and next generation sequencing

For all faecal and nasopharyngeal samples, DNA was isolated according to the modified protocol reported by Rodríguez-Nogales *et al.* [22] using Qiagen Allprep PowerFecal DNA kit (Qiagen, Germany). DNA was quantified using Qubit dsDNA HS assay kit (12640ES60, Yeason Biotechnology, Shanghai, China) and total DNA was amplified by targeting variable regions V3-V4 of the bacterial 16 S rRNA gene. Quality control of amplified products was achieved by running a high-throughput Invitrogen 96-well-E-gel (Thermo Fisher Scientific, Waltham, MA, USA). PCR products from the same samples were pooled in one plate and normalised with the high-throughput Invitrogen SequaPrep 96-well Plate kit. Pool samples were sequenced using an Illumina MiSeq.

Bioinformatic tools

Paired end reads quality was checked with FastQC software [23]. Trimming of adapters and filtering of low quality sequences was performed with Trimmomatic [24]. Filtered reads were further processed with QIIME2 software (open access, Northern Arizona University, Flagstaff, AZ, USA) by employing DADA2 software [25] to carry out denoising steps to obtain amplicon sequence variants (ASVs). SILVA reference database (138 99% full length) was used for taxonomic assignment [26]. Microbiota results were further analyze with the help of different R packages. Alpha and beta diversity as well as relative abundance were appraised with the *Phyloseq* package [27]. Beta diversity differences were obtained with a Permutational Multivariate Analysis of Variance (PERMANOVA) included in the *Vegan* package (*Eulerr* along with *microbiomeutilities* package [28] was utilised for constructing Venn diagrams [29]. Meanwhile, *microbial* package was employed to perform linear discriminant analysis (LDA) effect size (LEfSe) with an LDA score of 3 [29].

Statistical analysis

All data presented were analyzed and visualized using R software. For numerical clinical variables, normality and homoscedasticity were assessed using the `shapiro.test` and `leveneTest` functions from the *stats* and *car* package, respectively. Data were displayed as mean $\pm$ standard deviation (SD) for normally distributed variables, whereas median and interquartile range (IQR) were used for variables with non-normal distributions.

Statistical differences were evaluated using ANOVA for parametric data or Kruskal-Wallis test for non-parametric data, both available in the *stats* package. When significant differences were identified, post-hoc group comparisons were conducted using Tukey's method through the `TukeyHSD` function for ANOVA and Dunn's test via the `dunnTest` function from the *FSA* package for Kruskal-Wallis.

Categorical variables were expressed as percentages, and statistical significance was determined using Fisher's exact test with the `fisher.test` function from the *stats* package. Finally, for correlation analysis, Spearman's correlation coefficient was employed for non-normally distributed variables, while Pearson's correlation coefficient was used for normally distributed variables. The `cor` function from the *stats* package was used to calculate these coefficients, and the *corrplot* package was employed for visualization. Statistical significance was defined as a p-value of less than 0.05. To illustrate group differences, different letters (a, b or c) were used to denote statistically significant differences among groups. All data, models, and analytical output are on the linked GitHub repository <https://github.com/albarodnog/COVID-INMUNOBIOTICS-Group.git>.

Conclusion

This inter-individual variability between the COVID-19 patients could contribute to the different symptomatology observed. This study has identified a correlation between changes in the nasopharyngeal and stool microbiota with COVID-19 severity. A novel biomarker linked to severity of COVID-19 infection has been described based on changes in the abundance of bacterial species in nasopharyngeal and faecal samples. This knowledge can support the development of biomarkers to gauge disease severity and, also, the design of novel therapeutic strategies to mitigate adverse outcomes. Further investigations are imperative to explore how the association between nasopharyngeal and faecal microbiota can be modulated to uncover its role in enhancing immune health, preventing or treating SARS-CoV-2 infections, and fostering immunity.

Data availability statement

Participant data cannot be made publicly available due to the sensitive nature of the personal health data and privacy and confidentiality reasons. However, under certain conditions, these data could be accessible for statistical and scientific research. For further information, please contact the corresponding authors.

Acknowledgements

We acknowledge the collaboration of all the participants who voluntarily and selflessly participated in the study.

Additional information

Conflict of interest statement

All authors declare no interest.

Funding information

The research project was supported by Government of Andalusia (Spain) (CV20-99908).

Author contributions

Benita Martín-Castaño, Margarita Martínez-Zaldívar, Emilio Mota, Fernando Cobo, Concepcion Morales-García, Marta Alvarez-Estevéz, Federico García, Silvia Merlos, Paula García-Flores, Manuel Colmenero-Ruiz, José Hernández-Quero, María Nuñez were involved in the sample collection. Patricia Díez-Echave, Jorge García-García, Alba Rodríguez-Nogales, María Elena Rodríguez-Cabezas, Laura Hidalgo-García, Antonio Jesús Ruiz-Malagon, José Alberto Molina-Tijeras, María Jesús Rodríguez-Sojo and Anaïs Redruello were involved in the processing of samples and obtaining results. Rocio Morón and Emilio Fernández-Varón had access to the data and were involved in the conception and data analysis and interpretation. Alba Rodríguez-Nogales, Javier Martín, María Elena Rodríguez-Cabezas, Benita Martín-Castaño, Rocio Morón, Jorge García-García, Ángel Carazo and Julio Gálvez had access to the data and were involved in the conception and design of the work, data analysis and interpretation, critical revision of the article and final approval before submission. All authors reviewed the final manuscript and agreed to be accountable for all aspects of the work.

References

1. WHO (2023) **WHO COVID-19 dashboard** Geneva: World Health Organization
2. Harvey W.T., et al. (2021) **SARS-CoV-2 variants, spike mutations and immune escape** *Nat Rev Microbiol* **19**:409–424
3. Crook H., et al. (2021) **Long covid-mechanisms, risk factors, and management** *BMJ* **374**
4. Yang J., et al. (2020) **Prevalence of comorbidities and its effects in patients infected with SARS-CoV-2: a systematic review and meta-analysis** *Int J Infect Dis* **94**:91–95
5. Severe Covid, G.G., et al. (2020) **Genomewide Association Study of Severe Covid-19 with Respiratory Failure** *N Engl J Med* **383**:1522–1534
6. Bastard P., et al. (2020) **Autoantibodies against type I IFNs in patients with life-threatening COVID-19** *Science* **370**
7. Zhou P., et al. (2020) **A pneumonia outbreak associated with a new coronavirus of probable bat origin** *Nature* **579**:270–273
8. Peng L., et al. (2020) **SARS-CoV-2 can be detected in urine, blood, anal swabs, and oropharyngeal swabs specimens** *J Med Virol* **92**:1676–1680
9. Sun J., et al. (2020) **Isolation of infectious SARS-CoV-2 from urine of a COVID-19 patient** *Emerg Microbes Infect* **9**:991–993
10. Zhang H., et al. (2020) **Digestive system is a potential route of COVID-19: an analysis of single-cell coexpression pattern of key proteins in viral entry process** *Gut* **69**:1010–1018
11. Lamers M.M., et al. (2020) **SARS-CoV-2 productively infects human gut enterocytes** *Science* **369**:50–54
12. Zang R., et al. (2020) **TMPRSS2 and TMPRSS4 promote SARS-CoV-2 infection of human small intestinal enterocytes** *Sci Immunol* **5**
13. Candel S., et al. (2023) **The nasopharyngeal microbiome in COVID-19** *Emerg Microbes Infect* **12**
14. Ancona G., et al. (2023) **Gut and airway microbiota dysbiosis and their role in COVID-19 and long-COVID** *Front Immunol* **14**
15. De Bruyn A., et al. (2022) **Secondary infection in COVID-19 critically ill patients: a retrospective single-center evaluation** *BMC Infect Dis* **22**
16. Gu S., et al. (2020) **Alterations of the Gut Microbiota in Patients With Coronavirus Disease 2019 or H1N1 Influenza** *Clin Infect Dis* **71**:2669–2678
17. Zuo T., et al. (2020) **Alterations in Gut Microbiota of Patients With COVID-19 During Time of Hospitalization** *Gastroenterology* **159**:944–955

18. Wu Y., et al. (2021) **Altered oral and gut microbiota and its association with SARS-CoV-2 viral load in COVID-19 patients during hospitalization** *NPJ Biofilms Microbiomes* 7
19. Cao J., et al. (2021) **Integrated gut virome and bacteriome dynamics in COVID-19 patients** *Gut Microbes* 13:1–21
20. Mancabelli L., et al. (2023) **Disentangling the interactions between nasopharyngeal and gut microbiome and their involvement in the modulation of COVID-19 infection** *Microbiology Spectrum* 11:e02194–23
21. National Institutes of Health (2019) **NIH COVID-19 Treatment Guidelines**
22. Rodriguez-Nogales A., et al. (2017) **Differential intestinal anti-inflammatory effects of *Lactobacillus fermentum* and *Lactobacillus salivarius* in DSS mouse colitis: impact on microRNAs expression and microbiota composition** *Mol Nutr Food Res* 61
23. Bittencourt S. (2010) **FastQC: A quality control tool for high throughput sequence data** *Babraham Bioinformatics*
24. Bolger A.M., Lohse M., Usadel B (2014) **Trimmomatic: A flexible trimmer for Illumina Sequence Data** *Bioinformatics*
25. Callahan B.J., et al. (2016) **DADA2: High-resolution sample inference from Illumina amplicon data** *Nat Methods* 13:581–3
26. Kaehler B.D., et al. (2019) **Species abundance information improves sequence taxonomy classification accuracy** *Nat Commun* 10
27. Love MI, Anders S (2014) **Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2** *Genome Biology* 15
28. Sudarshan A., Shetty L.L. (2020) **microbiomeutilities: Utilities for Microbiome Analytics** *The microbiome R package relies on the independently developed*
29. R.C.T. and R: A language and environment for statistical computing (2013) **R Foundation for Statistical Computing**
30. Mutlu E.A., et al. (2014) **A compositional look at the human gastrointestinal microbiome and immune activation parameters in HIV infected subjects** *PLoS Pathog* 10
31. Guo X., et al. (2023) **Abnormal blood microbiota profiles are associated with inflammation and immune restoration in HIV/AIDS individuals** *mSystems* 8
32. Li J., et al. (2023) **Assessment of microbiota in the gut and upper respiratory tract associated with SARS-CoV-2 infection** *Microbiome* 11
33. Budden K.F., et al. (2017) **Emerging pathogenic links between microbiota and the gut-lung axis** *Nat Rev Microbiol* 15:55–63
34. Yamamoto S., et al. (2021) **The human microbiome and COVID-19: A systematic review** *PLoS One* 16
35. Di Stadio A., et al. (2020) **The Microbiota/Host Immune System Interaction in the Nose to Protect from COVID-19** *Life (Basel)* 10

36. Nguyen L.H., et al. (2023) **Metagenomic assessment of gut microbial communities and risk of severe COVID-19** *Genome Med* 15
37. Pijls B.G., et al. (2022) **Temporal trends of sex differences for COVID-19 infection, hospitalisation, severe disease, intensive care unit (ICU) admission and death: a meta-analysis of 229 studies covering over 10M patients** *F1000Res* 11
38. Mauvais-Jarvis F. (2020) **Aging, Male Sex, Obesity, and Metabolic Inflammation Create the Perfect Storm for COVID-19** *Diabetes* 69:1857–1863
39. Gomez-Pastora J., et al. (2020) **Hyperferritinemia in critically ill COVID-19 patients - Is ferritin the product of inflammation or a pathogenic mediator?** *Clin Chim Acta* 509:249–251
40. Rostami M., Mansouritorghabeh H. (2020) **D-dimer level in COVID-19 infection: a systematic review** *Expert Rev Hematol* 13:1265–1275
41. Zhou C., et al. (2020) **Increased Serum Levels of Hepcidin and Ferritin Are Associated with Severity of COVID-19** *Med Sci Monit* 26
42. Liu Q., et al. (2022) **Gut microbiota dynamics in a prospective cohort of patients with post-acute COVID-19 syndrome** *Gut* 71:544–552
43. Liu Q., et al. (2022) **Multi-kingdom gut microbiota analyses define COVID-19 severity and post-acute COVID-19 syndrome** *Nat Commun* 13
44. Yeoh Y.K., et al. (2021) **Gut microbiota composition reflects disease severity and dysfunctional immune responses in patients with COVID-19** *Gut* 70:698–706
45. Gao M., et al. (2021) **Characterization of the Human Oropharyngeal Microbiomes in SARS-CoV-2 Infection and Recovery Patients** *Adv Sci (Weinh)* 8
46. Gupta A., et al. (2022) **Nasopharyngeal microbiome reveals the prevalence of opportunistic pathogens in SARS-CoV-2 infected individuals and their association with host types** *Microbes Infect* 24
47. De Maio F., et al. (2020) **Nasopharyngeal Microbiota Profiling of SARS-CoV-2 Infected Patients** *Biol Proced Online* 22
48. Braun T., et al. (2021) **SARS-CoV-2 does not have a strong effect on the nasopharyngeal microbial composition** *Sci Rep* 11
49. Bose T., et al. (2023) **A cross-sectional study on the nasopharyngeal microbiota of individuals with SARS-CoV-2 infection across three COVID-19 waves in India** *Front Microbiol* 14
50. Wang M., et al. (2023) **The relationship between gut microbiota and COVID-19 progression: new insights into immunopathogenesis and treatment** *Front Immunol* 14
51. Nardelli C., et al. (2022) **Nasal Microbiome in COVID-19: A Potential Role of *Corynebacterium* in Anosmia** *Curr Microbiol* 80
52. Bozkurt H.S., Bilen O. (2021) **Oral booster probiotic bifidobacteria in SARS-COV-2 patients** *Int J Immunopathol Pharmacol* 35

53. Wei N., et al. (2023) **Characterization of oral bacterial and fungal microbiome in recovered COVID-19 patients** *BMC Microbiol* **23**
54. Bassis C.M., et al. (2014) **The nasal cavity microbiota of healthy adults** *Microbiome* **2**
55. Perez-Losada M., et al. (2023) **Nasal Bacteriomes of Patients with Asthma and Allergic Rhinitis Show Unique Composition, Structure, Function and Interactions** *Microorganisms* **11**
56. Garcia-Vidal C., et al. (2021) **Incidence of co-infections and superinfections in hospitalized patients with COVID-19: a retrospective cohort study** *Clin Microbiol Infect* **27**:83–88
57. Seibert B., et al. (2021) **Mild and Severe SARS-CoV-2 Infection Induces Respiratory and Intestinal Microbiome Changes in the K18-hACE2 Transgenic Mouse Model** *Microbiol Spectr* **9**
58. Schippa S., et al. (2020) **Nasal Microbiota in RSV Bronchiolitis** *Microorganisms* **8**
59. Ma S., et al. (2021) **Metagenomic analysis reveals oropharyngeal microbiota alterations in patients with COVID-19** *Signal Transduct Target Ther* **6**
60. Merenstein C., et al. (2021) **Signatures of COVID-19 Severity and Immune Response in the Respiratory Tract Microbiome** *mBio* **12**
61. Mizutani T., et al. (2022) **Correlation Analysis between Gut Microbiota Alterations and the Cytokine Response in Patients with Coronavirus Disease during Hospitalization** *Microbiol Spectr* **10**
62. Mankowska-Wierzbicka D., et al. (2023) **Alterations in Gut Microbiota Composition in Patients with COVID-19: A Pilot Study of Whole Hypervariable 16S rRNA Gene Sequencing** *Biomedicines* **11**
63. Ubeda C., et al. (2013) **Intestinal microbiota containing *Barnesiella* species cures vancomycin-resistant *Enterococcus faecium* colonization** *Infect Immun* **81**:965–73
64. Giacobbe D.R., et al. (2021) **Enterococcal bloodstream infections in critically ill patients with COVID-19: a case series** *Ann Med* **53**:1779–1786
65. Murphy E.C., Frick I.M. (2013) **Gram-positive anaerobic cocci--commensals and opportunistic pathogens** *FEMS Microbiol Rev* **37**:520–53
66. Kageyama Y., et al. (2022) **Lactobacillus plantarum induces innate cytokine responses that potentially provide a protective benefit against COVID-19: A single-arm, double-blind, prospective trial combined with an in vitro cytokine response assay** *Exp Ther Med* **23**
67. Chen J., et al. (2022) **Comparison of the respiratory tract microbiome in hospitalized COVID-19 patients with different disease severity** *J Med Virol* **94**:5284–5293
68. Haran J.P., et al. (2021) **Inflammation-type dysbiosis of the oral microbiome associates with the duration of COVID-19 symptoms and long COVID** *JCI Insight* **6**
69. Lu S., et al. (2023) **Metatranscriptomic analysis revealed *Prevotella* as a potential biomarker of oropharyngeal microbiomes in SARS-CoV-2 infection** *Front Cell Infect Microbiol* **13**

70. Seong H., et al. (2023) **Clinical implications of gut microbiota and cytokine responses in coronavirus disease prognosis** *Front Immunol* **14**
71. Ning Y., et al. (2022) **Comparative analysis of the gut microbiota composition between knee osteoarthritis and Kashin-Beck disease in Northwest China** *Arthritis Res Ther* **24**
72. Mengyi Z., et al. (2023) **Plasma metagenomics reveals regional variations of emerging and re-emerging pathogens in Chinese blood donors with an emphasis on human parvovirus B19** *One Health* **17**
73. Li S., et al. (2021) **Microbiome Profiling Using Shotgun Metagenomic Sequencing Identified Unique Microorganisms in COVID-19 Patients With Altered Gut Microbiota** *Front Microbiol* **12**
74. de Nies L., et al. (2023) **Altered infective competence of the human gut microbiome in COVID-19** *Microbiome* **11**
75. Xu X., et al. (2022) **Integrated analysis of gut microbiome and host immune responses in COVID-19** *Front Med* **16**:263–275
76. Lymberopoulos E., et al. (2022) **COVID-19 severity is associated with population-level gut microbiome variations** *Front Cell Infect Microbiol* **12**
77. Thissen J.B., et al. (2022) **Evaluation of co-circulating pathogens and microbiome from COVID-19 infections** *PLoS One* **17**
78. Sulaiman I., et al. (2021) **Microbial signatures in the lower airways of mechanically ventilated COVID-19 patients associated with poor clinical outcome** *Nat Microbiol* **6**:1245–1258
79. Devi P., et al. (2023) **Longitudinal study across SARS-CoV-2 variants identifies transcriptionally active microbes (TAMs) associated with Delta severity** *iScience* **26**
80. DeVoe C., et al. (2022) **Increased rates of secondary bacterial infections, including Enterococcus bacteremia, in patients hospitalized with coronavirus disease 2019 (COVID-19)** *Infect Control Hosp Epidemiol* **43**:1416–1423
81. Khosravi A., et al. (2014) **Gut microbiota promote hematopoiesis to control bacterial infection** *Cell Host Microbe* **15**:374–81
82. Moreira-Rosario A., et al. (2021) **Gut Microbiota Diversity and C-Reactive Protein Are Predictors of Disease Severity in COVID-19 Patients** *Front Microbiol* **12**
83. Zhou Y., et al. (2021) **Gut Microbiota Dysbiosis Correlates with Abnormal Immune Response in Moderate COVID-19 Patients with Fever** *J Inflamm Res* **14**:2619–2631
84. Ley R.E., et al. (2006) **Microbial ecology: human gut microbes associated with obesity** *Nature* **444**:1022–3

Author information

Benita Martin-Castaño[#]

Centro de Salud Las Gabias, Distrito Granada-Metropolitano, Granada, Spain, Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain

[#]These authors contributed equally to this work

Patricia Diez-Echave[#]

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Department of Pharmacology, Center for Biomedical Research (CIBM), University of Granada, Granada, Spain

[#]These authors contributed equally to this work

Jorge García-García

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Department of Pharmacology, Center for Biomedical Research (CIBM), University of Granada, Granada, Spain, Servicio Microbiología, Hospital Universitario Clínico San Cecilio, Granada, Spain
ORCID iD: [0000-0003-0925-9441](https://orcid.org/0000-0003-0925-9441)

For correspondence: jgarcia.51@ugr.es

Laura Hidalgo-García

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Department of Pharmacology, Center for Biomedical Research (CIBM), University of Granada, Granada, Spain

Antonio Jesús Ruiz-Malagon

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Department of Pharmacology, Center for Biomedical Research (CIBM), University of Granada, Granada, Spain

José Alberto Molina-Tijeras

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Department of Pharmacology, Center for Biomedical Research (CIBM), University of Granada, Granada, Spain

Maria Jesús Rodríguez-Sojo

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Department of Pharmacology, Center for Biomedical Research (CIBM), University of Granada, Granada, Spain

Anaïs Redruello

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain

Margarita Martínez-Zaldívar

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Centro de Salud “Salvador Caballero”, Distrito Granada-Metropolitano, Granada, Spain

Emilio Mota

Centro de Salud “Salvador Caballero”, Distrito Granada-Metropolitano, Granada, Spain

Fernando Cobo

Servicio Microbiología, Hospital Universitario Virgen de las Nieves, Granada, Spain

Marta Alvarez-Estevez

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Servicio Microbiología, Hospital Universitario Clínico San Cecilio, Granada, Spain, CIBER de Enfermedades Infecciosas (CIBER-Infecc), Instituto de Salud Carlos III, Madrid, Spain

Federico García

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Servicio Microbiología, Hospital Universitario Clínico San Cecilio, Granada, Spain, CIBER de Enfermedades Infecciosas (CIBER-Infecc), Instituto de Salud Carlos III, Madrid, Spain

Concepción Morales-García

Respiratory Medicine Department, Hospital Universitario Virgen de las Nieves, Granada, Spain

Silvia Merlos

Respiratory Medicine Department, Hospital Universitario Virgen de las Nieves, Granada, Spain

Paula García-Flores

Respiratory Medicine Department, Hospital Universitario Virgen de las Nieves, Granada, Spain

Manuel Colmenero-Ruiz

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Servicio de Medicina Intensiva, Hospital Universitario Clínico San Cecilio, Granada, Spain

José Hernandez-Quero

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Servicio de Enfermedades Infecciosas, Hospital Universitario Clínico San Cecilio, Granada, Spain

María Nuñez

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Servicio Farmacia Hospitalaria, Hospital Universitario Clínico San Cecilio, Granada, Spain, CIBER de Epidemiología y Salud Pública (CIBER-ESP), Instituto de Salud Carlos III, Madrid, Spain

Maria Elena Rodríguez-Cabezas

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Department of Pharmacology, Center for Biomedical Research (CIBM), University of Granada, Granada, Spain

Ángel Carazo

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Servicio Microbiología, Hospital Universitario Clínico San Cecilio, Granada, Spain

Javier Martín

Department of Cell Biology and Immunology, Institute of Parasitology and Biomedicine López-Neyra, CSIC, Granada, Spain

Rocío Morón

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Servicio Farmacia Hospitalaria, Hospital Universitario Clínico San Cecilio, Granada, Spain

For correspondence: rmoronr@gmail.com

Alba Rodríguez-Nogales[‡]

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Department of Pharmacology, Center for Biomedical Research (CIBM), University of Granada, Granada, Spain
ORCID iD: [0000-0003-1927-0628](https://orcid.org/0000-0003-1927-0628)

[‡]These authors also contributed equally to this work

Julio Gálvez[‡]

Instituto de Investigación Biosanitaria de Granada (ibs.GRANADA), Granada, Spain, Department of Pharmacology, Center for Biomedical Research (CIBM), University of Granada, Granada, Spain, CIBER de Enfermedades Hepáticas y Digestivas (CIBER-EHD), Instituto de Salud Carlos III, Madrid, Spain

[‡]These authors also contributed equally to this work

Editors

Reviewing Editor

Musa Ali

Hawassa University, Hawassa, Ethiopia

Senior Editor

K VijayRaghavan

National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bangalore, India

Reviewer #1 (Public review):

Summary:

The research study under review investigated the relationship between gut and identified potential biomarkers derived from the nasopharyngeal and gut microbiota-based that could aid predicting COVID-19 severity. The study reported significant changes in the richness and Shannon diversity index in nasopharyngeal microbiome associated with severe symptoms.

Strengths:

The study successfully identified differences in the microbiome diversity that could indicate or predict disease severity. Furthermore, the authors demonstrated a link between individual

nasopharyngeal organisms and the severity of SARS-CoV-2 infection. The density of the nasopharyngeal organism was shown to be a potential predictors of severity of COVID-19.

<https://doi.org/10.7554/eLife.95292.2.sa2>

Reviewer #3 (Public review):

Summary:

How the microbial composition of the human body is influenced by and influences disease progression is an important topic. For people with COVID-19, symptomatic progression and deterioration can be difficult to predict. This manuscript attempts to associate the nasal and fecal microbiomes of COVID-19 patients with the severity of disease symptoms, with the goal of identifying microbial markers that can predict disease outcomes.

Strengths:

Analysis of microbiomes from two distinct anatomical locations and across three distinct patient groups is a substantial undertaking. How these microbiomes influence and are influenced by COVID-19 disease progression is an important question. In particular, the putative biomarker identified here could be of clinical value with additional research.

Weaknesses:

The primary weaknesses of this analysis is the relatively low sample size for analyzing disease subsets and moderate correlation values observed for putative biomarkers. Regardless, this data can be used to inform future studies aiming to understand the contribution of multifactorial dysbiosis to COVID-19 disease progression.

<https://doi.org/10.7554/eLife.95292.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Public reviews:

Reviewer 1

We would like to express our gratitude to Reviewer 1 for providing a thorough summary of our work and highlighting its strengths. With regards to the weaknesses, we are committed to improve the manuscript by performing the necessary changes. First, we will specify the exact p-value in all cases.

Regarding the discussion section, we acknowledge the feedback regarding its potential confusion. In line with the reviewer's suggestion, we will reduce the literature review and highlight our findings.

Finally, for the preprint we did not include cofounders such as HIV infection and ethnicity as our study population did not exhibit viral infections and comprised only Hispanic individuals. We will make a more thorough description of the population of study and address these characteristics explicitly in both the methods section and the initial part of the results.

Reviewer 2

We appreciate and thank reviewer 2 for the commentaries. Although it is true that several papers have described the role of microbiome in COVID-19 severity, we firmly believe that our current work stands out. There is not much information related to this association in Mediterranean countries, especially in the south of Spain. In addition, most of the studies only describe microbiota composition in stool or nasopharyngeal samples separately, without investigating any potential relationships between them as we do.

(1) We agree with the reviewer idea of a limited sample size. We faced the challenge of collecting the samples during the peak of COVID-19 pandemic. Thus, doctors and nurses were overwhelmed and not always available for carrying out patient recruitment following the inclusion criteria. Despite these constraints, we ensured that all included samples met our specified inclusion criteria and were from subjects with confirmed symptomatology.

In addition, our main goal was to identify whether severity of the disease could be assessed through microbiota composition. Therefore we did not include a healthy group. Despite not having a large N, our results should be reproducible as they are supported by statistical analysis.

(2) We thank reviewer commentary, and since our original sentence may have lacked clarity, we intend to modify it to ensure it conveys the intended meaning more effectively.

Nonetheless, we remain confident in the significance of our findings. Not only have we found correlation between microbiota and COVID severity, but we have also described how specific bacteria from each condition is associated with key biochemical parameters of clinical COVID infection.

(3) We appreciate the feedback provided by the reviewer. In this case, we have performed 16S analysis due to its cost-effectiveness compared to metagenomic approaches. Furthermore, 16S analysis has undergone refinements that ensure comprehensive coverage and depth, along with standardized analysis protocols. Unlike 16S, metagenomic approaches lack software tools such as QIIME that facilitate standardization of analysis and, thus, reduce reproducibility of results.

(4) We sincerely appreciate this insightful suggestion. simply listing associations between both microbiomes and COVID-19 severity could not be enough, we intend to discuss how microbiota composition may be linked to the mechanisms underlying COVID-19 pathogenesis in our discussion.

(5) We are grateful for the constructive criticism and intend to rewrite our abstract to enhance clarity. Additionally, we will thoroughly review all figures and their descriptions to ensure accuracy and comprehensibility.

Reviewer 3

We acknowledge the annotations made by reviewer 3 and are committed to addressing all identified weaknesses to enhance the quality of our work. Our idea is to modify the methods section and figures to make them easier to understand.

Specifically, in the case of Figure 1, we recognize an error in the description of the Bray-Curtis test. We appreciate the commentary and we will make the necessary changes. Moreover, there is another observation related to Figure 1 description. We are going to modify it in order to gain accuracy.

For figure 2 we are planning to add a supplementary table showing the abundance of detected genus. Nevermind, we will also update the manuscript text to provide clarification

on how we obtained this result.

Regarding the clarification about "1% abundance," we want to emphasize that we are referring to relative abundance, where 1 represents 100%. To avoid confusion, we will explicitly state this in both the methods section and figure descriptions. Besides, it is true that the statistical test employed for the analysis is not mentioned in the figure description and we recognize that the image may be difficult to interpret. Therefore, we will modify the text and a supplementary table displaying the abundance and p values is going to be added.

Furthermore, we agree with the reviewer's suggestion to investigate whether the bacteria identified as potential biomarkers for each condition are specific to their respective severity index or if there is a threshold. Thus, we will reanalyze the data and include a supplementary table with the abundance of each biomarker for each condition. We will also place greater emphasis on these results in our discussion.

Finally, in response to the reviewer's suggestion, we are going to go through the nasopharyngeal-fecal axis part in the discussion. It is well described that COVID-19 induces a dysbiosis in both microbiomes. Consequently, we understand that the ratio we have described could be an interesting tool for assessing COVID severity development as it considers alterations in both environments. However, we acknowledge that there may be room for improvement in clarifying the significance of this intriguing finding and its implications.

<https://doi.org/10.7554/eLife.95292.2.sa0>