

Development of a new genotype–phenotype linked antibody screening system

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint version 1

May 3, 2024 (this version)

Posted to preprint server

March 8, 2024

Sent for peer review

February 20, 2024

Takashi Watanabe , **Hikaru Hata**, **Yoshiki Mochizuki**, **Fumie Yokoyama**, **Tomoko Hasegawa**, **Naveen Kumar**, **Tomohiro Kurosaki**, **Osamu Ohara**, **Hidehiro Fukuyama**

Laboratory for Integrative Genomics, RIKEN Center for Integrative Medical Sciences, Yokohama, 230-0045, Japan • Laboratory for Lymphocyte Differentiation, RIKEN Center for Integrative Medical Sciences, Yokohama, 230-0045, Japan • Laboratory for Integrated Bioinformatics, RIKEN Center for Integrative Medical Sciences, Yokohama, 230-0045, Japan • Laboratory of Lymphocyte Differentiation, WPI Immunology Frontier Research Center, Osaka University, Osaka, 565-0871, Japan • Department of Applied Genomics, Kazusa DNA Research Institute, Chiba 292-0818, Japan • Near-Infrared Photo-Immunotherapy Research Institute, Kansai Medical University, Hirakata, Osaka, 573-1010, Japan • INSERM EST, Strasbourg Cedex 2, 67200, France

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Antibodies are powerful tools for the therapy and diagnosis of various diseases. In addition to conventional hybridoma-based screening, recombinant antibody-based screening has become a common choice; however, its application is hampered by two factors: 1) screening starts only after Ig gene cloning and recombinant antibody production, and 2) the antibody is composed of paired chains, heavy and light, commonly expressed from two independent expression vectors. Here, we introduce a method for the rapid screening of recombinant monoclonal antibodies by establishing a Golden Gate-based dual-expression vector and *in vivo* expression of membrane-bound antibodies. Using this system, we demonstrated the efficient isolation of influenza cross-reactive antibodies with high affinity from mouse germinal center B cells over 4 days. This system is particularly useful for isolating therapeutic or diagnostic antibodies (e.g., during foreseen pandemics).

Impact Statement

A Golden Gate-based dual-expression vector enables rapid screening, facilitating efficient isolation of high-affinity cross-reactive antibodies for therapeutic or diagnostic use, offering a crucial advance for pandemic preparedness.

eLife assessment

The studies described here are **useful**; they are broadly applicable to all antibody discovery subfields but do not add significant improvement to techniques already published. The findings are **incomplete** with respect to the methodology since details that are crucial in order to repeat the experiment are lacking (such as a timestamp) and they do not take into account multiple recent papers that have tested similar strategies. These studies will be of interest to a specialized audience working on making antibodies to infectious agents.

Introduction

Technical improvements in the rapid isolation of monoclonal antibodies (mAbs) are critical for their diagnostic and therapeutic development. This need became apparent to the general public during the COVID-19 pandemic. Antibodies isolated from convalescent COVID-19 patients are valuable resources for the development of mAb therapeutics and rapid diagnostic tools. Human antibodies can be administered to other patients without the need to humanize in an emergency and with a slight modification of Fc receptor binding (1 [↗](#)) to increase the half-life or reduce the risk of antibody-dependent enhancement. Although production costs remain high, the efficiency of generating recombinant antibodies has dramatically improved with the use of mammalian cell lines, and the application of therapeutic antibodies is likely to expand in the near future (2 [↗](#)).

The gold standard method for isolating mAbs is the hybridoma technology developed by Kohler and Milstein in 1975 (3 [↗](#)), in which antibody-producing B cells are fused with immortal B cells, called myelomas, to produce long-lasting antibody-producing B cells. In addition to hybridoma technology, new methods have been developed to increase mAb screening efficiency. These include direct immortalization of B cells by gene reprogramming using the Epstein–Barr virus (4 [↗](#)) or retrovirus-mediated gene transfer (5 [↗](#)), cloning of variable region-encoding genes by single-cell PCR (6 [↗](#), 7 [↗](#)), single-cell culture screening (8 [↗](#)), and *in vitro* screening of recombinant antibody libraries (9 [↗](#)–13 [↗](#)). Some of these methods successfully yield high-affinity antibodies in various formats, including single variable domain on a heavy chain or single-chain fragment variable antibodies. Although hybridoma technology is highly reliable for the isolation of valuable antibodies from animal models, such as mice, rats, and hamsters, it is time-consuming and requires technical skill. The development of a new fusion partner cell line, SPYMEG, has enabled the production of human hybridomas and opened a new direction for the isolation of human mAbs (14 [↗](#)). Although many of these new technologies have increased the throughput of mAb isolation, the procedure still requires significant resources and time.

To dramatically enhance the efficiency of mAb isolation, we used next-generation sequencing (NGS) technology, which has changed the method for sequencing immunoglobulin (Ig) variable-region genes. For instance, tens of thousands of Ig genes specific to certain antigens can be identified by combining droplet-based single-cell isolation with DNA barcode antigen technology, followed by NGS (15 [↗](#)). Although Ig genes can be sequenced at a high throughput, there is no method for screening antibodies in a high-throughput format that is compatible with NGS technology.

In this study, we developed a new functional screening method that is compatible with NGS to rapidly identify antigen-specific clones. We first generated an Ig dual-expression vector using Golden Gate Cloning (16 [↗](#)), which enabled the linkage of heavy-chain variable and light-chain variable DNA fragments obtained from a single-sorted B cell, followed by the expression of

membrane-bound Ig. This single-step procedure enabled the enrichment of antigen-specific, high-affinity Igs by flow cytometry, which was significantly faster than conventional cloning-based methods that require sequential steps. To demonstrate the efficiency of our new method, we screened for potent broadly reactive antibodies (17–19) against the influenza virus in an experimental mouse model using our technology. Broad reactivity allows for influenza vaccines that work across seasonal variations in influenza strains, which can only be accomplished by screening a large number of candidate antibodies. We first raised cross-reactive B cells against various hemagglutinin (HA) antigens from the influenza virus by sequential immunization with heterotypic HA antigens from group 1 influenza. Using this technology, we obtained several mAbs that bind to other group 1 HA antigens and even to group 2 HA antigens of the influenza virus. Our technology can also be applied to human antibody screening and represents a new line of mAb screening that accelerates the isolation of therapeutic and diagnostic mAbs.

Results

Rapid membrane-bound dual Ig expression screening system

Cloning-based Ig screening involves several steps: cloning two Ig gene fragments encoding heavy and light chains independently from a single cell, co-expressing these Ig chains, and purifying individual recombinant antibodies. These steps are labor intensive. We established a new antibody screening system for the isolation of valuable mAbs (details in the method). Eventually, the antigen-binding Ig transformants were collected by sorting in a bulk fashion, and the unique CDR3 region and clones of interest were identified using the Ig-seq database (Fig. 1A).

We generated a dual-expression vector to express both Ig heavy and kappa/lambda in a single-expression vector (Fig. 1B). These plasmids were individually transfected transiently to the floating human FreeStyle 293 cell line. Igs were expressed on the cell surface in 2 days; their expression could be confirmed and normalized by Venus expression, since the Ig heavy chain is fused with Venus at the cytoplasmic domain tail (Fig. 1C). The advantage of this system is the rapid enrichment of clones of interest through bulk screening. In the current single-cell-based cloning/screening, plasmid DNA extracted from the collected antigen-binding bulk transformants was sequenced for the heavy chain CDR3 region. The CDR3 region is unique and can be used for the identification of a clone (20,21). This membrane-bound Ig expression system links antigen-binding features and genetic information of the Ig repertoire.

Building the Ig-seq database and analysis

As a model experiment, we obtained broadly reactive antibodies against influenza viruses using multiple HA probes. We prepared two HA proteins as probes: A/Puerto Rico/8/1934 (H1N1), designated as PR8, and A/Okuda/1957 (H2N2), designated as H2. A total of 380 IgG1⁺ germinal center B cells of either PR8⁺ (204 cells), H2⁺ (99 cells), or PR8⁺H2⁺ (77 cells) were collected in a single-cell fashion. We obtained sequences of 284 independent clones. The success rate of cloning the paired Ig fragments was 75.9%. An overview of heavy chain V-D-J and light chain V-J usage and repertoire clonality is shown in Supplementary Fig. 1A and B. Mutation rates and CDR3 lengths of the heavy chains of the three cell populations (PR8⁺, H2⁺, and PR8⁺H2⁺) were comparable (Supplementary Fig. 1C and D). These results indicate that broadly reactive antibodies do not require unique genetic traces to obtain breadth in our experimental setup.

Isolation of H2- and H1-reactive B cells

On day 2 after transfection, 287 transformants (Ig-expressing transformant libraries) were stained with HA probes (Fig. 2A). As shown in Fig. 2B, we selected three prominent populations, H1⁺H2⁺ (cross), H2⁺ (H2), and H1⁺ (H1), and collected them in bulk. In contrast to bacterial transfection, mammalian cell lines, such as FreeStyle 293 cells, can contain multiple plasmids in a




Figure 1.

System to link the antigen-binding feature with the Ig repertoire genetic information. **A.** Single cell-sorted B cells were subjected to two processes: Ig-seq database building and Ig-expressing transformant library preparation. Antigen-binding Ig transformants were collected by sorting in bulk and the sequences of the unique CDR3 regions and clones of interest were identified by referring to the Ig-seq database. **B.** To express both Ig heavy and kappa/lambda chains in a single-expression vector, we generated a dual-expression vector. Four gene fragments were assembled by the Golden Gate method using the BsaI restriction enzyme. They included: 1) the destination vector containing the Ig kappa constant gene and the ccdB Chloramphenicol cassette for negative selection, 2) fragments containing the Ig heavy constant gene fused to *Venus* derived from the donor vector, 3) the Ig heavy variable gene fragment, and 4) the Ig light variable gene fragment. **C.** Individually purified plasmids were transiently transfected to the floating human FreeStyle 293 cell line. Igs were expressed on the cell surface in 2 days, and expression levels were confirmed and normalized by *Venus* expression, since the Ig heavy chain was fused to *Venus* at the cytoplasmic domain tail.

cell and ectopically express multiple proteins. Therefore, we decided to mix the transformants instead of plasmids. We sorted 2,981 cells from these three populations and performed sequencing. Bulk Ig-seq data are referred to as Ig-seq data, as described above. All the transformants were examined individually. A total of 90 clones from the three populations were verified for their sequences and HA binding by flow cytometry analysis, as shown in [Fig. 2](#). Among these, 110 “H1⁺,” 67 “H2⁺,” and 13 “cross” clones were successfully isolated. Then, 138 transformants were verified for their binding to HAs by flow cytometry. Of these, 81 “H1⁺,” 48 “H2⁺,” and nine “cross” clones were successfully isolated. Finally, 8 of the 13 “cross” clones obtained in bulk were included in this group; the binding of one clone (D11p4) was so weak that it was not included in the gates sorted in bulk. Thus, we established a high-throughput method for sorting “cross” clones.

Characterization of “cross” clones for broad reactivity against the influenza virus

By generating a secretory form of mAbs for these nine cross clones, including C10p2, we characterized their affinity and broadness using six different HA antigens from strains A/Okuda/1957 (H2N2) named for H2, A/Puerto Rico/8/1934 (H1N1) for PR8, A/California/2009 (X-179A) [H1N1] Pdm09 for Cal, A/Texas/50/2012 (X-223) [H3N2] for H3, A/Egypt/N03072/2010(H5N1) for H5, and A/Brisbane/59/2007 [H1N1] ([19](#)) for Stem. We determined the affinities (K_d) of the antibodies for the six HA antigens using surface plasmon resonance analysis ([Table 1](#) and [Supplementary Fig. 2](#)), which ranged from 500 to 100 nM. With the highest affinity among them, the C10p2 antibody bound to Cal at $K_d \approx 5.66 \times 10^{10}$ (M). Surprisingly, the antibodies showed a broader spectrum of HA binding among the influenza strains. Seven of the nine mAbs bound to HA from the highly pathogenic avian influenza strain H5N1. Furthermore, the six antibodies that bound to H3 were categorized into group 2 ([22](#)). In addition, we observed that two of the broadly reactive antibodies recognized the stem region of the HA of A/Brisbane/59/2007. A possible scenario for other stem cell-negative antibodies is that they may either bind to the stem region of other strains or to the head region of HA. We tested whether some of these antibodies shared features with classic broadly neutralizing antibodies. C179, the first broadly neutralizing mouse antibody isolated in 1993 ([17](#)), reacts with the stem region of HAs from group 1 influenza virus strains ([22](#)). C179 binds to HAs from group 1 influenza viruses, including H1N1, H2N2, and H5N1. A competition assay for the binding of H2 and PR8 revealed that A6p4 competes with C179 ([Fig. 3](#)). In addition, our two-dimensional phylogeny map indicated that A6p4 was unique to all clones ([Supplementary Fig. 3](#)). Note that we used NSP2 as a negative control antibody, which binds specifically to the HA from A/California/2009 (X-179A) [H1N1] Pdm09 ([23](#)). These results demonstrate that we were able to isolate a broadly reactive mAb with an affinity comparable to that of a classic broadly reactive mAb (C179) using our technology.

Discussion

The discovery of therapeutic antibodies is accumulating, and their demand is increasing ([24](#)). Efficiently isolating one biologically significant antibody from the 10^{13} antibody repertoire is challenging. The hybridoma method has long been used for this purpose and remains reliable. However, the fusion rate of B and myeloma cells is $< 0.001\%$ in this method; therefore, the number of hybridomas obtained from one individual B cell is limited to several hundred clones, limiting its use in the production of mAbs. As an alternative choice, cloning-based antibody screening has gained popularity, especially for human antibody discovery, because of the lack of a human myeloma cell line similar to that used for murine cell fusion partners. The SPYMEG cell line can generate antigen-specific antibody-producing cells by fusing the murine SP2/0 myeloma cell line and human megakaryoblastic leukemia cell line MEG-01 with human PBMCs ([14](#), [25](#)). In addition to gene expression profiling, NGS facilitates antibody repertoire analysis. However,




Figure 2.

Isolation of H2 and H1-reactive B cells. **A.** Mixture of Ig-expressing transformant libraries stained with HA probes. **B.** Three prominent populations, H1⁺H2⁺ (cross), H2⁺ (H2), and H1⁺ (H1), were selected and collected in a bulk fashion. Then, the three prominent populations were sorted and sequenced. **C.** The bulk Ig-seq data were referred to as the single-cell Ig-seq data. Wells containing crossed antibodies were determined by comparing the sequence of the heavy chains of the populations included in H1⁺H2⁺ (cross) with the Ig-seq data of single cells. **D.** Cross antibodies obtained by bulk sorting were examined by single-cell presentation. **E.** A summary of the results to check whether a candidate mAb was cross-reactive.




Figure 3.

Competition assay using various mAbs for binding to H2 and PR8 was performed, including two previously reported antibodies, C179, a broadly-reactive mAb binding to the Stem region of HA, and NSP2 (Narita strain-specific mAb), a strain-specific mAb that binds to the head region of HA.

Surface plasmon resonance (Biacore)

						KD (M)
Group	1	1	1	2	1	1
	H2	PR8	Cal	H3	H5	Stem
A6p4	4.51E-08	1.41E-06	1.14E-05		2.91E-05	1.59E-08
B10p2	6.75E-09	2.3E-09	3.29E-09			
C10p2	1.66E-09	1.29E-09	5.66E-10			4.26E-08
A2p1	2.49E-06	1.42E-07	1.14E-05	1.25E-04	1.91E-05	
D11p4	7.93E-07	1.72E-06	1.04E-06	1.36E-05	1.05E-11*	
E11p2	4.4E-08	1.86E-08	1.67E-06	2.92E-07	2.37E-07	
G6p2	3.51E-08	3.53E-08	1.44E-08	1.78E-05	3.05E-08	
D4p4	1.1E-05	1.78E-05	6.4E-05	4.38E-05	2.47E-05	
G12p4	1.45E-07	1.52E-07	5.7E-07	9.44E-07	1.58E-07	

High
Middle
Low

Table 1.

Affinities (K_d) of the antibodies were determined to 6 HA antigens from strains A/Okuda/1957(H2N2) named for H2, A/Puerto Rico/8/1934(H1N1) for PR8, A/California/2009 (X-179A) [H1N1] Pdm09 for Cal, A/Texas/50/2012 (X-223) [H3N2] for H3, A/Egypt/N03072/2010(H5N1) for H5, and A/Brisbane/59/2007 (H1N1) for Stem using surface plasmon resonance analysis. H2, PR8, Cal, and H5 belong to group 1, and H3 belongs to group 2.

* The affinity of D11p4 for H5 was low and inaccurate (Supplementary Figure 2).

despite breakthroughs in sequencing technology, the process of linking antibody function and its genes is still long and labor intensive. To overcome this issue, we developed a new antibody screening system to directly link antigen–antibody binding (as a function) with a gene encoding the antibody by expressing its membrane-bound form. Among conventional cloning-based mAb isolation methods, this system demonstrates several advantages. 1) Membrane Ig expression can link the antigen-binding feature of membrane-expressed Ig, which can be linked to Ig DNA sequence information using our plasmid construct. Similar to the panning procedure used in phage display (26), antigen-binding cells carry the plasmid encoding the Ig genes through surface Ig so that the selection process in bulk format enriches the relevant plasmids for use in further experiments. 2) The dual Ig expression vector links heavy- and light-chain genes, which reduces the plasmid preparation time and stock by half. 3) Highly reliable Golden Gate Cloning technology using type IIs restriction enzymes can readily generate plasmid clones. This reduced the time required to generate an Ig plasmid library. 4) The population profile, defined by the fluorescence intensity during flow cytometry, directly reflected the affinity of a clone (Supplementary Fig. 4). Collectively, our technology has streamlined the isolation of mAbs for therapy and diagnosis.

Conclusion

Our findings indicate that the developed antibody presentation system facilitates antibody functional analysis and is well suited for the discovery of antibodies important for infectious diseases when combined with conventional NGS-based antibody repertoire analysis. Compared with droplet-based experimental systems, well-based systems are limited in the number of cells they can process. Furthermore, experiments involving infectious bacteria and viruses have imposed limitations on human experimentation. To solve these problems, the automation of experiments will become important in the future. By combining our screening system with robotic automation of experiments, it will be possible to obtain useful mAbs for various diseases quickly and in large quantities, which has broad implications for the development of vaccines against various diseases.

Materials and Methods

Protein purification of influenza HAs (H1 and H2) as antigens

A mixture of 25.7 µg of HA, comprising H1 strain A/Puerto Rico/8/1934 (PR8), H1 strain A/California/7/2009 (X-179A), H2 strain A/Okuda/1957 (H2), H3 strain A/Texas/50/2012 (X-223), or H5 strain A/Egypt/N03072/2010], 1.3 µg of NA (Strain-matched), and 3 µg of BirA-expressing plasmids were transfected into Expi293 cells using the Expifectamine 293 transfection kit (Thermo Fisher Scientific, Waltham, MA, USA) according to the manufacturer's instructions. The cells were then cultured in Expi293 Expression Medium (Thermo Fisher Scientific) supplemented with D-biotin at a final concentration of 100 µM for *in vivo* BirA biotinylation in a humidified incubator containing 8% CO₂ at 37°C and 125 rpm. If biotinylation was not necessary, transfection was performed without BirA, and cell culture was carried out without D-biotin. Culture supernatants were harvested at 3 days post-transfection and filtered through a 0.45 µm filter (Merck Millipore, Burlington, MA, USA). All proteins were purified with a Talon metal affinity resin (Takara Bio USA, Mountain View, CA, USA) and dialyzed against pyrogen-free PBS. The oligomeric state and purity were determined at 25°C by size-exclusion chromatography on a Superdex 200 10/300GL column (GE Healthcare Technologies, Chicago, IL).

HA (H1 and H2) protein immunization

Two individual BALB/c mice were intraperitoneally immunized sequentially, 2 weeks apart, with 15 µg of H1 (PR8) HA followed by 15 µg of H2 HA protein as an antigen and supplemented with AddaVax adjuvant (InvivoGen, San Diego, CA, USA). BALB/c mice were purchased from CLEA Japan (CLEA Japan, Inc., Tokyo, Japan). All animal experiments were performed using protocols approved by the Institutional Animal Care and Use Committee (IACUC) of the RIKEN Yokohama Branch.

Fluorescence-activated cell sorting

CD43-negative B cells were collected using AutoMACS (Miltenyi Biotec, Inc., Bergisch Gladbach, North Rhine-Westphalia, Germany) and stained with IgG1 (Bv510), non-biotinylated His-tagged purified recombinant H1 (PR8) protein, and biotinylated purified recombinant H2 protein on ice for 30 min. After washing three times, the cell suspension was incubated with Alexa Fluor 488-labeled anti-His antibodies on ice for 30 min. After washing three times, the cell suspension was incubated with Brilliant Violet 421-conjugated streptavidin on ice for 3 min. Next, the B cells were subjected to single-cell sorting using a BD FACSAria III (BD Biosciences, Franklin Lakes, NJ, USA) and collected into 96-well plates pre-loaded with lysis buffer. The single cells in the lysis buffer were immediately snap-frozen on powdered dry ice and stored at 80°C.

Amplification of a paired B cell repertoire amplicon from a single cell

Single cells were dropped into pre-loaded plates containing 4 µL/well of lysis buffer (2 U/µL RNase inhibitor, 0.5× PBS, 10 mM DTT) and 1 µL of 10 µM Oligo(dT)18 Primer. Cell lysates were incubated at 70°C for 90 s, followed by 35°C for 15 s, and then placed on ice for 2 min. For first-strand cDNA synthesis, 5 µL of RT mix (2× PCR buffer, 5 mM DTT, 2 mM dNTP Mix, 10 U RNase inhibitor, 100 U SuperScript III Reverse Transcriptase) was added to the cell lysate and incubated consecutively at 35°C for 5 min, 45°C for 20 min, 70°C for 10 min, and 4°C for chilling. For primer digestion, 5 µL of the primer digestion mix (1× ExoI Buffer, 7.5 U Exonuclease I) was prepared and incubated at 37°C for 30 min, 80°C for 20 min, and 4°C for chilling. For the poly-A tailing reaction, 5 µL of the poly-A tailing reaction mix (4× PCR buffer, 5 mM dATP, 0.1 U RNase H, 60 U Terminal transferase) was prepared and incubated at 37°C for 50 s, 65°C for 10 min, and 4°C for chilling. For the 2nd strand cDNA synthesis, 25 µL of the amplification mix (1× KAPA HiFi HotStart ReadyMix, 400 nM SMART_dT primer, 10 µL of poly-A tailing product, 1.5 µL of nuclease-free H₂O) was prepared and incubated at 95°C for 3 min, 40°C for 1 min, 65°C for 10 min, and 72°C for 5 min and chilled. For the cDNA amplification using an Ig-specific outer primer set, 20 µL of the outer amplification mix (1× KOD FX Neo buffer, 400 µM dNTPs, 1× UPM, 300 nM outer primer, 3.5 µL of the SMART product, 0.5 U KOD FX Neo polymerase) was prepared and incubated at 94°C for 2 min, followed by 25 cycles of 98°C for 10 s, 60°C for 30 s, and 68°C for 45 s and then chilled. For cDNA amplification with an Ig-specific inner primer set, 20 µL of the inner amplification mix (1× KOD FX Neo buffer, 400 µM dNTPs, 500 nM 5' matrix primer mix, 500 nM 3' matrix primer mix, 3.5 µL of the outer cDNA amplified product, 0.5 U KOD FX Neo polymerase) was prepared and incubated at 94°C for 2 min, followed by 25 cycles of 98°C for 10 s, 60°C for 30 s, and 68°C for 45 s and then chilled. Finally, 3 µL of the amplified inner cDNA product was used for further plate indexing by amplification. Then, 20 µL of the plate-indexing amplification mix (1× KOD FX Neo buffer, 400 µM dNTPs; 500 nM i5-Fwd or i7-Fwd, 500 nM i7-Rev or i5-Rev; 2 µL of the inner cDNA amplified product; 0.5 U KOD FX Neo polymerase) was prepared and incubated at 94°C for 2 min, followed by 25 cycles at 98°C for 10 s, 60°C for 30 s, and 68°C for 45 s and then chilled. All primers used are listed in **Supplementary Table 1** [↗](#) and **Supplementary Table 2** [↗](#).

Paired-end Illumina sequencing strategy and analysis

The libraries were sequenced on a MiSeq instrument (Illumina, San Diego, CA, USA) following the manufacturer's protocol. As the sequencing length was limited to 2×300 bp, we used the following strategy. Amplicons in both orientations were generated using an Illumina adapter sequence. Then, asymmetric 400+100 nt paired-end sequencing was performed to obtain high-quality 400 + 400 nt paired-end sequences of the original cDNA molecule. This approach provided a sufficient sequencing length to capture extra-long Ig variants. We followed Turchaninova's method (27 [↗](#)). Only reads that met the following four conditions were used as the resulting Ig sequences: 1) reads with a length of 260 nt or more; 2) cumulative clustered reads representing the top 5% of the entire reads; 3) 5'- and 3'-reads overlapping a length of 20 bases or more; and 4) a total length of 460 ± 50 bp for heavy chain or 390 ± 40 bp for light chain. All resulting sequence data were analyzed using IMGT/High V-QUEST (http://www.imgt.org/IMGT_vquest/vquest [↗](#)) for annotation (28 [↗](#)). Using the sequences and annotations, Ig database construction and visualization of the Ig repertoire were performed using the in-house BONSCI software.

Construction of a mouse IgG/IgK expression plasmid

The BsaI restriction sites were inserted into the paired B cell repertoire amplicon from a single cell using PCR. The destination vector contained the IL6 signal peptide, EF1a promoter, ccdB gene, and mouse IgK Fc region. The donor vector contained the mouse IgG1 Fc region, the *Venus* gene, and the EF1a promoter. The paired B cell repertoire amplicons, destination vector, and donor vector containing BsaI restriction sites were assembled using our assembly method. For this step, 10 μ L of the assembly mix (1 $\times$ T4 DNA ligase buffer, 1 $\times$ BSA, 1 U BsaI restriction enzyme, 40 U T4 DNA ligase, 100 ng heavy chain amplicon, 100 ng light chain amplicon, 100 ng destination vector, and 100 ng donor vector) was prepared and incubated for 25 cycles at 37°C for 3 min, 16°C for 4 min, 50°C for 5 min, and 80°C for 5 min, and then chilled. The antibody sequence that entered the construct was fused to the *Venus* sequence and expressed in membrane form.

Antibody display on FreeStyle 293 cultured cells

A mixture of 1 μ g of antibody-expressing plasmid was transfected into FreeStyle 293 cells using the 293fectin Transfection Reagent (Thermo Fisher Scientific) according to the manufacturer's instructions and cultured in FreeStyle 293 Expression Medium (Thermo Fisher Scientific) in a humidified incubator with 8% CO₂ at 37°C and 125 rpm. Antibodies were displayed on the surface of cultured cells, which enabled the determination of antigen specificity. Furthermore, the antibody-display cells were tested for binding activity with Alexa647-labeled H1 and Alexa568-labeled H2 using a BD FACSAria III (BD Biosciences).

Antibody production

To obtain a large amount of secretory antibodies, PCR fragments from the antibody-expressing plasmid were cloned into the pcDNA3.4-mIgG1 or pcDNA3.4-kappa vectors. A mixture of 15 μ g of pcDNA3.4-V-gene-mIgG1 vector and 15 μ g of pcDNA3.4-V-gene-kappa vector were transfected into Expi293 cells using the ExpiFectamine 293 Transfection Kit (Thermo Fisher Scientific), according to the manufacturer's instructions, and cultured in Expi293 Expression Medium (Thermo Fisher Scientific) in a humidified 8% CO₂ incubator at 37°C and 125 rpm. Then, 5 days post-transfection, the culture supernatants were harvested, and all proteins were purified with PureSpeed IMAC resin (Mettler Toledo, Columbus, OH, USA), according to the manufacturer's instructions.

Surface plasmon resonance

Kinetic analyses were performed at 25°C using a BIAcore 3000 machine (GE Healthcare Technologies). A2p1, B10p2, C10p2, E11p2, G6p2, A6p4, D4p4, D11p4, or G12p4 antibodies were immobilized on a CM5 sensor chip (GE Healthcare Technologies) with an amine-coupling kit

according to the manufacturer's instructions. HA probes were serially diluted at five different concentrations and injected at a flow rate of 30 μ L/min for 3 min with a dissociation time of 7 min in HBS-EP buffer (10 mM HEPES, pH 7.4, 150 mM NaCl, 3.4 mM EDTA, and 0.005% Surfactant P20). The chip was regenerated using 10 mM glycine (pH 2.5) as an HA probe.

Data access

All raw and processed sequencing data generated in this study have been submitted to the NCBI Gene Expression Omnibus (GEO; <https://www.ncbi.nlm.nih.gov/geo/>) under accession number GSE140720.

Acknowledgements

We thank Mr. Atsuo Kobayashi, Ms. Chieko Okamura, Dr. Seok-Won Kim, Mr. Maxime Hebrard, Dr. Todd D Taylor, and the RIKEN IMS Genome Platform for their technical support, and Dr. Christopher Nicholas for critical reading and English editing of the manuscript. We would like to thank Editage (www.editage.jp) for English language editing.

Funding

This study was supported in part by research grants from the RIKEN Research Program 'Single cell Project' (O.O., T.W., T.K., and H.F.) and Grants-in-Aid for Scientific Research (S) [26221306 to T.K. and H.F.]. The funding sources were not involved in the study design, data collection and interpretation, or paper submission decisions.

Conflict of interest

None declared.

Author contributions

T.W. and H.F. designed the study, the main conceptual ideas, and the proof outline. H.H., Y.M., F.Y., T.H., and N.K. collected the data. O.O. and T.K. supervised the project. T.W. wrote the manuscript with support from H.F. All authors discussed the results and commented on the manuscript.

Supplementary Tables

Supplementary Figures

Primer	Application	Sequence
First-strand cDNA synthesis		
Oligo(dT)18	Primer for cDNA synthesis	TTTTTTTTTTTTTTTTTT
Second-strand cDNA synthesis		
SMART_dT	TdT oligo	AAGCAGTGGTATCAACGCAGAGTAC(T) ₃₀ VN
Mouse IGH/IGK primer		
TdT_L_BsaI_heavy	mouse IgG heavy-chain cDNA	CTAGGGTCTCAAGCAGTGGTATCAACGCAGAGTAC
msVHE		GAGGTGCAGCTGCAGGAGTCTGG
W-gamma-outer		AGAAGGTGTGCACACCGCTGGAC
mC_G new2		ATCTCCACACACAGGGGCCAGTGGATAGAC
BsaI_IL6sp_L		CTAGGGTCTCAAGCAGATGAACTCCTTCTCCACAAGCG
mC_G_new2_BsaI		TCCTAGGTCTCCACACACAGGGGCCAGTGGATAGAC
TdT_L_BsaI_light	mouse IgK light-chain cDNA	CTAGGGTCTCAGTGGTATCAACGCAGAGTAC
mvkappa		GAYATTGTGMTSACMCARWCTMCA
mCkappa-outer		ACTGAGGCACCTCCAGATGTT
mC_K_new		AGGCACCTCCAGATGTAACTGCTCACTGG
BsaI_IL6sp_kappa_L		CTAGGGTCTCAGTGGTATGAACTCCTTCTCCACAAGCG
mC_K_new_BsaI		TCCTAGGTCTCCTCCAGATGTAACTGCTCACTGG

Supplementary Table 1.


Oligonucleotides

Primer	Applicati on	Sequence
Next Generation sequencing		
M13fwd_i5_mC_G_ new2 ^a	Well index	TGTA AACGACGGCCxxxxxxxxATCTCCACACAGGGGC CAGTGGATAGAC
M13fwd_i5_mC_K ^a		TGTA AACGACGGCCxxxxxxxxGCACCTCCAGATGTTAAC TG
M13fwd_i5_mC_L ^a		TGTA AACGACGGCCxxxxxxxxAGCTCCTCAGRGAAGGT G
M13rev_i7_msVHE ^b		GGAAACAGCTATGACyyyyyyyyGAGGTGCAGCTGCAGGAG TCTGG
M13rev_ i7_mv kappa ^b		GGAAACAGCTATGACyyyyyyyyGAYATTGTGMTSACMCAR WCTMCA
i5-Fwd ^c	Plate index	AATGATACGGCGACCA CCGAGATCTACACxxxxxxxx ACACTCTTTCCCTACACGACGCTCTTCCGATCTTGTA AAC GACGGCC
i7-Rev ^d		CAAGCAGAAGACGGCATA CAGAGATyyyyyyyy GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTGGAAAC AGCTATGAC
i5-Rev ^c		AATGATACGGCGACCA CCGAGATCTACACxxxxxxxx ACACTCTTTCCCTACACGACGCTCTTCCGATCTGGAAAC AGCTATGAC
i7-Fwd ^d		CAAGCAGAAGACGGCATA CAGAGATyyyyyyyy GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTTGTA AA ACGACGGCC

^axxxxxxxx: sequence showing the well index from D501 to D508. ^byyyyyyyy: sequence showing the well index from D701 to D712. ^cxxxxxxxx: sequence showing the plate index from D501 to D508. ^dyyyyyyyy: sequence showing the plate index from D701 to D712.


Supplementary Table 2.

Oligonucleotides




Supplementary Figure 1.

The Ig repertoires were visualized using the in-house software. **A**, Overview of heavy (V-D-J) and light (V-J) chain usages. **B**, the repertoire clonality from single cells. **C**, the mutation rates of heavy chains in three cell populations (PR8+, H2+, PR8+H2+) were compared. **D**, the lengths of CDR3 of heavy chains of three cell populations (PR8+, H2+, PR8+H2+) were compared.




Supplementary Figure 2.

Kinetics analysis was performed using a BIAcore 3000 machine (GE Healthcare). A6p4, B10p2, C10p2, A2p1, D11p4, E11p2, G6p2, D4p4 or G12p4 antibodies were immobilized on a CM5 sensor chip (GE Healthcare) using an amine-coupling kit according to the supplier's instructions. HA probes were serially diluted at five different concentrations in HBS-EP buffer.



Supplementary Figure 3.

Two hundred and eighty-four clones are plotted in a two-dimensional map of phylogeny trees for heavy and light chain genes. Nine mAb clones are overlaid in green. Six “cross” clones are located in a major cluster. B10p2 and C10p2 are co-localized and distant from the cluster. A6p4 is uniquely located in the map.



Supplementary Figure 4.

The affinity of antigen-antibody binding (in this case, probe and membrane-bound antibody expressing cell) was inferred from the fluorescence intensity (flow cytometry analysis).

References

1. Balsitis S. J., Williams K. L., et al. (2010) **Lethal antibody enhancement of dengue disease in mice is prevented by Fc modification** *PLoS Pathog* **6**
2. Fang X. T., Sehlin D., et al. (2017) **Efficient and inexpensive transient expression of multispecific multivalent antibodies in Expi293 cells** *Biol. Proced. Online* **19**:1–9
3. Köhler G., Milstein C. (1975) **Continuous cultures of fused cells secreting antibody of predefined specificity** *Nature* **256**:495–497
4. Traggiai E., Becker S., et al. (2004) **An efficient method to make human monoclonal antibodies from memory B cells: potent neutralization of SARS coronavirus** *Nat. Med* **10**:871–875
5. Kwakkenbos M. J., Diehl S. A., et al. (2010) **Generation of stable monoclonal antibody-Producing B cell receptor-positive human memory B cells by genetic programming** *Nat. Med* **16**:123–128
6. Wrammert J., Smith K., et al. (2008) **Rapid cloning of high-affinity human monoclonal antibodies against influenza virus** *Nature* **453**:667–671
7. Meijer P. J., Andersen P. S., et al. (2006) **Isolation of human antibody repertoires with preservation of the natural heavy and light chain pairing** *J. Mol. Biol* **358**:764–772
8. Kuraoka M., Schmidt A. G., et al. (2016) **Complex antigens drive permissive clonal selection in germinal centers** *Immunity* **44**:542–552
9. Clackson T., Hoogenboom H. R., et al. (1991) **Making antibody fragments using phage display libraries** *Nature* **352**:624–628
10. Feldhaus M. J., Siegel R. W., et al. (2003) **Flow-cytometric isolation of human antibodies from a nonimmune *Saccharomyces cerevisiae* surface display library** *Nat. Biotechnol* **21**:163–170
11. Harvey B. R., Georgiou G., et al. (2004) **Anchored periplasmic expression, a versatile technology for the isolation of high-affinity antibodies from *Escherichia coli*-expressed libraries** *Proc. Natl. Acad. Sci. U. S. A* **101**:9193–9198
12. Schaffitzel C., Hanes J., et al. (1999) **Ribosome display: an in vitro method for selection and evolution of antibodies from libraries** *J. Immunol. Methods* **231**:119–135
13. Mazor Y., Van Blarcom T., et al. (2007) **Isolation of engineered, full-length antibodies from libraries expressed in *Escherichia coli*** *Nat. Biotechnol* **25**:563–565
14. Kubota-Koketsu R., Mizuta H., et al. (2009) **Broad neutralizing human monoclonal antibodies against influenza virus from vaccinated healthy donors** *Biochem. Biophys. Res. Commun* **387**:180–185
15. Setliff I., Shiakolas A. R., et al. (2019) **High-throughput mapping of B cell receptor sequences to antigen specificity** *Cell* **179**:1636–1646

16. Kirchmaier S., Lust K., et al. (2013) **Golden GATeway cloning--a combinatorial approach to generate fusion and recombination constructs** *PLoS One* **8**
17. Okuno Y., Isegawa Y., et al. (1993) **A common neutralizing epitope conserved between the hemagglutinins of influenza A virus H1 and H2 strains** *J. Virol* **67**:2552–2558
18. Ekiert D. C., Kashyap A. K., et al. (2012) **Cross-neutralization of influenza A viruses mediated by a single antibody loop** *Nature* **489**:526–532
19. Impagliazzo A., Milder F., et al. (2015) **A stable trimeric influenza hemagglutinin stem as a broadly protective immunogen** *Science* **349**:1301–1306
20. D'Angelo S., Ferrara F., et al. (2018) **Many Routes to an Antibody Heavy-Chain CDR3: Necessary, Yet Insufficient, for Specific Binding** *Front. Immunol* **9**
21. Xu J. L., Davis M. M. (2000) **Diversity in the CDR3 region of VH is sufficient for most antibody specificities** *Immunity* **13**:37–45
22. Gamblin S. J., Skehel J. J. (2010) **Influenza hemagglutinin and neuraminidase membrane glycoproteins** *J. Biol. Chem* **285**:28403–28409
23. Adachi Y., Onodera T., et al. (2015) **Distinct germinal center selection at local sites shapes memory B cell response to viral escape** *J. Exp. Med* **212**:1709–1723
24. Kaplon H., Crescioli S., et al. (2023) **Antibodies to watch in 2023** *Mabs* **15**
25. Soni P., Yasuhara A., et al. (2018) **Evaluation of the fusion partner cell line SPYMEG for obtaining human monoclonal antibodies against influenza B virus** *J. Vet. Med. Sci* **80**:1020–1024
26. Petropoulos K (2012) **Phage display** *Methods Mol. Biol* **901**:33–51
27. Turchaninova M. A., Davydov A., et al. (2016) **High-quality full-length immunoglobulin profiling with unique molecular barcoding** *Nat. Protoc* **11**:1599–1616
28. Alamyar E., Duroux P., et al. (2012) **IMGT® tools for the nucleotide analysis of immunoglobulin (IG) and T cell receptor (TR) V-(D)-J repertoires, polymorphisms, and IG mutations: IMGT/V-QUEST and IMGT/HighV-QUEST for NGS** *Methods Mol. Biol* **882**:569–604

Article and author information

Takashi Watanabe

Laboratory for Integrative Genomics, RIKEN Center for Integrative Medical Sciences, Yokohama, 230-0045, Japan

For correspondence: twatanabe@riken.jp

ORCID iD: [0000-0002-6184-0375](https://orcid.org/0000-0002-6184-0375)

Hikaru Hata

Laboratory for Lymphocyte Differentiation, RIKEN Center for Integrative Medical Sciences, Yokohama, 230-0045, Japan

Yoshiki Mochizuki

Laboratory for Integrative Genomics, RIKEN Center for Integrative Medical Sciences,
Yokohama, 230-0045, Japan

Fumie Yokoyama

Laboratory for Integrative Genomics, RIKEN Center for Integrative Medical Sciences,
Yokohama, 230-0045, Japan

Tomoko Hasegawa

Laboratory for Integrative Genomics, RIKEN Center for Integrative Medical Sciences,
Yokohama, 230-0045, Japan

Naveen Kumar

Laboratory for Integrated Bioinformatics, RIKEN Center for Integrative Medical Sciences,
Yokohama, 230-0045, Japan

Tomohiro Kurosaki

Laboratory for Lymphocyte Differentiation, RIKEN Center for Integrative Medical Sciences,
Yokohama, 230-0045, Japan, Laboratory of Lymphocyte Differentiation, WPI Immunology
Frontier Research Center, Osaka University, Osaka, 565-0871, Japan

Osamu Ohara

Laboratory for Integrative Genomics, RIKEN Center for Integrative Medical Sciences,
Yokohama, 230-0045, Japan, Department of Applied Genomics, Kazusa DNA Research Institute,
Chiba 292-0818, Japan
ORCID iD: [0000-0002-3328-9571](https://orcid.org/0000-0002-3328-9571)

Hidehiro Fukuyama

Laboratory for Lymphocyte Differentiation, RIKEN Center for Integrative Medical Sciences,
Yokohama, 230-0045, Japan, Near-InfraRed Photo-Immunotherapy Research Institute, Kansai
Medical University, Hirakata, Osaka, 573-1010, Japan, INSERM EST, Strasbourg Cedex 2, 67200,
France

Copyright

© 2024, Watanabe et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/),
which permits unrestricted use and redistribution provided that the original author and
source are credited.

Editors

Reviewing Editor

Pramod Srivastava

UConn Health, Farmington, United States of America

Senior Editor

Tadatsugu Taniguchi

University of Tokyo, Tokyo, Japan

Reviewer #1 (Public Review):

Summary:

This paper by Watanabe et al described an expression system that can express the paired heavy and light chains of IgG antibodies from single cell B cells. In addition, they used FACS sorting for specific antigens to screen/select the specific populations for more targeted cloning of mAb genes. By staining with multiple antigens, they were able to zoom in to cross-reactive antibodies.

Strengths:

A highly efficient process that combines selection/screening with dual expression of both antibody chains. It is particularly suitable for the isolation of cross-reactive antibodies against conserved epitopes of different antigens, such as surface proteins of related viruses.

Weaknesses:

(1) The overall writing is very difficult to follow and the authors need to work on significant re-writing.

(2) The paper in its current form really lacks detail and it is NOT possible for readers to repeat or follow their methods. For example: a) It is not clear whether the authors checked the serum to see if the mice were producing antibodies before they sacrificed them to harvest spleen/blood i.e. using ELISA? b) How long after administration of the second dose were the mice sacrificed? c) What cell types are taken for single B cell sorting? Splenocytes or PBMC? These are just some of the questions which need to be addressed.

(3) According to the authors, 77 clones were sorted from the PR8+ and H2+ double positive quadrant. It is surprising that after transfection and re-analysing of bulk antibody presenting EXPI cells on FACS, only 13 clones (or 8 clones? - unclear) seemed to be truly cross-reactive. If that is the case, the approach is not as efficient as the authors claimed.

<https://doi.org/10.7554/eLife.95346.1.sa1>

Reviewer #2 (Public Review):

Summary:

Watanabe, Takashi, et al. investigated the use of the Golden Gate dual-expression vector system to enhance the modern standard for rapid screening of recombinant monoclonal antibodies. The presented data builds upon modern techniques that currently use multiple expression vectors to express heavy and light chain pairs. In a single vector, they express the linked heavy and light chain variable genes with a membrane-bound Ig which allows for rapid and more affordable cell-based screening. The final validation of H1 and H2 strain influenza screening resulted in 81 "H1+", 48 "H2+", and 9 "cross" reactive clones. The kinetics of some of the soluble antibodies were tested via SPR and validated with a competitive inhibition with classical well-characterized neutralizing clones.

Strengths:

In this study, Watanabe, Takashi, et al. further develop and refine the methodologies for the discovery of monoclonal antibodies. They elegantly merge newer technologies to speed up turnaround time and reduce the cost of antibody discovery. Their data supports the feasibility of their technique.

This study will have an impact on pandemic preparedness and antibody-based therapies.

Weaknesses:

A His tagged antigen was used for immunization and H1-his was used in all assays. Either the removal of His specific clones needs to be done before selection, or a different tag needs to be used in the subsequent assays.

This assay doesn't directly test the neutralization of influenza but rather equates viral clearance to competitive inhibition. The results would be strengthened with the demonstration of a functional antibody in vivo with viral clearance.

Limitations of this new technique are as follows: there is a significant loss of cells during FACs, transfection and cloning efficiency are critical to success, and well-based systems limit the number of possible clones (as the author discussed in the conclusions). Early enrichment of the B cells could improve efficiency, such as selection for memory B cells.

<https://doi.org/10.7554/eLife.95346.1.sa0>