

Targeting resident astrocytes attenuates neuropathic pain after spinal cord injury

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint version 1

April 15, 2024 (this version)

Posted to preprint server

January 23, 2024

Sent for peer review

January 23, 2024

Qing Zhao, Yanjing Zhu, Yilong Ren, Lijuan Zhao, Jingwei Zhao, Shuai Yin, Haofei Ni, Rongrong Zhu , Liming Cheng , Ning Xie 

Key Laboratory of Spine and Spinal Cord Injury Repair and Regeneration of Ministry of Education, Orthopaedic Department of Tongji Hospital, School of Medicine, School of Life Sciences and Technology, Tongji University, Shanghai, 200065 P. R. China • Division of Spine, Department of Orthopedics, Tongji Hospital, Tongji University School of Medicine, Tongji University, Shanghai, 200065 P. R. China • Clinical Center for Brain and Spinal Cord Research, Tongji University

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Astrocytes derive from different lineages and play a critical role in neuropathic pain after spinal cord injury (SCI). Whether selective eliminating these main origins of astrocytes in lumbar enlargement could attenuate SCI-induced neuropathic pain remains unclear. In this study, astrocytes in lumbar enlargement were lineage traced, targeted and selectively eliminated through transgenic mice injected with an adeno-associated virus vector and diphtheria toxin. Pain-related behaviors were measured with an electronic von Frey apparatus and a cold/hot plate after SCI. RNA sequencing, bioinformatics analysis, molecular experiment and immunohistochemistry were used to explore the potential mechanisms after astrocyte elimination. Through lineage tracing, we concluded the resident astrocytes but not ependymal cells were the main origins of astrocytes-induced neuropathic pain. SCI induced mice to obtain significant pain symptoms and astrocyte activation in lumbar enlargement. Selective resident astrocytes elimination in lumbar enlargement could attenuate neuropathic pain and activate microglia. Interestingly, the type I interferons (IFNs) signal was significantly activated after astrocytes elimination, and the most activated Gene Ontology terms and pathways were associated with the type I IFNs signal which was mainly activated in microglia and further verified in vitro and in vivo. Furthermore, different concentrations of interferon and Stimulator of interferon genes (STING) agonist could activate the type I IFNs signal in microglia. Our results elucidate that selectively eliminating resident astrocytes attenuated neuropathic pain associated with type I IFNs signal activation in microglia. Targeting type I IFNs signal is proven to be an effective strategy for neuropathic pain treatment after SCI.

eLife assessment

This **important** study demonstrated that ablation of astrocytes in the lumbar spinal cord not only reduced neuropathic pain but also caused microglia activation. The findings presented add considerable value to the current understanding of the role of astrocyte elimination in neuropathic pain, offering **convincing** evidence that supports existing hypotheses and insights into the interactions between astrocytes and microglial cells, likely through IFN-mediated mechanisms. This study may also offer a new therapeutic strategy for the treatment of debilitating neuropathic pain in patients with SCI.

Introduction

Spinal cord injury (SCI) can cause sensorimotor deficits and autonomic changes, and approximately 65% of SCI patients suffer chronic, severe neuropathic pain (Henwood and Ellis, 2004). Long-term unrelieved pain will hinder the patient's rehabilitation training, delay functional recovery, and seriously affect the patient's physical and mental health (Henwood and Ellis, 2004; Masri and Keller, 2012). Unfortunately, the effect of current treatments for neuropathic pain is limited (Dworkin et al., 2003). Exploring an effective treatment for neuropathic pain is a major clinical challenge. Astrocytes are the most abundant type of glial cell in the central nervous system (Dozio and Sanchez, 2018). Accumulating studies have shown that astrocytes play a key role in the induction and maintenance of persistent pain (Ji et al., 2019). Cytokines and chemokines produced by astrocytes can powerfully modulate excitatory and inhibitory synaptic transmission in pain circuits, leading to central sensitization, and a transition from acute to chronic neuropathic pain (Ji et al., 2013; Ji et al., 2019; Kawasaki et al., 2008). Inhibiting the proliferation of spinal astrocytes or glial fibrillary acidic protein (GFAP) expression could reduce neuropathic pain (Ji et al., 2019; Tsuda et al., 2011).

Astrocytes are mainly derived from resident astrocytes, NG2 progenitor and ependymal cells (Barnabé-Heider et al., 2010; Ren et al., 2017; Sofroniew, 2009). SCI leads to massive proliferation of resident astrocytes, which are the most abundant and mainly located around the scar (Barnabé-Heider et al., 2010; Sabelstrom et al., 2014). Ependyma is regarded as the main source of endogenous neural stem cells, which widely migrate to injury area and generate the majority of neuroprotective astrocytes after SCI (Barnabé-Heider et al., 2010; Meletis et al., 2008). However, we found that only a small number of ependymal cells could proliferate and migrate to the lesion score after SCI (Ren et al., 2017). NG2 cells, also known as oligodendrocyte progenitor cells, were previously considered to be the component of glial scar after SCI (Church et al., 2016; Hackett and Lee, 2016; Hesp et al., 2015). While, reports about their functions are still controversial, partly because there are several cell types that up-regulate proteoglycan NG2 after SCI, such as oligodendrocyte progenitor cells, pericytes and macrophages (Huang et al., 2020; Nishiyama et al., 2005). Until now, there is limited data on the main origins of astrocytes-induced neuropathic pain after SCI and whether selectively target these astrocytes could attenuate neuropathic pain is still unclear.

Cervical and thoracic SCI often induce below-level neuropathic pain (Gwak et al., 2012; Shiao and Lee-Kubli, 2018; Vierck, 2020), partly through activating signal transducer and activator of transcription 3 (STAT3)-dependent reactive astrocytes (Ono et al., 2020). The expression of tumor necrosis factor- α (TNF- α), interleukin-1 β (IL-1 β) and interleukin-6 (IL-6) in lumbar enlargement was closely associated with below-level neuropathic pain (Honjoh et al., 2019). Mainly because the ascending tracts and superficial dorsal horn (SDH) in lumbar enlargement are

closely associated with proprioception and pain signal processing of the lower limbs (Watson, 2012). Thus, targeting the astrocytes in lumbar enlargement may improve neuropathic pain after cervical or thoracic SCI.

In this study, we constructed a C57/BL6 mouse neuropathic pain model with spinal cord contusion at thoracic level 10 (T10). Pain behaviors were measured by an electronic von Frey apparatus and a cold/hot plate. Transgenic mice and lineage tracing were used to determine the main origins of astrocytes-induced neuropathic pain after SCI. To investigate if eliminating these astrocytes in lumbar enlargement could attenuate neuropathic pain after SCI, we used the transgenic mice, adeno-associated virus (AAV) vector and diphtheria toxin (DT). We identified a previously undescribed improvement effects of targeting these astrocytes on neuropathic pain. Combining with RNA sequencing, bioinformatics analysis, immunohistochemistry, molecular experiment and cultured microglial cells, we further explore the potential mechanisms of pain relief after selective astrocytes elimination.

Results

Resident astrocytes but not ependymal cells were the main origins of astrocytes-induced neuropathic pain after SCI

The construction strategies of transgenic mice were described in **Figure 1A**. The brain slices from C57BL/6-Foxj1^{em1(GFP-CreERT2-polyA)Smoc} mice showed that choroids plexus in both fourth ventricle and lateral ventricle had been labeled by reporter gene EGFP (**Figure 1B**). However, EGFP positive cells were not found in central canal of spinal cord under normal condition (data not shown), indicating that ependymal cells may not be activated. Therefore, C57BL/6-Foxj1^{em1(GFP-CreERT2-polyA)Smoc} mice were used to establish contusion SCI models and examine the activation and differentiation of ependymal cells. As we reported before (Ren *et al.*, 2017), we found ependymal cells did not transform to astrocytes in the lesion area (**Figure 1C**). S100 family protein p11 (S100A10) is previously regarded as a marker of neuroprotective astrocytes (Liddelow *et al.*, 2017), but not unique and expressed by various cell types (**Supplementary Figure 1A-C**). Instead, ependymal cells were observed to produce offspring of Tuj1 positive neurons, Oligo2 negative and S100A10 positive cells in epicenter (**Figure 1D-E**). Totally, ependymal cells did not transform to astrocytes as well as Oligo2 positive oligodendrocytes. Therefore, the resident astrocytes but not ependymal cells were the main origins of astrocytes-induced neuropathic pain after SCI.

Using transgenic mice and AAV vectors could selectively target resident astrocytes in lumbar enlargement

The experiment flowchart was described in **Figure 2A**. Compared with the sham group, the SCI group showed increased astrocytes activation (GFAP expression) in ventral spinothalamic tracts (VST, 65.08 ± 2.25 vs. 52.71 ± 1.02), SDH (59.62 ± 2.21 vs. 49.05 ± 1.30, **Figure 2B** and **Supplementary Figure 2A**), dorsal spinocerebellar tract (DSC, 81.64 ± 3.63 vs. 51.69 ± 1.22), gracile fasciculi (GR, 58.37 ± 1.89 vs. 49.63 ± 0.83) and ventral spinocerebellar tract (VSC, 75.73 ± 2.21 vs. 50.88 ± 1.29). Thus, astrocytes activation in the lumbar enlargement is associated with neuropathic pain after SCI. AAV2/5-GfaABC1D-Cre with a minimal GFAP promoter was used to selectively target resident astrocytes of *Gt(ROSA)26Sor^{em1(CAG-LSL-tdTomato-2A-DTR)Smoc}* mice. AAV2/5-GfaABC1D-Cre (1:3 dilution, ≥ 0.33E+13 V.G/ml) was injected into the spinal cord of mice (n=3, **Supplementary Figure 2B**). Two weeks after AAV2/5-GfaABC1D-Cre injection, immunofluorescence staining showed that a large number of astrocytes were labelled (**Supplementary Figure 2B**). Meanwhile, almost no tdTomato-positive cells were found in NEUN-positive neurons or Oligo2-positive oligodendrocytes (**Supplementary Figure 2B** and unshown data). To explore the optimal virus




Figure 1.

Ependymal cells did not transform to astrocytes in our transgenic mice.

(A), construction strategies of transgenic mice. (B), ependymal cells both in fourth ventricle (4V) and lateral ventricle (LV) had been labeled by reporter gene EGFP (green). $n = 6$ biological repeats. (C), ependymal cells did not produce astrocytes in the lesion area. $n = 3$ biological repeats. (D-E), ependymal cells were observed to produce offspring of neurons (Tuj1+, red), Oligo2 (white) negative and S100A10 (white in D or red in E) positive cells in epicenter. $n = 3$ biological repeats. 4V=fourth ventricle, LV=lateral ventricle, CB= Cerebellum. Scale bars had been indicated in pictures.

concentration, different dilutions including 1:10 ($\geq 1 \times 10^{12}$ V.G/ml), 1:100 ($\geq 1 \times 10^{11}$ V.G/ml), 1:250 ($\geq 0.4 \times 10^{11}$ V.G/ml), 1:500 ($\geq 0.2 \times 10^{11}$ V.G/ml), 1:1000 ($\geq 1 \times 10^{10}$ V.G/ml) and 1:10000 ($\geq 1 \times 10^9$ V.G/ml), were set for injection. Immunofluorescence results of spinal cord tissue and slices revealed that the dilution of 1:100 ($\geq 1 \times 10^{11}$ V.G/ml) was the optimal intervention concentration for concentrations below 1:100 can hardly target astrocytes (**Supplementary Figure 2B** and unshown data). Then AAV2/5-GfaABC1D-Cre ($\geq 1 \times 10^{11}$ V.G/ml) was injected into *Gt(ROSA)26Sor^{em1}(CAG-LSL-tdTomato-2A-DTR)Smoc* mice 2 weeks before T10 contusion. We found that the labeled resident astrocytes were mainly located in the dorsal side of lumbar enlargement (**Figure 1C**). It was also found that the labeled astrocytes by AAV2/5-GfaABC1D-Cre accounted for $39.03 \pm 1.97\%$ astrocytes in dorsal side of SDH (**Figure 1D**). All labeled astrocytes in SCI+AAV+DT group were eliminated after DT injection (**Figure 1C**) and the expression of GFAP was significantly decreased in dorsal side after resident astrocytes elimination (**Figure 1D**).

Selective resident astrocytes elimination in lumbar enlargement attenuated neuropathic pain

Pain-related behaviors were measured with an electronic von Frey apparatus and a cold/hot plate, including mechanical allodynia, thermal and cold hyperalgesia. The pain test was carried out as described in the experiment flowchart (**Figure 2A**). After contusion, the mice showed a complete hindlimb paralysis and weight loss, and these effects gradually recovered (**Figure 3A-B**). We found that the 5 g impactor and 6.25 mm height contusion could induce mice to obtain obvious pain symptoms (**Figure 3C-F**). Mechanical allodynia (reflected by paw withdrawal threshold) occurred at 14 dpi (day post injury, left hindlimb cut-offs vs. baseline; right hindlimb cut-offs vs. baseline), peaked at 28 dpi and sustained until 43 dpi (**Figure 3C-D**). Cold hyperalgesia (reflected by paw withdrawal frequency) and thermal hyperalgesia (reflected by paw withdrawal latency) also occurred at 14 dpi, peaked at 43 dpi (**Figure 3E-F**). There was no statistical significance in pain-related behaviors, weight and motor function between the sham group and the sham AAV group (**Figure 3A-F**).

A total of 36 mice with neuropathic pain were randomly selected to receive diphtheria toxin (DT) injection on 31, 32 and 34 dpi (male= 18, female= 18, SCI+AAV+DT group), and the 36 mice left were classified as control group (male= 18, female= 18, SCI+AAV group). Then pain tests were carried out on 35 dpi after DT injection. We observed that motor function (BMS score) and weight did not change significantly in mice of SCI+AAV+DT group after selective resident astrocytes elimination in the lumbar enlargement (**Figure 3A-B**). Mechanical allodynia of the left hindlimb attenuated after selective astrocytes elimination at 35 dpi (first day after astrocytes elimination, **Figure 3C**). The pain attenuation effects peaked at 41 dpi (seventh day after astrocytes elimination, **Figure 3C**). The mechanical allodynia of the right hindlimb was relieved after selective astrocytes elimination at 38 dpi (fourth day after astrocytes elimination), and the pain attenuation effects peaked at 41 dpi (**Figure 3D**). The attenuation of both cold and thermal pain occurred at 38 dpi, and the pain attenuation effects peaked at 41 dpi (**Figure 3E-F**). To further determine the effect of selective astrocytes elimination on hindlimb pain, we measured the pain parameters on left and right hindlimbs, and observed no obvious difference between them (data not shown). Furthermore, we explored the effect of sex on neuropathic pain, and observed no significant differences between male and female mice (**Supplementary Figure 3A-H**). In addition, we examined the effect of DT injection alone on neuropathic pain of wild-type mice after contusion. The contusion models were constructed as mentioned above. We observed that pain response was similar in wild-type mice with or without DT injection (**Supplementary Figure 4A**).




Figure 2.

Resident astrocytes in lumbar enlargement were selectively targeted and eliminated.

(A), the experiment flowchart. (B), the histogram and statistical results of GFAP fluorescence intensity. The location of three largest ascending tracts of lumbar enlargements, including (1) CU, GR; (2) LST, DLST, VST; (3) DSC, VSC, along with SDH and gray matter. CU=cuneate fasciculus, GR=gracile fasciculus, LST=lateral spinothalamic tract, DLST=dorsolateral spinothalamic tract, VST=ventral spinothalamic tract, DSC=dorsal spinocerebellar tract, VSC=ventral spinocerebellar tract, SDH=superficial dorsal horn. $n = 3$ biological repeats. Scale bar, 200 μ m. Values are the mean $\pm$ SEM. Statistical significance was determined by one-way ANOVA followed by Student Newman-Keuls post hoc test. $^{**}P < 0.01$, $^{***}P < 0.001$, Sham group vs. SCI Pain positive (+) group. (C), AAV2/5-GfaABC1D-Cre targeted astrocytes were mainly in dorsal side and reflected by tdTomato and GFAP double positive cells in SCI+AAV group, no tdTomato and GFAP double positive cells were found in SCI+AAV+DT group. $n = 3$ biological repeats. Scale bars had been indicated in pictures. (D), the efficiency of astrocytes labeling and elimination in dorsal side of spinal cord. Immunofluorescence analysis results of GFAP after selective astrocyte elimination. Values are the mean $\pm$ SEM. Statistical significance was determined by one-way ANOVA followed by Student Newman-Keuls post hoc test. $n = 3$ biological repeats. TdTomato positive cells indicated AAV. 2/5-GfaABC1D-Cre targeted astrocytes. $^{***}P < 0.001$.




Figure 3.

Selective astrocyte elimination in lumbar enlargement attenuated neuropathic pain.

Change of weight (**A**), BMS scores (**B**), mechanical allodynia (**C-D**), cold hyperalgesia (**E**) and thermal hyperalgesia (**F**) in mice after astrocyte elimination. Sham group, n=30; sham+AAV group, n=30; SCI+AAV group, n=36; SCI+AAV+DT group, n=36. Arrow, 500 ng DT injection was performed on 31, 32 and 34 dpi (days post-injury). Left, left hindlimbs; right, right hindlimbs. i.p., intraperitoneal injection. Values are the mean $\pm$ SEM. Statistical significance was determined by one-way ANOVA followed by Student Newman-Keuls post hoc test. *P < 0.05, **P < 0.01, ***P < 0.001, SCI+AAV+DT group vs. SCI+AAV group.

Microglia were activated after selective resident astrocytes elimination in lumbar enlargement

Immunofluorescence staining showed that increased activation of astrocytes and microglia and neuron loss were observed in the lesion area in SCI+AAV group and SCI+AAV+DT group after contusion (**Figure 4A**, **4C** and **Supplementary Figure 4B**). Compared with SCI+AAV group, enhanced expression of microglia marker Ionized calcium binding adaptor molecule 1 (IBA1) in lumbar enlargement was observed after selective resident astrocytes elimination in SCI+AAV+DT group (**Figure 4B-C**). Furthermore, compared with the SCI+AAV group, the heatmap of RNA-seq analysis results showed that the transcripts of pro-inflammatory microglial marker genes elevated in the SCI+AAV+DT group (**Figure 4D**). As previously reported (Watson, 2012), we selected the three largest ascending tracts, grey matter and SDH to analyze in the immunohistochemistry images. We found that selective resident astrocytes elimination led to enhanced IBA1 expression in the left DSC, left deep grey matter, left LST, left SDH, left VSC, right GR, right shallow grey matter and right SDH (**Figure 4E-H**). These results indicated that selective resident astrocytes elimination activated the microglia in lumbar enlargement.

Selective resident astrocytes elimination activated the type I IFNs signal in lumbar enlargement

To explore the potential mechanisms of pain attenuation after selective resident astrocytes elimination, RNA-seq and bioinformatics analysis were conducted. RNA-seq analysis revealed that there were few DEGs between the sham group and sham AAV group (**Supplementary Figure 4C**). Between SCI+AAV group and sham AAV group, 1872 differentially expressed genes (DEGs) were found in SCI+AAV group, including 1654 upregulated and 218 downregulated genes, as shown in volcano map (**Supplementary Figure 4D**). The Gene Ontology (GO) enrichment analysis and KEGG pathway analysis revealed that the abundance of DEGs enriched in several biological processes, which were associated with immune responses including immune system processes, inflammatory responses and innate immune responses (**Supplementary Figure 4E-F**). Compared with SCI+AAV group, the SCI+AAV+DT group (astrocytes elimination group) had 1120 DEGs, including 688 upregulated and 432 downregulated genes. We observed that the upregulated DEGs after selective resident astrocytes elimination were mainly enriched in regulating interferons (IFNs) production and signal transduction of type I IFNs (**Figure 5A**) (Borden et al., 2007; Owens et al., 2014; Rothhammer et al., 2016). Protein-protein interaction (PPI) network analysis showed the interaction of these DEGs (**Figure 5B**). Variability was observed between individual animals within the SCI+AAV and SCI+AAV+DT groups, but the expression differences of DEGs in these two groups were significant.

GO enrichment analysis showed that the dominant biological processes were associated with response to virus, immune and type I IFNs responses, including defense response to virus, response to virus, immune system process, cellular response to interferon-alpha/-beta, positive regulation of interferon-alpha/-beta production or secretion, type I interferon signaling pathway, response to interferon-beta, type I interferon biosynthetic process (**Figure 5C**). KEGG pathway analysis found that several signaling pathways were disturbed, including the influenza A, Epstein-Barr virus infection, Herpes simplex virus 1 infection, Measles, Hepatitis C and NOD-like receptor signaling pathways (**Figure 5D**). Heatmap showed significantly elevated transcripts level of critical genes involved in the type I IFNs signaling pathway, including *Jak1/2/3*, *Ifna16*, *Cgas*, *Tyk2*, *Ifnar1/2*, *Sting1*, *Irf3/7/9*, *Ifnb1*, and *Ifih1* (**Figure 5E**). Meanwhile, gene set enrichment analysis (GSEA) showed the disturbance of the Janus kinase (Jak)-STAT signaling pathway and cytokine-cytokine receptor interactions (**Figure 5F-G**). Thus, GO enrichment, KEGG pathway, and GSEA




Figure 4.

Microglia in lumbar enlargement were activated after selective astrocyte elimination.

(A-B), images of immunofluorescent staining using GFAP (green), Tuj1 (red) and IBA1 (white) as characteristic markers of astrocyte, neuron and microglia, respectively. $n = 3$ biological repeats. Scale bar, 200 μ m. (C), the histogram and statistical results of GFAP, IBA1 and Tuj1 fluorescence intensity in lesion area. (D), the heatmap showed that the transcripts of pro-inflammatory and anti-inflammatory microglial marker genes. $n = 3$ biological repeats. (E-H), the histogram and statistical results of IBA1 fluorescence intensity in lumbar enlargement. CU=cuneate fasciculus, GR=gracile fasciculus, LST=lateral spinothalamic tract, DLST=dorsolateral spinothalamic tract, VST=ventral spinothalamic tract, DSC=dorsal spinocerebellar tract, VSC=ventral spinocerebellar tract, SDH=superficial dorsal horn. Values are the mean $\pm$ SEM. Statistical significance was determined by one-way ANOVA followed by Student Newman-Keuls post hoc test. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, SCI+AAV+DT group vs. SCI+AAV group.




Figure 5.

Selective astrocyte elimination activated Type I IFNs signal.

(A), the heatmap showed the DEGs between SCI+AAV+DT group and SCI+AAV group. (B), PPI networks analysis showed the interaction of the DEGs between SCI+AAV+DT group and SCI+AAV group. (C), GO enrichment analysis revealed the dominant biological process of DEGs between SCI+AAV+DT group and SCI+AAV group. (D), KEGG pathway showed dysregulated signaling pathways between SCI+AAV+DT group and SCI+AAV group. (E-F), GSEA analysis showed the dysregulated signaling pathways between SCI+AAV+DT group and SCI+AAV group. (G), the heatmap showed that the critical genes expression involved in type I IFNs signal. n = 3 biological repeats.

analysis results were associated with the type I IFNs signaling pathway (Barrat et al., 2019 [↗](#); Ivashkiv and Donlin, 2014 [↗](#)). Totally, these results indicated that selective resident astrocytes elimination activated the type I IFNs signal in lumbar enlargement.

The type I IFNs signal was activated in microglia and associated with pain relief

In order to further verify the results of RNA-seq, RT-PCR was carried out and the results showed the mRNA expression of genes associated with the type I IFNs signaling pathway increased, including *IFIH1*(*MAD5*), *IRF3*, *IRF7*, *IF144*, *STAT1*, *STAT2*, *PARP14*, *IRF9*, *ISG15*, *OAS2*, *OAS3*, *ZBP1*, *IFIT1*, *USP18*, *DHX58*, and *Eifzak2* (Figure 6A [↗](#)). We also observed that mRNA expression of type I IFNs signaling genes was similar in wild-type mice with or without DT injection (Figure 6B [↗](#)). Furthermore, we selected three critical proteins involved in the type I IFNs signal with significantly increased expression at the transcriptional level, including IRF7, ISG15 and STAT1. Through immunofluorescence analysis, we found that the expression of IRF7, ISG15 and STAT1 proteins was also significantly increased after selective astrocytes elimination (Figure 6C-D [↗](#)).

Although various cells in the central nervous system can synthesize and secrete interferons, microglia are resident immune cells and are believed to be the main source of IFN- β within the central nervous system during inflammatory conditions (Owens et al., 2014 [↗](#)). Reinert and colleagues (Reinert et al., 2016 [↗](#)) identified microglia as the primary producers of type I IFNs. We also found that ISG15 and IRF7 were co-located with the microglial marker IBA1 (Figure 6E [↗](#)). Thus, ISG15 and IRF7 were mainly expressed by microglia. To further confirmed the expression of type I IFNs signaling factors were derived from microglia, the cultured and lipopolysaccharides (LPS)-activated microglia were used. Compared with the results for the control microglia (without LPS stimulation), the RT-PCR results revealed that 0.2 μ g/mL LPS mainly activated pro-inflammatory microglia, including the pro-inflammatory microglial markers TNF- α , IL-1 β , IL-6, and iNOS (Figure 6G [↗](#)). Anti-inflammatory microglial markers, including IL-4, TGF- β , IL-10, and Arg-1, were not significantly increased (Figure 6G [↗](#)). Additionally, RT-PCR results showed the upregulated mRNA expression of type I IFNs signaling critical genes in LPS-activated microglia, including *STING1*, *STAT2*, *STAT1*, *ISG15*, *IRF9*, and *IRF3* (Figure 6H [↗](#)). These results indicated that type I IFNs signal was activated in microglia.

In order to further verify the activation of type I IFNs signal in microglia, we used different concentrations of interferon agonist (Ploy:IC)(Khorrooshi et al., 2015 [↗](#)) and STING agonist (ADU-S100 and DMXAA) (Li et al., 2023 [↗](#)) to stimulate microglia at 12 and 24 hours. We found that the mRNA expression of *IFNb*, *ISG15*, *STAT2*, and *STAT1* increased significantly after Ploy:IC stimulation (Figure 7A-B [↗](#)). The mRNA expression levels of *IRF7*, *IFNb*, *STAT2*, *ISG15*, *STAT1*, and *IRF9* significantly increased after stimulation with ADU-S100 and DMXAA (Figure 7C-D [↗](#)). A previous study identified that type I IFNs signals can significantly reduce sensory neuron activation and pain level (Donnelly et al., 2021 [↗](#)). Thus, the pain relief after selective resident astrocytes elimination may be associated with the activation of type I IFNs signal in microglia after SCI.

Discussion

Astrogliosis is induced early in neuropathic pain models and sustained for exceptionally long durations, which covers both the acute-to-chronic pain transition and the maintenance phase of pain (Ji et al., 2019 [↗](#)). Cytokines and chemokines produced by astrocytes can powerfully modulate excitatory and inhibitory synaptic transmission in pain circuits, leading to central sensitization, and a transition from acute to chronic neuropathic pain (Ji et al., 2013 [↗](#); Ji et al., 2019 [↗](#); Kawasaki et al., 2008 [↗](#)). Recently, Kohro et al. (Kohro et al., 2020 [↗](#)) identified an SDH-localized Hes5+ astrocytes population that gated descending noradrenergic control of mechanosensory behaviour




Figure 6.

The type I IFNs signal was activated in microglia after selective resident astrocyte elimination.

(A-B), RT-PCR identified the effects of selective astrocyte elimination and DT injection alone on mRNA expression of type I IFNs signal genes. $n = 3$ biological repeats. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$. (C-D), images of immunofluorescent staining and statistical results using IRF7, ISG15 and STAT1 as critical proteins of type I IFNs signal and microglia marker IBA1. Scale bar, 200µm. $n = 3$ biological repeats. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, SCI+AAV+DT group vs. SCI+AAV group. (E), images of immunofluorescent staining using IRF7, ISG15 and IBA1. Scale bar, 200µm. $n = 3$ biological repeats. (F-G), the mRNA expression of pro- and anti-inflammatory microglial markers and type I IFNs signal critical genes after 0.2 µg/mL LPS stimulation. $n = 3$ biological repeats. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, Control group vs. LPS 0.2µg group. All values are the mean $\pm$ SEM. Statistical significance was determined by one-way ANOVA followed by Student Newman-Keuls post hoc test.




Figure 7.

Interferon agonist (Ploy:IC) an STING agonist (ADU-S100 and DMXAA) significantly activated the type I IFNs signal in microglia.

A-F, the mRNA expression levels of *IRF7*, *IRF3*, *IFNβ*, *STAT2*, *ISG15*, *STAT1*, and *IRF9* after stimulating with different concentrations of Ploy:IC, ADU-S100 and DMXAA. $n = 3$ biological repeats. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, treatment group vs. control group. All values are the mean $\pm$ SEM. Statistical significance was determined by one-way ANOVA followed by Student Newman-Keuls post hoc test.

via $\alpha 1A$ adrenergic receptors ($\alpha 1A$ -Ars). Inhibition of locus coeruleus noradrenergic signal transduction to Hes5 + astrocytes could effectively prevent mechanical hyperalgesia induced by capsaicin (Kohro *et al.*, 2020 [DOI](#)). In this work, we observed that astrocytes activation increased in the lumbar enlargement of mice with SCI-induced pain, including the three largest ascending tracts and the SDH. These findings further highlight the importance of astrocytes in processing pain signals and targeting astrocytes may improve neuropathic pain after SCI.

Resident astrocytes maintain stable under normal physiological state and rarely proliferates. SCI leads to massive proliferation of resident astrocytes especially in lesion area and re-express the neural progenitor cell markers, which are the most abundant and mainly located around the scar forming a region complementary to cells derived from ependymal cells (Barnabé-Heider *et al.*, 2010 [DOI](#); Sabelstrom *et al.*, 2014 [DOI](#)). Ependyma is regarded as the main source of endogenous neural stem cells, which widely migrate to injury area and generate the majority of neuroprotective astrocytes after SCI (Barnabé-Heider *et al.*, 2010 [DOI](#); Meletis *et al.*, 2008 [DOI](#)). However, we found that only a small number of ependymal cells could proliferate and migrate to the lesion score after SCI (Ren *et al.*, 2017 [DOI](#)). Through transgenic mice and lineage tracing, we examined the differentiation of ependymal cells and found that ependymal cells did not produce a large number of astrocytes. Instead, ependymal cells differentiated to Tuj1 positive neurons in epicenter. NG2 cells were previously considered to be the component of glial scar after SCI (Church *et al.*, 2016 [DOI](#); Hackett and Lee, 2016 [DOI](#); Hesp *et al.*, 2015 [DOI](#)). However, reports about their functions are still controversial, partly because there are many cell types that up regulate proteoglycan NG2 in the scar and lesion core after SCI (Nishiyama *et al.*, 2005 [DOI](#)). Totally, through lineage tracing, this study determined that the resident astrocytes but not ependymal cells were the main origins of astrocytes-induced neuropathic pain after SCI.

One of our novel findings was that selective resident astrocytes elimination in the lumbar enlargement could attenuate neuropathic pain after SCI. Previously, there were limited data on the association between local astrocytes depletion and neuropathic pain attenuation. We tried to target astrocytes in the spinal cord by using GFAP-Cre; *Gt(ROSA)26Sor^{em1}(CAG-LSL-tdTomato-2A-DTR)^{Smoc}* mice, but we identified that the mice were not suitable for labelling astrocytes (Supplementary Figure 5). The most reliable mouse strain for targeting astrocytes in vivo is *Aldh11-Cre/ERT2* mice (Srinivasan *et al.*, 2016 [DOI](#)). To selectively and locally target astrocytes in spinal cord, instead, selective astrocytes elimination in lumbar enlargement was performed by AAV2/5-GfaABC1D-Cre mediated local astrocytes transduction and DT injection. Several previous studies have reported astrocytes elimination in animal experiments. Through transgenical method, a previous study found that domain-specific depletion of astrocytes in dorsal spinal cord showed atrophy, reduction in the total number of astrocytes, loss of neuropil, and congested neurons (Tsai *et al.*, 2012 [DOI](#)). However, they did not observe increased inflammation, gliosis, or blood brain barrier (BBB) permeability in these mice (Tsai *et al.*, 2012 [DOI](#)). They speculated that remaining astrocytes were sufficient for structural maintenance (Tsai *et al.*, 2012 [DOI](#)). In contrast, domain-specific depletion of astrocytes in ventral spinal cord resulted in abnormal motor neuron synaptogenesis, which was not rescued by immigration of astrocytes from adjoining regions (Tsai *et al.*, 2012 [DOI](#)). With transgenically targeted DTR and AAV2/5-GfaABC1D-Cre injection, one another study found that DT-mediated ablation of chronic astrocytic scar could significantly prevent axon regrowth after severe SCI of mice (Anderson *et al.*, 2016 [DOI](#)). Two other studies also found that astrocytes elimination of the whole body impaired redox homeostasis and triggered neuronal loss in the central nervous system, accompanied by myelin sheath loss and motor function impairment (Allnoch *et al.*, 2019 [DOI](#); Schreiner *et al.*, 2015 [DOI](#)). Meanwhile, local astrocytes depletion through local DT injection did not affect astrocytes in other sites as shown in their supplemental information (Schreiner *et al.*, 2015 [DOI](#)). Our results also showed that selective resident astrocytes elimination in the lumbar enlargement does not have a significant impact on weight and motor function.

We also found the transcripts of pro-inflammatory microglial marker genes elevated and microglia could activate after selective resident astrocytes elimination. As the resident macrophages in the spinal cord, microglia play critical roles in the progress of neuropathic pain (Chen *et al.*, 2018 [DOI](#); Salter and Stevens, 2017 [DOI](#)). However, the role of microglia in chronic inflammatory pain is still controversial (Chen *et al.*, 2018 [DOI](#)). A previous study reported that P2×4 knockout mice exhibit unaltered inflammatory pain (Tsuda *et al.*, 2009 [DOI](#)). In contrast, Cx3cr1 knockout mice exhibit reduced inflammatory pain (Staniland *et al.*, 2010 [DOI](#)). Recently, Tansley S *et al.* (Tansley *et al.*, 2022 [DOI](#)) found that after peripheral nerve injury, microglia will degrade the extracellular matrix structure (neuronal peripheral network, PNN) in lamina I of the spinal dorsal horn, enhance the activity of projection neurons, and induce pain related behaviors. By comparison, Kohno *et al.* (Kohno *et al.*, 2022 [DOI](#)) described a spinal cord-resident pool of CD11c⁺ microglia that emerges during pain maintenance and contributes to the resolution of neuropathic pain in mice. Interesting, previous evidence indicated that direct activation of astrocytes using an optogenetic approach induces microglial reactivity and pain hypersensitivity (Nam *et al.*, 2016 [DOI](#)). Thus, microglia act as diverse roles in a pain development and resolution (Sideris-Lampretsas and Malcangio, 2022 [DOI](#)). Sex differences in the involvement of microglia in mediating chronic pain have been demonstrated (Sorge *et al.*, 2015 [DOI](#); Taves *et al.*, 2016 [DOI](#)). We also explored the effect of sex on neuropathic pain, and observed no significant differences between male and female mice. Thus, there were inconsistent association between sexes and neuropathic pain after SCI (Dominguez *et al.*, 2012 [DOI](#); Mapplebeck *et al.*, 2016 [DOI](#); McFarlane *et al.*, 2020 [DOI](#)). Furthermore, we also observed that the onset of cold/thermal hypersensitivity were delayed than mechanical allodynia, which may be related to the different processing of different types of pain signals.

Our study showed that among the most significant DEGs after selective resident astrocytes elimination were the genes involved in regulating type I IFNs production, and the signal transduction of type I IFNs was significantly upregulated. Molecular experiment results also found the mRNA expression of genes associated with the type I IFNs signal increased after selective astrocytes elimination. It has been verified that type I IFNs signal can significantly reduce sensory neuron activation and pain level (Donnelly *et al.*, 2021 [DOI](#)). Mice with type I IFNs signal deficiency were more sensitive to pain stimuli and had more obvious activation of intense sensory neurons (Donnelly *et al.*, 2021 [DOI](#)). Thus, we suspected that type I IFNs signal might be involved in the association between selective resident astrocytes elimination and neuropathic pain attenuation. Although interferons could derive from various cells, myeloid-lineage cells, microglia are resident immune cells in the central nervous system and are believed to be the main source of IFN-β within the CNS during inflammatory conditions (Owens *et al.*, 2014 [DOI](#)). Reinert and colleagues (Reinert *et al.*, 2016 [DOI](#)) identified microglia as the primary producers of type I IFNs, and the expression of cGAS and Sting1 were enriched in microglia (Chin, 2019 [DOI](#)). The association between selective astrocytes elimination, type I IFNs signal and microglia activation remains unclear. Considering that the death or ablation of numerous cell types in CNS including oligodendrocyte could spark activation of microglia (Gritsch *et al.*, 2014 [DOI](#)), more studies were needed to explore the effect of microglia activation on the association between astrocytes elimination and neuropathic pain attenuation.

There are also some limitations in this work. Although we determined that the selective resident elimination could significantly attenuate neuropathic pain, astrocyte elimination is difficult to achieve in clinical practice. Instead, we provided a strategy to potentially target and activate type I IFNs signal to improve pain after SCI. Additionally, astrocytes elimination may cause many pathophysiological changes in addition to activating type I IFNs signal, the observed effect may be a side effect of the elimination of astrocytes. For our work did not identify whether diphtheria toxin-induced cell apoptosis leads to the activation of the type I IFNs signaling pathway in microglia and is not induced only by astrocyte-specific clearance. Our experiments in cell culture induced by LPS were partly to give a possible mechanist explanation, the optimal methods should be performed in primary microglia obtained from SCI+AAV group or SCI+AAV+DT group and

compare to sham operated one. Finally, we also did not provide the rescue experiment, explore the homeostatic function of astrocytes and the roles of interferon agonist (Poly:IC) and STING agonist (ADU-S100 and DMXAA) during this work.

Conclusion

Above all, we determined that resident astrocytes but not ependymal cells are the main origins of astrocytes participate in neuropathic pain after SCI. The selective resident elimination in lumbar enlargement could significantly attenuate neuropathic pain, and the mechanisms were partly associated with type I IFNs signal activation in microglia (as the schematic diagram shown in **Figure 8**). Therefore, targeted intervention on type I IFNs signal may be an effective strategy for neuropathic pain treatment after SCI.

Materials and methods

Experimental models (organisms/strains)

The experimental procedures were designed to minimize the number of animals used as well as animal suffering. All animal experiments were carried out in accordance with the U.K. Animals (Scientific Procedures) Act, 1986 and associated guidelines, EU Directive 2010/63/EU for animal experiments, and approved by the Institutional Animal Care and Use Committee (IACUC) of the Tongji University School of Medicine (SYXK2019-0005). All transgenic and non-transgenic mice used were derived from in house breeding colonies maintained on C57/BL6 backgrounds. Mice were housed on a 12/12-h light/dark cycle and had food and water available ad libitum. The room temperature and humidity were appropriate. All behavioral testing was performed between 9:00 am and 5:00 pm. Transgenic mice were purchased from the Model Organisms Centre, Shanghai and the Jackson Laboratory. To target spinal cord ependymal cells, C57BL/6-Foxj1^{em1(GFP-CreERT2-polyA)Smoc} mice (Cat# NM-KI-200133, the Model Organisms Centre) were used in this experiment. C57BL/6-Foxj1^{em1(GFP-CreERT2-polyA)Smoc} mice were under the control of promoters with specific expression in ependymal cells (Foxj1), which have been widely used for identification of cells that express Foxj1 in the central nervous system and peripheral organs (Barnabé-Heider *et al.*, 2010; Ren *et al.*, 2017). *Gt(ROSA)26Sor^{em1(CAG-LSL-tdTomato-2A-DTR)Smoc}* mice were obtained from *Gt(ROSA)26Sor^{em1(CAG-LSL-RSR-tdTomato-2A-DTR)Smoc}* mice (Cat# NM-KI-190086, the Model Organisms Centre) crossed with C57BL/6J^{Smoc-Tg(CAG-Dre)Smoc} (Cat# NM-TG-00026, the Model Organisms Centre). *Gt(ROSA)26Sor^{em1(CAG-LSL-tdTomato-2A-DTR)Smoc}* mice carried the reporter gene tdTomato and diphtheria toxin receptor (DTR), which can be used for lineage tracing and selective elimination for specific cell population (Buch *et al.*, 2005; Madisen *et al.*, 2010), including astrocytes (Anderson *et al.*, 2016). GFAP-Cre; *Gt(ROSA)26Sor^{em1(CAG-LSL-tdTomato-2A-DTR)Smoc}* mice were obtained from *Gt(ROSA)26Sor^{em1(CAG-LSL-tdTomato-2A-DTR)Smoc}* mice crossed with FVB-Tg^{(GFAP-cre)25Mes/J} mice (Cat# JAX004600, the Jackson Laboratory). Wild-type C57BL/6J mice were obtained through reproduction. For genotyping, genomic DNA was isolated from ear biopsy for PCR. Transgenic mice and genotyping primers were listed in **Supplementary Table1**. B6.Cg-*Gt(ROSA)26Sor^{tm9(CAG-tdTomato)Hze/J}* mice (Cat# JAX007909, the Jackson Laboratory) were used for the AAV2/5-GfaABC1D-Cre test experiment. A total of 132 *Gt(ROSA)26Sor^{em1(CAG-LSL-tdTomato-2A-DTR)Smoc}* mice aged 2 months and older were randomly and equally divided into the following groups: the sham group (male=15, female=15), sham AAV group (male=15, female=15), SCI+AAV group (male=18, female=18), and SCI+AAV+DT group (male=18, female=18). Three or more independent experiments were performed.




Figure 8.

The potential schematic diagram of selective resident astrocytes elimination attenuated neuropathic pain after SCI.

(A), astrocytes in lumbar enlargement were targeted and selectively eliminated through transgenic mice injected with an adeno-associated virus vector (AAV2/5-GfaABC1D-Cre) and diphtheria toxin. Selective astrocyte elimination in lumbar enlargement could attenuate neuropathic pain after SCI, which were associated with type I IFNs signal and microglia activation. (B), the production of type I IFNs. Type I IFNs production is mainly caused by the contact of innate immune cells (mainly macrophages, microglia and astrocyte in CNS) surface or internal receptors (RIG-I receptor IFI1/MDA5, cGAS, etc.) with virus specific antigenic substances (DNA, RNA), then through intracellular signal molecule transmission (*STING*, *TBK*, *IKK*, etc.), and finally activate the transcription factor IRF3/7 to promote the expression of type I IFNs, including IFN- α and IFN- β . (C), the signal transduction of type I IFNs. Type I IFNs bind to the same two membrane spanning polypeptide chains type I IFNs receptor 1/2 (IFNAR1/2) (Borden *et al.*, 2007; Owens *et al.*, 2014) and lead to cross phosphorylation and activation of Tyk2/Jak1. Activation of Tyk2 and JAK1 phosphorylate STAT1/2 to form a heterodimer, which then translocate to the nucleus and associate with IRF9 to further form the heterotrimeric transcription factor complex IFN-stimulated gene factor-3 (ISGF3). Finally, ISGF3 translocate to the nucleus and bind to specific IFN-response elements (ISREs) to control the expression of IFNs-stimulated genes (ISGs) (Rothhammer *et al.*, 2016), including *ISG15*, *MX1*, *OAS1*, *OAS2*, *IFIT1*, *ZBP1*.

Spinal cord contusion model and motor function test

Thoracic spinal cord contusion, a widely used SCI model for neuropathic pain (Kramer et al., 2017), and the MASCIS Impactor Model III (W.M. Keck Center for Collaborative Neuroscience, Rutgers, the State University of New Jersey, USA) were applied in this study (Duan et al., 2021). Briefly, mice were weighed and deeply anaesthetized with isoflurane evaporated in a gas mixture containing 70% N₂O and 30% O₂ through a nose mask. The back skin was shaved and cleaned with 75% alcohol. A laminectomy at T10 was performed to remove the part of the vertebra overlying the spinal cord, exposing a circle of dura through an operating microscope (Zeiss, Germany) and rodent stereotaxic apparatus (RWD Life Science Co., Ltd, Shenzhen, China). The spinal column was stabilized using lateral clamps over the lateral processes at T9 and T11. Contusion was performed at T10 with a 5 g impactor and 6.25 mm height with a force of about 60kdyn, which could cause a moderate injury as previously reported (Wu et al., 2013; Wu et al., 2014), and then the wound was sutured. Mice had received natural illumination to keep warm before and after the surgery. The following symbols were indicators of a successful contusion model: (1) the impact point was located in the middle of T10, (2) paralysis of both hindlimbs occurred after awakening, (3) the cut-offs of mechanical pain were lower than their baselines. Unsuccessful models were excluded in the following experiment and analysis. Sham mice underwent laminectomy but not contusion. Urine was manually expressed from the bladders of the injured mice twice per day until autonomous urination recovered. Hindlimb motor function was tested in an open field chamber on days 0, 1 and 3 after SCI and weekly for up to 6 weeks. The Basso Mouse Scale (BMS) (Basso et al., 2006) was applied to quantify motor function by two investigators blinded to the genotypes and group assignments.

Measurement of mechanical sensitivity with an electronic von Frey apparatus

The mechanical allodynia of the hindlimbs was measured by the electronic von Frey apparatus (IITC Life Sciences, Woodland Hills, CA) as previously reported (Cunha et al., 2004; Martinov et al., 2013), which is a sensitive, objective and quantitative measurement tool for mechanical sensitivity in rodent models with pain (Cunha et al., 2004; Martinov et al., 2013). The electronic von Frey apparatus has a pressure-meter which consisted of a hand-held force transducer fitted with a 0.5 mm² polypropylene tip (Cunha et al., 2004). When positive response appeared, the intensity of the stimulus was automatically recorded. Our investigators were proficiency trained in advance to use the polypropylene tip vertically to the central of plantar. Briefly, mice were placed on a wire mesh platform in individual Plexiglas cubicles and allowed to acclimate to the Plexiglas restraint and testing room for 30 minutes. Before testing, the mice were quiet without exploratory behaviors or defecation, and not resting on their paws (Cunha et al., 2004). The average interstimulus interval was 2 minutes. Polypropylene tip was applied a vertical and gradual increase force to the plantar surface of the left and right hindlimbs for 3–4 seconds, or until a positive response appeared. Behaviours that were considered positive responses to the filament included brisk paw withdrawal, flinching, hunching of the back, licking of the stimulated area, and escape responses (Wu et al., 2016). All animals underwent 2 trials (5 measurements/trial) at 24 and 48 hours before contusion to calculate the average plantar paw withdrawal threshold of both hindlimbs at baseline. The measurement of post-contusion withdrawal thresholds was performed in a blinded fashion by two independent investigators at 2, 3, 4, 5, and 6 weeks after contusion. These timepoints were chosen for the nociceptive assays to calculate BMS scores was to avoid the issue of motor impairment for reflex-based assays, and these mice could maintain a standing posture from 2 weeks after contusion on.

Analgesia test for thresholds of cold and thermal pain

Cold and hot plate (Cat# YSL-21; Shanghai Yuyan Instruments Co., Ltd. China) was used to assess cold and thermal pain as previously reported (Carozzi *et al.*, 2013 [DOI](#); Wu *et al.*, 2016 [DOI](#)). Cold and thermal pain tests of the hindlimbs were conducted one day after mechanical allodynia measurement. Before testing, mice were also transferred into the testing room to acclimate to the environment for 30 minutes. The temperatures of 0°C (cold stimulus) and 50°C (hot stimulus) were used to measure the thresholds for noxious heat and cold, respectively (Wu *et al.*, 2016 [DOI](#)). To avoid tissue injury, temperatures of 0°C (cold stimulus) for 5 minutes and 50°C (hot stimulus) for 30 seconds were set as the cut-off points. The positive response included licking, jumping or hind paw withdrawal. The paw withdrawal latency (for hot stimuli) and paw withdrawal frequency (for cold stimuli) of positive responses were recorded. For hot stimuli, the paw withdrawal latency means the period from touching the hot plate to the emergence of positive response. For cold stimuli, the paw withdrawal frequency means the number of nociceptive responses observed in 5 minutes on cold plate. The tests were carried out 24 hours before contusion and 2, 3, 4, 5, and 6 weeks after contusion. The analgesia test for thresholds of cold and thermal pain were conducted by two investigators blinded to the groups and experimental conditions.

AAV2/5-GfaABC1D-Cre and DT injection

In this study, AAV2/5 with the GfaABC1D promoter (AAV2/5-GfaABC1D-Cre) was used to target Cre recombinase expression exclusively in astrocytes (Anderson *et al.*, 2016 [DOI](#)). The 681 bp GFAP promoter, GfaABC1D, was derived from the 2.2kb human GFAP promoter gfa2. Importantly, the GfaABC1D promoter is about twice as active as gfa2 with smaller size, which is better suited to viral vectors (Lee *et al.*, 2008 [DOI](#); Yu *et al.*, 2020 [DOI](#)). The expression of cargo genes under the GfaABC1D promoter in astrocytes is highly efficient and specific (Yu *et al.*, 2020 [DOI](#)). Previous study reported that AAV2/5 predominantly labeled astrocytes, displaying robust co-localization of reporter gene (eGFP) with the astrocytic marker, GFAP (Ortinski *et al.*, 2010 [DOI](#)). The AAV2/5 vector with a minimal GFAP promoter (AAV2/5-GfaABC1D-Cre) has been regarded as the most reliable viral approach and widely used to target astrocytes (Yu *et al.*, 2020 [DOI](#)). AAV2/5-GfaABC1D-Cre was purchased from Shanghai Taitool Bioscience Co., Ltd. (Cat# S0611-5, $\geq 1 \times 10^{13}$ V.G/ml). AAV2/5-GfaABC1D-Cre was injected into *Gt(ROSA)26Sor^{em1}(CAG-LSL--tdTomato-2A-DTR)Smoc* mice of SCI+AAV group and SCI+AAV+DT group 2 weeks before T10 contusion, and into mice of sham AAV group 2 weeks before T10 laminectomy. Sham group mice underwent L4-5 laminectomy but not AAV. Briefly, after precise location of L4-5 spinal cord as previously reported (Rigaud *et al.*, 2008 [DOI](#)), the back skin was shaved and cleaned with 75% alcohol. A laminectomy was performed at L4-5 as described above. AAV2/5-GfaABC1D-Cre was injected into the both sides of dorsal median of the lumbar enlargement (0.5 mm lateral to midline with a depth of 0.6 mm, 0.4 μ L per side, $\geq 1 \times 10^{11}$ V.G/ml in sterile saline), with a Hamilton 33-gauge (33 G) microinjection needle at a speed of 0.2 μ L/min. After injection, the needle was held for 5 minutes to avoid liquid reflux. Overall, no obvious tissue injury of spinal cord, no motor and sensory function impairment of hindlimbs were observed during the following two weeks. Spinal cord tissue fluorescence images were taken and handled with a Fluorescence-labeled Organism Bioimaging Instrument (FOBI, Neoscience Ltd., Suwon City, Rep. of Korea) after AAV2/5-GfaABC1D-Cre injection.

Diphtheria toxin (DT) injection

Mice received 500 ng diphtheria toxin (DT, Cat# 150, List Biological Laboratory) in 100 μ L distilled, deionized water (ddH₂O) through intraperitoneal injection (i.p.) (Binnewies *et al.*, 2019 [DOI](#)) on 31, 32 and 34 dpi (days post-injury), when the pain level peaked (the lowest paw withdrawal threshold of mechanical sensitivity). Then, the mice were sacrificed at 1 week after DT injection for RNA sequencing, bioinformatics analysis, immunohistochemistry and molecular experiment.

RNA sequencing (RNA-seq) and bioinformatics analysis

To further evaluate the role of selective astrocytes elimination in gene expression regulation and pain attenuation mechanisms, RNA-seq was used to examine the transcriptomes of lumbar enlargement tissues in sham group (n = 3), sham AAV group (n = 3), SCI+AAV group (n = 3), and SCI+AAV+DT group (n = 3). Previous findings revealed the sex-dependent difference in neuropathic pain after SCI (Fiore et al., 2022 [\[4\]](#); Stewart et al., 2021 [\[4\]](#)). Thus, the lumbar enlargement tissues of female mice were selected for RNA-seq experiment. Briefly, mice were euthanized after terminal anaesthesia by pentobarbital overdose. Spinal cord tissue was dissected from lumbar enlargements (L4-5) at 1 week after selective astrocytes elimination, with a dissecting microscope, and 2 mm segments were harvested and marked. A total of 12 samples were sent to the Beijing Genomics Institute (BGI) Company (Shenzhen, China) in solid carbon dioxide for further RNA-seq analysis, and sequencing was performed on a DNBSEQ platform at BGI Company. Bioinformatics analysis, including Gene Ontology (GO) enrichment, Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway, gene set enrichment analysis (GSEA) and protein-protein interaction (PPI) network analysis, was carried out with the online platform Dr. Tom (BGI Company). Only genes with transcripts per million (TPM) > 1 were analyzed. Differentially expressed genes (DEGs) were identified using the DEGseq method and screened with the criteria of q value ≤ 0.05 and log₂FC ≥ 0.6, where FC represents the fold change. False discovery rate (FDR) method was used to adjust for multiple comparisons to generate q-values in RNA-seq study. The normalized enrichment score (NES) was utilized to select the most activated and inactivated GO terms and pathways. The sequence and sample data have been deposited in the NCBI database under Sequence Read Archive (SRA) with Bioproject identification number PRJNA847704 (Accession number: SRR19611667 - SRR19611678). The single-cell RNA sequencing (scRNA-seq) results were analyzed through online data from injured mouse spinal cords (https://jaeleelab.shinyapps.io/sci_singlecell/ [\[4\]](#)) (Milich et al., 2021 [\[4\]](#)).

Immunofluorescence staining and analysis

Immunofluorescence staining procedures were conducted as described previously (Ren et al., 2017 [\[4\]](#)). After post-fixation and cryoprotection, a dissected 6 mm segment of spinal cord centred around the injury epicentre, or a 2 mm segment of the lumbar enlargement (L4-5), was coronally sectioned at 12 µm thickness and thaw-mounted onto Superfrost Plus slides (Citotest Labware Manufacturing Co., Ltd.). The primary antibodies used were as follows: glial fibrillary acidic protein (GFAP, Cat# ab4674, Abcam, 1:500), Neuronal nuclear antigen (NEUN, Cat# ab190565, Abcam, 1:500), Tubulin beta III (Tuj1, Cat# MAB1637, Millipore, 1:500), Ionized calcium binding adaptor molecule 1 (IBA1, Cat# 016-20001, Wako, 1:500), Oligodendrocyte Transcription Factor 2 (Olig2, Cat# AF2418, R&D system, 1:500), S100 Calcium Binding Protein A10 (S100A10, Cat#11250-1-AP, Protein Tech, 1:250), Interferon-stimulated Gene 15 (ISG15, Cat# sc-166755, Santa Cruz, 1:50), Interferon regulatory factor 7 (IRF7, Cat# sc-74471, Santa Cruz, 1:50), Signal transducer and activator of transcription 1 (STAT1, Cat# ab109461, Abcam, 1:150). The secondary antibodies were Alexa® Fluor 488 (Cat# abs20019A, Absin, 1:500), Alexa® Fluor 488 (Cat# A-32814, Invitrogen, 1:500), Alexa® Fluor 488 (Cat# A-32766, Invitrogen, 1:500), Alexa® Fluor 555 (Cat# A-32773, Invitrogen, 1:500), Alexa® Fluor 594 (Cat# ab150176, Abcam, 1:500), and Alexa® Fluor 647 (Cat# A-32795, Invitrogen, 1:500). The nuclei were stained with 4',6-diamidino-2-phenylindole (DAPI, Cat# C1002, Beyotime Institute of Biotechnology), and fluorescence images were taken by confocal microscopy (LSM 700, Carl Zeiss, Jena, Germany). Image acquisition was performed with ZEN 2.3 (blue edition, Carl Zeiss), and micrographs were assembled using Adobe Illustrator CC 2018. As previously reported (Watson, 2012 [\[4\]](#)), we selected four regions in the lumbar enlargement to analyze in the immunohistochemistry images, including three largest ascending tracts and SDH. The three largest ascending tracts were (1) the gracile and cuneate fasciculi (GR and CU, respectively); (2) the lateral, dorsolateral and ventral spinothalamic tracts (LST, DLST, and VST, respectively); and (3) the dorsal and ventral spinocerebellar tracts (DSC and VSC, respectively). These tracts are closely associated with proprioception and sensation (Watson, 2012 [\[4\]](#)). For

immunofluorescence analysis of lumbar enlargement, three sections were randomly selected to delineate the three tracts and SDH (3 areas per tract/SDH), then the 160 μm $\times$ 160 μm images were exported as single channel images through ZEN 2.3 (blue edition). Lesion area covered the lesion core, rostral and caudal injured spinal cords with a distance of 500 μm from lesion core, respectively. A total of three sections (8 areas of rostral and 8 of caudal injured spinal cords per section) were randomly selected per mice. Then the 160 μm $\times$ 160 μm images were exported by ZEN 2.3 (blue edition). Image-J software with customized macros was used to quantify the fluorescence intensity of proteins including the GFAP, Tuj1, IBA1, ISG15, IRF7 and STAT1 in the images mentioned above. For sham group and treatment groups, three mice per group were tested.

Microglial culture

To observe the production of type I interferon related factors by microglia after activation, we used activated microglia induced by lipopolysaccharides (LPS, Cat# Escherichia coli O111:B4, Sigma-Aldrich) in this study. Furthermore, different concentrations of interferon agonist (Ploy:IC (Cat# B5551, APEX BIO)) (Khoroshi *et al.*, 2015 [\[1\]](#)) and Stimulator of interferon genes (STING) agonist (ADU-S100 (Cat# HY-12885B, MCE) and DMXAA (Cat# HY-10964, MCE)) (Li *et al.*, 2023 [\[2\]](#)) were used to stimulate microglia for 12 and 24 hours. Mouse microglial SIM-A9 cells were purchased from ATCC (Cat# CRL-3265, ATCC) and cultured according to ATCC recommended growth conditions in DMEM-F12 (Cat# 11330057, Gibco) containing 10% heat-inactivated FBS (Cat# 16140-071, Gibco) and 5% heat-inactivated horse serum (Cat# 26050-088, Gibco) (Correia *et al.*, 2021 [\[3\]](#)). LPS at concentrations of 0.2 $\mu\text{g}/\text{mL}$ LPS (Jayakumar *et al.*, 2022 [\[4\]](#)) was used to stimulate SIM-A9 cells after they reached 60-70% confluency for 24 h. Microglia and culture medium were collected and centrifuged at 12,000 $\times$ g for 5 min to remove debris. The expression of pro-inflammatory and anti-inflammatory microglial marker genes were measured by RT-PCR. After the stimulation by Ploy:IC (2.5 μg , 12.5 μg , 25 μg , 50 μg , 100 μg , 200 μg), ADU-S100 (0.1 μg , 1 μg , 10 μg , 20 μg) and DMXAA (1 μg , 5 μg , 10 μg , 20 μg) for 12 and 24 hours, the microglia were also obtained for RT-PCR experiment. Control wells were also established, and the data were analyzed and compared with those of the control wells.

Quantitative real-time PCR (RT-PCR)

The total RNA of cells was isolated with RNAiso plus (Cat# 9108, Takara). The concentration and purity of RNA samples were measured using a Nanodrop ND-2000 (Thermo Science, MA, USA) for further experiments. Five hundred nanograms of RNA was converted to complementary DNA (cDNA), which was synthesized with a PrimeScript reverse transcriptase kit (Cat# RR037A, Takara). RT-PCR was performed using a TB Green TM Premix Ex Taq Kit (Cat# RR820A, Takara) on a Light Cycler Real-Time PCR System (480II, Roche). The primer sequences (Shanghai Generay Biotech Co., Ltd.) were designed through PrimerBank (<https://pga.mgh.harvard.edu/primerbank/> [\[5\]](#)) and are listed in **Tables S2-4**. The relative amounts of mRNA were calculated using the $\Delta\Delta\text{Ct}$ relative quantification method. GAPDH served as the control gene, and the mRNA levels of specific genes were normalized to GAPDH. Calculations and statistics were performed in Microsoft Excel version 16.36. Graphs were plotted in GraphPad Prism 8 version 8.4.3.

Statistical analysis

All experiments were conducted with three or more duplicates. All continuous data were shown as mean $\pm$ SEM. One-way ANOVA was performed followed by Student Newman-Keuls post hoc test for continuous data. P values < 0.05 were considered statistically significant. Data analyses were conducted using the Statistical Analysis System (SAS), version 9.4 (SAS Institute, Inc, Cary, NC, USA). Plots were generated using GraphPad Prism 8 software (GraphPad Software, San Diego, CA).

Acknowledgements

This work was supported by the International Cooperation Project of National Natural Science Foundation of China (Grant No. 81810001048), the National Natural Science Foundation of China (Grant Nos. 81974190, 82271419, 82225027, 81901902 and 81701217), the National Key R&D Program of China (Grant No. 2020YFC2008703), Shanghai Rising-Star Program (Grant No. 22QA1408200), the Fundamental Research Funds for the Central Universities (22120220555 and 22120230138), Programs in Emergency and Critical Care Medicine, Shanghai Municipal Health Commission (KPB1702), National Key Specialty Program, National Health Commission of China (GJ2301). We thank Gufa Lin, Bei Ma, Zhourui Wu and Xu Xu of our laboratory for help and advice. We also thank Pianpian Fan (Department of Pediatrics, West China second hospital, Sichuan University) for article grammar check and suggestions.

Author Contributions

Q.Z., Y.J.Z. and Y.L.R. contributed equally to this work. Q.Z., Y.J.Z. and Y.L.R. designed and performed experiments, analyzed the data and drafted the manuscript. L.J.Z, J.W.Z, S.Y. and H.F.N. participated in the establishment of spinal cord injury model and pain test. N.X., L.M.C., and R.R.Z. conceived and designed the experiments, supervised the overall project and revised the manuscript. All authors have read and approved the article. The authors also declare no conflict of interest.

Ethics

Animals (Scientific Procedures) Act, 1986 and associated guidelines, EU Directive 2010/63/EU for animal experiments, and approved by the Institutional Animal Care and Use Committee (IACUC) of the Tongji University School of Medicine (SYXK2019-0005).

Additional files

Supplementary files

Supplementary file 1.

Supplementary Figure 1. S100 family protein p11 (S100A10) expressed by various cell types.

Supplementary Figure 2. Astrocytes activation in the lumbar enlargement of neuropathic pain mice. The results of AAV2/5-GfaABC1D-Cre targeted astrocytes in lumbar enlargement.

Supplementary Figure 3. No differences in neuropathic pain were observed between male and female mice.

Supplementary Figure 4. Pain symptoms were similar in wild-type mice with or without DT injection; full fluorescence images of lesion area; RNA-seq analysis results between SCI+AAV group and sham AAV group.

Supplementary Figure 5. GFAP-Cre; *Gt(ROSA)26Sor^{em1}(CAG-LSL-tdTomato-2A-DTR)Smoc* mice were not suitable for labelling astrocytes.

Supplementary Table 1. Transgenic mice and genotyping primers.

Supplementary Table 2. Primers for type I interferon signal pathway genes.

Supplementary Table 3. Primers for pro- and anti-inflammatory microglial marker genes.

Supplementary Table 4. Primers for Type I IFN signal related genes.

Data availability

The sequence and sample data have been deposited in the NCBI database under Sequence Read Archive (SRA) with Bioproject identification number PRJNA847704 (Accession number: SRR19611667 - SRR19611678). Any information required to utilize and reanalyze the data reported in this paper is available from the corresponding author upon request (nxieprof18@tongji.edu.cn).

References

- Allnoch L., Baumgärtner W., Hansmann F (2019) **Impact of Astrocyte Depletion upon Inflammation and Demyelination in a Murine Animal Model of Multiple Sclerosis** *Int J Mol Sci* **20** <https://doi.org/10.3390/ijms20163922>
- Anderson M.A., Burda J.E., Ren Y., Ao Y., O'Shea T.M., Kawaguchi R., Coppola G., Khakh B.S., Deming T.J., Sofroniew M.V (2016) **Astrocyte scar formation aids central nervous system axon regeneration** *Nature* **532**:195–200 <https://doi.org/10.1038/nature17623>
- Barnabé-Heider F., Göritz C., Sabelström H., Takebayashi H., Pfriederger F.W., Meletis K., Frisén J (2010) **Origin of new glial cells in intact and injured adult spinal cord** *Cell Stem Cell* **7**:470–482 <https://doi.org/10.1016/j.stem.2010.07.014>
- Barrat F.J., Crow M.K., Ivashkiv L.B (2019) **Interferon target-gene expression and epigenomic signatures in health and disease** *Nat Immunol* **20**:1574–1583 <https://doi.org/10.1038/s41590-019-0466-2>
- Basso D.M., Fisher L.C., Anderson A.J., Jakeman L.B., McTigue D.M., Popovich P.G (2006) **Basso Mouse Scale for locomotion detects differences in recovery after spinal cord injury in five common mouse strains** *J Neurotrauma* **23**:635–659 <https://doi.org/10.1089/neu.2006.23.635>
- Binnewies M. *et al.* (2019) **Unleashing Type-2 Dendritic Cells to Drive Protective Antitumor CD4(+) T Cell Immunity** *Cell* **177**:556–571 <https://doi.org/10.1016/j.cell.2019.02.005>
- Borden E.C., Sen G.C., Uze G., Silverman R.H., Ransohoff R.M., Foster G.R., Stark G.R (2007) **Interferons at age 50: past, current and future impact on biomedicine** *Nat Rev Drug Discov* **6**:975–990 <https://doi.org/10.1038/nrd2422>
- Buch T., Heppner F.L., Tertilt C., Heinen T.J., Kremer M., Wunderlich F.T., Jung S., Waisman A (2005) **A Cre-inducible diphtheria toxin receptor mediates cell lineage ablation after toxin administration** *Nat Methods* **2**:419–426 <https://doi.org/10.1038/nmeth762>

- Carozzi V.A. *et al.* (2013) **Bortezomib-induced painful peripheral neuropathy: an electrophysiological, behavioral, morphological and mechanistic study in the mouse** *PLoS One* **8** <https://doi.org/10.1371/journal.pone.0072995>
- Chen G., Zhang Y.Q., Qadri Y.J., Serhan C.N., Ji R.R. (2018) **Microglia in Pain: Detrimental and Protective Roles in Pathogenesis and Resolution of Pain** *Neuron* **100**:1292–1311 <https://doi.org/10.1016/j.neuron.2018.11.009>
- Chin A.C. (2019) **Neuroinflammation and the cGAS-STING pathway** *Journal of neurophysiology* **121**:1087–1091 <https://doi.org/10.1152/jn.00848.2018>
- Church J.S., Kigerl K.A., Lerch J.K., Popovich P.G., McTigue D.M. (2016) **TLR4 Deficiency Impairs Oligodendrocyte Formation in the Injured Spinal Cord** *The Journal of neuroscience : the official journal of the Society for Neuroscience* **36**:6352–6364 <https://doi.org/10.1523/JNEUROSCI.0353-16.2016>
- Correia S.S. *et al.* (2021) **The CNS-penetrant soluble guanylate cyclase stimulator CYR119 attenuates markers of inflammation in the central nervous system** *J Neuroinflammation* **18** <https://doi.org/10.1186/s12974-021-02275-z>
- Cunha T.M., Verri W.A., Vivancos G.G., Moreira I.F., Reis S., Parada C.A., Cunha F.Q., Ferreira S.H. (2004) **An electronic pressure-meter nociception paw test for mice** *Brazilian journal of medical and biological research = Revista brasileira de pesquisas medicas e biologicas* **37**:401–407 <https://doi.org/10.1590/s0100-879x2004000300018>
- Dominguez C.A., Ström M., Gao T., Zhang L., Olsson T., Wiesenfeld-Hallin Z., Xu X.J., Piehl F. (2012) **Genetic and sex influence on neuropathic pain-like behaviour after spinal cord injury in the rat** *European journal of pain (London, England)* **16**:1368–1377 <https://doi.org/10.1002/j.1532-2149.2012.00144.x>
- Donnelly C.R. *et al.* (2021) **STING controls nociception via type I interferon signalling in sensory neurons** *Nature* **591**:275–280 <https://doi.org/10.1038/s41586-020-03151-1>
- Dozio V., Sanchez J.C. (2018) **Profiling the proteomic inflammatory state of human astrocytes using DIA mass spectrometry** *J Neuroinflammation* **15** <https://doi.org/10.1186/s12974-018-1371-6>
- Duan H., Pang Y., Zhao C., Zhou T., Sun C., Hou M., Ning G., Feng S. (2021) **A novel, minimally invasive technique to establish the animal model of spinal cord injury** *Ann Transl Med* **9**:881–881 <https://doi.org/10.21037/atm-21-2063>
- Dworkin R.H. *et al.* (2003) **Advances in neuropathic pain: diagnosis, mechanisms, and treatment recommendations** *Archives of neurology* **60**:1524–1534 <https://doi.org/10.1001/archneur.60.11.1524>
- Fiore N.T., Yin Z., Guneykaya D., Gauthier C.D., Hayes J.P., D'Hary A., Butovsky O., Moalem-Taylor G. (2022) **Sex-specific transcriptome of spinal microglia in neuropathic pain due to peripheral nerve injury** *Glia* **70**:675–696 <https://doi.org/10.1002/glia.24133>
- Gritsch S. *et al.* (2014) **Oligodendrocyte ablation triggers central pain independently of innate or adaptive immune responses in mice** *Nat Commun* **5** <https://doi.org/10.1038/ncomms6472>

Gwak Y.S., Kang J., Unabia G.C., Hulsebosch C.E (2012) **Spatial and temporal activation of spinal glial cells: role of gliopathy in central neuropathic pain following spinal cord injury in rats** *Exp Neurol* **234**:362–372 <https://doi.org/10.1016/j.expneurol.2011.10.010>

Hackett A.R., Lee J.K (2016) **Understanding the NG2 Glial Scar after Spinal Cord Injury** *Front Neurol* **7** <https://doi.org/10.3389/fneur.2016.00199>

Henwood P., Ellis J.A (2004) **Chronic neuropathic pain in spinal cord injury: the patient's perspective** *Pain Res Manag* **9**:39–45 <https://doi.org/10.1155/2004/863062>

Hesp Z.C., Goldstein E.Z., Miranda C.J., Kaspar B.K., McTigue D.M (2015) **Chronic oligodendrogenesis and remyelination after spinal cord injury in mice and rats** *The Journal of neuroscience : the official journal of the Society for Neuroscience* **35**:1274–1290 <https://doi.org/10.1523/JNEUROSCI.2568-14.2015>

Honjoh K., Nakajima H., Hirai T., Watanabe S., Matsumine A (2019) **Relationship of Inflammatory Cytokines From M1-Type Microglia/Macrophages at the Injured Site and Lumbar Enlargement With Neuropathic Pain After Spinal Cord Injury in the CCL21 Knockout (plt) Mouse** *Front Cell Neurosci* **13** <https://doi.org/10.3389/fncel.2019.00525>

Huang W., Bai X., Meyer E., Scheller A (2020) **Acute brain injuries trigger microglia as an additional source of the proteoglycan NG2** *Acta Neuropathol Commun* **8** <https://doi.org/10.1186/s40478-020-01016-2>

Ivashkiv L.B., Donlin L.T (2014) **Regulation of type I interferon responses** *Nat Rev Immunol* **14**:36–49 <https://doi.org/10.1038/nri3581>

Jayakumar P., Martínez-Moreno C.G., Lorenson M.Y., Walker A.M., Morales T (2022) **Prolactin Attenuates Neuroinflammation in LPS-Activated SIM-A9 Microglial Cells by Inhibiting NF- κ B Pathways Via ERK1/2** *Cell Mol Neurobiol* **42**:2171–2186 <https://doi.org/10.1007/s10571-021-01087-2>

Ji R.R., Berta T., Nedergaard M (2013) **Glia and pain: is chronic pain a gliopathy?** *Pain* **154**:S10–S28 <https://doi.org/10.1016/j.pain.2013.06.022>

Ji R.R., Donnelly C.R., Nedergaard M (2019) **Astrocytes in chronic pain and itch** *Nat Rev Neurosci* **20**:667–685 <https://doi.org/10.1038/s41583-019-0218-1>

Kawasaki Y., Zhang L., Cheng J.K., Ji R.R (2008) **Cytokine mechanisms of central sensitization: distinct and overlapping role of interleukin-1 β , interleukin-6, and tumor necrosis factor- α in regulating synaptic and neuronal activity in the superficial spinal cord** *J Neurosci* **28**:5189–5194 <https://doi.org/10.1523/jneurosci.3338-07.2008>

Khorooshi R., Mørch M.T., Holm T.H., Berg C.T., Dieu R.T., Dræby D., Issazadeh-Navikas S., Weiss S., Lienenklaus S., Owens T (2015) **Induction of endogenous Type I interferon within the central nervous system plays a protective role in experimental autoimmune encephalomyelitis** *Acta Neuropathol* **130**:107–118 <https://doi.org/10.1007/s00401-015-1418-z>

Kohno K. *et al.* (2022) **A spinal microglia population involved in remitting and relapsing neuropathic pain** *Science* **376**:86–90 <https://doi.org/10.1126/science.abf6805>

Kohro Y. *et al.* (2020) **Spinal astrocytes in superficial laminae gate brainstem descending control of mechanosensory hypersensitivity** *Nature Neuroscience* <https://doi.org/10.1038/s41593-020-00713-4>

Kramer J.L., Minhas N.K., Jutzeler C.R., Erskine E.L., Liu L.J., Ramer M.S (2017) **Neuropathic pain following traumatic spinal cord injury: Models, measurement, and mechanisms** *J Neurosci Res* **95**:1295–1306 <https://doi.org/10.1002/jnr.23881>

Lee Y., Messing A., Su M., Brenner M (2008) **GFAP promoter elements required for region-specific and astrocyte-specific expression** *Glia* **56**:481–493 <https://doi.org/10.1002/glia.20622>

Li N., Wang C., Zhao Y., Wang Y., Gao T., Yu Y., Wang G., Zhang L (2023) **STING controls opioid-induced itch and chronic itch via spinal tank-binding kinase 1-dependent type I interferon response in mice** *J Neuroinflammation* **20** <https://doi.org/10.1186/s12974-023-02783-0>

Liddel S.A. *et al.* (2017) **Neurotoxic reactive astrocytes are induced by activated microglia** *Nature* **541**:481–487 <https://doi.org/10.1038/nature21029>

Madisen L. *et al.* (2010) **A robust and high-throughput Cre reporting and characterization system for the whole mouse brain** *Nat Neurosci* **13**:133–140 <https://doi.org/10.1038/nn.2467>

Mapplebeck J.C.S., Beggs S., Salter M.W (2016) **Sex differences in pain: a tale of two immune cells** *Pain* **157**:S2–s6 <https://doi.org/10.1097/j.pain.0000000000000389>

Martinov T., Mack M., Sykes A., Chatterjea D (2013) **Measuring changes in tactile sensitivity in the hind paw of mice using an electronic von Frey apparatus** *J Vis Exp* <https://doi.org/10.3791/51212>

Masri R., Keller A (2012) **Chronic pain following spinal cord injury** *Adv Exp Med Biol* **760**:74–88 https://doi.org/10.1007/978-1-4614-4090-1_5

McFarlane K., Otto T.E., Bailey W.M., Veldhorst A.K., Donahue R.R., Taylor B.K., Gensel J.C (2020) **Effect of Sex on Motor Function, Lesion Size, and Neuropathic Pain after Contusion Spinal Cord Injury in Mice** *J Neurotrauma* **37**:1983–1990 <https://doi.org/10.1089/neu.2019.6931>

Meletis K., Barnabe-Heider F., Carlen M., Evergren E., Tomilin N., Shupliakov O., Frisen J (2008) **Spinal cord injury reveals multilineage differentiation of ependymal cells** *PLoS Biol* **6** <https://doi.org/10.1371/journal.pbio.0060182>

Milich L.M., Choi J.S., Ryan C., Cerqueira S.R., Benavides S., Yahn S.L., Tsoulfas P., Lee J.K (2021) **Single-cell analysis of the cellular heterogeneity and interactions in the injured mouse spinal cord** *Journal of Experimental Medicine* **218** <https://doi.org/10.1084/jem.20210040>

Nam Y. *et al.* (2016) **Reversible Induction of Pain Hypersensitivity following Optogenetic Stimulation of Spinal Astrocytes** *Cell Rep* **17**:3049–3061 <https://doi.org/10.1016/j.celrep.2016.11.043>

Nishiyama A., Yang Z., Butt A (2005) **Astrocytes and NG2-glia: what's in a name?** *Journal of anatomy* **207**:687–693 <https://doi.org/10.1111/j.1469-7580.2005.00489.x>

Ono T., Kohro Y., Kohno K., Tozaki-Saitoh H., Nakashima Y., Tsuda M (2020) **Mechanical pain of the lower extremity after compression of the upper spinal cord involves signal transducer and activator of transcription 3-dependent reactive astrocytes and interleukin-6** *Brain Behav Immun* **89**:389–399 <https://doi.org/10.1016/j.bbi.2020.07.025>

- Ortinski P.I., Dong J., Mungenast A., Yue C., Takano H., Watson D.J., Haydon P.G., Coulter D.A (2010) **Selective induction of astrocytic gliosis generates deficits in neuronal inhibition** *Nat Neurosci* **13**:584–591 <https://doi.org/10.1038/nn.2535>
- Owens T., Khoroshchi R., Wlodarczyk A., Asgari N (2014) **Interferons in the central nervous system: a few instruments play many tunes** *Glia* **62**:339–355 <https://doi.org/10.1002/glia.22608>
- Reinert L.S. *et al.* (2016) **Sensing of HSV-1 by the cGAS-STING pathway in microglia orchestrates antiviral defence in the CNS** *Nat Commun* **7** <https://doi.org/10.1038/ncomms13348>
- Ren Y. *et al.* (2017) **Ependymal cell contribution to scar formation after spinal cord injury is minimal, local and dependent on direct ependymal injury** *Sci Rep* **7** <https://doi.org/10.1038/srep41122>
- Rigaud M., Gemes G., Barabas M.E., Chernoff D.I., Abram S.E., Stucky C.L., Hogan Q.H (2008) **Species and strain differences in rodent sciatic nerve anatomy: implications for studies of neuropathic pain** *Pain* **136**:188–201 <https://doi.org/10.1016/j.pain.2008.01.016>
- Rothhammer V. *et al.* (2016) **Type I interferons and microbial metabolites of tryptophan modulate astrocyte activity and central nervous system inflammation via the aryl hydrocarbon receptor** *Nat Med* **22**:586–597 <https://doi.org/10.1038/nm.4106>
- Sabelstrom H., Stenudd M., Frisen J (2014) **Neural stem cells in the adult spinal cord** *Exp Neurol* **260**:44–49 <https://doi.org/10.1016/j.expneurol.2013.01.026>
- Salter M.W., Stevens B (2017) **Microglia emerge as central players in brain disease** *Nat Med* **23**:1018–1027 <https://doi.org/10.1038/nm.4397>
- Schreiner B. *et al.* (2015) **Astrocyte Depletion Impairs Redox Homeostasis and Triggers Neuronal Loss in the Adult CNS** *Cell Reports* **12**:1377–1384 <https://doi.org/10.1016/j.celrep.2015.07.051>
- Shiao R., Lee-Kubli C.A (2018) **Neuropathic Pain After Spinal Cord Injury: Challenges and Research Perspectives** *Neurotherapeutics : the journal of the American Society for Experimental NeuroTherapeutics* **15**:635–653 <https://doi.org/10.1007/s13311-018-0633-4>
- Sideris-Lampretsas G., Malcangio M (2022) **Pain-resolving microglia** *Science* **376**:33–34 <https://doi.org/10.1126/science.abo5592>
- Sofroniew M.V (2009) **Molecular dissection of reactive astrogliosis and glial scar formation** *Trends in neurosciences* **32**:638–647 <https://doi.org/10.1016/j.tins.2009.08.002>
- Sorge R.E. *et al.* (2015) **Different immune cells mediate mechanical pain hypersensitivity in male and female mice** *Nat Neurosci* **18**:1081–1083 <https://doi.org/10.1038/nn.4053>
- Srinivasan R., Lu T.Y., Chai H., Xu J., Huang B.S., Golshani P., Coppola G., Khakh B.S (2016) **New Transgenic Mouse Lines for Selectively Targeting Astrocytes and Studying Calcium Signals in Astrocyte Processes In Situ and In Vivo** *Neuron* **92**:1181–1195 <https://doi.org/10.1016/j.neuron.2016.11.030>

- Staniland A.A., Clark A.K., Wodarski R., Sasso O., Maione F., D'Acquisto F., Malcangio M (2010) **Reduced inflammatory and neuropathic pain and decreased spinal microglial response in fractalkine receptor (CX3CR1) knockout mice** *J Neurochem* **114**:1143–1157 <https://doi.org/10.1111/j.1471-4159.2010.06837.x>
- Stewart A.N., Lowe J.L., Glaser E.P., Mott C.A., Shahidehpour R.K., McFarlane K.E., Bailey W.M., Zhang B., Gensel J.C (2021) **Acute inflammatory profiles differ with sex and age after spinal cord injury** *J Neuroinflammation* **18** <https://doi.org/10.1186/s12974-021-02161-8>
- Tansley S. *et al.* (2022) **Microglia-mediated degradation of perineuronal nets promotes pain** *Science* **377**:80–86 <https://doi.org/10.1126/science.abl6773>
- Taves S., Berta T., Liu D.L., Gan S., Chen G., Kim Y.H., Van de Ven T., Laufer S., Ji R.R. (2016) **Spinal inhibition of p38 MAP kinase reduces inflammatory and neuropathic pain in male but not female mice: Sex-dependent microglial signaling in the spinal cord** *Brain Behav Immun* **55**:70–81 <https://doi.org/10.1016/j.bbi.2015.10.006>
- Tsai H.H. *et al.* (2012) **Regional astrocyte allocation regulates CNS synaptogenesis and repair** *Science* **337**:358–362 <https://doi.org/10.1126/science.1222381>
- Tsuda M. *et al.* (2011) **JAK-STAT3 pathway regulates spinal astrocyte proliferation and neuropathic pain maintenance in rats** *Brain* **134**:1127–1139 <https://doi.org/10.1093/brain/awr025>
- Tsuda M., Kuboyama K., Inoue T., Nagata K., Tozaki-Saitoh H., Inoue K. (2009) **Behavioral phenotypes of mice lacking purinergic P2X4 receptors in acute and chronic pain assays** *Mol Pain* **5** <https://doi.org/10.1186/1744-8069-5-28>
- Vierck C (2020) **Mechanisms of Below-Level Pain Following Spinal Cord Injury (SCI)** *J Pain* **21**:262–280 <https://doi.org/10.1016/j.jpain.2019.08.007>
- Watson C (2012) **The Somatosensory System** *In The Mouse Nervous System* :563–570 <https://doi.org/10.1016/b978-0-12-369497-3.10021-4>
- Wu J., Renn C.L., Faden A.I., Dorsey S.G (2013) **TrkB.T1 contributes to neuropathic pain after spinal cord injury through regulation of cell cycle pathways** *J Neurosci* **33**:12447–12463 <https://doi.org/10.1523/jneurosci.0846-13.2013>
- Wu J., Zhao Z., Sabirzhanov B., Stoica B.A., Kumar A., Luo T., Skovira J., Faden A.I (2014) **Spinal cord injury causes brain inflammation associated with cognitive and affective changes: role of cell cycle pathways** *J Neurosci* **34**:10989–11006 <https://doi.org/10.1523/jneurosci.5110-13.2014>
- Wu J., Zhao Z., Zhu X., Renn C.L., Dorsey S.G., Faden A.I (2016) **Cell cycle inhibition limits development and maintenance of neuropathic pain following spinal cord injury** *Pain* **157**:488–503 <https://doi.org/10.1097/j.pain.0000000000000393>
- Yu X., Nagai J., Khakh B.S (2020) **Improved tools to study astrocytes** *Nature Reviews Neuroscience* **21**:121–138 <https://doi.org/10.1038/s41583-020-0264-8>

Article and author information

Qing Zhao

Key Laboratory of Spine and Spinal Cord Injury Repair and Regeneration of Ministry of Education, Orthopaedic Department of Tongji Hospital, School of Medicine, School of Life Sciences and Technology, Tongji University, Shanghai, 200065 P. R. China, Division of Spine, Department of Orthopedics, Tongji Hospital, Tongji University School of Medicine, Tongji University, Shanghai, 200065 P. R. China
ORCID iD: [0000-0003-3437-533X](https://orcid.org/0000-0003-3437-533X)

Yanjing Zhu

Key Laboratory of Spine and Spinal Cord Injury Repair and Regeneration of Ministry of Education, Orthopaedic Department of Tongji Hospital, School of Medicine, School of Life Sciences and Technology, Tongji University, Shanghai, 200065 P. R. China

Yilong Ren

Key Laboratory of Spine and Spinal Cord Injury Repair and Regeneration of Ministry of Education, Orthopaedic Department of Tongji Hospital, School of Medicine, School of Life Sciences and Technology, Tongji University, Shanghai, 200065 P. R. China, Division of Spine, Department of Orthopedics, Tongji Hospital, Tongji University School of Medicine, Tongji University, Shanghai, 200065 P. R. China

Lijuan Zhao

Key Laboratory of Spine and Spinal Cord Injury Repair and Regeneration of Ministry of Education, Orthopaedic Department of Tongji Hospital, School of Medicine, School of Life Sciences and Technology, Tongji University, Shanghai, 200065 P. R. China, Division of Spine, Department of Orthopedics, Tongji Hospital, Tongji University School of Medicine, Tongji University, Shanghai, 200065 P. R. China

Jingwei Zhao

Key Laboratory of Spine and Spinal Cord Injury Repair and Regeneration of Ministry of Education, Orthopaedic Department of Tongji Hospital, School of Medicine, School of Life Sciences and Technology, Tongji University, Shanghai, 200065 P. R. China, Division of Spine, Department of Orthopedics, Tongji Hospital, Tongji University School of Medicine, Tongji University, Shanghai, 200065 P. R. China

Shuai Yin

Key Laboratory of Spine and Spinal Cord Injury Repair and Regeneration of Ministry of Education, Orthopaedic Department of Tongji Hospital, School of Medicine, School of Life Sciences and Technology, Tongji University, Shanghai, 200065 P. R. China, Division of Spine, Department of Orthopedics, Tongji Hospital, Tongji University School of Medicine, Tongji University, Shanghai, 200065 P. R. China

Haofei Ni

Key Laboratory of Spine and Spinal Cord Injury Repair and Regeneration of Ministry of Education, Orthopaedic Department of Tongji Hospital, School of Medicine, School of Life Sciences and Technology, Tongji University, Shanghai, 200065 P. R. China, Division of Spine, Department of Orthopedics, Tongji Hospital, Tongji University School of Medicine, Tongji University, Shanghai, 200065 P. R. China

Rongrong Zhu

Key Laboratory of Spine and Spinal Cord Injury Repair and Regeneration of Ministry of Education, Orthopaedic Department of Tongji Hospital, School of Medicine, School of Life Sciences and Technology, Tongji University, Shanghai, 200065 P. R. China

For correspondence: rrzhu@tongji.edu.cn

Liming Cheng

Key Laboratory of Spine and Spinal Cord Injury Repair and Regeneration of Ministry of Education, Orthopaedic Department of Tongji Hospital, School of Medicine, School of Life Sciences and Technology, Tongji University, Shanghai, 200065 P. R. China, Division of Spine, Department of Orthopedics, Tongji Hospital, Tongji University School of Medicine, Tongji University, Shanghai, 200065 P. R. China, Clinical Center for Brain and Spinal Cord Research, Tongji University

For correspondence: limingcheng@tongji.edu.cn

Ning Xie

Key Laboratory of Spine and Spinal Cord Injury Repair and Regeneration of Ministry of Education, Orthopaedic Department of Tongji Hospital, School of Medicine, School of Life Sciences and Technology, Tongji University, Shanghai, 200065 P. R. China, Division of Spine, Department of Orthopedics, Tongji Hospital, Tongji University School of Medicine, Tongji University, Shanghai, 200065 P. R. China

For correspondence: nxieprof18@tongji.edu.cn

Copyright

© 2024, Zhao et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Bamidele Owoyele

University of Ilorin, Ilorin, Nigeria

Senior Editor

Michael Taffe

University of California, San Diego, San Diego, United States of America

Reviewer #1 (Public Review):

Summary:

In this study the authors demonstrated that ablation of astrocytes in lumbar spinal cord not only reduced neuropathic pain but also caused microglia activation. Furthermore, RNA sequencing and bioinformatics revealed an activation of STING/type I IFNs signal pathway in spinal cord microglia after astrocyte ablation.

Strengths:

The findings are novel and interesting and provide new insights into astrocyte-microglia interaction in neuropathic pain. This study may also offer a new therapeutic strategy for the treatment of debilitating neuropathic pain in patients with SCI.

Weaknesses:

More details are needed to justify the sample size, statistics, and sex of animals.

<https://doi.org/10.7554/eLife.95672.1.sa1>

Reviewer #2 (Public Review):

Summary:

In the manuscript, Zhao et al. have carried out a thorough examination of the effects of targeted ablation of resident astrocytes on behavior, cellular responses, and gene expression after spinal cord injury. Employing transgenic mice models alongside pharmacogenetic techniques, the authors have successfully achieved the selective removal of these resident astrocytes. This intervention led to a notable reduction in neuropathic pain and induced a shift in microglial cell reactivation states within the spinal cord, significantly altering transcriptome profiles predominantly associated with interferon (IFN) signaling pathways.

Strengths:

The findings presented add considerable value to the current understanding of the role of astrocyte elimination in neuropathic pain, offering convincing evidence that supports existing hypotheses and valuable insights into the interactions between astrocytes and microglial cells, likely through IFN-mediated mechanisms. This contribution is highly relevant and suggests that further exploration in this direction could yield meaningful results.

Weaknesses:

The methodology and evidence underpinning the study are solid, yet some areas would benefit from further clarification, particularly concerning methodological details and the choice of statistical analyses. Additionally, the manuscript's organization and clarity could be improved, as certain figures and schematics appear inconsistent or misleading.

<https://doi.org/10.7554/eLife.95672.1.sa0>