

Perturbations in eIF3 subunit stoichiometry alter expression of ribosomal proteins and key components of the MAPK signaling pathways

Reviewed Preprint

v2 • October 18, 2024

Revised by authors

Reviewed Preprint

v1 • March 25, 2024

Anna Herrmannová , Jan Jelínek, Klára Pospíšilová, Farkas Kerényi, Tomáš Vomastek, Kathleen Watt, Jan Brábek, Mahabub Pasha Mohammad, Susan Wagner, Ivan Topisirovic, Leoš Shivaya Valášek 

Laboratory of Regulation of Gene Expression, Institute of Microbiology of the Czech Academy of Sciences, Prague, Videnska 1083, 142 20, the Czech Republic • Laboratory of Bioinformatics, Institute of Microbiology of the Czech Academy of Sciences, Prague, Videnska 1083, 142 20, the Czech Republic • Laboratory of Cell Signaling, Institute of Microbiology of the Czech Academy of Sciences, Prague, Videnska 1083, 142 20, the Czech Republic • Science for Life Laboratory, Department of Oncology-Pathology, Karolinska Institutet, Solna, Sweden • Laboratory of Cancer Cell Invasion, Faculty of Science, Charles University, Prague, Vinicna 7, 128 43, Czech Republic • Lady Davis Institute, Gerald Bronfman Department of Oncology, Department of Biochemistry, Division of Experimental Medicine, McGill University, Montréal, H3T 1E2, QC Canada

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Protein synthesis plays a major role in homeostasis and when dysregulated leads to various pathologies including cancer. To this end, imbalanced expression of eukaryotic translation initiation factors (eIFs) is not only a consequence but also a driver of neoplastic growth. eIF3 is the largest, multi-subunit translation initiation complex with a modular assembly, where aberrant expression of one subunit generates only partially functional subcomplexes. To comprehensively study the effects of eIF3 remodeling, we contrasted the impact of eIF3d, eIF3e or eIF3h depletion on the translome of HeLa cells using Ribo-seq. Depletion of eIF3d or eIF3e, but not eIF3h reduced the levels of multiple components of the MAPK signaling pathways. Surprisingly, however, depletion of all three eIF3 subunits increased MAPK/ERK pathway activity. Depletion of eIF3e and partially eIF3d also increased translation of TOP mRNAs that encode mainly ribosomal proteins and other components of the translational machinery. Moreover, alterations in eIF3 subunit stoichiometry were often associated with changes in translation of mRNAs containing short uORFs, as in the case of the proto-oncogene MDM2 and the transcription factor ATF4. Collectively, perturbations in eIF3 subunit stoichiometry exert specific effect on the translome comprising signaling and stress-related transcripts with complex 5' UTRs that are implicated in homeostatic adaptation to stress and cancer.

eLife Assessment

This study demonstrates mRNA-specific regulation of translation by subunits of the eukaryotic initiation factor complex 3 (eIF3) using **convincing** methods, data, and analyses. The investigations have generated **important** information that will be of interest to biologists studying translation regulation. However, the physiological significance of the gene expression changes that were observed is not clear.

<https://doi.org/10.7554/eLife.95846.2.sa4>

Introduction

Translational control contributes significantly to the overall regulation of gene expression. Indeed, regulation at the level of translation provides quicker adaptive and usually reversible responses to environmental cues than transcriptional regulation as it acts on pre-existing mRNAs by simply turning their expression on or off (Sonenberg & Hinnebusch, 2009 [↗](#)). As such, translational control critically contributes to the maintenance of overall cellular homeostasis, suggesting that its dysregulation has detrimental effects on cell proliferation and, more broadly, on health (Scheper *et al*, 2007 [↗](#)). In a rapidly growing number of cases, mutations and/or impaired expression of many translational components have been shown to either directly lead to or significantly contribute to various pathological conditions, including cancer (reviewed in (de la Parra *et al*, 2018b [↗](#); Robichaud & Sonenberg, 2017 [↗](#); Spilka *et al*, 2013 [↗](#); Sriram *et al*, 2018 [↗](#))).

Translation can be divided into four stages with the initiation stage playing the most critical role with respect to translational control. It begins with the formation of the 43S pre-initiation complex (PIC) composed of the 40S ribosomal subunit, the eIF2*GTP*Met-tRNA_i^{Met} ternary complex (eIF2-TC), and eIFs 1, 1A, 3, and 5 (reviewed in (Valášek, 2012 [↗](#))). The next step is the mRNA recruitment to form the 48S PIC, which is facilitated by the eIF4F cap-binding complex in concert with eIF3. Once the mRNA is loaded, the 48S PIC starts scanning the mRNA's 5' untranslated region (UTR) until the start codon has been recognized by base-pairing with the Met-tRNA_i^{Met} anticodon. This drives conformational changes in the 48S PIC, co-operatively mediated by eIF1, eIF1A, eIF2, eIF3 and eIF5, resulting in the closure of the 40S mRNA binding channel and subsequent ejection of the eIF2*GDP*eIF5 subcomplex to allow subunit joining to commence (reviewed in (Hinnebusch, 2017 [↗](#))).

Cellular stresses primarily inhibit cap-dependent translation initiation via the eIF4F and eIF2-TC complexes. The former is under the control of two major stress-signaling hubs – the anabolic kinase – the mechanistic target of rapamycin complex 1 (mTORC1) and the mitogen-activated protein kinases (MAPKs). They promote growth under nutrient-rich and stress-free conditions and deactivate in response to limited supply of amino acids, oxygen, ATP, growth factors or nutrients and play a role in the unfolded protein response (Bhat *et al*, 2015 [↗](#); Lindqvist *et al*, 2018 [↗](#); Roux & Topisirovic, 2018 [↗](#)). These pathways are hyperactive in the vast majority of cancers, mainly due to activating mutations of RAS, BRAF and PI3K or loss-of-function mutations of the tumor suppressor PTEN, which are among the most common genetic perturbations in cancer (Kandoth *et al*, 2013 [↗](#)). The latter – eIF2-TC – is a major target of the pro-survival, adaptive pathway called integrated stress response (ISR) encompassing four members of the eIF2 α kinase family (PERK, PKR, GCN2 and HRI) that respond to a multitude of stimuli including nutrient limitation and endoplasmic reticulum stress by rapidly shutting down general translation (Pakos-Zebrucka *et al*, 2016 [↗](#)).

Mitogen-activated protein kinase (MAPK) cascades have been shown to play a key role in the transduction of extracellular signals into cellular responses. Three families of MAPK signaling pathways have been well characterized in mammalian cells: extracellular signal-regulated kinase (ERK), c-Jun N-terminal kinase/stress-activated protein kinase (JNK/SAPK), and p38 kinase (Cargnello & Roux, 2011 [↗](#); Zhang & Liu, 2002 [↗](#)). The activation of MAPKs is a multistep process. In case of the best described Raf-MEK-ERK pathway, the essential linkers of epidermal growth factor receptors and other receptor tyrosine kinases with the MAPK cascade include adaptor protein Grb2, a Son of Sevenless (SOS) family of guanine nucleotide exchange factors, and a small GTP binding protein Ras (**Figure 1A** [↗](#)). The MAPK cascade begins with MAPKKK (represented by c-Raf) which phosphorylates and activates MEK1 and MEK2 (MAPKKs). MEKs eventually phosphorylate, and thereby activate p44 MAPK and p42 MAPK, also known as ERK1 and ERK2, respectively (Lavoie *et al*, 2020 [↗](#); Stokoe *et al*, 1994 [↗](#)). The activated ERKs then translocate to the nucleus where they phosphorylate and transactivate various transcription factors such as Elk-1, c-Jun and c-Fos, thereby altering gene expression to promote growth, differentiation, or mitosis. The ERK/MAPK pathway also directly regulates translational machinery, for example by stimulating the activity of both ribosomal S6 protein kinases (RSKs; RSK1-4 in humans) (Carriere *et al*, 2008 [↗](#)) and MAPK-interacting kinases (MNKs; MNK1 and 2 in humans) (Buxade *et al*, 2008 [↗](#); Fukunaga & Hunter, 1997 [↗](#); Waskiewicz *et al*, 1997 [↗](#)). MNKs interact with eIF4G and phosphorylate eIF4E at Ser209, a site that increases the oncogenic potential of eIF4E (Topisirovic *et al*, 2004 [↗](#)) and facilitates translation of specific mRNAs related to migration, metastasis and inflammation (Furic *et al*, 2010 [↗](#); Pyronnet *et al*, 1999 [↗](#); Robichaud *et al*, 2015 [↗](#)). Following the stimulation of the ERK pathway, activated RSKs phosphorylate rpS6, eIF4B, eEF2K, and PDCD4, which are all important translational regulators (reviewed in (Bhat *et al*, 2015 [↗](#); Robichaud & Sonenberg, 2017 [↗](#))). RSKs have also been implicated in the regulation of other signaling pathways involved in growth control, stress-response and cancer including the regulation of mTORC1 (reviewed in (Roux & Topisirovic, 2018 [↗](#))). For example, RSKs have been shown phosphorylate and inhibit tuberous sclerosis complex 2 (TSC2) that acts as the negative regulator of mTORC1 signaling (Roux *et al*, 2004 [↗](#)) (**Figure 1A** [↗](#)).

It has been shown that alterations in genomic sequences, protein levels, stoichiometry, and/or activity of eIFs and other components of the translational apparatus and relevant signaling cascades, as well as in cis-regulatory elements of mRNAs, can circumvent the stress-induced translational shutdown that these pathways impose by allowing selective recruitment of mRNAs, rewiring translational control of mRNA cis-regulatory elements, and/or enabling alternative start codon selection (Kovalski *et al*, 2022 [↗](#); Robichaud *et al*, 2019 [↗](#); Sriram *et al*, 2018 [↗](#)). Naturally, this leads to various pathologies. Overexpression of eIF4F complex components, for example, has been observed in various cancer entities (Bhat *et al*, 2015 [↗](#); De Benedetti & Harris, 1999 [↗](#); Lazaris-Karatzas *et al*, 1990 [↗](#); Ruggero, 2013 [↗](#); Shuda *et al*, 2000 [↗](#); Truitt & Ruggero, 2016 [↗](#)). In addition to eIF4F components, dysregulation of the multi-subunit eIF3 complex is also involved in oncogenesis. Ten of the twelve eIF3 subunits have been linked to human cancers, many due to overexpression (a, b, c, d, g, h, m or i), some due to underexpression (e and f) (reviewed in (de la Parra *et al*, 2018b [↗](#); Gomes-Duarte *et al*, 2018 [↗](#); Hershey, 2015 [↗](#); Valasek *et al*, 2017 [↗](#); Zhang *et al*, 2007 [↗](#))). In this regard, it is worth stressing that some eIF3 subunits have been shown to directly interact with specific mRNAs (Lee *et al*, 2015 [↗](#)), eIF3d together with DAP5 was shown to facilitate alternate form of cap-dependent mRNA translation (de la Parra *et al*, 2018a [↗](#)), and eIF3k was recently proposed to serve as a regulator of the balanced ribosome content (Duan *et al*, 2023 [↗](#)). Despite all these observations, the exact mechanism(s) by which eIF3 affects various pathologies is virtually unknown. Moreover, previous studies that investigated the effect of eIF3 in specific diseases focus on altered levels of only one of the eIF3 subunits, while neglecting the need to consider the context of altered expression of the entire eIF3 complex.

Here, we draw attention to this unfortunate trend. eIF3 is the largest and most complex eIF of all eIFs, it is modular in nature (**Figure 1B** [↗](#)) (reviewed in (Valasek *et al*, 2017 [↗](#); Wolf *et al*, 2020 [↗](#))), and plays multiple roles not only in translation initiation, but also in termination, ribosomal




Figure 1.

Schematic models of the core of MAPK/ERK pathway and human eIF3 complex.

(A) The MAPK/ERK pathway activation by growth factors and receptor tyrosine kinases (RTKs). RTKs signal through GRB2-Sos and Ras to activate Raf-MEK-ERK signaling cascade. Active ERK phosphorylates numerous substrates, both nuclear and cytoplasmic, to affect gene transcription, protein translation, cell growth, proliferation and survival. Both RTKs and Ras also activate PI3K-AKT pathway that affect cell survival and growth. Conventional PAKs (PAK1-3) phosphorylate c-Raf and contribute to its activation. Created with Biorender.com. (B) eIF3 subunits forming the PCI/MPN octamer are indicated by the grey background. The rectangle marks the seven α -helices involved in formation of the 7-helix bundle. The Yeast-Like Core (YLC) comprising the eIF3 subunits a, b, g and i is depicted, and so is the eIF3-associated factor eIF3j. Arrows indicate subunits targeted by siRNA and subjected to Ribo-Seq in this study; eIF3 subcomplexes generated in individual knock-downs are boxed (adapted from (Wagner *et al.*, 2016)).

recycling and stop codon readthrough (Beznosková *et al*, 2013 [↗](#); Beznosková *et al*, 2015 [↗](#); Pisarev *et al*, 2007 [↗](#); Poncova *et al*, 2019 [↗](#)). Each of these roles may be regulated differently by individual subunits and/or their combinations, making it practically impossible to study them in isolation. Eight eIF3 subunits (a, c, e, f, h, k, l and m) form a structural scaffold called the PCI (Proteasome, COP9, eIF3) / MPN (Mpr1-Pad1 N-terminal) octamer, whereas the remaining four non-octameric subunits (b, d, g and i) are likely to be more flexible. The essential subunits b, g, and i form a separate module that is connected to the octamer *via* the C-terminal domain (CTD) of eIF3a; together they are called YLC for Yeast Like Core (Wagner *et al*, 2016 [↗](#)). The eIF3d subunit is located on the eIF3 periphery and is attached to the octamer primarily via eIF3e, but partly also *via* eIF3a and c (Bochler *et al*, 2020 [↗](#); Brito Querido *et al*, 2020). On the 40S subunit, the octamer is positioned close to the mRNA exit site, whereas the mobile YLC shuffles between the mRNA entry site and the subunit interface during the AUG selection process (des Georges *et al*, 2015 [↗](#); Llacer *et al*, 2021 [↗](#); Simonetti *et al*, 2020 [↗](#)).

We recently reported a comprehensive *in vivo* analysis of the modular dynamics of the human eIF3 complex (Wagner *et al*, 2020 [↗](#); Wagner *et al*, 2014 [↗](#); Wagner *et al*, 2016 [↗](#)). Using a systematic individual downregulation strategy, we showed that the expression of all 12 eIF3 subunits is interconnected such that perturbation of the expression of one subunit results in the down-regulation of entire modules leading to the formation of partial eIF3 subcomplexes with limited functionality (Herrmannová *et al*, 2020 [↗](#)). eIF3d is the only exception in this respect, as its downregulation does not influence expression of any other eIF3 subunit. Here we took advantage of this knowledge to examine the specific effect of selected eIF3 subcomplexes on the translation efficiency, transcriptome-wide, in HeLa cells.

Methods

Cell lines, culture conditions and transfection

HeLa cells (ATCC: CCL2) were grown at 37 °C and 5% CO₂ in Ø 15 cm dishes in DMEM (Sigma, cat #D6429) supplemented with 10% FBS (Gibco, cat # 10270-106). Twenty-four hours after seeding, cells were transfected with the ON-TARGETplus siRNA cocktail system from Dharmacon at a final concentration of 5 nM. Catalog numbers for all siRNAs used in this study are listed in Supplementary Table S1. INTERFERin (Polyplus, cat # 101000016) was used as a transfection reagent, 100 µl per dish. Cells were harvested 72 hours after transfection at approximately 80% confluency and cytoplasmatic lysates were prepared as described previously (Herrmannová *et al*, 2020 [↗](#)). For monitoring levels of phosphorylated c-Jun protein, cytoplasmatic and nuclear lysates were prepared using NE-PER™ Nuclear and Cytoplasmic Extraction Reagents from Thermo Scientific™.

Ribosome profiling (Ribo-Seq) and RNA-seq library preparation

For **footprint (FP) libraries** cycloheximide was added to a final concentration of 100 µg/ml 1 min prior to harvesting. Cells were washed with ice cold 1x PBS and lysed in buffer A (10 mM HEPES [pH 7.5], 62.5 mM KCl, 2.5 mM MgCl₂, 1 mM DTT, 1 mM PMSF, 1 µg/ml Aprotinin, 1 µg/ml Leupeptin, 1 µg/ml Pepstatin, Complete Mini EDTA-free [Roche, cat # 11836170001] – 1 tablet/5 ml, 1% Triton X-100, 100 µg/ml Cycloheximide) on the dish. The resulting lysate was collected and cleared by centrifugation. Aliquots of twelve absorbance units (AU) OD₂₆₀ of WCE were flash frozen in liquid nitrogen and used for control polysome profile and footprint isolation. An aliquot of 1 mg total protein was taken for control western blot, and another 200 µl aliquot was mixed with 1 ml of RNA Blue (TopBio, cat # R013) and taken for RNA isolation and control qPCR. When the lysate passed all three control experiments (polysome profile, western blot, qPCR), one aliquot of 12 AU OD₂₆₀ was thawed on ice, digested with 5 µl RNaseI (Ambion, cat # AM2295) for 30 min at 24 °C with mild shaking (300 rpm) and inactivated with SuperaseIN (Ambion, cat #AM2696). 80S ribosomes were separated by high-velocity sedimentation through a 5% to 45% sucrose gradient at

39,000 rpm for 2.5 h using the SW41Ti rotor. The gradients were scanned at 254 nm to visualize the ribosomal species and fractions containing 80S ribosomes were collected and mixed with RNA Blue. RNA was isolated according to vendor's instructions, FP RNA was size selected on 15% TBE-Urea PAGE (Biorad, cat # 3540092) in range 20-40 nt and footprint library was prepared according to (Ingolia *et al.*, 2012 [DOI](#)). FP libraries were sequenced on NovaSeq6000 (Illumina) with read length 50 nt.

For **mRNA libraries** the growth media was aspirated and cells were harvested into 5 ml of RNA Blue per Ø 15 cm dish. RNA was isolated according to vendor's instructions. mRNA was isolated from the total RNA using Poly(A)purist MAG kit (Ambion, cat # AM1922), fragmented using RNA Fragmentation Reagents (Ambion, cat # AM8740) and size selected on 15% TBE-Urea PAGE in range 50-100 nt. Size selected mRNA was then dephosphorylated and library was prepared using SMARTer smRNA-Seq Kit (TAKARA, cat # 635029) according to vendor's instructions. mRNA libraries were sequenced on NovaSeq6000 (Illumina) with read length 150 nt.

RNA isolation, reverse transcription and qPCR

Total RNA was isolated using RNA Blue reagent (TopBio, cat # R013) 72 h post-transfection according to the manufacturer's instructions. After DNase I digestion (NEB, cat # M0303L), cDNA was synthesized from 1 µg RNA using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems, # 4368813). qPCR was performed in triplicates using 5× HOT FIREPol EvaGreen qPCR Mix Plus (Solis BioDyne # 08-25-00020). The obtained data were normalized to ALAS1 mRNA levels and non-transfected control. All qPCR primers are listed in Supplementary Table S2.

Western blotting

All samples were resolved using SDS-PAGE followed by western blotting. All primary antibodies used in this study are listed in Supplementary Table S3. The signal was developed using SuperSignal West Femto Maximum Sensitivity Substrate (Thermo Scientific, cat # 34096) and detected in a G-Box imager (Syngene) using a series of varying exposure times. Signals were processed with Quantity One (BioRad). Only signals from the same strips and with the same exposure times were compared.

Polysome profiles

One aliquot of 12 AU OD₂₆₀ was thawed on ice and separated by high-velocity sedimentation through a 5% to 45% sucrose gradient at 39,000 rpm for 2.5 h using the SW41Ti rotor. The gradients were scanned at 254 nm to visualize the ribosomal species.

60S/40S ratio calculation

Cell lysates were prepared as above by on dish lysis in buffer A. To 6 AU OD₂₆₀ EDTA was added to final concentration of 50mM and the lysate was separated by high-velocity sedimentation through a 5% to 50% sucrose gradient at 39,000 rpm for 3 h using the SW41Ti rotor. The gradients were scanned at 260 nm to visualize the ribosomal species. Area under the curve was quantified to calculate the 60S:40S ratio to check for ribosome biogenesis defect.

Ribosome content profiles

Cells were counted by Corning Automated Cell Counter and 10 million cells were pelleted in microcentrifuge tube and lysed in buffer A (as above). EDTA was added to the lysate to final concentration of 50mM and the lysate was separated by high-velocity sedimentation through a 5% to 50% sucrose gradient at 39,000 rpm for 3.5 h using the SW41Ti rotor. The gradients were scanned at 260 nm to visualize the ribosomal species. Area under the curve was quantified to calculate the total amount of ribosomes between samples.

Analysis of Sequencing Data

We processed the ribosome **footprint libraries** as follows. Adapter sequence “CTGTAGGCACCATCAAT” and all sequences 3’ of the adapter were trimmed from the 3’ end of each read by Cutadapt v. 3.5 (Martin, 2011 [↗](#)). Flanking Ns were trimmed from both 5’ and 3’ ends of each read. Reads without adapter sequence and reads shorter than 15 nt after trimming were discarded. Next, rRNA and tRNA reads were discarded after alignment with Bowtie2 v. 2.4.4 (Langmead & Salzberg, 2012 [↗](#)). For rRNA, we used RNA5S1, RNA5-8SN5, RNA18SN5 and RNA28SN5 from NCBI (Sayers *et al.*, 2022 [↗](#)), and ENSG00000211459 and ENSG00000210082 from Ensembl (Howe *et al.*, 2021 [↗](#)). For tRNA we used all human tRNA sequences from tRNAdb (Juhling *et al.*, 2009 [↗](#)) and all high confidence hg38 tRNA sequences from GtRNAdb (Chan & Lowe, 2016 [↗](#)). Remaining reads were aligned to genome and transcriptome from Ensembl (GRCh38.p13, annotation release 104) by the STAR aligner v. 2.7.9a_2021-06-25 (Dobin *et al.*, 2013 [↗](#)) using local alignment with 10% mismatches, and with indels and soft clipping allowed for transcriptomic alignment. Gene counts were counted from genomic alignment using Samtools index v. 1.13 (Danecek *et al.*, 2021 [↗](#)) and Htseq-count v. 0.13.5 (Anders *et al.*, 2015 [↗](#)) within CDS regions. Transcriptomic alignments were further processed using MANE Select set from MANE project v. 0.95 (Morales *et al.*, 2022 [↗](#)) and custom programs “mane2ensembl_gtf”, “filter_reverse_reads”, “filter_ambiguous_genes” and “select_transcripts”. See Supplementary Spreadsheet S1 for detailed overview of raw read processing.

The **RNA-seq libraries** were processed the same as above, except that raw read processing by Cutadapt had to be adapted to different library construction method (SMARTer smRNA-Seq Kit, TAKARA). In particular, first 3 nucleotides on the 5’ end were removed, and poly(A) sequence (10xA) was trimmed from the 3’ end.

Quality checks were done by FastQC v. 0.11.9 (Andrews, 2010 [↗](#)) and Ribo-seQC v. 1.1 (Calviello *et al.*, 2019). Triplet periodicity was checked on transcriptomic alignments using custom modified riboWaltz (Lauria *et al.*, 2018 [↗](#)) and a custom script “Triplet_periodicity.R”. Gene counts were used to check correlation between replicates using a custom script “Correlation-samples.R”. Read length counts were used to check read length distribution using a custom script “Lengths.R”.

PCA plot, sample distances clustering and the differential expression analysis were performed on gene counts by DESeq2 (Love *et al.*, 2014 [↗](#)) with biomaRt library (Durinck *et al.*, 2009 [↗](#)) used to download gene annotations. Differential translation efficiency (TE) values between NT and the knock-downs were calculated as described in (Chothani *et al.*, 2019 [↗](#)). The p_{adj} threshold for differentially translated transcripts was set to 0.05. For details see custom script “Differential analysis.R”. Results of differential expression analysis were further used to generate heatmap and scatterplots of differentially expressed genes and genes with defined p-adjusted value were used to analyze dependency between mRNA characteristics and changes in TE using custom scripts “Significant_TE_heatmap.R”, “Significant_TE_intersection.R” and “Correlation-features.R”. To get mRNA characteristics we used MANE Select set to select a single representative transcript for each protein-coding gene. uORF database uORFdb (Manske *et al.*, 2023 [↗](#)) was used to get number of ATG uORFs per mRNA.

Metagene analysis was done as described in (Lin *et al.*, 2020 [↗](#)), but 5’ ends of reads were used instead of P-sites. Samtools and custom programs “transcripts_startstop_positions”, “read_counts” and “region_readcounts” were used to get read counts relative to the start codon. These were normalized to a number of reads in a region starting 75 bases after the start codon and ending 15 bases before the stop codon and averaged over the samples. Only transcripts that met the following parameters were considered: CDS longer than 450 nt and RPKM higher than 10. Weighted counts of biological replicates were further used to compute average read counts for each sample. Moving average method was used with a 21 bases window size.

For footprint coverage plots of individual mRNAs, MANE transcripts of examined genes (ENST00000674920-ATF4; ENST00000258149-MDM2; ENST00000251849-RAF1) were selected. Reads were counted for each position using Samtools and a custom script read_counts and normalized to number of read counts over all positions of transcript. Weighted counts of biological replicates were further used to compute average read counts for each sample. Graphs were smoothened with a sliding window of 30nt. All used custom scripts are available on Github.

KEGG/GO pathway enrichment analysis

We analyzed enrichment of the differentially translated transcripts in KEGG pathways (Kanehisa & Sato, 2020 [↗](#)) and Gene Ontology (Ashburner *et al.*, 2000 [↗](#); Gene Ontology, 2021) using Enrichr (Kuleshov *et al.*, 2016 [↗](#)), ShinyGO v0.741 (Ge *et al.*, 2020 [↗](#)) and KEGG Mapper Search (Kanehisa *et al.*, 2022 [↗](#)). Venn diagrams were created using InteractiVenn (Heberle *et al.*, 2015 [↗](#)).

Analysis of Gene Signatures

To assess changes in the regulation of known gene signatures in the eIF3 KDs, empirical cumulative distribution functions (ECDFs) of $\log_2$ fold changes in FP and total mRNA were plotted independently. Fold change ECDFs for genes belonging to a particular signature were compared to those for all other genes, and differences in distributions were calculated at the quantiles. Significant directional shifts between the signatures and the background were identified using the Wilcoxon rank-sum test.

Results

High quality sequencing libraries prepared from HeLa cells expressing different eIF3 subcomplexes

To investigate the impact of different human eIF3 subunits and partial eIF3 sub-complexes, which arise in cells as a result of unbalanced eIF3 expression, on translational efficiency transcriptome-wide, we performed Ribo-Seq together with RNA-Seq, which allowed us to draw conclusions about gene expression at the transcriptional and translational levels and their interdependence. In particular, using the siRNA technology, we examined HeLa cells with reduced levels of eIF3d, eIF3e or eIF3h subunits and cells treated with non-targeting (NT) siRNA. These eIF3 subunits were selected because the eIF3d depletion leads to a reduction in protein levels of only eIF3d and leaves the rest of the eIF3 complex intact, whereas downregulation of eIF3e or eIF3h generates partial eIF3 subcomplexes as previously described (Wagner *et al.*, 2016 [↗](#)). Specifically, targeting eIF3e with siRNA results in reduced protein levels of eIF3e, eIF3d, eIF3k and eIF3l, whereas downregulation of eIF3h produces a partial eIF3 complex lacking subunits eIF3h, eIF3k and eIF3l (**Figure 1B** [↗](#)). In terms of impact on cell growth, both eIF3d^{KD} and eIF3e^{KD} strongly reduce proliferation, while the effect of eIF3h^{KD} is only moderate (Supplementary Figure 1A) (Wagner *et al.*, 2016 [↗](#)).

We first performed several control experiments with aliquots of cell lysates made to prepare individual Ribo-Seq libraries to document an efficient downregulation of the targeted eIF3 subunit, as well as the expected phenotype (Wagner *et al.*, 2016 [↗](#)). Using qPCR, we measured mRNA levels of the siRNA targeted eIF3 subunits (eIF3d, eIF3e, eIF3h), as well as of eIF3b as a negative control and a housekeeping gene ALAS1 for normalization purposes. Routinely, a difference of 4 to 5 cycles was achieved for each downregulated subunit, corresponding to ~16–32 fold decrease in mRNA levels; the mRNA levels of the control subunit eIF3b remained unchanged (Supplementary Figure 1B); note that Supplementary Figure 1B summarizes qPCR results for all samples used to generate Ribo-Seq and RNA-Seq libraries.

Next, we determined protein levels of selected eIF3 subunits by Western blotting. Protein levels of all eIF3 subunits targeted individually by siRNA were strongly reduced (Supplementary Figure 1C). Depletion of eIF3e also induced a marked decrease in the eIF3d, eIF3k and eIF3l protein levels (Supplementary Figure 1C and data not shown), as previously reported (Wagner *et al.*, 2016). Finally, we performed polysome profiling to assess the impact on translation and obtained essentially the same results as before (Wagner *et al.*, 2016). The eIF3d^{KD} showed the most pronounced defect in translation initiation, marked by a robust increase of the 80S monosome peak and a concomitant decrease of polysome content, similar to eIF3e^{KD}, while eIF3h^{KD} had only a modest effect on polysome formation (Supplementary Figure 1D).

Ribo-Seq libraries were generated from each knock-down and NT control in three biological replicates as described in (Ingolia *et al.*, 2012). RNA-Seq libraries were prepared in four biological replicates using the SMARTer smRNA-Seq Kit to normalize for mRNA levels and calculate translational efficiency (TE). Both Ribo-Seq and RNA-seq libraries were highly reproducible as shown by the Spearman correlation coefficient (Supplementary Figure 2A). All Ribo-Seq libraries showed a similar read length distribution with a prominent peak at 30 – 32 nt and an expected 3 nt periodicity in coding sequences (CDS) (Supplementary Figure 2B-C and Supplementary Figure 3A). As expected, Ribo-seq libraries were strongly enriched for CDS reads compared to RNA-Seq libraries (Supplementary Figure 3B – C). The PCA plot and hierarchical clustering (Figure 2A and Supplementary Figure 4A) showed clustering of the samples into two main groups: Ribo-Seq and RNA-seq, and also into two subgroups; NT and eIF3h^{KD} samples clustered on one side and eIF3e^{KD} and eIF3d^{KD} samples on the other. These results suggest that the eIF3h depletion has a much milder impact on the translome than depletion of eIF3e or eIF3d, which agrees with the growth phenotype and polysome profile analyses (Supplementary Figure 1A and 1D).

eIF3 subcomplexes alter translation efficiency of a large proportion of genes

Using the R DESeq2 package (Love *et al.*, 2014), we identified transcripts with significantly altered TE ($P_{\text{adjusted}} < 0.05$) for each knock-down compared to the NT control (Figure 2B – D, Supplementary Spreadsheet S2), henceforth referred to as Differential Translation Efficiency Genes (DTEGs) (Chothani *et al.*, 2019). In total, we identified 21 DTEGs in eIF3h^{KD} (19 downregulated and 2 upregulated), 1180 DTEGs in eIF3e^{KD} (622 down and 558 up), and 1971 DTEGs in eIF3d^{KD} (1040 down and 931 up) out of a total of 17520 genes that were assigned TE values.

The number of DTEGs in each knock-down correlated well with the severity of the overall phenotype and with the impact on global translation, as judged from the polysome profiles. eIF3d^{KD} with the strongest decrease in polysomes had the highest number of DTEGs, whereas eIF3h^{KD} with only 21 DTEGs had a marginal effect on polysome formation (Supplementary Figure 1D). Consistently, the overall distribution of fold-changes in TE showed most pronounced changes in eIF3d^{KD} (Figure 2E). It should be noted, however, that in contrast to polysome profiling, differential expression analysis of Ribo-Seq data normalizes out global changes in translational rates, and reveals only specific shifts of ribosome localization on individual mRNAs between the NT and KD datasets. Therefore, the fact that all three knock-downs showed an increase in TE (not just a decrease) for approximately half of the affected genes, clearly points to an impairment of a specific mode of regulation rather than simply to a non-specific global translational shutdown.

Interestingly, there is a considerable overlap between the knock-downs studied, such that only 1045 DTEGs are unique for eIF3d^{KD} and the rest (926) are shared with other two knock-downs; in particular 923 with eIF3e^{KD} and 21 with eIF3h^{KD} – in the latter case, the overlap reaches 100% (Figure 2F, Supplementary Figure 5A – C). Remarkably, eIF3d^{KD} affects neither the expression of




Figure 2.

DTEGs identified in eIF3d^{KD}, eIF3e^{KD} and eIF3h^{KD} largely overlap.

(A) Principal component analysis of read count per gene of the RNA-Seq and Ribo-Seq libraries. (B - D) Volcano plots of significant ($P_{\text{adjusted}} < 0.05$) DTEGs in individual knock-downs: eIF3d^{KD} (B), eIF3e^{KD} (C) and eIF3h^{KD} (D). The volcano plots show the Log₂ fold-change of TE (x-axis) versus the significance -Log₁₀ p-adjusted value (y-axis). Genes without significant p-adjusted value are plotted in grey, downregulated DTEGs are plotted in blue and upregulated DTEGs in orange. For each plot, the top 10 most significant DTEGs are labeled with gene names. The Volcano plots were generated using modified script from Galaxy (Blankenberg *et al.*, 2014). (E) Heatmap and dendrogram resulting from hierarchical clustering analysis of significant TE changes observed in eIF3d^{KD}, eIF3e^{KD} and eIF3h^{KD} cells. (F - H) Venn diagrams depicting the overlap in all DTEGs (F), in upregulated DTEGs (G), and in downregulated DTEGs (H) among eIF3d^{KD}, eIF3e^{KD} and eIF3h^{KD}. Based on the overlaps, DTEGs were divided in several subgroups for further analysis: eIF3d^{KD} "all", eIF3d^{KD} "unique", eIF3e^{KD} "all", eIF3e^{KD} "unique", eIF3d^{KD}/eIF3e^{KD} "common" and eIF3h^{KD} "all".

any other eIF3 subunit nor the integrity of the eIF3 complex, yet it has the most pronounced impact on translation (Wagner *et al.*, 2016 [\[4\]](#)), correlating well with the highest number of DTEGs identified. In case of eIF3e^{KD}, only 257 DTEGs are unique and the rest (923) are shared with eIF3d^{KD}, 18 of which are also shared with eIF3h^{KD} (Figure 2F [\[4\]](#), Supplementary Figure 5A – C). The eIF3 subcomplex formed in eIF3e^{KD} lacks subunits e, k and l and partially also eIF3d (Wagner *et al.*, 2016 [\[4\]](#)) – the significantly reduced protein levels of eIF3d are evident in Supplementary Figure 1C. Thus, it is not surprising that the DTEGs identified in eIF3e^{KD} largely overlap with those identified in eIF3d^{KD}; in fact, this result supports the robustness and accuracy of this approach. The fact that the magnitude of the knock-down effect is bigger in eIF3d^{KD} over eIF3e^{KD} is discussed below.

eIF3d^{KD} and eIF3e^{KD} increase TE of mRNAs encoding proteins associated with membrane organelles

To analyze enrichment in KEGG pathways and GO Biological Processes in a meaningful way, we analyzed the DTEGs divided not only into upregulated and downregulated groups, but also into specific subgroups – eIF3d^{KD} “all”, eIF3e^{KD} “all”, eIF3d^{KD} “unique”, eIF3e^{KD} “unique” and eIF3d^{KD}/eIF3e^{KD} “common” (Figure 2F [\[4\]](#) – H). To disentangle the effects of the changes in mRNA abundance vs. translational regulation, we also investigated translationally exclusive DTEGs whose mRNA abundance did not change significantly and therefore the change in TE can be attributed solely to translational regulation; eIF3d^{KD} “translation only” and eIF3e^{KD} “translation only”, as well as translationally buffered DTEGs whose mRNA abundance changes in opposite direction than translational regulation, i.e. mRNA increases but TE decreases so the net protein production is not changed or even decreased; eIF3d^{KD} “buffered” and eIF3e^{KD} “buffered” (Supplementary Figure 5D – F). For schematic illustrating classification of genes based on fold changes of footprints (FP), mRNA, and TE, see Supplementary Figure 5G – H (adopted from (Chothani *et al.*, 2019 [\[4\]](#)) with permission).

Analysis of DTEGs upregulated in eIF3d^{KD} and eIF3e^{KD} showed similar results for almost all groups analyzed (eIF3d^{KD} “all”, “unique”, and “translation only”, eIF3d^{KD}/eIF3e^{KD} “common”, as well as eIF3e^{KD} “all” and “translation only”) (Figure 3A [\[4\]](#), B, and Supplementary Figure 6A), with the strongest enrichment in the KEGG “Lysosome” and “Protein processing in endoplasmic reticulum” pathways. The majority of DTEGs in these pathways were solely translationally regulated meaning that their mRNA levels stayed unchanged while the TE increased. For example, for the KEGG pathway “Lysosome”, more than 70% (i.e. 37 out of the 51) DTEGs upregulated in the eIF3d^{KD} “all”, fall into the eIF3d^{KD} “translation only” group. This suggests that when translation initiation is compromised by eIF3d/e^{KD}, cells may increase protein degradation and turnover by translationally upregulating lysosomal and ER proteins. Interestingly and in accord with our results, Lin *et al.* detected increased lysosomal load in MCF-10A cells downregulated for eIF3e (Lin *et al.*, 2020 [\[4\]](#)). They speculated that this is due to the need to clear defective mitochondria. In support, we also found enrichment of mitochondria-associated DTEGs, especially mitochondrial ribosomal proteins (see below).




Figure 3.

KEGG pathway enrichment analysis of the eIF3d^{KD}- and eIF3e^{KD}- associated upregulated DTEGs reveals upregulation of ribosomal proteins.

(A) Venn diagram of the 10 most significantly enriched KEGG pathways for eIF3d^{KD} “all” and eIF3e^{KD} “all” groups of upregulated DTEGs, highlighting that most of the pathways are common to both knock-downs. Lysosome and Protein processing in ER are highlighted in bold. The complete results of the KEGG enrichment analysis with corresponding *p*-values can be found in Supplementary Figure 5A. (B) Venn diagram as in A but displaying eIF3d^{KD} “translation only” and eIF3e^{KD} “translation only” groups of the upregulated DTEGs. (C) The bar chart shows the top 10 enriched KEGG terms for eIF3e^{KD} “unique” upregulated DTEGs. Orange bars correspond to terms with significant *p*-values (< 0.05), grey bars correspond to terms with not significant *p*-values (> 0.05). The *p*-values were calculated by Enrichr gene set search engine. (D) The list of genes pre-selected for western blot analysis and a heatmap showing their respective log2 fold-change values from differential expression analysis of FP, TE and mRNA in eIF3e^{KD} and eIF3d^{KD}. Positive values indicating significant upregulation are in shades of orange. ns = not significant *p*-adjusted value. (E) Western blot analysis of selected ribosomal proteins performed in the indicated knock-downs and the NT control. GAPDH was used as a loading control. (F) Relative protein levels of selected ribosomal proteins normalized to GAPDH; NT control = 100%. Dots represent results from individual biological replicates. One sample t-test was used for statistical evaluation, *P*-values: * = *P*<0.05, ** = *P*<0.01, ns = not significant. (G) eIF3d^{KD}, eIF3e^{KD} and eIF3h^{KD} do not influence the balanced ribosomal subunits production. The 60S/40S ratio was calculated from polysome profiles carried out in the presence of 50mM EDTA. Dots represent results from individual biological replicates. Paired t test was used for statistical evaluation, all downregulations were individually compared to NT. ns = not significant *P*-value. (H) Ribosomal content is increased in eIF3d^{KD} and eIF3e^{KD}. One representative polysome profile, carried out in the presence of 50mM EDTA, made from 10 million cells is shown in the upper panel. Relative ribosomal content normalized to NT control set to 100% is shown in the lower panel. Individual biological replicates are depicted as dots. One sample t-test was used for statistical evaluation, *P*-values: * = *P*<0.05, ** = *P*<0.01. All plots in (F-H) were created in GraphPad Prism version 8.4.3 for Windows.

Among the eIF3d^{KD} “unique upregulated” DTEGs, we identified one interesting KEGG pathway, the ABC transporters, which did not show up in other gene groups (Supplementary Figure 6A, in green). A total of 12 different ABC transporters had elevated TE (9 of them are unique to eIF3d^{KD}, while 3 were also found in eIF3e^{KD}), 6 of which (ABCC1-5, ABCC10) belong to the C subfamily, known to confer multidrug resistance with alternative designation as multidrug resistance protein (MRP1-5, MRP7) (Sodani *et al*, 2012 [\[1\]](#)). Interestingly, all six of these ABCC transporters were upregulated solely at the translational level (Supplementary Spreadsheet S2).

eIF3e^{KD} increases production of ribosomal proteins

Interestingly, the KEGG pathway “Ribosome” and GO terms connected with translation and ribosome biogenesis were enriched amongst DTEGs upregulated specifically in eIF3e^{KD} “unique upregulated” DTEGs (Figure 3C [\[2\]](#) and Supplementary Figure 6B). We then compared the significance of enrichment for “Ribosome” KEGG term between eIF3e^{KD} and eIF3d^{KD} “all upregulated” groups. This was the only case where a term or a pathway was more significantly enriched in eIF3e^{KD} (*P* = 5.199E-10) than in eIF3d^{KD} (NS, *P* = 0.1183) (Supplementary Figure 6A, in green). Intrigued by this result, we filtered out all ribosomal proteins (RPs) from our data set (according to HGNC database, Tweedie *et al*, 2021 [\[3\]](#)), regardless of the TE changes, which consisted of 89 cytoplasmic and 77 mitochondrial RPs, and analyzed them separately (Supplementary Spreadsheet S2). In total, we identified 32 RPs with significantly changed TE in eIF3e^{KD} (14 cytoplasmic and 18 mitochondrial) and only 16 in eIF3d^{KD} (9 cytoplasmic and 7 mitochondrial). Moreover, all the cytoplasmic RPs with increased TE in eIF3e^{KD} (and most mitochondrial ones) were upregulated solely at the translational level and their mRNA levels were not changed. We selected several RPs with different combinations of changes in TE or FP or mRNA

levels in both eIF3e^{KD} and eIF3d^{KD} (Figure 3D) and subjected them to western blotting. All ribosomal proteins tested showed visibly increased protein levels in both eIF3e^{KD} and eIF3d^{KD} (Figure 3E and F), while their mRNA levels remained unchanged (Supplementary Figure 6C).

As the production of the ribosomes seemed to be increased, we next examined the 60S/40S subunit ratio to control for unbalanced ribosome biogenesis. We performed polysome profiling in the presence of EDTA that splits all ribosomes into 40S and 60S subunits and showed that the 60S/40S ratio remained unaffected (Figure 3G). Furthermore, we performed a similar experiment with exactly 10 million cells *per* each sample and confirmed that the total amount of ribosomes in cells is indeed increased in eIF3e^{KD} and eIF3d^{KD} (Figure 3H).

Overall, our results suggest that eIF3e and, to some extent, its binding partner eIF3d maintain control over production of RPs in the cell. A similar role has recently been attributed to eIF3k, which has been shown to specifically repress the synthesis of the small ribosomal protein RPS15A, and cells depleted of eIF3k showed increased ribosomal content (Duan *et al.*, 2023). As mentioned above, targeting eIF3e with siRNA results in co-downregulation of not only eIF3e and eIF3d, but also of eIF3k and eIF3l (Wagner *et al.*, 2016) which may at least partially explain these findings.

In addition to mitochondrial RPs, we also observed increased TE for several other mitochondrial proteins from the Complex I, Complex IV and Complex V (KEGG term Oxidative phosphorylation hsa00190; rank 17; #of genes 12 out of 134; $p = 0.0003431$) in eIF3e^{KD}, but many of these transcripts, mainly from the Complex I, are encoded by mitochondrial DNA and translated in mitochondria directly, so these results should be verified by Mitoribosome profiling, as the traditional Ribo-Seq method is not suitable for this (Pearce *et al.*, 2021).

Because RPs and other components of the translational machinery frequently harbor 5' terminal oligopyrimidine (TOP) motifs that render their translation highly sensitive to mTORC1 (Philippe *et al.*, 2020; Stolovich *et al.*, 2002), we first investigated whether mTORC1 translation is affected in eIF3d^{KD} or eIF3e^{KD} but found no significant change in TE for any mTORC1 components. Next, we examined the effect of depletion of eIF3 subunits on translation of all known TOP mRNAs. We observed that transcripts harboring TOP motifs (Philippe *et al.*, 2020) were significantly translationally activated in eIF3d^{KD} and eIF3e^{KD}, but not in eIF3h^{KD} (Figure 4A - C). Interestingly, the effect of eIF3d^{KD} and eIF3e^{KD} on translation of non-TOP mRNAs that are mTORC1-sensitive (Gandin *et al.*, 2016) differed from what was observed for TOP mRNAs, as non-TOP transcripts were predominantly translationally offset (buffered), such that translation efficiency was increased to maintain a constant level of polysome association despite decreased total mRNA abundance (Supplementary Figure 7A - C). Together, these findings suggest that eIF3e and eIF3d may play a role in tempering translation of TOP mRNAs under normal cellular conditions, and that the loss of these eIF3 subunits has distinct impact on the translome as compared to mTORC1.

eIF3d^{KD} and eIF3e^{KD} decrease translation efficiency of the key components of the MAPK/ERK and other signaling pathways implicated in cancer

The eIF3d^{KD} or eIF3e^{KD} “all downregulated” and eIF3d^{KD}/eIF3e^{KD} “common downregulated” groups were enriched for a similar set of KEGG pathways (Figure 5A and Supplementary Figure 8A). The effect is generally stronger for eIF3d^{KD}, which simply means that this knock-down has a higher number of DTEGs in the same pathway than eIF3e^{KD} (for example “MAPK signaling pathway” has 42 DTEGs in eIF3d^{KD} vs. 30 DTEGs in eIF3e^{KD}). Therefore, we focused primarily on DTEGs identified in eIF3d^{KD} “all” (Figure 5A and Supplementary Figure 8A). Most of the top ten KEGG results are related to and/or encompass the “MAPK signaling pathway” term




Figure 4.

Loss of eIF3 subunits leads to translational activation of mRNAs with 5'UTR TOP motifs.

(A-C) Scatterplots from transcriptome analysis of eIF3d^{KD} (A), eIF3e^{KD} (B), and eIF3h^{KD} (C) with the location of transcripts harboring 5' UTR TOP motifs (Philippe *et al.*, 2020) colored (left panels). Middle and right panels show the empirical cumulative distribution functions of log₂ fold changes in FP and total mRNA for the transcripts with TOP motifs. The background constituting all other transcripts are shown as grey curves. Significant differences between the distributions were identified using the Wilcoxon rank-sum test. Differences between the distributions at each quantile are indicated. Shift to the right indicates increased expression, while shift to the left indicates decreased expression. eIF3d^{KD} and eIF3e^{KD} show very significant increase of FPs of TOP mRNAs, suggesting mainly translational upregulation.

(Supplementary Figure 8B), like “Chronic myeloid leukemia” and the “Neurotrophin signaling pathway”. Depletion of eIF3d affected multiple components of the MAPK signaling pathways, which plays a critical role in the regulation of cell proliferation (Zhang & Liu, 2002 [↗](#)) and cancer development and progression (reviewed in (Dhillon *et al.*, 2007 [↗](#))). These include the docking protein Grb2 (encoded by *GRB2*) and guanine nucleotide exchange factor Sos (*SOS1*), through kinases c-Raf (*RAF1*), MEK1 (*MAP2K1*), Erk2 (*MAPK1*), MNK1 (*MKNK1*), Rsk2 (*RPS6KA3*), to the transcription factor CREB (*ATF4*). In addition, Rac1 and PAK1, that activate c-Raf by increasing phosphorylation of Ser338 (King *et al.*, 1998 [↗](#)) were also significantly affected (**Figure 1A** [↗](#), Supplementary Figure 8C and Supplementary Spreadsheet S2). Consistent with the observed translational suppression of MNK1, which phosphorylates eIF4E (Waskiewicz *et al.*, 1997 [↗](#)), mRNAs that critically depend on phospho-eIF4E levels (Karampelias *et al.*, 2022 [↗](#)) were translationally offset in eIF3d^{KD} and eIF3e^{KD} cells, such that translation efficiency was decreased to maintain a constant level of polysome association despite increased total mRNA abundance (Supplementary Figure 9A - B).




Figure 5.

KEGG pathway enrichment analysis of the eIF3d^{KD}- and eIF3e^{KD}-associated downregulated DTEGs reveals downregulation of MAPK signaling pathways components.

(A) Venn diagram of the 10 most significantly enriched KEGG pathways for eIF3d^{KD} “all” and eIF3e^{KD} “all” groups of downregulated DTEGs, highlighting that most of the pathways are common to both knock-downs. The MAPK signaling pathway is highlighted in bold. The complete results of the KEGG enrichment analysis with corresponding *p*-values can be found in Supplementary Figure 7A. (B) The list of genes pre-selected for western blot analysis and a heatmap showing their respective log2 fold-change values from differential expression analysis of FP, TE and mRNA in eIF3d^{KD}. Negative values indicating a significant downregulation are in shades of blue, while positive values showing significant upregulation are in shades of orange. ns = not significant *p*-adjusted value. (C) Western blot analysis of selected proteins constituting the MAPK/ERK pathway performed in the indicated knock-downs and the NT control. GAPDH was used as a loading control. (D) Relative protein levels of selected MAPK/ERK pathway proteins normalized to GAPDH; NT control = 100%. Dots represent results from individual biological replicates. One sample *t*-test was used for statistical evaluation, *P*-values: * = *P* < 0.05, ** = *P* < 0.01, *** = *P* < 0.001, ns = not significant. (E) Western blot analysis of the phosphorylation status of the ERK1/2 proteins (left) and its relative quantification normalized to NT = 1 (right). One-sample *t*-test was used for statistical evaluation, **** = *P* < 0.0001. (F) Western blot analysis of the phosphorylation status of the c-Jun transcription factor in cytoplasmatic and nuclear fractions in the indicated knock-downs and the NT control. The protein loading was 50ug of total protein for cytoplasmatic lysate and 30ug of total protein for nuclear lysate, as indicated. GAPDH was used as cytoplasmatic loading control and Lamin B1 was used as nuclear loading control. This experiment was repeated 3 times with similar results. All plots in (D-E) were created in GraphPad Prism version 8.4.3 for Windows.

Consistently, Gene Ontology (GO) analysis revealed strong association of downregulated DTEGs with biological processes like “protein phosphorylation” and “regulation of transcription” and the molecular function term “protein serine/threonine kinase activity” (Supplementary Figure 10A – B). Key integrin-mediated signaling proteins that were implicated in cancer cell plasticity, adhesion, migration and survival (reviewed in (Cooper & Giancotti, 2019 [\[1\]](#))) also had significantly decreased TE, specifically, proto-oncogenes Crk (*CRK*) and Crk-L (*CRKL*), *PRKCA* – a catalytic subunit of PI3-kinase, Rac1 GTPase (*RAC1*), JNK2 (*MAPK9*), as well as Abl1 kinase (*ABL1*) and BCR activator of Rho-GEF (*BCR*). The TGF-beta signaling, important for a wide range of cellular processes as well as for tumor cell invasion, metastasis, and resistance to chemotherapy (reviewed in (Colak & Ten Dijke, 2017 [\[2\]](#))), was affected at the level of the intracellular transducer Smad4 (*SMAD4*) and receptor activated SMADs, namely Smad3 (*SMAD3*) and Smad5 (*SMAD5*) and TGF beta (*TGFB3*) itself. The NFkB pathway, a regulator of immune and inflammatory responses (Liu *et al*, 2017 [\[3\]](#)), also implicated in cancer cell proliferation, survival, angiogenesis and plasticity (reviewed in (Xia *et al*, 2014 [\[4\]](#))), was downregulated at the level of NFkB (*NFKB1*) itself and IKK kinase (*CHUK*). Notably, a direct link between activation of NFkB and eIF3 has very recently been described (She *et al*, 2023 [\[5\]](#)). Finally, the canonical Wnt signaling pathway, which shapes the stem cells-like properties of cancer cells and boosts proliferation (reviewed in (Zhang & Wang, 2020 [\[6\]](#))), was downregulated at the level of a protein Dishevelled (*DVL3*) and GSK3 beta kinase (*GSK3B*). Physiological relevance of all these hits remains to be seen; for the complete list of all DTEGs, please see Supplementary Spreadsheet S2.

eIF3d^{KD} and eIF3e^{KD} reduce protein levels of key components of the activated MAPK/ERK pathway

Interestingly, more than half (26 of the 42) downregulated DTEGs belonging to the “MAPK signaling pathway” can be classified as translationally buffered; i.e., their mRNA levels increase while their TE decreases (Kusnadi *et al.*, 2022 [DOI](#)). However, this is not a typical case of translational buffering because mRNA upregulation is not only nullified by translational repression, but completely overruled, so that the total number of FPs is also reduced. To verify that the reduction in TE and FP counts is indeed manifested by a reduction in cellular protein levels, we selected a set of genes with different combinations of changes in TE or FP or mRNA levels, including translationally exclusive and translationally buffered genes, as well as genes with unchanged TE, and subjected them to western blotting (**Figure 5B - [DOI](#)**).

We found, that AKT1, GRB2, MAP2K1 (MEK1), PAK1, RPS6KA1 (RSK1), RPS6KA3 (RSK2), RAF1 and SOS1 had visibly decreased protein levels in eIF3d^{KD} and most of them also in eIF3e^{KD}. An interesting exception is the transcription factor c-Jun, which showed increased mRNA and FP levels, unchanged TE, but robust increases in protein levels in all three knock-downs (**Figure 5C - [DOI](#)**). The mRNA levels of all selected genes were verified by qPCR, showing a good correlation with our sequencing data; whereas PAK1 showed significant decrease, JUN mRNA was significantly increased in both eIF3d^{KD} and eIF3e^{KD} (Supplementary Figure 10C). To assess the functional consequences of the downregulation of MAPK/ERK pathway components, we examined the phosphorylation status of effector kinases ERK1/2, which directly phosphorylate several transcription factors including c-Jun (Leppa *et al.*, 1998 [DOI](#)). As with other MAPK proteins, ERK1/2 expression was reduced in eIF3d^{KD} and eIF3e^{KD}, but its phosphorylation at activation sites Thr202/Tyr204 was unexpectedly increased in all three knock-downs (**Figure 5E [DOI](#)**). Next, we examined the phosphorylation status of the upregulated c-Jun protein. As phosphorylated c-Jun translocates to nucleus to activate transcription of specific genes, we performed western blot of both cytoplasmatic and nuclear fractions. In accord with increased ERK1/2 phosphorylation, c-Jun was found to be activated by phosphorylation at Ser63 and largely accumulated in the nucleus (**Figure 5F [DOI](#)**). Thus, despite reduced MAPK/ERK pathway protein levels, MAPK/ERK signaling appears to be activated when eIF3 function in translation is compromised.

Intact eIF3 contributes to translational regulation of ATF4 and MDM2

Knocking down eIF3h generates an eIF3 subcomplex lacking eIF3h and two nonessential subunits eIF3k and eIF3l (**Figure 1B [DOI](#)**), having only negligible impact on overall translation. Accordingly, we identified 19 transcripts whose TE was decreased and only 2 transcripts whose TE was increased compared to the NT control (**Figure 2D [DOI](#)**). Given the low number of DTEGs, we observed no significant enrichment of any KEGG pathway within the upregulated DTEGs and only a few significant enrichments for the eIF3h^{KD} “all downregulated” DTEGs in pathways like “Prostate cancer”, “Viral carcinogenesis” or “PI3-Akt signaling pathway”, all of which included only two common hits – MDM2 and ATF4 (**Figure 6A [DOI](#)**, Supplementary Table S4). Additionally, we also noticed enrichment of GO terms associated with ubiquitination (**Figure 6B [DOI](#)**, Supplementary Table S5), which was mainly due to the MDM2, WAC, RNF10, RMND5A, USP38 and ZRANB1 genes, all of which showed decreased TE. MDM2, WAC, RNF10 and RMND5A are all E3 ubiquitin ligases while USP38 and ZRANB1 are de-ubiquitinases. Note that all of the above downregulated DTEGs are common for all three knock-downs.

Perhaps the most interesting of these genes is the proto-oncogene MDM2, an E3 ubiquitin ligase that acts as a negative regulator of the p53 protein (Marine & Lozano, 2010 [DOI](#)), whose overexpression has been reported in many different cancer types (Rayburn *et al.*, 2005 [DOI](#)). We did confirm that the MDM2 protein levels were reduced in all three knock-downs (**Figure 6C [DOI](#)**).




Figure 6.

KEGG pathway and GO enrichment analysis of the eIF3h^{KD}-associated DTEGs reveals downregulation of proto-oncogene MDM2 and a defective stress-induced upregulation of ATF4.

(A) The bar chart shows the top 10 enriched KEGG terms for eIF3h^{KD} “all downregulated” DTEGs. Blue bars correspond to terms with significant p -values (< 0.05), grey bars correspond to terms with not significant p -values (> 0.05). The p -values were calculated by Enrichr gene set search engine. (B) The bar chart shows the top 10 enriched GO Biological Process terms for eIF3h^{KD} “all downregulated” DTEGs. Blue bars correspond to terms with significant p -values (< 0.05). The p -values were calculated by Enrichr gene set search engine. PCP; protein catabolic process. (C) Western blot analysis of the MDM2 expression performed in the indicated knock-downs and NT control. GAPDH was used as a loading control; NT control = 100%. Dots represent results from individual biological replicates. One sample t-test was used for statistical evaluation, P -values: * = $P < 0.05$, ** = $P < 0.01$. Plots was created in GraphPad Prism version 8.4.3 for Windows. (D) Western blot analysis of the stress-induced upregulation of ATF4 expression performed in the indicated knock-downs and NT control. Before harvesting, cells were incubated for 3 hours with 1 μ M Thapsigargin to induce ER stress or with DMSO as a stress-free control. GAPDH was used as a loading control. (E) Normalized ribosomal footprint coverage along the ATF4 mRNA (first 450 nucleotides). Schematic with regulatory elements is shown at the bottom. Average of all three replicates is shown. Footprint coverage was normalized to all footprints mapping to the ATF4 mRNA. (F) Same as in (E) only for MDM2 mRNA.




Figure 8.

Differential TE transcripts in eIF3d^{KD} and eIF3e^{KD} show negative correlation with the UTR length and uORF content while they positively correlate with GC content of their 3' UTRs and coding sequences (CDS).

(A) Bar plot showing the Spearman correlation between the observed ΔTE values for all genes with assigned adjusted p -values in each knock-down and different mRNA features (5' and 3' UTR length or GC content, uORF content in 5' UTR). *** = $P < 10^{-20}$, ** = $P < 10^{-10}$, * = $P < 0.005$. (B-C) Box and whisker plots comparing UTR lengths or GC content (in UTRs or CDS) among mRNAs with TE significantly increased (red, upregulated), decreased (blue, downregulated), or unchanged (white) in eIF3d^{KD} (B) and eIF3e^{KD} (C); *** = $P_{adj} < 10^{-10}$, ** = $P_{adj} < 10^{-5}$, * = $P_{adj} < 0.05$, color indicates comparison set. (D) Same as in (B-C) but for the number of AUG-initiated uORFs in 5' UTRs in the eIF3d, eIF3e and eIF3h knock-downs. Outliers are not shown for better clarity. (E) Histogram of the frequency of AUG-initiated uORFs per 5' UTR in all transcripts listed in the uORFdb database that were assigned a p -value in this study ($n=11450$), for downregulated translation only DTEGs in eIF3d^{KD} ($n=1027$) and for downregulated translation only DTEGs in eIF3e^{KD} ($n=618$). (F) Venn diagram summarizing the main results of this study. Groups of significant DTEGs identified in given knock-downs (encircled – color-coded) are indicated by a blue arrow for downregulated or an orange arrow for upregulated. Results confirmed by western blotting are shown in bold. Ribosomal proteins were placed on the borderline between eIF3d^{KD} and eIF3e^{KD} because they were identified as eIF3e^{KD} “unique upregulated” DTEGs, but their protein levels were elevated in both eIF3e^{KD} and eIF3d^{KD}, as shown by western blotting. The “MAPK/ERK signaling” is shown in bold italics because it is an independent phenomenon that could be detected only by western blotting.

An equally interesting DTEG is ATF4, which is among the top 10 most significant DTEGs in all three knock-downs (**Figure 2B - D**). In eIF3h^{KD}, it can be classified as a translationally buffered gene because it has strongly upregulated mRNA levels, but this increase is not accompanied by a concomitant increase in the FP count, thereby resulting in decreased TE. In eIF3d^{KD} and eIF3e^{KD}, it is even significantly translationally repressed (FPs are decreased). Using qPCR, we confirmed ATF4 mRNA upregulation in all three knock-downs (Supplementary Figure 11A). Because ATF4 protein levels are on the verge of the western blot detection limit under non-stress conditions, we examined whether all three knock-downs tested had any effect on ATF4 induction upon 3 hour-long ER stress provoked by 1μM Thapsigargin (Harding *et al*, 2000). Indeed, stress-mediated ATF4 induction was almost abolished in eIF3d^{KD} and eIF3e^{KD}, and strongly decreased in eIF3h^{KD} (**Figure 6D**). This differential effect correlates well with our sequencing data, where the decrease in TE was also stronger in eIF3d^{KD} and eIF3e^{KD}. The ATF4 protein levels in DMSO control are shown as a no stress reference. Overall, these findings further confirm that an intact eIF3 complex is required for proper translational control of ATF4, as we and others have previously suggested (Guan *et al*, 2017; Hronova *et al*, 2017; Mukhopadhyay *et al*, 2023; Shu *et al*, 2022; Wagner *et al.*, 2020), and as discussed below.

As both *ATF4* and *MDM2* are known to contain uORFs in their 5' UTR that regulate their translation under stress vs. non-stress conditions (Akulich *et al*, 2019; Dey *et al*, 2010; Lu *et al*, 2004; Smirnova *et al*, 2024; Vatter & Wek, 2004), we examined the footprint coverage in the 5' UTR of these genes in more detail. In the case of *ATF4*, a markedly increased accumulation of footprints was observed at the most 5' Start-stop (St-st) element, specifically in eIF3d^{KD} and eIF3e^{KD}, while coverage of uORF1 decreased, and so did also the basal level of uORF2 (modestly) and *ATF4* (**Figure 6E**). This agrees well with our observations that ribosomes are stalled at this St-st element (Rendleman *et al*, 2023; Wagner *et al.*, 2020) and further suggests that both eIF3e and eIF3d are required to clear these St-st-trapped ribosomes. Regarding *MDM2*, footprint coverage was increased specifically for uORF1 in all three knock-downs (**Figure 6F**), consistent with its several-fold stronger inhibitory role on *MDM2* expression compared to uORF2 (Jin *et al*, 2003), which seems to be magnified in the absence of fully functional eIF3. Collectively, these results are in line with the well-established role of eIF3 in translation reinitiation (reviewed in (Valasek *et al.*, 2017)). In addition, we found similarly accumulated footprints also at uORF1 and uORF2 of the *RAF1* kinase, one of the downregulated components of the MAPK pathway (Supplementary Figure 11B).

Last but not least, eIF3d was shown to mediate mechanism of alternative cap-dependent translation initiation in conjunction with DAP5 (p97/NAT1) (Alard *et al*, 2023; de la Parra *et al.*, 2018a; Volta *et al*, 2021) and, importantly, DAP5-dependent mRNAs also frequently contain uORFs (David *et al*, 2022). Consistent with this, the effects of depletion of DAP5 on the translome were comparable to those observed in eIF3d-depleted cells (Supplementary Figure 12A - C); i.e., no changes in mRNA levels and the FP coverage moving in the same direction, either up or down. Since similar effects were also observed with eIF3e^{KD} and eIF3h^{KD}, it seems that the DAP5-dependent mechanism may rely upon not only eIF3d but also on the entire eIF3 complex.

Presence of uORFs, UTR length and GC content correlate with the change in translation efficiency

eIF3 is critically required for mRNA recruitment to the 40S ribosomal subunit (reviewed in (Valasek *et al.*, 2017)) and several eIF3 subunits have been proposed to regulate translation of specific subsets of mRNAs (Lee *et al.*, 2015; Lee *et al*, 2016). We therefore asked whether the changes in TE that were observed in the eIF3d, eIF3e or eIF3h knock-downs correlate with specific mRNA properties. Indeed, in all three knock-downs, the change in TE was negatively correlated with the length of 5' and 3' UTR, and with the presence of uORFs in 5' UTR (Spearman correlation, $P < 1 \times 10^{-45}$) (Figure 7A). Moreover, we found a significant positive Spearman correlation between TE changes and the GC content in the coding region (CDS) and 3' UTRs.

To explore these correlations in more detail, we focused only on those mRNAs that are exclusively regulated at the translational level in each knock-down and therefore are more likely to contain distinct features that facilitate their translational regulation. We divided them into those with significantly changed TE (decreased and increased) and those with unchanged TE and searched for significant differences in the median values of the tested traits (Figure 7B - C). In accordance with the Spearman correlation, in eIF3d^{KD} and eIF3e^{KD}, mRNAs of downregulated DTEGs had longer 5' UTRs, while upregulated DTEGs possessed shorter 5' UTRs, when compared to mRNAs with unchanged TE. The same trend is seen for the 3' UTR length in eIF3e^{KD}, but with lower significance. In case of eIF3d^{KD}, only the downregulated DTEGs showed a significant change in 3' UTR length, but not DTEGs that were upregulated.

In terms of GC content, while we observed little significant difference in GC content in the 5' UTR of eIF3d^{KD} and eIF3e^{KD}, there was a highly significant difference in the CDS and 3' UTR of these knock-downs in both downregulated and upregulated DTEGs, indicating that mRNAs with increased TE tend to have more GC-rich 3' UTRs and, surprisingly, CDS and vice versa. For eIF3h^{KD}, we only observed differences of low significance due to the low number of DTEGs that were only translationally regulated (only 8 downregulated and 1 upregulated) (data not shown).

In summary, mRNAs with decreased TE in cells wherein eIF3d and eIF3e were depleted exhibited on average longer 5' and 3' UTRs and lower GC content in the CDS and 3' UTRs. In turn, mRNAs with increased TE have on average shorter 5' UTRs and a higher GC content in the CDS and 3' UTRs.

We next asked whether 5' UTRs of upregulated or downregulated DTEGs contain any specific sequence motifs or short upstream open reading frames (uORFs). We did not identify any specific sequence motif, but expectedly, the 5' UTRs of “downregulated translation only” DTEGs have a strong tendency to contain uORFs starting with AUG (Figure 7D - E). These findings suggest that the intact eIF3 complex is primarily required for translational control of subset of mRNAs containing short uORFs in their 5' UTR even under normal cellular conditions, like for example *MDM2*, *ATF4* or *RAF1*, which is in line with its role in translation reinitiation (Valasek *et al.*, 2017), and further discussed below.

Discussion

In this study, we interrogated the effects of altering composition of eIF3 by depleting three eIF3 subunits in HeLa cells, followed by RNA-Seq and Ribo-Seq to elucidate their role in translational regulation of specific mRNAs by differential expression analysis. We focused on differential translation efficiency genes (DTEGs) and identified several distinct groups of DTEGs associated with individual eIF3 knock-downs.

Interestingly, while DTEGs identified in eIF3d^{KD} and eIF3e^{KD} largely overlapped, considerably more DTEGs were identified in eIF3d^{KD} than in eIF3e^{KD}, correlating well with the severity of the growth phenotype of both knock-downs, with the former being more detrimental despite not affecting eIF3 integrity (Wagner *et al.*, 2016). The higher number of DTEGs identified in the same pathways in eIF3d^{KD} vs. eIF3e^{KD} could be explained by differences in eIF3d protein levels between the two knock-downs. Despite being markedly co-downregulated in eIF3e^{KD}, eIF3d levels were more dramatically reduced in eIF3d^{KD} compared to eIF3e^{KD}, as expected. Furthermore, eIF3d is a peripheral subunit that binds to the head of the 40S subunit (mainly via the small ribosomal protein Rps16 (Herrmannová *et al.*, 2020)) and is connected to the eIF3 octamer not only via eIF3e but also via the eIF3a and c subunits (Bochler *et al.*, 2020). Therefore, we hypothesize that eIF3d may remain partially operational even in the absence of eIF3e, and therefore the more

severe phenotype in eIF3d^{KD} is likely to stem from lower levels of the eIF3d protein compared to eIF3e^{KD}. Following this logic, phenotypes that are common to eIF3d^{KD} and eIF3e^{KD} are likely a consequence of reduction in eIF3d protein levels.

Perhaps the most intriguing finding is that eIF3d^{KD} and, to a large extent, also eIF3e^{KD} result in translational repression of a marked number of the MAPK pathways constituents, significantly reducing their protein levels, despite the apparent compensatory mechanisms that increase their mRNA levels. Somewhat unexpectedly, despite the downregulation of a number of its components, eIF3d^{KD} and eIF3e^{KD} but also eIF3h^{KD} appear to activate MAPK/ERK pathway as evidenced by the elevated ERK1/2 and c-Jun phosphorylation. Notably, the eIF3d influence on the MAPK/ERK pathway was previously demonstrated in HTR-8/SVneo cells, where overexpression of eIF3d led to hypo-phosphorylation of MEK1 and ERK1/2, while downregulation had the opposite effect; however, no significant effect on the protein levels of its components was detected in this study (Li *et al.*, 2021 [\[1\]](#)). Importantly, since the increased ERK1/2 phosphorylation was observed not only in eIF3d^{KD} and eIF3e^{KD} but also in eIF3h^{KD}, where expression of MAPK pathways proteins is unaffected, we conclude that activation of the MAPK/ERK pathway is a general consequence of the compromised eIF3 function.

Expression of the transcription factor c-Jun is partially controlled by the MAPK/ERK pathway and, consistently, we found that it is strongly upregulated in all three knock-downs. Moreover, we observed that all three knockdowns activated c-Jun as evidenced by its increased nuclear translocation. Although *JUN* mRNA was previously shown to interact directly with eIF3 through a stable stem loop occurring in its 5' UTR, which positively stimulated its translation (Lee *et al.*, 2015 [\[2\]](#)), in this study we observed upregulation of the *JUN* mRNA abundance, whereby the depletion of eIF3d, e, or h had no conspicuous effect on its translational efficiency. This can be explained by a recent report showing that eIF3d binds to *JUN* mRNA with increased affinity during chronic ER stress, suggesting that eIF3d-dependent translational regulation may be triggered by stress (Mukhopadhyay *et al.*, 2023 [\[3\]](#)). Because c-Jun is a component of the AP1 transcription factor complex, wherein it either homodimerizes or heterodimerizes with other AP1 proteins, we examined the effects of eIF3 subunit depletion on mRNA, FP and TE trends of other AP1 family members. Similarly to c-Jun, only FOSL1 and ATF3 showed increased mRNA levels and FP counts with no net changes in TE, whereas other members of the JUN (JUNB and JUND) and FOS (FOS, FOSB, FOSL2) families remained unaltered. In addition, the ATF family members ATF2 and ATF4 were even downregulated (Supplementary Spreadsheet S2). Notwithstanding that our data suggest that impairment of eIF3 function may be compensated by the activation of the MAPK/ERK pathway, further research is required to unravel the full spectrum of changes associated with the MAPK pathways in eIF3-impaired cells.

Recently, Lin *et al.* carried out Ribo-seq analysis in MCF-10A cells downregulated for eIF3e (Lin *et al.*, 2020 [\[4\]](#)). They identified 240 downregulated and 220 upregulated eIF3e^{KD}-specific DTEGs that overlap very poorly with our DTEGs (data not shown). In addition, they identified a set of 2683 eIF3e-dependent mRNAs encoding many proteins associated with mitochondrial and membrane functions that showed increased ribosome density between codons 25 and 75 in eIF3e^{KD}. Those mRNAs were therefore proposed to require eIF3e to recruit chaperons to promote their early elongation. While this set of genes displayed a significant overlap with our DTEGs (see below), we did not observe a similar phenomenon (a specifically increased ribosome density early on) in our datasets. In metagene analysis, we did observe some accumulation of footprints at the beginning of the CDS, but this was present in all samples, including NT, whereby peaks were located around codon 20 downstream of the translation initiation sites (Supplementary Figure 13A).

Nonetheless, still considering that the “accumulation effect” observed by Lin *et al.*, most probably resulting from ribosome stalling and/or slower elongation rates, could artificially increase FPs and TE without actually increasing synthesis of the corresponding proteins in our datasets, we trimmed the first 225 nt (75 codons) from all mRNA's CDSs, and repeated the differential

expression analysis in the same way. When compared, the two DE analyses showed approximately 80% - 90% identical DTEGs and high correlation (Spearman correlation coefficient of eIF3d^{KD}: 0.924, eIF3e^{KD}: 0.925, eIF3h^{KD}: 0.834), suggesting that our DE analysis is not biased by ribosome accumulation in the first 75 codons and reflects true changes in translation efficiency.

Despite this “accumulation effect” discrepancy, comparing our DTEGs with the Lin et al. eIF3e-dependent data, we found a very significant overlap of 446 genes between the eIF3e-dependent mRNAs and our eIF3d^{KD} “all upregulated” dataset ($P < 1 \times 10^{-53}$), while the overlap with eIF3e^{KD} “all upregulated” – 251 genes – was surprisingly smaller ($P < 1 \times 10^{-24}$). Moreover, our KEGG enrichment analysis of eIF3e-dependent mRNAs from Lin et al. revealed the same top 2 KEGG pathways (data not shown) as in case of eIF3d^{KD} and eIF3e^{KD} “all upregulated” groups; i.e. Lysosome and Protein processing in ER (**Figure 3A** and Supplementary Figure 6A). Considering that eIF3d was co-depleted with eIF3e in the eIF3e^{KD} of Lin et al., it is conceivable that not eIF3e but the eIF3d subunit (or cooperatively both subunits together) ensures loading of quality control factors to promote elongation of transmembrane proteins. In any case, resolving this issue will require further research.

Based on our analysis, eIF3e seems to be primarily involved in controlling the balanced production of mature ribosomes, as its loss resulted in significant number of “unique upregulated” DTEGs with strong enrichment for the KEGG Ribosome and GO Translation and Ribosome biogenesis terms (**Figure 3C** and Supplementary Figure 6B). Focusing directly on RPs, nearly twice as many RPs with significantly altered TEs were identified in eIF3e^{KD} compared to eIF3d^{KD} (32 vs. 16) and a similar pattern was also evident at the FP level (91 vs. 64). However, despite this obvious difference, RPs were increased similarly in both eIF3d^{KD} and eIF3e^{KD} (**Figure 3E - F** and **3H**). This discrepancy between our Ribo-Seq and Western blot results in eIF3e^{KD} versus eIF3d^{KD} (**Figure 3D vs. 3E**) might be explained by the enhanced stabilization of RPs by the previously described compensatory dosage mechanism at the protein level (Ishikawa, 2021). In particular, to preserve the stoichiometry of the ribosome when numerous RPs are translationally upregulated, the rest of the RPs would be protected from degradation. This speculation can be supported by the impact of all knock-downs on translation efficiency of different E3 ubiquitin ligases, as described for eIF3h^{KD} (Supplementary Table S5). Similar effect of increased RPs production was recently reported also for eIF3k^{KD} (Duan *et al.*, 2023). Of note, eIF3e^{KD} also decreases protein levels of eIF3k and eIF3l and thus this RP-specific effect could be at least partially attributed to the loss of eIF3k. However, increased RP production was also observed in eIF3d^{KD}, where eIF3k is normally present, but was not observed in eIF3h^{KD}, where eIF3k and eIF3l are also co-depleted. Thus, although it now seems well-established that the eIF3 complex exerts control over RP production, the exact contribution of its individual subunits to this regulation awaits further analysis.

An important hint could be that all cytoplasmic RPs with increased TE in eIF3e^{KD} (and most of mitochondrial RPs) were upregulated solely at the translational level (their mRNA levels did not change), which may well correspond to the fact that RP mRNAs belong to the TOP mRNAs family. These mRNAs contain a polypyrimidine tract in their 5' UTR and were suggested to be regulated solely at the translational level in response to stress or unfavorable growth conditions through mTORC1 or PI3-kinase pathways (Cockman *et al.*, 2020; Philippe *et al.*, 2020; Stolovich *et al.*, 2002). Besides RPs, this family also includes some translational initiation and elongation factors (for the list of TOP mRNAs see Supplementary Spreadsheet S2). Accordingly, we found that TOP transcripts were significantly translationally activated in eIF3d^{KD} and eIF3e^{KD} (**Figure 4A - C**), suggesting a role of eIF3e and eIF3d in repressing translation of TOP mRNAs under normal cellular conditions. Interestingly, since mTOR-activated translation of non-TOP transcripts was predominantly translationally offset (buffered) in eIF3d^{KD} and eIF3e^{KD} cells (Supplementary Figure 7A - C), it appears that both effects are independent of each other; i.e., mTOR does not generally operate *via* eIF3 and eIF3 does not require mTOR activation to fine tune RP production.

It is also noteworthy that upregulation of cytosolic ribosomal proteins has previously been reported in association with the eIF3m downregulation – albeit only at transcriptional level (Smekalova *et al.*, 2020 [DOI](#)), or with downregulation of eEF2 (Gerashchenko *et al.*, 2020 [DOI](#)).

In addition, as we were unable to find any specific sequence motif in the 5' UTR's of the mRNAs whose TE was affected in eIF3 subunit-depleted cells, we cross-referenced our eIF3d^{KD} “all” and eIF3e^{KD} “all” groups with transcripts that physically interact with eIF3 as reported by (Lee *et al.*, 2015 [DOI](#)). We found that out of the 402 transcripts with a putative eIF3 binding sites in their 5' UTRs, 92 and 65 had significantly altered TE in the eIF3d^{KD} and eIF3e^{KD}, respectively (51 were common for both eIF3d^{KD} and eIF3e^{KD}) (Supplementary Figure 13B). This notable overlap involving for example the MAPK1 gene further underscores the robustness of our screening approach.

Among the few downregulated DTEGs in eIF3h^{KD}, we found *ATF4* and *MDM2* mRNAs, whose translation is tightly regulated by the presence of uORFs in their 5' UTRs in response to specific stress conditions (Akulich *et al.*, 2019 [DOI](#); Dey *et al.*, 2010 [DOI](#); Jin *et al.*, 2003 [DOI](#); Smirnova *et al.*, 2024 [DOI](#); Vattam & Wek, 2004 [DOI](#)). Here, we showed that these mRNAs were translationally downregulated in our knock-downs under non-stress conditions and, accordingly, their most 5' short uORFs (or St-st in the case of *ATF4*) accumulated FPs, most probably slowing down the flow of ribosomes downstream. The fact that transcripts of these genes contain uORFs in their 5' UTRs also correlates well with our finding that 5' UTRs of “downregulated translation only” DTEGs have a strong tendency to contain AUG-initiated uORFs (Figure 7D - E). Noteworthy, the impaired *ATF4* induction in our eIF3 knock-downs under ER stress (Figure 6D [DOI](#)), was reported before also for knock-downs of eIF3a, eIF3c, eIF3d, eIF3g, but not eIF3l (Guan *et al.*, 2017 [DOI](#)). Moreover, eIF3h and specifically modified eIF3a have been directly shown to modulate resumption of scanning from *ATF4*'s uORF1 for efficient reinitiation downstream; i.e. one of the key steps in the *ATF4* translational control (Hronova *et al.*, 2017 [DOI](#); Shu *et al.*, 2022 [DOI](#)). Therefore, we propose that the intact eIF3 complex is required for stimulation of reinitiation on uORF-containing mRNAs, such as *ATF4* and *MDM2* to ensure their efficient translation on demand. Interestingly, the impact on translation of *MDM2* mRNA could indicate that eIF3 is also linked to the key tumor suppressor p53 (Marine & Lozano, 2010 [DOI](#)).

Among a few upregulated DTEGs in eIF3h^{KD}, we noticed PRRC2A, which was significantly upregulated in all three knock-downs and was recently identified by the Teleman's group as a protein interacting with the translational machinery, including eIF3. Interestingly, they found that PRRC2 proteins affected translation initiation by promoting leaky scanning (Bohlen *et al.*, 2023 [DOI](#)) and that the triple knock-down of all PRRC2 proteins (PRRC2A, PRRC2B, PRRC2C) specifically decreased translation efficiency of mRNAs harboring uORFs in their 5' UTRs, including *ATF4* and *RAF1* studied also here, similar to our eIF3 knock-downs. They concluded that uORFs are a key element determining PRRC2-dependence. Thus, we speculate that by increased expression of PRRC2A (and also PRRC2B and PRRC2C in eIF3d^{KD} and eIF3e^{KD}), cells attempt to compensate for ineffective translation of mRNAs with short uORFs, which often encode regulatory proteins. Intriguingly, all PRRC2 proteins themselves have a highly translated overlapping-uORF in their 5' UTR, suggesting that they might be a subject to an autoregulatory mechanism.

Although our results presented here and summarized in Figure 7F are limited to HeLa cells and future work is needed to confirm their universal validity in other cells and tissues, a growing body of evidence suggests that eIF3 plays a critical role in translational regulation of specific mRNAs encoding regulatory proteins, very often utilizing uORFs and/or specific secondary structures in their 5' UTRs. Here, we demonstrate that alterations in the eIF3 subunit stoichiometry and/or eIF3 subcomplexes have distinct effects on the translome; for example, they affect factors that play a prominent (either positive or negative) role in cancer biology (e.g., *MDM2* and *cJUN*), but the resulting impact is unclear so far. Considering the complex interactions between these factors as well as the complexity of the eIF3 complex *per se*, future studies are required to delineate the

specific oncogenic and tumor suppressive pathways that play a predominant role in mediating the effects of perturbations in the eIF3 complex in the context of neoplasia. Taking into account that malignant cells exhibit augmented activity of many components of the translation machinery and are inherently believed to become “addicted” to elevated protein synthesis (Ruggero, 2013 [↗](#); Silvera *et al*, 2010 [↗](#)), this type of research is important because it is increasingly apparent that targeting general components of the canonical translational machinery, such as eIF3, which acts upstream of effector proteins directly involved in tumorigenesis, may hold promise for overcoming the major obstacle associated with intratumor heterogeneity.

Acknowledgements

We are thankful to Olga Krýdová for technical and administrative assistance and to all lab members for fruitful discussions. We are also thankful to Jonathan Bohlen for sharing and discussing results of PRRC2 proteins downregulation. This work was supported by a Grant of Excellence in Basic Research (EXPRO 2019) provided by the Czech Science Foundation (19-25821X), the Praemium Academiae grant provided by the Czech Academy of Sciences, and CZ.02.01.01/00/22_008/0004575 RNA for therapy by ERDF and MEYS (all to L.S.V.), Czech Science Foundation grant 19-08013S (to T.V), ELIXIR CZ research infrastructure project (MEYS Grant No: LM2023055) including access to computing and storage facilities, by National Institute for Cancer Research (Programme EXCELES, ID Project No. LX22NPO5102) funded by the European Union - Next Generation EU (to J.B.) and by Fonds de Recherche du Québec – Santé (FRQS) Senior Investigator award (awarded to I.T.).

Authors contributions

A.H. and L.S.V. conceived and designed the project.

A.H., J.J., K.P., F.K., T.V., K.W. and M.P.M. carried out all experiments and analyzed the data.

A.H., J.J. and L.S.V. interpreted results and wrote the paper with input from K.P., F.K., T.V., J.B., M.P.M., K.W., I.T. and S.W.

Declaration of interest

The authors declare no competing interests.

Supplemental information

Supplemental Information can be found online at

Data availability

The datasets generated during this study were deposited in the GEO database with the following accession number: GSE216967. All custom programs source code are published at <https://github.com/cas-bioinf/RibosomeProfiling> [↗](#).

References

1. Akulich KA, Sinitcyn PG, Makeeva DS, Andreev DE, Terenin IM, Anisimova AS, Shatsky IN, Dmitriev SE (2019) **A novel uORF-based regulatory mechanism controls translation of the human MDM2 and eIF2D mRNAs during stress** *Biochimie* **157**:92–101
2. Alard A, Katsara O, Rios-Fuller T, Parra C, Ozerdem U, Ernlund A, Schneider RJ (2023) **Breast cancer cell mesenchymal transition and metastasis directed by DAP5/eIF3d-mediated selective mRNA translation** *Cell Rep* **42**
3. Anders S, Pyl PT, Huber W (2015) **HTSeq--a Python framework to work with high-throughput sequencing data** *Bioinformatics* **31**:166–169
4. Andrews S (2010) **FastQC: A quality control tool for high throughput sequence data**
5. Ashburner M *et al.* (2000) **Gene ontology: tool for the unification of biology. The Gene Ontology Consortium** *Nat Genet* **25**:25–29
6. Beznosková P, Cuchalová L, Wagner S, Shoemaker CJ, Gunišová S, Von der Haar T, Valášek LS (2013) **Translation initiation factors eIF3 and HCR1 control translation termination and stop codon read-through in yeast cells** *PLoS Genet* **9**
7. Beznosková P, Wagner S, Jansen ME, von der Haar T, Valášek LS (2015) **Translation initiation factor eIF3 promotes programmed stop codon readthrough** *Nucleic Acids Res* **43**:5099–5111
8. Bhat M, Robichaud N, Hulea L, Sonenberg N, Pelletier J, Topisirovic I (2015) **Targeting the translation machinery in cancer** *Nat Rev Drug Discov* **14**:261–278
9. Blankenberg D, Von Kuster G, Bouvier E, Baker D, Afgan E, Stoler N, Galaxy T, Taylor J, Nekrutenko A (2014) **Dissemination of scientific software with Galaxy ToolShed** *Genome Biol* **15**
10. Bochler A, Querido JB, Prilepskaja T, Soufari H, Simonetti A, Del Cistia ML, Kuhn L, Ribeiro AR, Valasek LS, Hashem Y (2020) **Structural Differences in Translation Initiation between Pathogenic Trypanosomatids and Their Mammalian Hosts** *Cell Rep* **33**
11. Bohlen J, Roiuk M, Neff M, Teleman AA (2023) **PRRC2 proteins impact translation initiation by promoting leaky scanning** *Nucleic Acids Res* **51**:3391–3409
12. Querido J Brito, Sokabe M, Kraatz S, Gordiyenko Y, Skehel JM, Fraser CS, Ramakrishnan V (2020) **Structure of a human 48S translational initiation complex** *Science* **369**:1220–1227
13. Buxade M, Parra-Palau JL, Proud CG (2008) **The Mnks: MAP kinase-interacting kinases (MAP kinase signal-integrating kinases)** *Front Biosci* **13**:5359–5373
14. Calviello L, WSydow D, Harnett D, Ohler U (2019) **Ribo-seQC: comprehensive analysis of cytoplasmic and organellar ribosome profiling data** *bioRxiv*
15. Cargnello M, Roux PP (2011) **Activation and function of the MAPKs and their substrates, the MAPK-activated protein kinases** *Microbiol Mol Biol Rev* **75**:50–83

16. Carriere A, Ray H, Blenis J, Roux PP (2008) **The RSK factors of activating the Ras/MAPK signaling cascade** *Front Biosci* **13**:4258–4275
17. Chan PP, Lowe TM (2016) **GtRNAdb 2.0: an expanded database of transfer RNA genes identified in complete and draft genomes** *Nucleic Acids Res* **44**:D184–189
18. Chothani S, Adami E, Ouyang JF, Viswanathan S, Hubner N, Cook SA, Schafer S, Rackham OJL (2019) **deltaTE: Detection of Translationally Regulated Genes by Integrative Analysis of Ribo-seq and RNA-seq Data** *Curr Protoc Mol Biol* **129**
19. Cockman E, Anderson P, Ivanov P (2020) **TOP mRNPs: Molecular Mechanisms and Principles of Regulation** *Biomolecules*
20. Colak S, Ten Dijke P (2017) **Targeting TGF-beta Signaling in Cancer** *Trends Cancer* **3**:56–71
21. Cooper J, Giancotti FG (2019) **Integrin Signaling in Cancer: Mechanotransduction, Stemness, Epithelial Plasticity, and Therapeutic Resistance** *Cancer Cell* **35**:347–367
22. Danecek P *et al.* (2021) **Twelve years of SAMtools and BCFtools** *Gigascience*
23. David M *et al.* (2022) **DAP5 drives translation of specific mRNA targets with upstream ORFs in human embryonic stem cells** *RNA* **28**:1325–1336
24. De Benedetti A, Harris AL (1999) **eIF4E expression in tumors: its possible role in progression of malignancies** *Int J Biochem Cell Biol* **31**:59–72
25. de la Parra C, Ernlund A, Alard A, Ruggles K, Ueberheide B, Schneider RJ (2018) **A widespread alternate form of cap-dependent mRNA translation initiation** *Nat Commun* **9**
26. de la Parra C, Walters BA, Geter P, Schneider RJ (2018) **Translation initiation factors and their relevance in cancer** *Curr Opin Genet Dev* **48**:82–88
27. des Georges A, Dhote V, Kuhn L, Hellen CU, Pestova TV, Frank J, Hashem Y (2015) **Structure of mammalian eIF3 in the context of the 43S preinitiation complex** *Nature* **525**:491–495
28. Dey S, Baird TD, Zhou D, Palam LR, Spandau DF, Wek RC (2010) **Both transcriptional regulation and translational control of ATF4 are central to the integrated stress response** *J Biol Chem* **285**:33165–33174
29. Dhillon AS, Hagan S, Rath O, Kolch W (2007) **MAP kinase signalling pathways in cancer** *Oncogene* **26**:3279–3290
30. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR (2013) **STAR: ultrafast universal RNA-seq aligner** *Bioinformatics* **29**:15–21
31. Duan H *et al.* (2023) **eIF3 mRNA selectivity profiling reveals eIF3k as a cancer-relevant regulator of ribosome content** *EMBO J*
32. Durinck S, Spellman PT, Birney E, Huber W (2009) **Mapping identifiers for the integration of genomic datasets with the R/Bioconductor package biomaRt** *Nat Protoc* **4**:1184–1191
33. Fukunaga R, Hunter T (1997) **MNK1, a new MAP kinase-activated protein kinase, isolated by a novel expression screening method for identifying protein kinase substrates** *EMBO J* **16**:1921–1933

34. Furic L *et al.* (2010) **eIF4E phosphorylation promotes tumorigenesis and is associated with prostate cancer progression** *Proc Natl Acad Sci U S A* **107**:14134–14139
35. Gandin V *et al.* (2016) **mTORC1 and CK2 coordinate ternary and eIF4F complex assembly** *Nat Commun* **7**
36. Ge SX, Jung D, Yao R (2020) **ShinyGO: a graphical gene-set enrichment tool for animals and plants** *Bioinformatics* **36**:2628–2629
37. Ontology C Gene (2021) **The Gene Ontology resource: enriching a GOld mine** *Nucleic Acids Res* **49**:D325–D334
38. Gerashchenko MV *et al.* (2020) **Translation elongation factor 2 depletion by siRNA in mouse liver leads to mTOR-independent translational upregulation of ribosomal protein genes** *Sci Rep* **10**
39. Gomes-Duarte A, Lacerda R, Menezes J, Romao L (2018) **eIF3: a factor for human health and disease** *RNA Biol* **15**:26–34
40. Guan BJ *et al.* (2017) **A Unique ISR Program Determines Cellular Responses to Chronic Stress** *Mol Cell* **68**:885–900
41. Harding HP, Novoa I, Zhang Y, Zeng H, Wek R, Schapira M, Ron D (2000) **Regulated translation initiation controls stress-induced gene expression in mammalian cells** *Mol Cell* **6**:1099–1108
42. Heberle H, Meirelles GV, da Silva FR, Telles GP, Minghim R (2015) **InteractiVenn: a web-based tool for the analysis of sets through Venn diagrams** *BMC Bioinformatics* **16**
43. Herrmannová A, Prilepskaja T, Wagner S, Sikrova D, Zeman J, Poncova K, Valasek LS (2020) **Adapted formaldehyde gradient cross-linking protocol implicates human eIF3d and eIF3c, k and l subunits in the 43S and 48S pre-initiation complex assembly, respectively** *Nucleic Acids Res* **48**:1969–1984
44. Hershey JW (2015) **The role of eIF3 and its individual subunits in cancer** *Biochim Biophys Acta* **1849**:792–800
45. Hinnebusch AG (2017) **Structural Insights into the Mechanism of Scanning and Start Codon Recognition in Eukaryotic Translation Initiation** *Trends Biochem Sci* **42**:589–611
46. Howe KL *et al.* (2021) **Ensembl 2021** *Nucleic Acids Res* **49**:D884–D891
47. Hronova V, Mohammad MP, Wagner S, Panek J, Gunisova S, Zeman J, Poncova K, Valasek LS (2017) **Does eIF3 promote reinitiation after translation of short upstream ORFs also in mammalian cells?** *RNA Biol* <https://doi.org/10.1080/15476286.15472017.11353863>
48. Ingolia NT, Brar GA, Rouskin S, McGeachy AM, Weissman JS (2012) **The ribosome profiling strategy for monitoring translation in vivo by deep sequencing of ribosome-protected mRNA fragments** *Nat Protoc* **7**:1534–1550
49. Ishikawa K (2021) **Multilayered regulation of proteome stoichiometry** *Curr Genet* **67**:883–890

50. Jin X, Turcott E, Englehardt S, Mize GJ, Morris DR (2003) **The two upstream open reading frames of oncogene mdm2 have different translational regulatory properties** *J Biol Chem* **278**:25716–25721
51. Juhling F, Morl M, Hartmann RK, Sprinzl M, Stadler PF, Putz J (2009) **tRNADB 2009: compilation of tRNA sequences and tRNA genes** *Nucleic Acids Res* **37**:D159–162
52. Kandoth C *et al.* (2013) **Mutational landscape and significance across 12 major cancer types** *Nature* **502**:333–339
53. Kanehisa M, Sato Y (2020) **KEGG Mapper for inferring cellular functions from protein sequences** *Protein Sci* **29**:28–35
54. Kanehisa M, Sato Y, Kawashima M (2022) **KEGG mapping tools for uncovering hidden features in biological data** *Protein Sci* **31**:47–53
55. Karampelias C *et al.* (2022) **MNK2 deficiency potentiates beta-cell regeneration via translational regulation** *Nat Chem Biol* **18**:942–953
56. King AJ, Sun H, Diaz B, Barnard D, Miao W, Bagrodia S, Marshall MS (1998) **The protein kinase Pak3 positively regulates Raf-1 activity through phosphorylation of serine 338** *Nature* **396**:180–183
57. Kovalski JR, Kuzuoglu-Ozturk D, Ruggero D (2022) **Protein synthesis control in cancer: selectivity and therapeutic targeting** *EMBO J* **41**
58. Kuleshov MV *et al.* (2016) **Enrichr: a comprehensive gene set enrichment analysis web server 2016 update** *Nucleic Acids Res* **44**:W90–97
59. Kusnadi EP, Timpone C, Topisirovic I, Larsson O, Furic L (2022) **Regulation of gene expression via translational buffering** *Biochim Biophys Acta Mol Cell Res* **1869**
60. Langmead B, Salzberg SL (2012) **Fast gapped-read alignment with Bowtie 2** *Nat Methods* **9**:357–359
61. Lauria F, Tebaldi T, Bernabo P, Groen EIJ, Gillingwater TH, Viero G (2018) **riboWaltz: Optimization of ribosome P-site positioning in ribosome profiling data** *PLoS Comput Biol* **14**
62. Lavoie H, Gagnon J, Therrien M (2020) **ERK signalling: a master regulator of cell behaviour, life and fate** *Nat Rev Mol Cell Biol* **21**:607–632
63. Lazaris-Karatzas A, Montine KS, Sonenberg N (1990) **Malignant transformation by a eukaryotic initiation factor subunit that binds to mRNA 5' cap** *Nature* **345**:544–547
64. Lee AS, Kranzusch PJ, Cate JH (2015) **eIF3 targets cell-proliferation messenger RNAs for translational activation or repression** *Nature* **522**:111–114
65. Lee AS, Kranzusch PJ, Doudna JA, Cate JH (2016) **eIF3d is an mRNA cap-binding protein that is required for specialized translation initiation** *Nature* **536**:96–99
66. Leppa S, Saffrich R, Ansorge W, Bohmann D (1998) **Differential regulation of c-Jun by ERK and JNK during PC12 cell differentiation** *EMBO J* **17**:4404–4413

67. Li X, Wang Z, Liu G, Guo J (2021) **EIF3D promotes the progression of preeclampsia by inhibiting of MAPK/ERK1/2 pathway** *Reprod Toxicol* **105**:166–174
68. Lin Y *et al.* (2020) **eIF3 Associates with 80S Ribosomes to Promote Translation Elongation, Mitochondrial Homeostasis, and Muscle Health** *Mol Cell* **79**:575–587
69. Lindqvist LM, Tandoc K, Topisirovic I, Furic L (2018) **Cross-talk between protein synthesis, energy metabolism and autophagy in cancer** *Curr Opin Genet Dev* **48**:104–111
70. Liu T, Zhang L, Joo D, Sun SC (2017) **NF-kappaB signaling in inflammation** *Signal Transduct Target Ther* **2**
71. Llacer JL, Hussain T, Dong J, Villamayor L, Gordiyenko Y, Hinnebusch AG (2021) **Large-scale movement of eIF3 domains during translation initiation modulate start codon selection** *Nucleic Acids Res* **49**:11491–11511
72. Love MI, Huber W, Anders S (2014) **Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2** *Genome Biol* **15**
73. Lu PD, Harding HP, Ron D (2004) **Translation reinitiation at alternative open reading frames regulates gene expression in an integrated stress response** *J Cell Biol* **167**:27–33
74. Manske F, Ogoniak L, Jurgens L, Grundmann N, Makalowski W, Wethmar K (2023) **The new uORFdb: integrating literature, sequence, and variation data in a central hub for uORF research** *Nucleic Acids Res* **51**:D328–D336
75. Marine JC, Lozano G (2010) **Mdm2-mediated ubiquitylation: p53 and beyond** *Cell Death Differ* **17**:93–102
76. Martin M (2011) **Cutadapt removes adapter sequences from high-throughput sequencing reads** *EMBnetjournal* **17**:10–12
77. Morales J *et al.* (2022) **A joint NCBI and EMBL-EBI transcript set for clinical genomics and research** *Nature* **604**:310–315
78. Mukhopadhyay S, Amodeo ME, Lee ASY (2023) **eIF3d controls the persistent integrated stress response** *Mol Cell* **83**:3303–3313
79. Pakos-Zebrucka K, Koryga I, Mnich K, Ljujic M, Samali A, Gorman AM (2016) **The integrated stress response** *EMBO Rep* **17**:1374–1395
80. Pearce SF, Cipullo M, Chung B, Brierley I, Rorbach J (2021) **Mitoribosome Profiling from Human Cell Culture: A High Resolution View of Mitochondrial Translation** *Methods Mol Biol* **2192**:183–196
81. Philippe L, van den Elzen AMG, Watson MJ, Thoreen CC (2020) **Global analysis of LARP1 translation targets reveals tunable and dynamic features of 5' TOP motifs** *Proc Natl Acad Sci U S A* **117**:5319–5328
82. Pisarev AV, Hellen CUT, Pestova TV (2007) **Recycling of Eukaryotic Posttermination Ribosomal Complexes** *Cell* **131**:286–299

83. Poncova K, Wagner S, Jansen ME, Beznoskova P, Gunisova S, Herrmannova A, Zeman J, Dong J, Valasek LS (2019) **uS3/Rps3 controls fidelity of translation termination and programmed stop codon readthrough in co-operation with eIF3** *Nucleic Acids Res* **47**:11326–11343
84. Pyronnet S, Imataka H, Gingras AC, Fukunaga R, Hunter T, Sonenberg N (1999) **Human eukaryotic translation initiation factor 4G (eIF4G) recruits Mnk1 to phosphorylate eIF4E** *EMBO Journal* **18**:270–279
85. Rayburn E, Zhang R, He J, Wang H (2005) **MDM2 and human malignancies: expression, clinical pathology, prognostic markers, and implications for chemotherapy** *Curr Cancer Drug Targets* **5**:27–41
86. Rendleman J *et al.* (2023) **Regulatory start-stop elements in 5' untranslated regions pervasively modulate translation** *bioRxiv*
87. Robichaud N *et al.* (2015) **Phosphorylation of eIF4E promotes EMT and metastasis via translational control of SNAIL and MMP-3** *Oncogene* **34**:2032–2042
88. Robichaud N, Sonenberg N (2017) **Translational control and the cancer cell response to stress** *Curr Opin Cell Biol* **45**:102–109
89. Robichaud N, Sonenberg N, Ruggero D, Schneider RJ (2019) **Translational Control in Cancer** *Cold Spring Harb Perspect Biol*
90. Roux PP, Ballif BA, Anjum R, Gygi SP, Blenis J (2004) **Tumor-promoting phorbol esters and activated Ras inactivate the tuberous sclerosis tumor suppressor complex via p90 ribosomal S6 kinase** *Proc Natl Acad Sci U S A* **101**:13489–13494
91. Roux PP, Topisirovic I (2018) **Signaling Pathways Involved in the Regulation of mRNA Translation** *Mol Cell Biol*
92. Ruggero D (2013) **Translational control in cancer etiology** *Cold Spring Harb Perspect Biol*
93. Sayers EW *et al.* (2022) **Database resources of the national center for biotechnology information** *Nucleic Acids Res* **50**:D20–D26
94. Scheper GC, van der Knaap MS, Proud CG (2007) **Translation matters: protein synthesis defects in inherited disease** *Nat Rev Genet* **8**:711–723
95. She R, Luo J, Weissman JS (2023) **Translational fidelity screens in mammalian cells reveal eIF3 and eIF4G2 as regulators of start codon selectivity** *Nucleic Acids Res* **51**:6355–6369
96. Shu XE, Mao Y, Jia L, Qian SB (2022) **Dynamic eIF3a O-GlcNAcylation controls translation reinitiation during nutrient stress** *Nat Chem Biol* **18**:134–141
97. Shuda M *et al.* (2000) **Enhanced expression of translation factor mRNAs in hepatocellular carcinoma** *Anticancer Res* **20**:2489–2494
98. Silvera D, Formenti SC, Schneider RJ (2010) **Translational control in cancer** *Nat Rev Cancer* **10**:254–266
99. Simonetti A, Guca E, Bochler A, Kuhn L, Hashem Y (2020) **Structural Insights into the Mammalian Late-Stage Initiation Complexes** *Cell Rep* **31**

100. Smekalova EM *et al.* (2020) **In Vivo RNAi-Mediated eIF3m Knockdown Affects Ribosome Biogenesis and Transcription but Has Limited Impact on mRNA-Specific Translation** *Mol Ther Nucleic Acids* **19**:252–266
101. Smirnova AM *et al.* (2024) **Stem-loop-induced ribosome queuing in the uORF2/ATF4 overlap fine-tunes stress-induced human ATF4 translational control** *Cell Rep* **43**
102. Sodani K, Patel A, Kathawala RJ, Chen ZS (2012) **Multidrug resistance associated proteins in multidrug resistance** *Chin J Cancer* **31**:58–72
103. Sonenberg N, Hinnebusch AG (2009) **Regulation of Translation Initiation in Eukaryotes: Mechanisms and Biological Targets** *Cell* **136**:731–745
104. Spilka R, Ernst C, Mehta AK, Haybaeck J (2013) **Eukaryotic translation initiation factors in cancer development and progression** *Cancer Lett* **340**:9–21
105. Sriram A, Bohlen J, Teleman AA (2018) **Translation acrobatics: how cancer cells exploit alternate modes of translational initiation** *EMBO Rep*
106. Stokoe D, Macdonald SG, Cadwallader K, Symons M, Hancock JF (1994) **Activation of Raf as a result of recruitment to the plasma membrane** *Science* **264**:1463–1467
107. Stolovich M, Tang H, Hornstein E, Levy G, Cohen R, Bae SS, Birnbaum MJ, Meyuhas O (2002) **Transduction of growth or mitogenic signals into translational activation of TOP mRNAs is fully reliant on the phosphatidylinositol 3-kinase-mediated pathway but requires neither S6K1 nor rpS6 phosphorylation** *Mol Cell Biol* **22**:8101–8113
108. Topisirovic I, Ruiz-Gutierrez M, Borden KL (2004) **Phosphorylation of the eukaryotic translation initiation factor eIF4E contributes to its transformation and mRNA transport activities** *Cancer Res* **64**:8639–8642
109. Truitt ML, Ruggero D (2016) **New frontiers in translational control of the cancer genome** *Nat Rev Cancer* **16**:288–304
110. Tweedie S, Braschi B, Gray K, Jones TEM, Seal RL, Yates B, Bruford EA (2021) **Genenames.org: the HGNC and VGNC resources in 2021** *Nucleic Acids Res* **49**:D939–D946
111. Valášek LS (2012) **‘Ribozoomin’ – Translation Initiation from the Perspective of the Ribosome-bound Eukaryotic Initiation Factors (eIFs)** *Curr Protein Pept Sci* **13**:305–330
112. Valasek LS, Zeman J, Wagner S, Beznoskova P, Pavlikova Z, Mohammad MP, Hronova V, Herrmannova A, Hashem Y, Gunisova S (2017) **Embraced by eIF3: structural and functional insights into the roles of eIF3 across the translation cycle** *Nucleic Acids Res* **45**:10948–10968
113. Vattam KM, Wek RC (2004) **Reinitiation involving upstream ORFs regulates ATF4 mRNA translation in mammalian cells** *Proc Natl Acad Sci U S A* **101**:11269–11274
114. Volta V, Perez-Baos S, de la Parra C, Katsara O, Ernlund A, Dornbaum S, Schneider RJ (2021) **A DAP5/eIF3d alternate mRNA translation mechanism promotes differentiation and immune suppression by human regulatory T cells** *Nat Commun* **12**

115. Wagner S, Herrmannová A, Hronová V, Gunišová S, Sen ND, Hannan RD, Hinnebusch AG, Shirokikh NE, Preiss T, Valášek LS (2020) **Selective Translation Complex Profiling Reveals Staged Initiation and Co-translational Assembly of Initiation Factor Complexes** *Mol Cell* **79**:546–560
116. Wagner S, Herrmannová A, Malik R, Peclínová L, Valasek LS (2014) **Functional and Biochemical Characterization of Human Eukaryotic Translation Initiation Factor 3 in Living Cells** *Mol Cell Biol* **34**:3041–3052
117. Wagner S, Herrmannová A, Sikrová D, Valasek LS (2016) **Human eIF3b and eIF3a serve as the nucleation core for the assembly of eIF3 into two interconnected modules: the yeast-like core and the octamer** *Nucleic Acids Res* **44**:10772–10788
118. Waskiewicz AJ, Flynn A, Proud CG, Cooper JA (1997) **Mitogen-activated protein kinases activate the serine/threonine kinases Mnk1 and Mnk2** *EMBO J* **16**:1909–1920
119. Wolf DA, Lin Y, Duan H, Cheng Y (2020) **eIF-Three to Tango: emerging functions of translation initiation factor eIF3 in protein synthesis and disease** *J Mol Cell Biol* **12**:403–409
120. Xia Y, Shen S, Verma IM (2014) **NF-kappaB, an active player in human cancers** *Cancer Immunol Res* **2**:823–830
121. Zhang L, Pan X, Hershey JW (2007) **Individual overexpression of five subunits of human translation initiation factor eIF3 promotes malignant transformation of immortal fibroblast cells** *J Biol Chem* **282**:5790–5800
122. Zhang W, Liu HT (2002) **MAPK signal pathways in the regulation of cell proliferation in mammalian cells** *Cell Res* **12**:9–18
123. Zhang Y, Wang X (2020) **Targeting the Wnt/beta-catenin signaling pathway in cancer** *J Hematol Oncol* **13**

Editors

Reviewing Editor

Sandeep Eswarappa

Indian Institute of Science Bangalore, Bengaluru, India

Senior Editor

Jonathan Cooper

Fred Hutchinson Cancer Research Center, Seattle, United States of America

Reviewer #1 (Public review):

Summary:

In this manuscript, Herrmannová et al explore changes in translation upon individual depletion of three subunits of the eIF3 complex (d, e and h) in mammalian cells. The authors provide a detailed analysis of regulated transcripts, followed by validation by RT-qPCR and/or Western blot of targets of interest, as well as GO and KEGG pathway analysis. The authors confirm prior observations that eIF3, despite being a general translation initiation factor, functions in mRNA-specific regulation, and that eIF3 is important for translation re-initiation. They show that global effects of eIF3e and eIF3d depletion on translation and cell growth are

concordant. Their results support and extend previous reports suggesting that both factors control translation of 5'TOP mRNAs. Interestingly, they identify MAPK pathway components as a group of targets coordinately regulated by eIF3 d/e. The authors also discuss discrepancies with other reports analyzing eIF3e function.

Strengths:

Altogether, a solid analysis of eIF3 d/e/h-mediated translation regulation of specific transcripts. The data will be useful for scientists working in the Translation field.

Weaknesses:

The authors could have explored in more detail some of their novel observations, as well as their impact on cell behavior.

The manuscript has improved with the new corrections. I appreciate the authors' attention to the minor comments, which have been fully solved. The authors have not, however, provided additional experimental evidence that uORF-mediated translation of Raf-1 mRNA depends on an intact eIF3 complex, nor have they addressed the consequences of such regulation for cell physiology. While I understand that this is a subject of follow-up research, the authors could have at least included their explanations/ speculations regarding major comments 2-4, which in my opinion could have been useful for the reader.

<https://doi.org/10.7554/eLife.95846.2.sa3>

Reviewer #2 (Public review):

Summary:

mRNA translation regulation permits cells to rapidly adapt to diverse stimuli by fine tuning gene expression. Specifically, the 13-subunit eukaryotic initiation factor 3 (eIF3) complex is critical for translation initiation as it aids in 48S PIC assembly to allow for ribosome scanning. In addition, eIF3 has been shown to drive transcript-specific translation by binding mRNA 5' cap structures through the eIF3d subunit. Dysregulation of eIF3 has been implicated in oncogenesis, however the precise eIF3 subunit contributions are unclear. Here, Herrmannová et al. aim to investigate how eIF3 subcomplexes, generated by knock down (KD) of either eIF3e, eIF3d or eIF3h, affect the global translome. Using Ribo-seq and RNA-seq, the authors identified a large number of genes that exhibit altered translation efficiency upon eIF3d/e KD, while translation defects upon eIF3h KD were mild. eIF3d/eKD share multiple dysregulated transcripts, perhaps due to both subcomplexes lacking eIF3d. Both eIF3d/e KD increase translation efficiency (TE) of transcripts encoding lysosomal, ER and ribosomal proteins, suggesting a role of eIF3 in ribosome biogenesis and protein quality control. Many transcripts encoding ribosomal proteins harbor a TOP motif, and eIF3d KD and eIF3e KD cells exhibit a striking induction of these TOP-modified transcripts. On the other hand, eIF3d KD and eIF3e KD leads to a reduction of MAPK/ERK pathway proteins. Despite this downregulation, eIF3d KD and eIF3e KD activates MAPK/ERK signaling as ERK1/2 and c-Jun phosphorylation was induced. Finally, in all three knockdowns, MDM2 and ATF4 protein levels are reduced. This is notable because MDM2 and ATF4 both contain short uORFs upstream of the start codon, and further supports a role of eIF3 in reinitiation. Altogether, Herrmannová et al. have gained key insights to precise eIF3-mediated translational control as it relates to key signaling pathways implicated in cancer.

Strengths:

The authors have provided a comprehensive set of data to analyze RNA and ribosome footprinting upon perturbation of eIF3d, eIF3e, and eIF3h. As described above in the summary, these data present many interesting starting points to understand additional roles of the eIF3 complex and specific subunits in translational control.

Weaknesses:

- The differences between eIF3e and eIF3d knockdown are difficult to reconcile, especially since eIF3e knockdown leads to reduction in eIF3d levels.
- The paper would be strengthened by experiments directly testing what RNA determinants allow for transcript-specific translation regulation by the eIF3 complex. This would allow the paper to be less descriptive.
- The paper would have more biological relevance if eIF3 subunits were perturbed to mimic naturally occurring situations where eIF3 is dysregulated. For example, eIF3e is aberrantly upregulated in certain cancers, and therefore an overexpression and profiling experiment would have been more relevant than a knockdown experiment.

The first review is unchanged as no additional experiments were provided to address the first review.

<https://doi.org/10.7554/eLife.95846.2.sa2>

Reviewer #3 (Public review):**Summary:**

In this article, Herrmannová et al catalog the changes in ribosome association with mRNAs when the multisubunit eukaryotic translation initiation factor 3 is disrupted by knocking down individual subunits. They find that RNAs relying on TOP motifs for translation, such as ribosomal protein RNAs, and RNAs encoding modification enzymes in the ER and components of the lysosome are upregulated. In contrast, proteins encoding components of MAP kinase cascades are downregulated when subunits of eIF3 are knocked down, but retain elevated levels of activity.

Strengths:

The authors use ribosome profiling of well-characterized mutants lacking subunits of eIF3 and assess the changes in translation that take place. They supplement the ribosome association studies with western blotting to determine protein level changes of affected transcripts. They analyze what transcripts undergo translation changes, which is important for understanding more broadly how translation initiation factor levels affect cancer cell translomes. Changes observed by both ribosome profiling and western blotting supports their claims that eIF3 functions in mRNA-specific control of translation.

Weaknesses:

- (1) The paper would be strengthened if there were a clear model tying the various effects together or linking individual subunit knockdown to cancerous phenotypes. It is noted that the authors plan to address such outcomes of eIF3 dysregulation in future work, which will be of interest.
- (2) The paper could also be strengthened if some of the experiments were performed in at least one other cell type to determine whether changes observed are general or cell-type specific. The authors discuss this issue and provide a literature citation to support a more general mechanism.

<https://doi.org/10.7554/eLife.95846.2.sa1>

Author response:

The following is the authors' response to the current reviews.

Reviewer #1 (Public review):

Summary:

In this manuscript, Herrmannová et al explore changes in translation upon individual depletion of three subunits of the eIF3 complex (d, e and h) in mammalian cells. The authors provide a detailed analysis of regulated transcripts, followed by validation by RT-qPCR and/or Western blot of targets of interest, as well as GO and KEGG pathway analysis. The authors confirm prior observations that eIF3, despite being a general translation initiation factor, functions in mRNA-specific regulation, and that eIF3 is important for translation re-initiation. They show that global effects of eIF3e and eIF3d depletion on translation and cell growth are concordant. Their results support and extend previous reports suggesting that both factors control translation of 5'TOP mRNAs. Interestingly, they identify MAPK pathway components as a group of targets coordinately regulated by eIF3 d/e. The authors also discuss discrepancies with other reports analyzing eIF3e function.

Strengths:

Altogether, a solid analysis of eIF3 d/e/h-mediated translation regulation of specific transcripts. The data will be useful for scientists working in the Translation field.

Weaknesses:

The authors could have explored in more detail some of their novel observations, as well as their impact on cell behavior.

The manuscript has improved with the new corrections. I appreciate the authors' attention to the minor comments, which have been fully solved. The authors have not, however, provided additional experimental evidence that uORF-mediated translation of Raf-1 mRNA depends on an intact eIF3 complex, nor have they addressed the consequences of such regulation for cell physiology. While I understand that this is a subject of follow-up research, the authors could have at least included their explanations/ speculations regarding major comments 2-4, which in my opinion could have been useful for the reader.

Our explanations/speculations regarding major comments 2 and 3 were included in the Discussion. We apologize for this misunderstanding as we thought that we were supposed to explain our ideas only in the responses. We did not discuss the comment 4, however, as we are really not sure what is the true effect and did not want to go into wild speculations in our manuscript. We thank this reviewer for his insightful comments and understanding.

The following is the authors' response to the original reviews.

Reviewer #1 (Recommendations For The Authors):

Major comments:

(1) The authors report the potential translational regulation of Raf kinase by re-initiation. It would be interesting to show that Raf is indeed regulated by uORF-mediated translation, and that this is dependent on an intact eIF3 complex. Analyzing the potential consequences of Raf1 regulation for cancer cell proliferation or apoptosis would be a plus.

We agree that this is an interesting and likely possibility. In fact, another clue that translation of Raf1 is regulated by uORFs comes from Bohlen et al. 2023 (PMID: 36869665) where they

showed that RAF1 translation is dependent on PRRC2 proteins (that promote leaky scanning through these uORFs). We noted in the discussion that our results from eIF3d/e/hKD and the PRRC2A/B/CKD partly overlap. It is a subject of our follow-up research to investigate whether eIF3 and PRRC2 co-operate together to regulate translation of this important mRNA.

(2) The authors show that eIF3 d/e -but not 3h- has an effect on cell proliferation. First, this indicates that proliferation does not fully correlate with eIF3 integrity. Depletion of eIF3d does not affect the integrity of eIF3, yet the effects on proliferation are similar to those of eIF3e. What is the possibility that changes in proliferation reflect functions of eIF3d outside the eIF3 complex? What could be the real consequences of disturbing eIF3 integrity for the mammalian cell? Please, discuss.

Yes, proliferation does not fully correlate with eIF3 integrity. Downregulation of eIF3 subunits that lead to disintegration of eIF3 YLC core (a, b, c, g, i) have more detrimental effect on growth and translation than downregulation of the peripheral subunits (e, k, l, f, h, m). Our previous studies (Wagner et al. 2016, PMID: 27924037 and Herrmannová et al. 2020, PMID: 31863585) indicate that the YLC core of eIF3 can partially support translation even without its peripheral subunits. In this respect eIF3d (as a peripheral subunit) is an amazing exception, suggesting it may have some specialized function(s). Whether this function resides outside of the eIF3 complex or not we do not know, but do not think so. Mainly because in the absence of eIF3e – its interaction partner, eIF3d gets rapidly degraded. Therefore, it is not very likely that eIF3d exists alone outside of eIF3 complex with moonlighting functions elsewhere. We think that eIF3d, as a head-interacting subunit close to an important head ribosomal protein RACK1 (a landing pad for regulatory proteins), is a target of signaling pathways, which may make it important for translation of specific mRNAs. In support of these thoughts, eIF3d (in the context of entire eIF3) together with DAP5 were shown to promote translation by an alternate capdependent (eIF4F-independent) mechanism (Lee et al. 2016, PMID: 27462815; de la Parra et al. 2018, PMID:30076308). In addition, the eIF3d function (also in the context of entire eIF3) was proved to be regulated by stress-triggered phosphorylation (Lamper et al. 2020, PMID: 33184215).

(3) Figure 6D: Surprisingly, reduced levels of ERK1/2 upon eIF3d/e-KD are compensated by increased phosphorylation of ERK1/2 and net activation of c-Jun. Please comment on the functional consequences of buffering mechanisms that the cell deploys in order to counteract compromised eIF3 function. Why would the cell activate precisely the MAPK pathway to compensate for a compromised eIF3 function?

This we do not know. We can only speculate that when translation is compromised, cells try to counteract it in two ways: 1) they produce more ribosomes to increase translational rates and 2) activate MAPK signaling to send pro-growth signals, which can in the end further boost ribosome biogenesis.

(4) Regarding DAP-sensitive transcripts, can the authors discuss in more detail the role of eIF3d in alternative cap-dependent translation versus re-initiation? Are these transcripts being translated by a canonical cap- and uORF-dependent mechanism or by an alternative capdependent mechanism?

This is indeed not an easy question. On one hand, it was shown that DAP5 facilitates translation re-initiation after uORF translation in a canonical cap-dependent manner. This mechanism is essential for translation of the main coding sequence (CDS) in mRNAs with structured 5' leaders and multiple uORFs. (Weber et al. 2022, PMID: 36473845; David et al., 2022, PMID: 35961752). On the other hand, DAP5 was proposed to promote alternative, eIF4F-independent but cap-dependent translation, as it can substitute the function of the eIF4F complex in cooperation with eIF3d (de la Parra et al., 2018, PMID: 30076308; Volta et al., 2021


34848685). Overall, these observations paint a very complex picture for us to propose a clear scenario of what is going on between these two proteins on individual mRNAs. We speculate that both mechanisms are taking place and that the specific mechanism of translation initiation differs for differently arranged mRNAs.

Minor comments:

(5) Figure S2C: why is there a strong reduction of the stop codon peak for 3d and 3h KDs?

We have checked the Ribowaltz profiles of all replicates (in the Supplementary data we are showing only a representative replicate I) and the stop codon peak differs a lot among the replicates. We think that this way of plotting was optimized for calculation and visualization of P-sites and triplet periodicity and thus is not suitable for this type of comparison among samples. Therefore, we have performed our own analysis where the 5' ends of reads are used instead of P-sites and triplicates are averaged and normalized to CDS (see below please), so that all samples can be compared directly in one plot (same as Fig. S13A but for stop codon). We can see that the stop codon peak really differs and is the smallest for eIF3hKD. However, these changes are in the range of 20% and we are not sure about their biological significance. We therefore refrain from drawing any conclusions. In general, reduced stop codon peak may signal faster termination or increased stop codon readthrough, but the latter should be accompanied by an increased ribosome density in the 3'UTR, which is not the case. A defect in termination efficiency would be manifested by an increased stop codon peak, instead.

Author response image 1.



(6) Figures 5 and S8: Adding a vertical line at 'zero' in all cumulative plots will help the reader understand the author's interpretation of the data.

We have added a dashed grey vertical line at zero as requested. However, for interpretation of these plots, the reader should focus on the colored curve and whether it is shifted in respect to the grey curve (background) or not. Shift to the right indicates increased expression, while shift to the left indicates decreased expression. The reported p-value then indicates the statistical significance of the shift.

(7) The entire Figure 2 are controls that can go to Supplementary Material. The clustering of Figure S3B could be shown in the main Figure, as it is a very easy read-out of the

consistent effects of the KDs of the different eIF3 subunits under analysis.

We have moved the entire Figure 2 to Supplementary Material as suggested (the original panels can be found as Supplementary Figures 1B, 1C and 3A). Figure S3B is now the main Figure 2E.

(8) *There are 3 replicates for Ribo-Seq and four for RNA-Seq. Were these not carried out in parallel, as it is usually done in Ribo-seq experiments? Why is there an extra replicate for RNASeq?*

Yes, the three replicates were carried out in parallel. We have decided to add the fourth replicate in RNA-Seq to increase the data robustness as the RNA-Seq is used for normalization of FP to calculate the TE, which was our main analyzed metrics in this article. We had the option to add the fourth replicate as we originally prepared five biological replicates for all samples, but after performing the control experiments, we selected only the 3 best replicates for the Ribo-Seq library preparation and sequencing.

(9) *Please, add another sheet in Table S2 with the names of all genes that change only at the translation (RPF) levels.*

As requested, we have added three extra sheets (one for each downregulation) for differential FP with Padjusted <0.05 in the Spreadsheet S2. We also provide a complete unfiltered differential expression data (sheet named “all data”), so that readers can filter out any relevant data based on their interest.

(10) *Page 5, bottom: '...we showed that the expression of all 12 eIF3 subunits is interconnected such that perturbation of the expression of one subunit results in the down-regulation of entire modules...'. This is not true for eIF3d, as shown in Fig1B and mentioned in Results.*

This reviewer is correct. By this generalized statement, we were trying to summarize our previous results from Wagner et al., 2014, PMID: 24912683; Wagner et al., 2016, PMID: 27924037 and Herrmannová et al., 2020, PMID: 31863585. The eIF3d downregulation is the only exception that does not affect expression of any other eIF3 subunit. Therefore, we have rewritten this paragraph accordingly: “We recently reported a comprehensive *in vivo* analysis of the modular dynamics of the human eIF3 complex (Wagner *et al.*, 2020; Wagner *et al.*, 2014; Wagner *et al.*, 2016). Using a systematic individual downregulation strategy, we showed that the expression of all 12 eIF3 subunits is interconnected such that perturbation of the expression of one subunit results in the down-regulation of entire modules leading to the formation of partial eIF3 subcomplexes with limited functionality (Herrmannová *et al.*, 2020). eIF3d is the only exception in this respect, as its downregulation does not influence expression of any other eIF3 subunit.”

(11) *Page 10, bottom: 'The PCA plot and hierarchical clustering... These results suggest that eIF3h depletion impacts the translome differentially than depletion of eIF3e or eIF3d.' This is already obvious in the polysome profiles of Figure S2C.*

We agree that this result is surely not surprising given the polysome profile and growth phenotype analyses of eIF3hKD. But still, we think that the PCA plot and hierarchical clustering results represent valuable controls. Nonetheless, we rephrased this section to note that this result agrees with the polysome profiles analysis: “The PCA plot and hierarchical clustering (Figure 2A and Supplementary Figure 4A) showed clustering of the samples into two main groups: Ribo-Seq and RNA-seq, and also into two subgroups; NT and eIF3hKD samples clustered on one side and eIF3eKD and eIF3dKD samples on the other. These results

suggest that the eIF3h depletion has a much milder impact on the translome than depletion of eIF3e or eIF3d, which agrees with the growth phenotype and polysome profile analyses (Supplementary Figure 1A and 1D)."

(12) Page 12: 'As for the eIF3dKD "unique upregulated" DTEGs, we identified one interesting and unique KEGG pathway, the ABC transporters (Supplementary Figure 5A, in green).' This sentence is confusing, as there are more pathways that are significant in this group, so it is unclear why the authors consider it 'unique'.

The eIF3dKD "unique upregulated" group comprises genes with increased TE only in eIF3dKD but not in eIF3eKD or eIF3hKD (500 genes, Fig 2G). All these 500 genes were examined for enrichment in the KEGG pathways, and the top 10 significant pathways were reported (Fig S6A). However, 8 out of these 10 pathways were also significantly enriched in other gene groups examined (e.g. eIF3d/eIF3e common). Therefore, the two remaining pathways ("ABC transporters" and "Other types of O-glycan biosynthesis") are truly unique for eIF3dKD. We wanted to highlight the ABC transporters group in particular because we find it rather interesting (for the reasons mentioned in the article). We have corrected the sentence in question to avoid confusion: "Among the eIF3dKD "unique upregulated" DTEGs, we identified one interesting KEGG pathway, the ABC transporters, which did not show up in other gene groups (Supplementary Figure 6A, in green). A total of 12 different ABC transporters had elevated TE (9 of them are unique to eIF3dKD, while 3 were also found in eIF3eKD), 6 of which (ABCC1-5, ABCC10) belong to the C subfamily, known to confer multidrug resistance with alternative designation as multidrug resistance protein (MRP1-5, MRP7) (Sodani *et al*, 2012).

Interestingly, all six of these ABCC transporters were upregulated solely at the translational level (Supplementary Spreadsheet S2)."

(13) Note typo ('Various') in Figure 4A.

Corrected

(14) The introduction could be shortened.

This is a very subjective requirement. In fact, when this manuscript was reviewed in NAR, we were asked by two reviewers to expand it substantially. Because a number of various research topics come together in this work, e.g. translational regulation, the eIF3 structure and function, MAPK/ERK signaling, we are convinced that all of them demand a comprehensive introduction for non-experts in each of these topics. Therefore, with all due respect to this reviewer, we did not ultimately shorten it.

Reviewer #2 (Recommendations For The Authors):

- In Figure 2, it would be useful to know why eIF3d is destabilized by eIF3e knockdown - is it protein degradation and why do the eIF3d/e knockdowns not more completely phenocopy each other when there is the same reduction to eIF3d as in the eIF3d knockdown sample?

Yes, we do think that protein degradation lies behind the eIF3d destabilization in the eIF3eKD, but we have not yet directly demonstrated this. However, we have shown that eIF3d mRNA levels are not altered in eIF3eKD and that Ribo-Seq data indicate no change in TE or FP for eIF3d-encoding mRNA in eIF3eKD. Nonetheless, it is important to note (and we discuss it in the article) that eIF3d levels in eIF3dKD are lower than eIF3d levels in eIF3eKD (please see Supplementary Figure 1C). In fact, we believe that this is one of the main reasons for the eIF3d/e knockdowns differences.

- The western blots in Figures 4 and 6 show modest changes to target protein levels and would be strengthened by quantification.

We have added the quantifications as requested by this reviewer and the reviewer 3.

- For Figure 4, this figure would be strengthened by experiments showing if the increase in ribosomal protein levels is correlated with actual changes to ribosome biogenesis.

As suggested, we performed polysome profiling in the presence of EDTA to monitor changes in the 60S/40S ratio, indicating a potential imbalance in the biogenesis of individual ribosome subunits. We found that it was not affected (Figure 3G). In addition, we performed the same experiment, normalizing all samples to the same number of cells (cells were carefully counted before lysis). In this way, we confirmed that eIF3dKD and eIF3eKD cells indeed contain a significantly increased number of ribosomes, in agreement with the western blot analysis (Figure 3H).

- In Figure 6, there needs to be a nuclear loading control.

This experiment was repeated with Lamin B1 used as a nuclear loading control – it is now shown as Fig. 5F.

- For Figure 8, these findings would be strengthened using luciferase reporter assays where the various RNA determinants are experimentally tested. Similarly, 5' TOP RNA reporters would have been appreciated in Figure 4.

This is indeed a logical continuation of our work, which represents the current work in progress of one of the PhD students. We apologize, but we consider this time- and resource-demanding analysis out of scope of this article.

Reviewer #3 (Recommendations For The Authors):

(1) Within the many effects observed, it is mentioned that eIF3d is known to be overexpressed while eIF3e is underexpressed in many cancers, but knockdown of either subunit decreases MDM2 levels, which would be expected to increase P53 activity and decrease tumor cell transformation. In contrast, they also report that 3e/3d knockdown dramatically increases levels of cJUN, presumably due to increased MAPK activity, and is expected to increase protumor gene expression. Additional discussion is needed to clarify the significance of the findings, which are a bit confusing.

This is indeed true. However, considering the complexity of eIF3, the largest initiation factor among all, as well as the broad portfolio of its functions, it is perhaps not so surprising that the observed effects are complex and may seem even contradictory in respect to cancer. To acknowledge that, we expanded the corresponding part of discussion as follows: “Here, we demonstrate that alterations in the eIF3 subunit stoichiometry and/or eIF3 subcomplexes have distinct effects on the translome; for example, they affect factors that play a prominent (either positive or negative) role in cancer biology (e.g., MDM2 and cJUN), but the resulting impact is unclear so far. Considering the complex interactions between these factors as well as the complexity of the eIF3 complex *per se*, future studies are required to delineate the specific oncogenic and tumor suppressive pathways that play a predominant role in mediating the effects of perturbations in the eIF3 complex in the context of neoplasia.”

(2) There are places in the text where the authors refer to changes in transcriptional control when RNA levels differ, but transcription versus RNA turnover wasn't tested, e.g.

page 16 and Figure S10, qPCR does not confirm "transcriptional upregulation in all three knockdowns" and page 19 "despite apparent compensatory mechanisms that increase their transcription."

This is indeed true, the sentences in question were corrected. The term “increased mRNA levels” was used instead of transcriptional upregulation (increased mRNA stabilization is also possible).

(3) Similarly, the authors suggest that steady-state LARP1 protein levels are unaffected based on ribosome footprint counts (page 21). It is incorrect to assume this, because ribosome footprints can be elevated due to stalling on RNA that isn't being translated and doesn't yield more protein, and because levels of translated RNA/synthesized proteins do not always reflect steady-state protein levels, especially in mutants that could affect lysosome levels and protein turnover. Also page 12, 1st paragraph suggests protein production is down when ribosome footprints are changed.

Yes, we are well-aware of this known limitation of Ribo-seq analysis. Therefore, the steady-state protein levels of our key hits were verified by western blotting. In addition, we have removed the sentence about LARP1 because it was based on Ribo-Seq data only without experimental evaluation of the steady-state LARP1 protein levels.

(4) The translation buffering effect is not clear in some Figures, e.g. S6, S8, 8A, and B. The authors show a scheme for translationally buffered RNAs being clustered in the upper right and lower left quadrants in S4H (translation up with transcript level down and v.v.), but in the FP versus RNA plots, the non-TOP RNAs and 4E-P-regulated RNAs don't show this behavior, and appear to show a similar distribution to the global changes. Some of the right panels in these figures show modest shifts, but it's not clear how these were determined to be significant. More information is needed to clarify, or a different presentation, such as displaying the RNA subsets in the left panels with heat map coloring to reveal whether RNAs show the buffered translation pattern defined in purple in Figure S4H, or by reporting a statistical parameter or number of RNAs that show behavior out of total for significance. Currently the conclusion that these RNAs are translationally buffered seems subjective since there are clearly many RNAs that don't show changes, or show translation-only or RNA-only changes.

We would like to clarify that S4H does not indicate a necessity for changes in FPs in the buffered subsets. Although opposing changes in total mRNA and FPs are classified as buffering, often we also consider the scenario where there are changes to the total mRNA levels not accompanied by changes in ribosome association.

In figure S6, the scatterplots indicate a high density of genes shifted towards negative fold changes on the x-axis (total mRNA). This is also reflected in the empirical cumulative distribution functions (ecdfs) for the log2 fold changes in total mRNA in the far right panels of A and B, and the lack of changes in log2 fold change for FPs (middle panels). Similarly, in figure S8, the scatterplots indicate a density of genes shifted towards positive fold changes on the x-axis for total mRNA. The ecdfs also demonstrate that there is a significant directional shift in log2 fold changes in the total mRNA that is not present to a similar degree in the FPs, consistent with translational offsetting. It is rightly pointed out that not all genes in these sets follow the same pattern of regulation. We have revised the title of Supplementary Figure S6 (now S7) to reflect this. However, we would like to emphasize that these figures are not intended to communicate that all genes within these sets of interest are regulated in the same manner, but rather that when considered as a whole, the predominant effect seen is that of translational offsetting (directional shifts in the log2 fold change distribution of total mRNA that are not accompanied by similar shifts in FP mRNA log2 fold changes).

The significance of these differences was determined by comparing the ecdfs of the log2 fold changes for the genes belonging to a particular set (e.g. non-TOP mTOR-sensitive, p-eIF4E-sensitive) against all other expressed genes (background) using a Wilcoxon rank sum test. This allows identification of significant shifts in the distributions that have a clear directionality (if there is an overall increase, or decrease in fold changes of FPs or total mRNA compared to background). If log2 fold changes are different from background, but without a clear directionality (equally likely to be increased or decreased), the test will not yield a significant result. This approach allows assessment of the overall behavior of gene signatures within a given dataset in a manner that is completely threshold-independent, such that it does not rely on classification of genes into different regulatory categories (translation only, buffering, etc.) based on significance or fold-change cut-offs (as in S4H). Therefore, we believe that this unbiased approach is well-suited for identifying cases when there are many genes that follow similar patterns of regulation within a given dataset.

(5) Page 10-*"These results suggest that eIF3h depletion impacts the translome differentially than depletion of eIF3e or eIF3d" ...These results suggest that eIF3h has less impact on the translome, not that it does so differently. If it were changing translation by a different mechanism, I would not expect it to cluster with control.*

This sentence was rewritten as follows: "The PCA plot and hierarchical clustering (Figure 2A and Supplementary Figure 4A) showed clustering of the samples into two main groups: RiboSeq and RNA-seq, and also into two subgroups; NT and eIF3hKD samples clustered on one side and eIF3eKD and eIF3dKD samples on the other. These results suggest that the eIF3h depletion has a much milder impact on the translome than depletion of eIF3e or eIF3d, which agrees with the growth phenotype and polysome profile analyses (Supplementary Figure 1A and 1D)."

Other minor issues:

(1) *There are some typos: Figure 2 leves, Figure 4 variou,*

Corrected.

(2) *Figure 3, font for genes on volcano plot too small*

Yes, maybe, however the resolution of this image is high enough to enlarge a certain part of it at will. In our opinion, a larger font would take up too much space, which would reduce the informativeness of this graph.

(3) *Figure S5, highlighting isn't defined.*

The figure legend for S5A (now S6A) states: "Less significant terms ranking 11 and below are in grey. Terms specifically discussed in the main text are highlighted in green." Perhaps it was overlooked by this reviewer.

(4) *At several points the authors refer to "the MAPK signaling pathway", suggesting there is a single MAPK that is affected, e.g in the title, page 3, and other places when it seems they mean "MAPK signaling pathways" since several MAPK pathways appear to be affected.*

We apologize for any terminological inaccuracies. There are indeed several MAPK pathways operating in cells. In our study, we focused mainly on the MAPK/ERK pathway. The confusion probably stems from the fact that the corresponding term in the KEGG pathway database is labeled "MAPK signaling pathway" and this term, although singular, includes all MAPK

pathways. We have carefully reviewed the entire article and have corrected the term used accordingly to either: 1) MAPK pathways in general, 2) the MAPK/ERK pathway for this particular pathway, or 3) "MAPK signaling pathway", where the KEGG term is meant.

(5) Some eIF3 subunit RNAs have TOP motifs. One might expect 3e and 3h levels to change as a function of 3d knockdown due to TOP motifs but this is not observed. Can the authors speculate why the eIF3 subunit levels don't change but other TOP RNAs show TE changes? Is this true for other translation factors, or just for eIF3, or just for these subunits? Could the Western blot be out of linear range for the antibody or is there feedback affecting eIF3 levels differently than the other TOP RNAs, or a protein turnover mechanism to maintain eIF3 levels?

This is indeed a very interesting question. In addition to the mRNAs encoding ribosomal proteins, we examined all TOP mRNAs and added an additional sheet to the S2 supplemental spreadsheet with all TOP RNAs listed in (Philippe et al., 2020, PMID: 32094190). According to our Ribo-Seq data, we could expect to see increased protein levels of eIF3a and eIF3f in eIF3dKD and eIF3eKD, but this is not the case, as judged from extensive western blot analysis performed in (Wagner et. al 2016, PMID: 27924037). Indeed, we cannot rule out the involvement of a compensatory mechanism monitoring and maintaining the levels of eIF3 subunits at steady-state – increasing or decreasing them if necessary, which could depend on the TOP motif-mediated regulation. However, we think that in our KDs, all non-targeted subunits that lose their direct binding partner in eIF3 due to siRNA treatment become rapidly degraded. For example, co-downregulation of subunits d, k and l in eIF3eKD is very likely caused by protein degradation as a result of a loss of their direct binding partner – eIF3e. Since we showed that the yeast eIF3 complex assembles co-translationally (Wagner et. al 2020, PMID: 32589964), and there is no reason to think that mammalian eIF3 differs in this regard, our working hypothesis is that free subunits that are not promptly incorporated into the eIF3 complex are rapidly degraded, and the presence or absence of the TOP motif in the 5' UTR of their mRNAs has no effect. As for the other TOP mRNAs, translation factors eEF1B2, eEF1D, eEF1G, eEF2 have significantly increased FPs in both eIF3dKD and eIF3eKD, but we did not check their protein levels by western blotting to conclude anything specific.

<https://doi.org/10.7554/eLife.95846.2.sa0>