

Actin dynamics switches two distinct modes of endosomal fusion in yolk sac visceral endoderm cells

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint version 1

March 20, 2024 (this version)

Sent for peer review

February 2, 2024

Posted to preprint server

December 18, 2023

Seiichi Koike, Masashi Tachikawa, Motosuke Tsutsumi, Takuya Okada, Tomomi Nemoto, Kazuko Keino-Masu, Masayuki Masu 

Graduate School of Comprehensive Human Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan • Department of Molecular Neurobiology, Institute of Medicine, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan • Laboratory of Molecular and Cellular Biology, Graduate School of Science and Engineering for Research, University of Toyama, 3190 Gofuku, Toyama 930-8555, Toyama, Japan • Graduate School of Nanobioscience, Yokohama City University, 22-2 Seto, Kanazawa-ku, Yokohama, Kanagawa 236-0027, Japan • Exploratory Research Center on Life and Living Systems (ExCELLS), National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji, Okazaki, Aichi 444-8787, Japan • National Institute for Physiological Sciences, National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji, Okazaki, Aichi 444-8787, Japan

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Membranes undergo various patterns of deformation during vesicle fusion, but how this membrane deformation is regulated and contributes to fusion remains unknown. In this study, we developed a new method of observing the fusion of individual late endosomes and lysosomes by using yolk sac visceral endoderm cells that have huge endocytic vesicles. We found that there were two distinct fusion modes that were differently regulated. In homotypic fusion, two late endosomes fused quickly, whereas in heterotypic fusion they fused to lysosomes slowly. Mathematical modeling showed that vesicle size is a critical determinant of these fusion types and that membrane fluctuation forces can overcome the vesicle size effects. We found that actin filaments were bound to late endosomes and forces derived from dynamic actin remodeling were necessary for quick fusion during homotypic fusion. Furthermore, cofilin played a role in endocytic fusion by regulating actin turnover. These data suggest that actin promotes vesicle fusion for efficient membrane trafficking in visceral endoderm cells.

eLife assessment

This study presents **valuable** findings on two distinct modes of endosomal fusion and the roles of actin dynamics in this process. The evidence supporting the authors' claims is **solid**, although the underlying molecular mechanisms and whether the proposed fusion modes are applicable in other cell types remain unclear. The work will be of interest to cell biologists and biophysicists working on the cytoskeleton and organelles.

Introduction

Eukaryotic cells contain numerous organelles that are separated from the cytoplasm by a lipid bilayer. Exchange of proteins, lipids, and metabolites between the organelles occurs through vesicle budding and fusion. To date, theoretical and computer biophysics have modeled several steps in the process of membrane fusion (Chernomordik and Kozlov, 2008 [DOI](#); Lu and Guo, 2019 [DOI](#)). The priming of membranes is followed by the merger of the outer leaflet of the membranes to establish a hemifusion stalk. The stalk then expands into a hemifusion diaphragm and the merger of the inner leaflets leads to the formation of a fusion pore, which expands and allows content mixing. Recently, the hemifusion stage was observed in live cells, providing strong evidence of the hemifusion model (Zhao et al., 2016 [DOI](#)).

Studies utilizing time-lapse microscopic observation of live cells, however, have revealed that membrane behavior after pore formation is unexpectedly divergent. For example, in giant endosomal vesicles in cells overexpressing constitutively active Rab5 protein, two different types of membrane fusion were observed (Roberts et al., 1999 [DOI](#)). In “explosive” fusion, which is commonly seen in cells, the fusion pore expanded quickly. In contrast, in “bridge” fusion, the pore did not expand; instead, one vesicle gradually shrank and disappeared over time as the result of the transfer of its vesicle content into another vesicle. In homotypic vacuole fusion in *Saccharomyces cerevisiae*, vacuole membranes fused at the vertices instead of a central pore expanding radially, and then a boundary membrane between the two docked vacuoles was released into the vacuole lumen (Wang et al., 2002 [DOI](#)). Because pore dynamics determine the rate of content mixing, they have significant effects on the trafficking efficiency of cargo molecules. However, how different modes of membrane deformation are induced and whether the distinct fusion modes are exchangeable remain unknown.

Membrane deformation is generally controlled by membrane-associating proteins as well as by several physical parameters including the vesicle volume, membrane area, bending energy, and osmotic energy in cells (Farsad and De Camilli, 2003 [DOI](#); Lipowsky, 1991 [DOI](#)). Actin and its regulating proteins, via interaction with the membrane, exert physical forces on and cause fluctuation in the membrane, which eventually leads to membrane deformation by modification of the physical parameters (Salbreux et al., 2012 [DOI](#)). Actin produces forces in two different manners. First, assembly of branched actin filaments, nucleated from the membrane surface by actin nucleators such as the actin-related protein 2/3 (Arp2/3) complex and formins, provides pushing forces on the membrane (Goley and Welch, 2006 [DOI](#); Svitkina, 2018 [DOI](#)). Second, actin generates pulling forces through myosin-driven sliding of actin filaments (Svitkina, 2018 [DOI](#)). Both forces play roles in induction of membrane deformation during vesicle fusion (Ebrahim et al., 2019 [DOI](#); Ma et al., 2020 [DOI](#); Tran et al., 2015 [DOI](#); Uenishi et al., 2013 [DOI](#)). Many experiments have been carried out to elucidate the roles of actin in vesicle fusion through manipulation of actin dynamics by use of specific drugs or gene modifications. These experiments gave rise to different results depending on the experimental conditions. For example, polymerization or stabilization of actin enhanced membrane fusion in some experiments but inhibited it in others (reviewed in Eitzen, 2003 [DOI](#)). These different results imply that the regulation mechanisms of vesicle fusion by actin filaments is more complex than previously assumed. Moreover, how the actin forces contribute to the fusion modes remains unknown, as does how the force generated by actin polymerization or by actin-dependent motor proteins contributes to membrane deformation during fusion.

In this study, we utilized mouse yolk sac visceral endoderm (VE) cells as a model system for addressing the above questions. VE cells play a critical role in the maternofetal exchange of nutrients before the establishment of a chorioallantoic placenta: they vigorously endocytose maternal proteins, hydrolyze the proteins in lysosomes, and supply the resultant products to the developing embryo (Bielinska et al., 1999 [DOI](#); Jollie 1990 [DOI](#)). Reflecting this high endocytic activity,

VE cells are characterized by huge endosomal vesicles (Bielinska et al., 1999 [↗](#); Kawamura et al., 2012 [↗](#); Koike et al., 2009 [↗](#)). These properties of VE cells enable us to observe endosomal dynamics including vesicle fusion. Previously, by using a whole embryo culture system combined with specific pharmacologic inhibitors for intracellular signaling molecules, we showed that actin dynamics regulated by lysophosphatidic acid (LPA) receptors and the Rho-ROCK-LIM kinase pathway are required for the formation of huge lysosomes (Koike et al., 2009 [↗](#)).

Here, we show that late endosomes of VE cells exhibit different modes of fusion with target organelles depending on their targets: they undergo homotypic fusion with late endosomes in an explosive fusion-like manner and heterotypic fusion with lysosomes in a bridge fusion-like manner. Our mathematical modeling predicts that the vesicle size is a critical determinant of the fusion types and that membrane perturbation overcomes the size effects. We provide evidence that actin filaments are associated with late endosomes and that actin dynamics generate forces promoting vesicle fusion of endosomes for efficient membrane trafficking.

Results

Endocytic vesicles in mouse yolk sac VE cells

To visualize endosomal vesicles in yolk sac VE cells, an embryonic day 8.5 (E8.5) whole embryo was labeled with a fluorescent membrane probe, FM1-43, and subsequently stained with organelle markers. The embryo was put on a glass-bottom dish, and VE cells were observed by use of confocal microscopy (Fig. 1A [↗](#)). These observations revealed that different types of endosomal vesicles formed three layers from the apical to the basal area in the supranuclear portion of VE cells (Fig. 1, B [↗](#) and C [↗](#)). In the most superficial portion (within about 1 μm of the apical surface; designated as zone 1), fuzzy FM1-43 staining was observed (Fig. 1B [↗](#)). The small vesicles in this zone were positive for EEA1 (Fig. 1B [↗](#)), indicating that they were early endosomes. Immediately beneath zone 1, many medium-sized round vesicles were observed (zone 2). Electroporation of GFP-Rab7 revealed that these vesicles were positive for Rab7, suggesting that they were late endosomes. In the deep portion (zone 3), large amorphous lysosomes, positive for LysoTracker Red and LAMP1, were observed (Fig. 1B [↗](#); Koike et al., 2009 [↗](#)).

To examine whether these vesicles constitute endocytic pathways, we traced vesicle trafficking in VE cells. After E8.5 whole embryos were pulse-labeled with Alexa 488-transferrin for 5 min, fluorescently labeled endocytic vesicles were tracked over time. The fluorescent signals were initially detected in small vesicles (area < 2.0 μm^2) in zone 1; 10–20 min after labeling, the signals were mainly detected in medium-sized vesicles (2.0–25 μm^2) in zone 2; 40 min after labeling, the signals were strongly observed in large lysosomes (> 25 μm^2) in zone 3 (Fig. 1, D [↗](#) and 1E [↗](#)). The time course of the numbers of labeled vesicles in each zone showed that endocytosed transferrin was rapidly transported from early endosomes to lysosomes via late endosomes in VE cells, as reported previously (Richardson et al., 2000 [↗](#)). In addition, rhodamine-labeled dextran, which was transported by pinocytosis pathways, was also transported to lysosomes in a similar fashion (Fig. S1).

Two distinct modes of late endosomal fusion

To examine the dynamic changes of endocytic vesicles, we performed time-lapse imaging of Alexa 488-transferrin- or FM 1-43-labeled vesicles. We focused on late endosomes in zone 2 because these vesicles were large enough and suitable for microscopic observation of membrane fusion processes, as described below.

First, from 5 min after pulse labeling, fusion of two neighboring late endosomes was observed frequently (Fig. 2, A [↗](#) and B [↗](#); Video S1). The fusion showed an explosive mode, in which a membrane pore quickly expanded and two vesicles fused to become a single vesicle (Fig. 2B [↗](#)).




Figure 1.

Endocytic vesicles in VE cells

(A) Schematic diagram of the imaging setup. An E8.5 whole mouse embryo was observed with a laser confocal microscope. **(B)** Confocal images of VE cells stained with FM 1-43, LysoTracker Red, and anti-EEA1 antibody, or electroporated with GFP-Rab7. The yellow dotted lines indicate the cell boundaries. The yellow arrows show some vesicles that were positive for LysoTracker Red in zone 2. **(C)** For descriptive purposes, the supranuclear portion of VE cells is divided into 3 zones. Zone 1, zone 2, and zone 3 contain early endosomes (EE), late endosomes (LE), and lysosomes, respectively. **(D)** Confocal images of VE cells labeled with Alexa 488-transferrin. Images taken 1, 10, 20, and 40 min after pulse labeling are shown. The yellow dotted lines indicate the cell boundaries. **(E)** Time-course analysis of the Alexa 488-transferrin-containing vesicles in VE cells. $n = 4$. The scale bars indicate $4 \mu\text{m}$ (B) and $10 \mu\text{m}$ (D).

The size distribution of the late endosomes that underwent fusion ($7.3 \pm 6.6 \mu\text{m}^2$, $n = 34$) was similar to that of the total late endosomes in the observed area at the start of recording ($5.7 \pm 6.4 \mu\text{m}^2$, $n = 265$; **Fig. 2C**), indicating that late endosomes of any size underwent homotypic fusion. The fusion was completed within 20–50 s (the time required from membrane fusion to single spherical vesicle formation was 31.9 ± 0.2 s, $n = 37$), and the time was weakly correlated with the vesicle size ($r = 0.60$, **Fig. 2D**).

Second, from 15 min after pulse labeling, some late endosomes shrank gradually and disappeared from the focal plane (**Fig. 3A** and Video S2). This disappearance appeared to result from the fusion of late endosomes with the underlying lysosomes. This type of fusion, so-called bridge fusion, described in a previous study (Nicot et al., 2006; Roberts et al., 1999), occurred as a result of the slow mixing of vesicular contents through a narrow fusion pore. However, it was previously reported that the late endosomes of E6.5 mouse VE cells were engulfed by lysosomes (Kawamura et al., 2012). We therefore observed the disappearance of late endosomes in more detail using different approaches. First, we used differential labeling of late endosomes and lysosomes. When whole embryos were incubated with rhodamine-dextran for 20 min and subsequently pulse-labeled with Alexa 488-transferrin for 5 min, lysosomes and late endosomes were labeled with dextran and transferrin, respectively, at the time of observation (15 min after transferrin labeling). In a diminishing late endosome, the fluorescence intensity of transferrin in the vesicle increased initially and then decreased quickly (**Fig. 3, B–D**). At the same time, the rhodamine-dextran in the adjacent lysosome entered the late endosome (**Fig. 3, B–D**), indicating that the vesicular contents of the late endosomes and lysosomes were mixed. Second, using electron microscopy, we observed pore formation between a late endosome and a lysosome (**Fig. 3E**). These data suggest that the two vesicles underwent fusion but not engulfment in E8.5 embryos. Heterotypic fusion was slower than homotypic fusion and was completed within 90–250 s (154.8 ± 1.4 s, $n = 25$), and the fusion time (the time required from membrane fusion to disappearance of the vesicle) did not correlate with the vesicle size (**Fig. 3F**). The size of the labeled late endosomes at this time point ($10.1 \pm 8.1 \mu\text{m}^2$, $n = 256$) was larger than that at 5 min after labeling ($6.7 \pm 6.2 \mu\text{m}^2$, $n = 242$), indicating that larger late endosomes tended to undergo heterotypic fusion (**Fig. 3G**, see also below).

Vesicle size is an important factor for the transition of fusion modes

To examine whether the differences in membrane deformation of late endosomes depend on their fusion targets, we adopted a mathematical modeling approach that describes the membrane deformation processes after pore formation. According to thermodynamics, the deformation of a physical object proceeds as its free energy decreases. The free energy of a vesicle consists of two terms: bending energy and osmotic energy. The bending energy, b , is the integral of the square of the mean curvature all over the surface (Helfrich, 1973). The osmotic energy is the product of the osmotic pressure difference between the inside and the outside of the vesicle, Lp , and the vesicle volume, V . Thus, the free energy is given by

$$F = b - \Delta p \cdot V.$$

Note that the bending energy depends only on its shape; in other words, two membranes with the same shape but with different sizes have the same bending energy. On the other hand, the volume depends on both the size and the shape. Therefore, the vesicle size intrinsically affects the balance of both energy sources and may be a critical factor for membrane deformation during fusion.

We focused on a closed membrane having a constricted neck region and introduced two shape parameters, the neck width, w , and the area ratio between both sides of the neck, a (**Fig. 4A**), so that the basic processes of the two fusion types are described by the changes in w and a just after




Figure 2.

Homotypic fusion between late endosomes

(A) Time-lapse imaging of endocytic vesicles in VE cells. After 5 min of pulse-labeling with Alexa 488-transferrin, homotypic fusion of late endosomes (arrows) was frequently observed in zone 2. **(B)** By labeling of VE cells with FM1-43, the fusion process of the cell membranes in zone 2 was observed. **(C)** Histograms showing the size distribution of total late endosomes (black line, $n = 265$) and the late endosomes that underwent homotypic fusion (gray line, $n = 37$). **(D)** Correlation between the size of the late endosomes that underwent homotypic fusion and the time required for completion of fusion. The sizes of the larger of the fused vesicles are plotted. Thirty-seven fusion events were measured. The scale bars indicate 5 μm (A, top) and 1.5 μm (A, bottom; B).




Figure 3.

Heterotypic fusion between late endosomes and lysosomes

(A) Time-lapse imaging of VE cells. After 15 min of pulse-labeling with Alexa 488-transferrin, heterotypic fusion of late endosomes was observed: they shrank gradually and disappeared from the focal plane (arrows). **(B)** Time-lapse imaging of late endosomes in VE cells. Late endosomes and lysosomes were labeled with Alexa 488-transferrin (Tfn, green) and rhodamine-dextran (Dex, red), respectively. The bottom figures show optical sectioning through the dotted line in the upper figure. **(C and D)** Time course of the fluorescence intensity. In a late endosome undergoing fusion (ROI 1 in [B-C]), the fluorescence intensity of transferrin (green) slightly increased and then quickly decreased after membrane fusion. The fluorescence signal of dextran (red) in an underlying lysosome leaked into the fusing late endosome. The white box in (C) indicates the area shown in (B). The gray line in (D) indicates the diameter of the transferrin-positive late endosomes. The blue dotted lines in (D) indicate the moment the fusion pore was formed in ROI 1. ROI 2 is a negative control in a lysosome that did not undergo fusion. **(E)** Electron microscopic image showing pore formation between a late endosome (LE) and a lysosome (Ly). The inset shows a magnified picture of the boxed region. **(F)** Correlation between the size of late endosomes that underwent heterotypic fusion and the time required for completion of fusion. Twenty-five fusion events were measured. **(G)** Histograms showing the size distribution of late endosomes at 5 min (gray line, $n = 256$) and 15 min (black line, $n = 242$) after labeling with Alexa 488-transferrin. The average size of the vesicles was larger at 15 min than at 5 min ($**P < 0.01$, unpaired t-test). The scale bars indicate 5 μm (A, top), 1.5 μm (A, bottom), 2 μm (B), 3 μm (C), and 1 μm (E).

pore formation, $w = 0$. The relaxation of the neck in explosive fusion is expressed by the increase in w . On the other hand, the absorption of smaller vesicles by the larger one in bridge fusion is the decrease in w while w remains small.

When calculating the free energy as a function of w and a , the deformation is expressed as the process of descending the free-energy landscape in the space. Because the shape giving the minimum free energy is most likely to be realized, we first specified the minimum-free-energy shape and then calculated the free energy value for each parameter set.

We considered two extreme conditions: the membrane size is sufficiently small or large. In the former case, the osmotic energy is negligible and the free energy is approximated to

$$F \simeq F^{\text{small}} = b.$$

The free-energy landscape on the shape parameter space (w, a) is shown in **Fig. 4B**. In the latter case, although the osmotic energy is dominant, the bending energy is also not negligible because the bending energy at the neck region becomes considerably large.

Thus, the free energy for the large membrane case is

$$F \simeq F^{\text{large}} = B^{\text{neck}} - \Delta p \cdot V,$$

where B^{neck} is the bending energy at the neck. The free-energy landscape for a vesicle with the projected area of $10 \text{ } [\mu\text{m}^2]$ is shown in **Fig. 4C**.

These free-energy landscapes (**Fig. 4, B** and **C**) clearly differ from each other, implying that they drive different deformation processes. In particular, the contour of the F^{small} landscape around $w \sim 0$ is perpendicular to the w axis, whereas the contour of the F^{large} landscape is parallel to the w axis. Typical orbits of the shape deformations from the initial condition $(w, a) = (0.0, 0.3)$ generated by the gradients of the landscape are shown as gray arrows in **Fig. 4, B** and **C**. The orbit in the F^{small} landscape is characterized by a smooth increase in w ; i.e., the neck expands quickly (**Fig. 4D**), resulting in explosive fusion. On the other hand, in the F^{large} landscape, a increases through the orbit, whilst w remains small, which evokes bridge fusion (**Fig. 4E**).

The deformation processes observed in our modeling seem to be qualitatively consistent with our experimental results showing that the size of the fusion target changes the fusion modes (**Fig. 2** and **3**). Next, we made a rough estimate of how much of the vesicle size the mathematical model expects to switch the types of fusion. The typical bending energy of a (spherical) vesicle is $b \approx 2 \times 10^{-18} \text{ } [\text{J}]$ (Evans and Rawicz, 1990). Referring to the density differences in the dominant ions of endosomes and cytoplasm (Scott and Gruenberg, 2011), we assume that the total ion difference is in the range of $1\text{--}10 \text{ } \mu\text{m}$, and the osmotic pressure difference is $\Delta p \approx 2.5\text{--}25 \text{ } [\text{Pa}]$. From these assumptions, $0.22\text{--}1.0 \text{ } [\mu\text{m}^2]$ would be the vesicle size to switch fusion modes. However, the size of the observed late endosomes ($5\text{--}20 \text{ } \mu\text{m}^2$ in **Fig. 3F**) is much larger than the predicted size, suggesting that homotypic fusion between late endosomes should show the bridge fusion mode.

To resolve these inconsistencies, we considered the fluctuation effects. Intracellular membranes receive various fluctuations including thermal noise and active fluctuation from the cytoskeletons and motor proteins (Biswas et al., 2017; Chen et al., 2009). Although orbits descending the F^{large} landscape are energetically restricted to keep w small, these fluctuations may force the orbits to deviate from the restricted path (the dashed arrow in **Fig. 4C**); i.e., they are expected to induce explosive fusion. Then, we performed Monte-Carlo simulations to generate orbits descending the free-energy landscape with various magnitudes of fluctuations and determined the probabilities of orbits that deviate from the path (**Fig. 4F**). When we considered only the




Figure 4.

Mathematical modeling analysis of membrane deformation after connection of 2 vesicles

(A) Definition of shape parameters. **(B)** Free-energy landscape for small vesicles. The gray arrow (D) indicates a typical orbit of membrane deformation from the initial condition $(w, a) = (0.0, 0.3)$. **(C)** Free-energy landscape for large vesicles. The solid gray arrow (E) indicates the typical orbit of membrane deformation, whereas the dashed gray arrow (D') indicates the typical orbit of membrane deformation under fluctuated conditions. The inset is a blowup of the region $w = 0 \sim 0.15$, $a = 0.25 \sim 0.35$. **(D-E)** Time series of shape variation on the typical orbits indicated in (B) and (C), respectively. Note that the exact time scale cannot be estimated in this analysis. **(F)** Probability of the explosive fusions in the F_{large} landscape (the orbit of the dashed gray arrow in [C]) for various vesicle sizes, osmotic pressure differences, and fluctuation energies, estimated by Monte-Carlo simulations. $\Gamma = 2\pi\kappa\sqrt{A}/R_0$ with the bending modulus $k = 10[kBT]$ and the minimum bending radius $R_0 = 100[nm]$. The green shaded region indicates the condition: $\Delta p \approx 2.5 \sim 25[Pa]$, $V = 4A\sqrt{A}/3\sqrt{\pi}$ with $A = 2 \sim 25[\mu m^2]$ and the fluctuation energy $= 1[k_B T]$ (thermal fluctuation at room temperature). The green shaded region corresponds to late-endosome-sized vesicles ($2.0 \sim 25[\mu m^2]$) under estimated osmotic pressure difference ($\Delta p \approx 2.5 \sim 25[Pa]$) and the thermal noise at room temperature. The blue-shaded region indicates the vesicles under larger fluctuations.

thermal noise with room temperature with the estimated osmotic pressure difference ($\Delta p \approx 2.5\text{--}25$ [Pa]), as for late-endosome-sized vesicles ($2.0\text{--}25$ [μm^2]), the simulation predicts that vesicles will show bridge fusion (the green-shaded region in **Fig. 4F**). However, if we take into account the fact that a larger fluctuation allows explosive fusion (the blue-shaded region in **Fig. 4F**), fluctuations generated by the cytoskeleton and motor proteins, much larger than thermal noise, may induce explosive fusion.

Actin filaments are bound to late endosomes in VE cells

Previously we found that lysosome biogenesis was regulated by actin dynamics in VE cells (Koike et al., 2009). To examine the relationship between actin filaments and endocytic vesicles, actin filaments were visualized by use of Alexa 488-phalloidin. Endocytic vesicles were visualized by immunostaining with anti-mouse IgG because IgG is endocytosed from the maternal tissue *in utero* and can be stained without incubation of embryos *in vitro* for the pulse labeling of endosomes. In VE cells, strong signals were detected to be associated with the cell membrane (**Fig. 5, A–C**). In addition, in zone 1, fuzzy cytoplasmic staining was observed (**Fig. 5A**), which appeared to correspond to the actin meshwork in the terminal web. In zone 2, small punctates and short filaments were detected in the cytoplasm (**Fig. 5B**) and were bound to the surface of late endosomes (**Fig. 5, B, B'**, and **B''**) and infrequently to lysosomes in zone 3 (**Fig. 5, C, C'**, and **C''**). Optical cross-sectional images and 3D reconstruction showed that several rod-like actin filaments, extending from the apical side of VE cells into the cytoplasm, surrounded and held the late endosomes (**Fig. 5, D–F**). These data indicate that most actin punctates observed in the x-y plane correspond to the cross-sectional images of actin filaments extending in the apical-to-basal direction. In addition, when a fusion protein of an actin-bundling protein, fascin-1, with a red fluorescent protein was expressed in VE cells by electroporation, fascin-1 was highly colocalized with the actin punctates, suggesting that most of the actin punctates surrounding the late endosomes were bundled (**Fig. 5G**).

To investigate the endosome-associated actin filaments in more detail, we performed high-resolution imaging analysis using a deconvolution technique, which had resolution comparable to super-resolution microscopy. It demonstrated that fine actin filaments extended radially from the endosomal membranes in the x-y plane, in addition to the actin filaments extending in the apical-basal axis (**Fig. 5H**).

To examine the dynamics of actin filaments associated with late endosomes, E7.5 whole embryos were electroporated with a pEGFP-actin vector, and the EGFP signals in VE cells were observed by use of confocal microscopy 24 h after electroporation. Because high levels of actin overexpression led to the deterioration of cell functions (see white arrows in **Fig. S3A**), cells expressing low levels of EGFP-actin were selected for analysis: those selected cells were normal in morphology and endocytosis (see white arrowheads in **Fig. S3A**). Punctate signals of EGFP-actin were seen on the late endosomal membrane similarly to those observed in Alexa 488-phalloidin labeling. Time-lapse imaging showed that some new spots had appeared, others had disappeared, and two actin spots were unified on the membrane surface of endocytic vesicles (**Fig. 6A**). To examine the turnover rate of actin spots in VE cells, we performed fluorescence recovery after a photobleaching (FRAP) assay. After EGFP-actin on a late endosome was photobleached, recovery of its fluorescence was measured (**Fig. 6B**). Recovery of actin spots on late endosomes occurred more rapidly than the control recovery measured at the cell surface ($t_{1/2} = 29.0$ s; **Fig. 6, C** and **D**). Especially, the recovery of actin spots in the apical area of late endosomes was faster than in the basal area ($t_{1/2} = 11.5$ s in the apical area vs. $t_{1/2} = 18.9$ s in the basal area; **Fig. 6C**). These results indicated that actin spots on the surface of late endosomes were dynamically regulated, especially in the apical area.

The time-lapse images showed that when late endosomes moved in an x-y plane in zone 2, spots of EGFP-actin were always at the rear-end of the endosomes along the moving direction (**Fig. 6E**, $n = 13$). Because it is known that Arp2/3-dependent actin polymerization on the membrane acts as




Figure 5.

Actin filaments are associated with late endosomes in VE cells

(A–C) Confocal images of E8.5 mouse embryos stained with Alexa 488-phalloidin and anti-mouse IgG. Mouse IgG was incorporated into late endosomes and used as a marker for endosomes. Punctate cytoplasmic staining of phalloidin (green) was associated with the IgG-containing endocytic vesicles (red) in zone 2 (B–B'') and at lower frequencies with the lysosomes in zone 3 (arrows, C–C''). (D) Optical sectioning of VE cells along the dotted line in the upper panel. Actin filaments, extending from the apical surface into the cytoplasm (white arrowheads), were closely associated with the endocytic vesicles (red). (E and F) Three-dimensional surface rendering of actin filaments and a late endosome. The side view (E) and basal view (F) are shown. (G) Alexa 488-phalloidin staining of mRFP-fascin-1-expressing VE cells. Actin punctates were highly colocalized with fascin-1, whereas fine actin filaments extending radially in the x-y plane from endosomes were negative for mRFP-fascin-1 (white arrow heads). The fluorescence intensity plot along the white arrow in the middle panel shows colocalization of fascin-1 and phalloidin in the large spot. (H) Deconvolution images of late endosomes and actin. In addition to strong phalloidin-positive spots (white arrow heads), which were observable by use of conventional confocal microscopy, actin filaments extending from the endosomal surface (orange arrow heads) were visualized in the x-y axis image. In the x-z axis image, a late endosome was surrounded by actin filaments, which extended from the apical cell surface to the basal side in the cytoplasm. The scale bars indicate 3 μm (A–D), 5 μm (G), and 1 μm (E–F, H).




Figure 6.

Dynamic regulation of actin filaments surrounding late endosomes

(A) Representative time-lapse images of GFP-actin spots on a late endosome. VE cells were electroporated with GFP-actin at E7.5, and time-lapse analysis was performed on the next day. The arrow indicates the fusion of actin spots. The arrowhead indicates the appearance of a new actin spot. **(B)** The turnover rate of actin was analyzed by use of a FRAP assay. The area ($4\ \mu\text{m} \times 4\ \mu\text{m}$) covering a single late endosome in VE cells expressing GFP-actin was bleached, and the fluorescence recovery was observed by use of time-lapse microscopy. The yellow boxes indicate the laser-irradiated area. The time laser irradiation was ended is indicated as 0. **(C-D)** Graphs showing the fluorescence recovery of GFP-actin surrounding late endosomes in the apical region of zone 2 (light blue in [C], $n = 4$) and the basal region of zone 2 (dark blue in [C], $n = 3$) or at the cell surface of VE cells (D, $n = 8$). Fluorescence intensity was normalized to prebleaching intensities. **(E)** GFP-actin localization during vesicle movement. Endocytic vesicles (red) visualized by use of rhodamine-dextran in zone 2 are shown. When a vesicle (arrow) moved in a certain direction (upward in this figure), the actin patch (arrowhead) was observed at the rear end of the movement direction. The dotted line indicates the initial position of the vesicle. The scale bars indicate $1\ \mu\text{m}$ (A) and $3\ \mu\text{m}$ (B, E).

the driving force of the motility of intracellular pathogens (Welch and Way, 2013 [↗](#)) and endosomes/lysosomes in *Xenopus* egg extracts (Taunton et al., 2000 [↗](#)), we examined the localization of Arp2/3 in VE cells to test the possible role of Arp2/3 in endosome migration. When a fusion protein of Arp3-EGFP was expressed in VE cells by electroporation, the EGFP signals were distributed in dots on late endosomes and partially colocalized with phalloidin spots in the x-y plane in zone 2 (Fig. S3B). In addition, Arp3-EGFP signals were always seen at the rear-end of the endosomes along the moving direction (Fig. S3C). Taken together, these data suggest that the radial branched filaments are temporally polymerized to provide the propulsive force on the membrane for intracellular movement.

Actin dynamics regulate homotypic fusion of late endosomes

To examine the roles of actin in late endosomal fusion, we tested whether perturbation of actin dynamics caused defects in vesicle fusion. Treatment with cytochalasin D (a cell-permeable inhibitor of actin polymerization) for 5 min reduced the frequency of homotypic fusion in a dose-dependent manner (Fig. 7, A [↗](#) and B [↗](#); Video S3). In addition, it altered the speed and mode of the fusion. Treatment with 0.1 or 1.0 μ M cytochalasin D slowed down homotypic fusion (Fig. 7, C [↗](#) and D [↗](#); Fig. S4A). After membrane fusion, two endosomes remained in a peanut shape for a long time, indicating slower neck expansion than that in the control cells (Fig. S4A). FM 1-43 labeling revealed that the membranes were rapidly fused, whereas it took a long time for the gourd-shaped vesicles to form a large round vesicle (Fig. S4A). Furthermore, upon 1.0 μ M cytochalasin D treatment, about one-third of homotypic fusion showed bridge fusion (Fig. 7, C [↗](#) and E [↗](#); Video S4). Therefore, cytochalasin D slowed down the fusion processes and changed the fusion mode to bridge fusion at higher concentrations. These data are consistent with our mathematical model showing that larger fluctuation is necessary for explosive fusion in homotypic fusion of late endosomes (Fig. 4F [↗](#)). Treatment with jasplakinolide (a cell-permeable stabilizer of actin filaments) also reduced the frequency of homotypic fusion in a dose-dependent manner (Fig. 7, A [↗](#) and B [↗](#); Video S5), indicating that dynamic actin turnover is required for homotypic fusion.




Figure 7.

Actin dynamics regulate homotypic fusion in VE cells

(A) Confocal images of endocytic vesicles in zone 2. E8.5 mouse embryos pretreated with 0.1 μ M cytochalasin D (CD) or 0.1 μ M jasplakinolide (Jasp) were labeled with Alexa 488-transferrin for 5 min. Time-lapse images were taken for 400 s from 5 min after labeling. Arrows indicate homotypic fusion. **(B)** Frequency of homotypic fusion calculated from the number of fusions that occurred in 400 s. Data are represented as means $\pm$ SEMs of 4–9 independent experiments; individual data are shown by circular dots. Data were analyzed by use of one-way ANOVA with the Tukey multiple comparison test (** P <0.01; *** P <0.001). **(C)** Quantitation of types of homotypic fusion. Fusion types are classified into normal explosive fusion, slow explosive fusion (more than 60 s), and bridge fusion. Cytochalasin D treatment slowed down the fusion process in a dose-dependent manner and induced bridge fusion at 1 μ M. Independent experiment numbers are shown on the top of the bars. **(D)** Quantitation of the time for completion of homotypic fusion. Treatment with 0.1 μ M or 1 μ M cytochalasin D or with 20 μ M CK666, but not with 0.1 μ M jasplakinolide, resulted in a significant increase in the fusion time. Data are represented as means $\pm$ SEMs of 4–30 independent experiments; individual data are shown by circular dots. Data were analyzed by use of one-way ANOVA with the Tukey multiple comparison test (** P <0.01; *** P <0.001). **(E)** Representative images of homotypic fusion of late endosomes in VE cells treated with 1 μ M cytochalasin D. Endosomes were visualized by incubation with Alexa 488-transferrin. The arrows indicate the vesicles undergoing homotypic fusion in the bridge fusion mode. **(F)** Time-lapse imaging of GFP-actin localization during homotypic fusion of late endosomes. Localization of GFP-actin (green) and endocytic vesicles (red), visualized by use of rhodamine-dextran, were observed. When a pair of endosomes (arrows) fused, GFP-actin was newly polymerized at the opposite positions of the docking site and its fluorescence intensity was increased (arrowheads). **(G)** Distribution of GFP-actin during homotypic fusion of late endosomes. The fluorescence intensities of GFP-actin (dark gray, n = 5) and FM 1-43 (light gray, n = 5) were measured during homotypic fusion. The percentages of their signals in the proximal (docking site, shown as 1), lateral (shown as 2), and distal (shown as 3) regions at the predocking, docking, rounding, and postfusion stages are shown. Data are represented as means $\pm$ SEMs (* P <0.05, unpaired t-test). **(H)** The time course of changes in the fluorescence intensity of GFP-actin around a pair of late endosomes during homotypic fusion. Time 0 s indicates the moment the fusion pore of two late endosomes was formed. The graph indicates the average of 5 independent experiments. The scale bars indicate 10 μ m (A), 2 μ m (E), and 3 μ m (F).

We next observed the distribution of actin filaments during homotypic fusion. When late endosomes underwent fusion, EGFP-actin punctates were excluded from the membrane docking sites and highly polymerized transiently at both ends of the fusing vesicles (opposite to the docking region) during the rounding period after membrane fusion (**Fig. 7F**). Fluorescence intensity of EGFP-actin on endosomes was quantitated in the proximal, lateral, and distal regions at the predocking, docking, rounding, and postfusion stages. This analysis indicated that actin filament distribution was spatially regulated during fusion (**Fig. 7G**, dark gray bar), whereas FM 1-43 distribution, used as an unchanged control, was uniform on the membrane throughout the rounding stages (**Fig. 7G**, light gray bar). Furthermore, the total fluorescence intensity of EGFP-actin on late endosomes significantly increased transiently after pore formation (**Fig. 7H**), indicating that actin filaments were polymerized temporarily, not simply distributed differently. Moreover, accumulation of Arp3-EGFP in the distal region during the rounding stage was observed similarly to EGFP-actin accumulation (**Fig. S4B**), and treatment with CK-666, an Arp2/3 inhibitor, resulted in a decrease in the frequency of homotypic fusion (**Fig. 7B**) and slowed down homotypic fusion (**Fig. 7, C** and **D**). These results suggest that branched actin filaments nucleated by the Arp2/3 complex on the distal surface of late endosomes are necessary to transform the vesicle into a large round sphere quickly after membrane fusion of late endosomes.

Homotypic and heterotypic fusions are differentially regulated by actin

We next examined the role of actin filaments in heterotypic fusion. Cytochalasin D treatment led to a significant reduction in the frequency of heterotypic fusion in a dose-dependent manner (**Fig. 8, A–C** and Video S6). In contrast, jasplakinolide treatment did not inhibit heterotypic fusion and somehow increased the frequency of heterotypic fusion with lysosomes as early as 5 min after pulse labeling (**Fig. 8B** and Video S7). In addition, CK-666 treatment had no effects, suggesting that Arp2/3 was not required for heterotypic fusion (**Fig. 8C**). These data suggest that actin played distinct roles in homotypic fusion and heterotypic fusion and that stable actin filaments are implicated in driving heterotypic fusion.

Myosins are a superfamily of actin-based motor proteins and their members are grouped into many classes according to their structures and functions (Hartman and Spudich, 2012). To determine which class of myosins is important for vesicle fusion and motility, we examined the effects of inhibitors for several myosin classes that are known to be involved in vesicle transport. We tested inhibitors of myosin I (pentachloropseudilin, PCIP), nonmuscle myosin II (NM II) (blebbistatin), myosin V (MyoVin-1), and myosin VI (2,4,6-triiodophenol, TIP; Bond et al., 2013). Heterotypic fusion frequency was reduced to 20% by MyoVin-1, to a lesser extent but significantly by blebbistatin, but was not affected by PCIP or TIP (**Fig. 8D**). In contrast, homotypic fusion frequency was significantly reduced by PCIP, MyoVin-1, and TIP (**Fig. S5**). EGFP-myosin IIA signals overlapped with phalloidin-positive spots on late endosomes (**Fig. 8E**), whereas EGFP-myosin Va signals were juxtaposed to phalloidin-positive spots on late endosomes (**Fig. 8E**).

Cofilin-dependent actin dynamics are involved in late endosome fusion

Late endosomes undergo homotypic fusion almost exclusively at the beginning and then switch to heterotypic fusion over time after endocytosis (**Fig. 9A**). Given that actin plays a different role in homotypic fusion and heterotypic fusion and that pharmacologic changes in actin dynamics shift the fusion mode as shown above, it is suggested that actin dynamics are a critical factor for the transition of fusion modes. Thus, we next decided to look into the regulation of actin dynamics in two different fusion modes. Previously, we found that lysosome biogenesis was regulated by actin dynamics through cofilin downstream of the autotaxin–LPA receptor–Rho–ROCK–LIM kinase pathway in VE cells (Koike et al., 2009). When VE cells were incubated with the S3 peptide, which contains the N-terminal 16 amino acid sequence of cofilin fused to a cell permeation sequence and is used as a competitive inhibitor for LIM kinase and serves as a cofilin activator (Maekawa et al., 1999; Aizawa et al., 2001), lysosomes were fragmented in VE cells and the fluorescence intensity of phalloidin in VE cells was reduced (Koike et al., 2009).

We thus focused on cofilin in endosomal fusion in VE cells. YFP-cofilin showed small punctate staining that colocalized with phalloidin-positive spots in the x-y plane and extended along the phalloidin signal (**Fig. 9, B and C**). When the fluorescence intensity of Alexa 546-phalloidin- and YFP-cofilin-positive filaments was quantitated along the apical-basal axis of late endosomes in zone 2, cofilin was slightly abundant in the apical region, whereas the phalloidin signal was stronger in the basal region (**Fig. 9D**). A FRAP assay of EGFP-actin on a late endosome showed that treatment of E7.5 whole embryos with the S3 peptide (15 µg/ml) for 24 h resulted in faster turnover of actin filaments of late endosomes when compared with treatment with the RV peptide (Aizawa et al., 2001), which contains the reverse sequence of cofilin and is used as a negative control ($t_{1/2} = 10.9$ s with the S3 peptide, $n = 9$ vs $t_{1/2} = 18.3$ s with the RV peptide, $n = 9$; **Fig. 9E**). These results indicate that cofilin distributes with actin filaments on late endosomes and regulates actin dynamics on late endosomes. Finally, the roles of cofilin in late endosomal fusion were investigated. After treatment with the S3 peptide for 24 h, Alexa 488-transferrin-labeled vesicles were observed by use of time-lapse imaging. When compared with the RV peptide, the S3 peptide




Figure 8.

Actin stabilization induces heterotypic fusion of late endosomes with lysosomes

(A) Confocal images of endocytic vesicles in zone 2. E8.5 mouse embryos pretreated with 0.1 μM cytochalasin D (CD) or with 0.01 μM jasplakinolide (Jasp) were labeled with Alexa488-transferrin for 5 min. Time-lapse images were taken for 400 s from 15 min after labeling. The arrows indicate heterotypic fusion with lysosomes. **(B)** Frequency of heterotypic fusion calculated from the number of fusions that occurred in 400 s from 5 min after labeling. Data are represented as means $\pm$ SEMs of 4–7 independent experiments; individual data are shown by circular dots. Data were analyzed by use of one-way ANOVA with the Tukey multiple comparison test (** $P<0.01$, *** $P<0.001$). **(C)** Quantitation of heterotypic fusion frequency at 15 min after labeling of VE cells pretreated with cytochalasin D (CD), jasplakinolide (Jasp), or CK666. Data are represented as means $\pm$ SEMs of 4–8 independent experiments; individual data are shown by circular dots. Data were analyzed by use of one-way ANOVA with the Tukey multiple comparison test (* $P<0.05$, ** $P<0.01$). **(D)** Effects of myosins on heterotypic fusion. Frequency of heterotypic fusion was calculated at 15 min after labeling. Data are represented as means $\pm$ SEMs of 5–8 independent experiments; individual data are shown by circular dots. Data were analyzed by use of one-way ANOVA with the Tukey multiple comparison test (* $P<0.05$, *** $P<0.001$). **(E)** Colocalization of EGFP-myosin IIA or EGFP-myosin Va with phalloidin in zone 2 of VE cells. The insets show the magnified images of the boxed area. Following electroporation of the EGFP-fused myosin expression plasmids and 24-h culture, whole embryos were stained with Alexa546-phalloidin. EGFP-myosin IIA signals overlapped with phalloidin-positive spots on late endosomes, whereas most of the EGFP-myosin Va signals were juxtaposed with phalloidin-positive spots on late endosomes (see the schemas and fluorescence intensity plots). The scale bars indicate 10 μm (A) and 3 μm (D).




Figure 9.

Transition from homotypic to heterotypic fusion

(A) Time course of frequencies of homotypic (blue line) and heterotypic fusion (red line). The graph indicates the means of 6 independent experiments. **(B)** Distribution of cofilin in zone 2 of VE cells. After whole embryos were electroporated with YFP-cofilin and cultured for 1 d, they were stained with Alexa 546-phalloidin. The lower graph plotting the fluorescence intensity along the white arrow in the upper image indicates colocalization of actin filaments and cofilin. **(C)** Optical sectioning images (right) of VE cells along the dotted line in the x-y axial panel (left). Embryos, which were electroporated with YFP-cofilin and cultured for 1 d, were stained with Alexa 546-phalloidin. The actin filaments, extending from the apical to the basal region, were highly colocalized with cofilin. Note that the white arrows and blue arrowheads indicate an actin filament that surrounds late endosomes and the cell membrane, respectively. **(D)** Quantification of the fluorescence intensity of phalloidin and cofilin along the apical-basal axis. The intensity of spots colocalized with phalloidin and cofilin around late endosomes observed in the x-y images was quantified along the z-axis. Fluorescence intensity was normalized to the intensities in the most apical plane (8 late endosomes from 3 VE cells). Data were analyzed by use of one-way ANOVA with the Tukey multiple comparison test and represented as means $\pm$ SEMs (** $P < 0.01$, *** $P < 0.001$). **(E)** FRAP assay to evaluate the dynamics of actin at the apical region of zone 2 in VE cell pretreated with the S3 peptide. After treatment with the S3 or RV peptide overnight, the area ($4 \mu\text{m} \times 4 \mu\text{m}$) covering a single late endosome in VE cells expressing EGFP-actin was bleached and the fluorescence recovery was observed by time-lapse microscopy. Three independent experiments were performed. **(F–G)** Frequency of homotypic (F) and heterotypic fusion (G) in embryos pretreated with 15 $\mu\text{g}/\text{ml}$ S3 or RV peptides. Frequencies of homotypic and heterotypic fusions were calculated according to the numbers of fusions in 400 s from 5 min (F) and 15 min (G) after labeling with Alexa 488-transferrin. Data are represented as means $\pm$ SEMs of 3–4 independent experiments; individual data are shown by circular dots. Data were analyzed by use of one-way ANOVA with the Tukey multiple comparison test (* $P < 0.05$). The scale bars indicate 8 μm (B) and 2 μm (C).

strongly inhibited homotypic fusion (**Fig. 9F**) but not heterotypic fusion (**Fig. 9G**). These data suggest that spatial differentiation of actin dynamics along the apical-basal direction, partially contributed by cofilin, may be a critical factor for the transition of fusion modes.

Discussion

In this study, we developed a novel imaging method of analyzing the motile behavior of late endosomes in VE cells. By using the technique, we found that VE cells have two different types of vesicle fusion: homotypic fusion between late endosomes and heterotypic fusion between late endosomes and lysosomes. We found that actin filaments were associated with late endosomes and that the forces generated by actin were required for inducing quick fusion of late endosomes and for facilitating transport of late endosomes to lysosomes. We also showed that cofilin plays a role in late endosome fusion via actin turnover (**Fig. 10**).

Actin filaments are associated with endosomal vesicles

We found two types of actin filaments associated with late endosomes in VE cells: those that extended along the apical-basal axis and those that extended radially from the endosomes (**Fig. 5**). Our findings resemble the previous observation that early endosomes were distributed along with actin cables and that Arp2/3-dependent small cloud-like patches protruded from the membrane (Morel et al., 2009). Actin is reportedly distributed around vesicles in different patterns. In addition to the well-known interaction of vesicles with actin rails in directional vesicle transport (Titus, 2018), actin was localized at the vertex ring in vacuole fusion in yeast (Wang et al., 2002) and entirely surrounded phagosomes in macrophages (Liebl and Griffiths, 2009). Advanced microscopy techniques have clarified the fine structure of actin, which is assembled into polyhedron-like lattices around vesicles in exocrine glands (Ebrahim et al., 2019).

We observed strong actin signals as patches on late endosomes in the focal planes in confocal microscopy. Three-dimensional reconstruction showed that most of these patches represent the cross-section of rod-shaped actin filaments extending from the apical side into the cytoplasm in VE cells. In addition to these patches, some signals seem to represent actin filaments extending radially from membranes. Although it is difficult to distinguish between these two types of patches by use of conventional confocal microscopy, we successfully observed finer actin filaments extending radially from the endosomal membranes by use of high-resolution imaging analysis. Furthermore, we showed that Arp3-positive actin spots appeared in the site opposite to that of endosome movement (**Fig. 6E**, Fig. S3C) like a comet tail (Goley and Welch, 2006), that Arp3-positive actin spots were asymmetrically distributed and temporarily polymerized in the distal region during endosomal fusion (**Fig. 7, F** and **G**; Fig. S4B), and that Arp2/3 inhibition reduced homotypic fusion (**Fig. 7B**). These results suggest that radial actin filaments polymerized by Arp2/3 generate force for homotypic fusion in VE cells. This notion is compatible with the findings of previous reports that the radial filaments were nucleated and extended with Arp2/3-dependent branches on the lateral membrane of endosomes to push vesicles (Goley and Welch, 2006; Svitkina, 2018). Given that nucleation and Arp2/3 activities are regulated by nucleation-promoting factors (NPFs; Goley and Welch, 2006), the activity of NPFs and membrane fusion processes are likely tightly linked to accurate membrane fusion. Differential regulation of NPF activity may give rise to different patterns of actin surrounding vesicles and membrane dynamics. Future investigation is necessary to elucidate how the NPF activity is spatially and temporally regulated on membranes.




Figure 10.

Summary of late endosome fusion in VE cells

In VE cells, late endosomes (LE) show two different types of fusion with different targets. In homotypic fusion, once a membrane pore is formed between two late endosomes, the pore quickly expands and a single fused vesicle is formed. In contrast, in heterotypic fusion, the pore does not expand; instead, the late endosome gradually shrinks and disappears over time as the result of a transfer of its vesicle content into lysosomes. Two different types of actin are associated with the late endosomal membrane in VE cells: the actin filaments that extend along the apical-to-basal axis and the filaments that are radially polymerized from the membrane. Cofilin, an actin-binding protein that severs and depolymerizes actin filaments activates actin turnover. In the apical region, late endosomes receive squeezing forces via active actin modification by cofilin and Arp2/3, which leads to homotypic fusion in an explosive mode. In contrast, in the basal region, actin is more static and late endosomes receive sliding forces via myosins.

Actin regulates membrane behavior after pore formation

In this study, we found that the membranes of late endosomes behave differently depending on the fusion mode. In homotypic fusion, the fusion pore formed between docked endosomal vesicles expanded quickly and the fusion was completed within 20–30 s (explosive fusion, **Fig. 2**). In contrast, in heterotypic fusion with lysosomes, the pore opened very slowly and late endosomes gradually shrank in diameter and disappeared over 1–2 min (bridge fusion, **Fig. 3**). Using mathematical modeling, we showed that vesicle size is an important factor determining the fusion mode. When the size of the vesicle becomes larger, the free energy for neck relaxation decreases and the pore is kept small. Our modeling predicted that the late endosomes are large enough to allow bridge fusion to occur, although they exhibit explosive fusion *in vivo*. Similarly, homotypic fusion in an explosive mode was observed in very large vesicles, such as vacuoles in yeast, endosomal vesicles in coelomocytes of *C. elegans*, and secretory granules in the salivary glands (Liu et al., 2016; Masedunskas et al., 2011; Wang et al., 2002). These observations indicate that vesicle size is not the sole factor determining the fusion mode. Lipid composition, a well-known parameter affecting its bending modulus, is also involved but the effect on fusion mode may be less dominant when compared with the vesicle size effects (Simon et al., 1998; Faizi et al., 2019; Ridolfi et al., 2021). One of the critical factors other than vesicle size that determine the fusion mode is membrane fluctuation. We found that homotypic fusion was converted to bridge-like fusion after cytochalasin D treatment (**Fig. 7C**), which suggests that actin dynamics around vesicles cause fluctuation and affect the fusion modes. This finding can be successfully explained by introducing minor modifications in our mathematical modeling so that a larger fluctuation can shift the fusion mode from bridge to explosive fusion.

VE cells vigorously endocytose maternal proteins and supply the digested products to the developing embryo to support embryonic development. This seems to be why VE cells have such huge endosomal and lysosomal vesicles. However, larger vesicle size leads to the reduction in the vesicle fusion speed and subsequent nutrient trafficking efficiency. If vesicle size were the only factor determining the fusion mode, this disadvantage would occur, but if fluctuation is taken into account, it can be overcome, as shown in our mathematical modeling, and the forces generated by actin are thought to be responsible for increasing the rate of vesicular trafficking. This idea is supported by previous reports that knockout mice of actin-related genes (e.g., Arc and NM IIA) showed abnormal VE cells and embryonic lethality (Conti et al., 2004; Liu et al., 2000).

In heterotypic fusion, late endosomes fused with lysosomes in a bridge-like manner. This may be because lysosomes are too large in comparison with late endosomes and actin accumulation on the lysosomal membrane is scarce, resulting in insufficient force of actin for explosive fusion. Bridge fusion was previously observed in live cells only when giant vesicles were artificially generated by gene modification, for example, in endosomes enlarged by expression of constitutively-active Rab5 (Roberts et al., 1999) and lysosomes enlarged in the intestinal cells of *ppk-3*-deleted *C. elegans* (Nicot et al., 2006). These findings suggest that the enlargement of these vesicles exceeds the intrinsic forces of actin to induce explosive fusion.

Like homotypic fusion, heterotypic fusion is also dependent on actin, as shown by reduction in fusion frequency upon cytochalasin D treatment. Intriguingly, in heterotypic fusion, NM II, myosin V, and stabilized actin filaments, but not Arp2/3, were required, indicating a distinct contribution of actin to heterotypic fusion when compared with homotypic fusion. We showed that, consistent with the well-known roles of myosin V in trafficking of lysosomes, ER, melanosomes, and synaptic vesicles (Hammer III and Sellers, 2011), myosin V facilitates the directional transport of late endosomes towards lysosomes along stabilized actin filaments that serve as a rail. Moreover, our results indicate that NM II, which is primarily responsible for generating contractile force (Svitkina, 2018), partially contributes to heterotypic fusion. It is well established that NM II facilitates the constriction of the membrane into the plasma membrane after membrane pore formation in exocytosis (Ebrahim et al., 2019) in cooperation with actin depolymerization by cofilin-1 (Miklavc et al., 2015). In contrast, our data showed that actomyosin appeared not to

form the lattice on the surface of endosomes for the contraction and that cofilin does not show a robust influence on heterotypic fusion. Rather, actin stabilization tended to promote heterotypic fusion. These data suggest that NM II may contribute to heterotypic fusion through a different mechanism from that of exocytosis. NM II is also implicated in the movement of the nucleus along with cell migration, which is akin to a “conveyor-belt” by the sliding of the nuclear membrane along the actin filament (Gomes et al., 2005 [↗](#)). Further investigation is necessary to elucidate the mechanisms by which NM II transports late endosomes along actin filaments.

Determination of fusion targets

Late endosomes, which fuse frequently with neighboring late endosomes, shift the target of fusion towards lysosomes. What triggers the transition of the fusion targets? We found that larger endosomes tend to fuse with lysosomes (Fig. 3G [↗](#)). Thus, one possibility is that the larger endosomes located in the basal position have a higher opportunity to fuse with lysosomes than do the smaller endosomes in the more apical portion. However, upon treatment with jasplakinolide, even smaller endosomes distributing in the more apical area underwent heterotypic fusion with lysosomes (Fig. 8, A [↗](#) and B [↗](#)), thus suggesting rather that the size and the position of vesicles are not the sole factors and that the different dynamics of actin shift the fusion modes. Furthermore, we showed weaker actin polymerization/depolymerization activity in the basal area as compared with that in the apical area (Fig. 6C [↗](#)), a decrease in the frequency of homotypic fusion but not of heterotypic fusion induced by S3 peptide treatment (Fig. 9, F-G [↗](#)), and an increase in actin filaments on endosomes in the basal area as compared with in the apical area (Fig. 9D [↗](#)). These findings support the notion that cofilin-dependent regulation of actin polymerization/depolymerization may play a critical role in determination of the fusion target. In parallel with VE cells, it was shown that cofilin was specifically distributed in the apical area of epithelial cells and was highly active under the plasma membrane (Morris et al., 2011 [↗](#); Vitriol et al., 2013 [↗](#)). Cofilin regulates actin remodeling for lamellipodia formation and endosomal trafficking (Harrington et al., 2011 [↗](#); Vitriol et al., 2013 [↗](#)). Furthermore, early endosomes located close to the plasma membrane fuse frequently when compared with late endosomes located at the perinuclear region (Gruenberg et al., 1989 [↗](#)). Taken together, higher dynamics of actin filaments regulated by cofilin at the cell peripheral area may increase the fusion frequency in many cells.

In this study, we developed a method of observing and manipulating late endosomes in VE cells. This method enables, for the first time, the observation of fusion of single vesicles with two different targets and is very useful to examine the molecular mechanisms that determine the fusion targets of vesicles and regulate different fusion modes.

Materials and Methods

Animal experiments

All the experiments using animals were approved by the Animal Care and Use Committee of the University of Tsukuba and performed under its guidelines. Noon of the day on which a vaginal plug was observed was taken as embryonic day 0.5 (E0.5). Time-pregnant ICR mice were purchased from CLEA Japan and Japan SLC.

Immunohistochemistry and phalloidin staining

For whole-mount immunohistochemistry, E8.5 embryos were fixed with 4% paraformaldehyde (PFA) in PBS at 4°C for 3 h and incubated with FITC-conjugated anti-EEA1 (1:100, BD Bioscience) or Cy3-labeled anti-mouse IgG (1:500, Jackson ImmunoResearch). For phalloidin staining, the embryos were fixed with 4% PFA in PBS at 37°C for 15 min and incubated with 0.2 U/ml Alexa Fluor 488- or Alexa Fluor 546-conjugated phalloidin (Molecular Probes) for 60 min.

Ex vivo whole embryo culture

The whole embryo culture was performed as described previously (Koike et al., 2009). In brief, embryos were dissected at E7.5 and cultured in 100% rat serum (Charles River) supplemented with 2 mg/ml glucose in a culture bottle placed in a rotation drum culture system (Ikemoto Rika) at 37°C under 5% O₂, 5% CO₂, and 90% N₂ for 24 h. In the pharmacologic experiments, embryos were incubated with 15 µg/ml S3 and RV peptides (a kind gift of Dr Kensaku Mizuno, Tohoku University).

Endocytic uptake assays and intracellular vesicle tracing

Whole embryos with intact yolk sacs dissected out at E8.5 or embryos cultured for 1 d from E7.5 were transferred into DMEM (Sigma-Aldrich) supplemented with 0.1% BSA. Embryos were incubated in 100 nM LysoTracker Red (Molecular Probes) or 12 µM FM 1-43 (Molecular Probes) in 0.1% BSA/DMEM at 37°C for 5 min. For tracing endocytic vesicles, embryos were incubated with 20 µg/ml Alexa Fluor 488-labeled transferrin (Molecular Probes) or 100 µg/ml dextran rhodamine B 70,000 MW (Molecular Probes) at 37°C for 5 min. For differential labeling of late endosomes and lysosomes, whole embryos were initially cultured in 100% rat serum supplemented with 2 mg/ml glucose and 75 µg/ml dextran rhodamine B 70,000 MW at 37°C for 20 min, and then incubated with 20 µg/ml Alexa Fluor 488-labeled transferrin for 5 min. To test the effects of actin inhibitors, embryos were incubated with 0.01–1 µM cytochalasin D (Sigma-Aldrich), 0.01–1 µM jasplakinolide (Calbiochem), 20 µM CK666 (Calbiochem), 10 µM pentachloropseudilin (AOBIOUS), 10 µM blebbistatin (Sigma-Aldrich), 10 µM MyoVin-1 (Calbiochem), and 10 µM 2,4,6-triiodophenol (Sigma-Aldrich) for 5 min, and the same drugs were added to the medium during observation.

Image acquisition and analysis

An embryo was placed on a glass-bottom dish (No.1S thickness, Matsunami Glass), and confocal fluorescent images were obtained by use of a LSM510 laser scanning microscope (Carl Zeiss) equipped with a C-APOCHROMAT 63x/1.2 W Korr objective lens (Carl Zeiss) and a pinhole set to 1.0 AU. Time-lapse images were obtained by collecting 40 images with time intervals of 5 or 10 s at room temperature. The series of images of the confocal z-stacks were obtained with a lateral resolution of 1024 x 1024 pixels and a scanning depth of 6–8 µm with 13–17 layers and 0.5-µm intervals.

For high-resolution imaging, the samples on the glass-bottom dish were mounted with ProLong Glass (Thermo Fisher Scientific). Images were captured by use of a TCS SP8 laser confocal microscope (Leica Microsystems) equipped with an HC PL APO CS2 100x/1.40 OIL objective (Leica). Alexa Fluor 488 and Alexa Fluor 594 dyes were excited at 488 and 594 nm of a supercontinuum white light laser, respectively. The emitted fluorescence was sequentially detected by use of HyD-SMD detectors through a spectral separation module (505–550 nm for AlexaFluor 488 and 605–700 nm for Alexa Fluor 594). The pinhole size was set at 0.40 Airy units (60.7 µm). The pixel size and z-step were set at 45 nm and 182 nm, respectively. The acquired images were processed using the Lightning deconvolution algorithm on LAS-X software (Leica). Three-dimensional reconstructions of the images were carried out on NIS-Elements software (Nikon) or Imaris software (Bitplane).

Electroporation of whole embryos

The electroporation of whole embryos was described previously (Koike et al., 2009). An E7.5 embryo was inserted into a small hole of an agarose mold (2% agarose/PBS, 7 x 7 mm) placed in Hanks buffered saline solution (HBSS). The solution bathing the embryos was then replaced with HBSS containing 2 mg/ml DNA. Three square electric pulses (30 V, 50 ms duration, 1 pulse/s) were delivered to the agarose mold by use of an electroporator (CUY21, Nepa Gene) and a 10-mm electrode (CUY650P10, Nepa Gene). After electroporation, the embryo was rinsed with HBSS and cultured as described in the previous section. Rab7, nonmuscle myosin IIA, and myosin Va subcloned into a pEGFP-C2 expression vector (Clontech), Arp3 subcloned into a pEGFP-N1

expression vector (Clontech), fascin-1 subcloned into a pmRFP-C1 expression vector (Clontech), pEGFP-Actin (BD Biosciences), and YFP-cofilin (kind gifts of Dr Kensaku Mizuno) subcloned into the pEGFP-C1 vector (Clontech) were used.

Photobleaching assay

Photobleaching analysis was performed with the LSM510 laser scanning microscope. The EGFP-actin-electroporated embryo was observed by use of a 63x objective lens. For the bleaching, an ROI was drawn around a late endosome (4 μm x 4 μm) or cell boundary and was bleached by use of a 488-nm laser beam for 8 bleach iterations at 100% intensity, and fluorescence recovery into the bleached area was monitored. To calculate the percentage of recovery after photobleaching, the intensity profiles for the actin filaments in the bleached area, a background region, and the whole cells were collected. The background was subtracted from all the data, and the intensity of the whole cells was used to correct for fluorescence loss during image acquisition.

Electron microscopy

E8.5 mouse embryos were fixed with 2% glutaraldehyde and 2% PFA in 0.1 M phosphate buffer (pH 7.4) overnight. After 3 washes with 0.1 M phosphate buffer, pH 7.4, the embryos were postfixed with 1.0% osmium tetroxide in 0.1 M phosphate buffer for 2 h, dehydrated, and embedded in Epon 812 according to a standard procedure. Ultrathin sections were stained with uranyl acetate and lead citrate and observed by use of an electron microscope (H-7100, Hitachi).

Acknowledgements

This work was supported in part by KAKENHI (17024006) and the 21st Century COE program from the MEXT, Japan, KAKENHI (JP22H04926), Grant-in-Aid for Transformative Research Areas – Platforms for Advanced Technologies and Research Resources “Advanced Bioimaging Support” from JSPS, Japan, and Takeda Science Foundation. We thank Drs. K. Mizuno, K. Ohashi, and N. Ohsumi, for reagents, protocols, and advice; Y. Sato (Carl Zeiss, Japan) for help in 3D reconstruction; and H. Bito and F. Miyamasu for useful comments on the manuscript.

Author contributions

Conceptualization, S.K., T.O., K.K.-M., and M.M.; Methodology, S.K., M.Ta., M.Ts., and T.N.; Software, M.Ta.; Formal analysis, S.K. and M.Ta.; Investigation, S.K., T.O., K.K.-M., M.Ts., T.N., and M.M.; Writing—Original Draft, S.K. and M.M.; Writing—Review & Editing, S.K., M.Ta., T.O., K.K.-M., and M.M.; Visualization, S.K. and M.Ta.; Supervision, M.M.; Funding acquisition, S.K. and M.M.

Declaration of interests

The authors declare no competing interests.

References

- Aizawa H. *et al.* (2001) **Phosphorylation of cofilin by LIM-kinase is necessary for semaphorin 3A-induced growth cone collapse** *Nat. Neurosci* **4**:367–373 <https://doi.org/10.1038/86011>
- Bielinska M., Narita N., Wilson D.B. (1999) **Distinct roles for visceral endoderm during embryonic mouse development** *Int. J. Dev. Biol* **43**:183–205
- Biswas A., Alex A., Sinha B. (2017) **Mapping Cell Membrane Fluctuations Reveals Their Active Regulation and Transient Heterogeneities** *Biophys. J* **113**:1768–1781 <https://doi.org/10.1016/j.bpj.2017.08.041>
- Bond L. M., Tumbarello D.A., Kendrick-Jones J., Buss F. (2013) **Small-molecule inhibitors of myosin proteins** *Future Med. Chem* **5**:41–52 <https://doi.org/10.4155/fmc.12.185>
- Chen C.H., Tsai F.C., Wang C.C., Lee C.H. (2009) **Three-dimensional characterization of active membrane waves on living cells** *Phys. Rev. Lett* **103** <https://doi.org/10.1103/PhysRevLett.103.238101>
- Chernomordik L.V., Kozlov M.M. (2008) **Mechanics of membrane fusion** *Nat. Struct. Mol. Biol* **15**:675–683 <https://doi.org/10.1038/nsmb.1455>
- Conti M.A., Even-Ram S., Liu C., Yamada K.M., Adelstein R.S. (2004) **Defects in cell adhesion and the visceral endoderm following ablation of nonmuscle myosin heavy chain II-A in mice** *J. Biol. Chem* **279**:41263–41266 <https://doi.org/10.1074/jbc.C400352200>
- Ebrahim S. *et al.* (2019) **Dynamic polyhedral actomyosin lattices remodel micron-scale curved membranes during exocytosis in live mice** *Nat. Cell Biol* **21**:933–939 <https://doi.org/10.1038/s41556-019-0365-7>
- Eitzen G. (2003) **Actin remodeling to facilitate membrane fusion** *Biochim. Biophys. Acta* **1641**:175–181 [https://doi.org/10.1016/s0167-4889\(03\)00087-9](https://doi.org/10.1016/s0167-4889(03)00087-9)
- Evans E., Rawicz W. (1990) **Entropy-driven tension and bending elasticity in condensed-fluid membranes** *Phys. Rev. Lett* **64**:2094–2097 <https://doi.org/10.1103/PhysRevLett.64.2094>
- Faizi H. A., Frey S.L., Steinkuhler J., Dimova R., Vlahovska P.M. (2019) **Bending rigidity of charged lipid bilayer membranes** *Soft Matter* **15**:6006–6013 <https://doi.org/10.1039/c9sm00772e>
- Farsad K., De Camilli P. (2003) **Mechanisms of membrane deformation** *Curr. Opin. Cell Biol* **15**:372–381 [https://doi.org/10.1016/s0955-0674\(03\)00073-5](https://doi.org/10.1016/s0955-0674(03)00073-5)
- Goley E.D., Welch M.D. (2006) **The ARP2/3 complex: an actin nucleator comes of age** *Nat. Rev. Mol. Cell Biol* **7**:713–726 <https://doi.org/10.1038/nrm2026>
- Gomes E.R., Jani S., Gundersen G.G. (2005) **Nuclear movement regulated by Cdc42, MRCK, myosin, and actin flow establishes MTOC polarization in migrating cells** *Cell* **121**:451–463 <https://doi.org/10.1016/j.cell.2005.02.022>

- Gruenberg J., Griffiths G., Howell K.E. (1989) **Characterization of the early endosome and putative endocytic carrier vesicles in vivo and with an assay of vesicle fusion in vitro** *J. Cell Biol* **108**:1301–1316 <https://doi.org/10.1083/jcb.108.4.1301>
- Hammer J.A., Sellers J.R. (2011) **Walking to work: roles for class V myosins as cargo transporters** *Nat. Rev. Mol. Cell Biol* **13**:13–26 <https://doi.org/10.1038/nrm3248>
- Harrington A.W., St Hillaire C., Zweifel L.S., Glebova N.O., Philippidou P., Halegoua S., Ginty D.D. (2011) **Recruitment of actin modifiers to TrkA endosomes governs retrograde NGF signaling and survival** *Cell* **146**:421–434 <https://doi.org/10.1016/j.cell.2011.07.008>
- Hartman M.A., Spudich J.A. (2012) **The myosin superfamily at a glance** *J. Cell Sci* **125**:1627–1632 <https://doi.org/10.1242/jcs.094300>
- Helfrich W. (1973) **Elastic properties of lipid bilayers: theory and possible experiments** *Z. Naturforsch. C* **28**:693–703 <https://doi.org/10.1515/znc-1973-11-1209>
- Jollie W.P. (1990) **Development, morphology, and function of the yolk-sac placenta of laboratory rodents** *Teratology* **41**:361–381 <https://doi.org/10.1002/tera.1420410403>
- Kawamura N., Sun-Wada G.H., Aoyama M., Harada A., Takasuga S., Sasaki T., Wada Y. (2012) **Delivery of endosomes to lysosomes via microautophagy in the visceral endoderm of mouse embryos** *Nat. Commun* **3** <https://doi.org/10.1038/ncomms2069>
- Koike S., Keino-Masu K., Ohto T., Sugiyama F., Takahashi S., Masu M. (2009) **Autotaxin/lysophospholipase D-mediated lysophosphatidic acid signaling is required to form distinctive large lysosomes in the visceral endoderm cells of the mouse yolk sac** *J. Biol. Chem* **284**:33561–33570 <https://doi.org/10.1074/jbc.M109.012716>
- Liebl D., Griffiths G. (2009) **Transient assembly of F-actin by phagosomes delays phagosome fusion with lysosomes in cargo-overloaded macrophages** *J. Cell Sci* **122**:2935–2945 <https://doi.org/10.1242/jcs.048355>
- Lipowsky R. (1991) **The conformation of membranes** *Nature* **349**:475–481 <https://doi.org/10.1038/349475a0>
- Liu D., Bei D., Parmar H., Matus A. (2000) **Activity-regulated, cytoskeleton-associated protein (Arc) is essential for visceral endoderm organization during early embryogenesis** *Mech. Dev* **92**:207–215 [https://doi.org/10.1016/S0925-4773\(99\)00340-8](https://doi.org/10.1016/S0925-4773(99)00340-8)
- Liu K. *et al.* (2016) **Negative regulation of phosphatidylinositol 3-phosphate levels in early-to-late endosome conversion** *J. Cell Biol* **212**:181–198 <https://doi.org/10.1083/jcb.201506081>
- Lu T., Guo H.X. (2019) **How the Membranes Fuse: From Spontaneous to Induced** *Adv. Theor. Simul* **2** <https://doi.org/10.1002/adts.201900032>
- Ma W., Chang J., Tong J., Ho U., Yau B., Kebede M.A., Thorn P. (2020) **Arp2/3 nucleates F-actin coating of fusing insulin granules in pancreatic beta cells to control insulin secretion** *J. Cell Sci* **133** <https://doi.org/10.1242/jcs.236794>
- Maekawa M., Ishizaki T., Boku S., Watanabe N., Fujita A., Iwamatsu A., Obinata T., Ohashi K., Mizuno K., Narumiya S. (1999) **Signaling from rho to the actin cytoskeleton through protein kinases ROCK and LIM-kinase** *Science* **285**:895–898 <https://doi.org/10.1126/science.285.5429.895>

- Masedunskas A., Sramkova M., Parente L., Sales K.U., Amornphimoltham P., Bugge T.H., Weigert R. (2011) **Role for the actomyosin complex in regulated exocytosis revealed by intravital microscopy** *Proc. Natl. Acad. Sci. USA* **108**:13552–13557 <https://doi.org/10.1073/pnas.1016778108>
- Miklavc P., Ehinger K., Sultan A., Felder T., Paul P., Gottschalk K.E., Frick M. (2015) **Actin depolymerisation and crosslinking join forces with myosin II to contract actin coats on fused secretory vesicles** *J. Cell Sci* **128**:1193–203 <https://doi.org/10.1242/jcs.165571>
- Morel E., Parton R.G., Gruenberg J. (2009) **Annexin A2-Dependent Polymerization of Actin Mediates Endosome Biogenesis** *Dev. Cell* **16**:445–457 <https://doi.org/10.1016/j.devcel.2009.01.007>
- Morris K., Ihnatovych I., Ionetz E., Reed J., Braundmeier A., Strakova Z. (2011) **Cofilin and Slingshot Localization in the Epithelium of Uterine Endometrium Changes During the Menstrual Cycle and in Endometriosis** *Reprod. Sci* **18**:1014–1024 <https://doi.org/10.1177/1933719111401663>
- Nicot A.S., Fares H., Payraastre B., Chisholm A.D., Labouesse M., Laporte J. (2006) **The phosphoinositide kinase PIKfyve/Fab1p regulates terminal lysosome maturation in *Caenorhabditis elegans*** *Mol. Biol. Cell* **17**:3062–3074 <https://doi.org/10.1091/mbc.e05-12-1120>
- Richardson S.J., Southwell B.R., Jaworowski A. (2000) **The rat visceral yolk sac internalizes maternal transferrin and secretes hydrolyzed products towards the fetus** *Comp. Biochem. Phys. B* **125**:29–36 [https://doi.org/10.1016/S0305-0491\(99\)00150-9](https://doi.org/10.1016/S0305-0491(99)00150-9)
- Ridolfi A., Caselli L., Baldoni M., Montis C., Mercuri F., Berti D., Valle F., Brucale M. (2021) **Stiffness of Fluid and Gel Phase Lipid Nanovesicles: Weighting the Contributions of Membrane Bending Modulus and Luminal Pressurization** *Langmuir* **37**:12027–12037 <https://doi.org/10.1021/acs.langmuir.1c01660>
- Roberts R.L., Barbieri M.A., Pryse K.M., Chua M., Stahl P.D. (1999) **Endosome fusion in living cells overexpressing GFP-rab5** *J. Cell Sci* **112**:3667–3675 <https://doi.org/10.1242/jcs.112.21.3667>
- Salbreux G., Charras G., Paluch E. (2012) **Actin cortex mechanics and cellular morphogenesis** *Trends Cell Biol* **22**:536–545 <https://doi.org/10.1016/j.tcb.2012.07.001>
- Scott C.C., Gruenberg J. (2011) **Ion flux and the function of endosomes and lysosomes: pH is just the start** *Bioessays* **33**:103–110 <https://doi.org/10.1002/bies.201000108>
- Simson R., Wallraff E., Faix J., Niewohner J., Gerisch G., Sackmann E. (1998) **Membrane bending modulus and adhesion energy of wild-type and mutant cells of *Dictyostelium* lacking talin or cortexillins** *Biophys. J* **74**:514–22 [https://doi.org/10.1016/S0006-3495\(98\)77808-7](https://doi.org/10.1016/S0006-3495(98)77808-7)
- Svitkina T.M. (2018) **Ultrastructure of the actin cytoskeleton** *Curr. Opin. Cell Biol* **54**:1–8 <https://doi.org/10.1016/j.ceb.2018.02.007>
- Taunton J., Rowning B.A., Coughlin M.L., Wu M., Moon R.T., Mitchison T.J., Larabell C.A. (2000) **Actin-dependent propulsion of endosomes and lysosomes by recruitment of N-WASP** *J. Cell Biol* **148**:519–530 <https://doi.org/10.1083/jcb.148.3.519>
- Titus M.A. (2018) **Myosin-driven intracellular transport** *Cold Spring Harb. Perspect. Biol* **10** <https://doi.org/10.1101/cshperspect.a021972>

- Tran D.T., Masedunskas A., Weigert R., Ten Hagen K.G. (2015) **Arp2/3-mediated F-actin formation controls regulated exocytosis in vivo** *Nat. Commun* **6** <https://doi.org/10.1038/ncomms10098>
- Uenishi E., Shibasaki T., Takahashi H., Seki C., Hamaguchi H., Yasuda T., Tatebe M., Oiso Y., Takenawa T., Seino S. (2013) **Actin Dynamics Regulated by the Balance of Neuronal Wiskott-Aldrich Syndrome Protein (N-WASP) and Cofilin Activities Determines the Biphasic Response of Glucose-induced Insulin Secretion** *J. Biol. Chem* **288**:25851–25864 <https://doi.org/10.1074/jbc.M113.464420>
- Vitriol E.A., Wise A.L., Berginski M.E., Bamburg J.R., Zheng J.Q. (2013) **Instantaneous inactivation of cofilin reveals its function of F-actin disassembly in lamellipodia** *Mol. Biol. Cell* **24**:2238–2247 <https://doi.org/10.1091/mbc.E13-03-0156>
- Wang L., Seeley E.S., Wickner W., Merz A.J. (2002) **Vacuole fusion at a ring of vertex docking sites leaves membrane fragments within the organelle** *Cell* **108**:357–369 [https://doi.org/10.1016/S0092-8674\(02\)00632-3](https://doi.org/10.1016/S0092-8674(02)00632-3)
- Welch M.D., Way M. (2013) **Arp2/3-Mediated Actin-Based Motility: A Tail of Pathogen Abuse** *Cell Host Microbe* **14**:242–255 <https://doi.org/10.1016/j.chom.2013.08.011>
- Zhao W.D., Hamid E., Shin W., Wen P.J., Krystofiak E.S., Villarreal S.A., Chiang H.C., Kachar B., Wu L.G. (2016) **Hemi-fused structure mediates and controls fusion and fission in live cells** *Nature* **534**:548–552 <https://doi.org/10.1038/nature18598>

Article and author information

Seiichi Koike

Graduate School of Comprehensive Human Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan, Department of Molecular Neurobiology, Institute of Medicine, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan, Laboratory of Molecular and Cellular Biology, Graduate School of Science and Engineering for Research, University of Toyama, 3190 Gofuku, Toyama 930-8555, Toyama, Japan
ORCID ID: [0000-0001-8203-4498](https://orcid.org/0000-0001-8203-4498)

Masashi Tachikawa

Graduate School of Nanobioscience, Yokohama City University, 22-2 Seto, Kanazawa-ku, Yokohama, Kanagawa 236-0027, Japan

Motosuke Tsutsumi

Exploratory Research Center on Life and Living Systems (ExCELLS), National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji, Okazaki, Aichi 444-8787, Japan, National Institute for Physiological Sciences, National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji, Okazaki, Aichi 444-8787, Japan

Takuya Okada

Graduate School of Comprehensive Human Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan, Department of Molecular Neurobiology, Institute of Medicine, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan

Tomomi Nemoto

Exploratory Research Center on Life and Living Systems (ExCELLS), National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji, Okazaki, Aichi 444-8787, Japan, National Institute for Physiological Sciences, National Institutes of Natural Sciences, 5-1 Higashiyama, Myodaiji, Okazaki, Aichi 444-8787, Japan

Kazuko Keino-Masu

Graduate School of Comprehensive Human Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan, Department of Molecular Neurobiology, Institute of Medicine, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan

Masayuki Masu

Graduate School of Comprehensive Human Sciences, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan, Department of Molecular Neurobiology, Institute of Medicine, University of Tsukuba, 1-1-1 Tennodai, Tsukuba, Ibaraki 305-8575, Japan

For correspondence: mmasu@md.tsukuba.ac.jp

ORCID iD: [0000-0002-4726-7429](https://orcid.org/0000-0002-4726-7429)

Copyright

© 2024, Koike et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Li Yu

Tsinghua University, Beijing, China

Senior Editor

Felix Campelo

Institute of Photonic Sciences, Barcelona, Spain

Reviewer #1 (Public Review):

Summary:

This manuscript employs yolk sac visceral endoderm cells as a novel model for studying endosomal fusion, observing two distinct fusion behaviors: quick homotypic fusion between late endosomes, and slower heterotypic fusion between late endosomes and lysosomes. The mathematical modeling suggests that vesicle size critically influences the mode of fusion. Further investigations reveal that actin filaments are dynamically associated with late endosomal membranes, and are oriented in the x-y plane and along the apical-basal axis. Actin and Arf2/3 were shown to appear at the rear end of the endosomes along the moving direction suggesting polymerization of actin may provide force for the movement of endosomes. Additionally, the authors found that actin dynamics regulate homotypic and heterotypic fusion events in a different manner. The authors also provide evidence to suggest that Cofilin-dependent actin dynamics are involved in late endosome fusion.

Strengths:

The unique feature of this study is that the authors use yolk sac visceral endoderm cells to study endosomal fusion. Yolk sac visceral endoderm cells have huge endocytic vesicles, endosomes, and lysosomes, offering an excellent system to explore endosomal fusion dynamics and the assembly of cellular factors on membranes. The manuscript provides a valuable and convincing observation of the modes of endosomal fusion and the roles of actin dynamics in this process, and the conclusions of the study are justified by the data.

Weaknesses:

While the study offers compelling observations, it falls short of delivering clear mechanistic insights. Key questions remain unaddressed, such as the functional significance of actin filaments that extend apically in positioning late endosomes, the ways in which actin dynamics influence fusion events, and the functional implications of the slower bridge fusion process.

<https://doi.org/10.7554/eLife.95999.1.sa2>

Reviewer #2 (Public Review):

Summary:

Seiichi Koike et al. studied two fusion models, explosive fusion, and bridge fusion, utilizing yolk sac visceral endoderm cells. They elucidated these two fusion models in vivo by employing mathematical modeling and incorporating fluctuations derived from actin dynamics as a key regulator for rapid homotypic fusion between late endosomes.

Strengths:

This study uncovered the role of actin dynamics in regulating the transition of fusion models in homotypic fusion between late endosomes and introduced a method for observing the fusion of single vesicles with two different targets. The role of actin dynamics in vesicle fusion in other systems has been extensively studied. This study could offer useful insights for research on vesicle fusion.

Weaknesses:

The physiological significance of different fusion models is lacking.

<https://doi.org/10.7554/eLife.95999.1.sa1>

Reviewer #3 (Public Review):

Summary:

The authors found two endosomal fusion modes by live cell imaging of endosomes in yolk sac lateral endoderm cells of 8.5-day-old embryonic mice and described the fusion modes by mathematical models and simulations. They also showed that actin polymerization is involved in the regulation of one of the fusion modes.

Strengths:

The strength of this study is that the authors' claims are well supported by beautiful live cell images and theoretical models. By using specialized cells, yolk sac visceral endoderm cells, the live images of endosomal fusion, localization of actin-related molecules, and validation data from multiple inhibitor experiments are clear.

Weaknesses:

This study does not include any assessment of whether the two types of endosome fusions claimed by the authors occur in general cells, so the article is limited to showing a phenomenon specific to yolk sac lateral endoderm cells. Also, the study does not show the physiological importance of the two types of fusion. There are some unclear points in the method of image analysis and some of the descriptions in the text are not logical.

<https://doi.org/10.7554/eLife.95999.1.sa0>