

De novo-designed minibinders expand the synthetic biology sensing repertoire

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

[About eLife's process](#)

Reviewed preprint version 1

April 19, 2024 (this version)

Sent for peer review

February 2, 2024

Posted to preprint server

January 15, 2024

Zara Y. Weinberg , Sarah S. Soliman, Matthew S. Kim, Irene P. Chen, Melanie Ott, Hana El-Samad 

Department of Biochemistry & Biophysics, UCSF, San Francisco CA • Tetrad Graduate Program, UCSF, San Francisco CA • Cell Design Institute, San Francisco CA • Gladstone Institutes, San Francisco CA • Department of Medicine, UCSF, San Francisco CA • Chan Zuckerberg Biohub–San Francisco, San Francisco CA • Altos Labs, San Francisco CA

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Synthetic and chimeric receptors capable of recognizing and responding to user-defined antigens have enabled “smart” therapeutics based on engineered cells. These cell engineering tools depend on antigen sensors which are most often derived from antibodies. Advances in the *de novo* design of proteins have enabled the design of protein binders with the potential to target epitopes with unique properties and faster production timelines compared to antibodies. Building upon our previous work combining a *de novo*-designed minibinder of the Spike protein of SARS-CoV-2 with the synthetic receptor synNotch (SARSNotch), we investigated whether minibinders can be readily adapted to a diversity of cell engineering tools. We show that the Spike minibinder LCB1 easily generalizes to a next-generation proteolytic receptor SNIPR that performs similarly to our previously reported SARSNotch. LCB1-SNIPR successfully enables the detection of live SARS-CoV-2, an improvement over SARSNotch which can only detect cell-expressed Spike. To test the generalizability of minibinders to diverse applications, we tested LCB1 as an antigen sensor for a chimeric antigen receptor (CAR). LCB1-CAR enabled CD8⁺ T cells to cytotoxically target Spike-expressing cells. Our findings suggest that minibinders represent a novel class of antigen sensors that have the potential to dramatically expand the sensing repertoire of cell engineering tools.

eLife assessment

This study presents a **useful** investigation to test *de novo*-designed mini binders against the Spike protein of SARS-CoV-2 within two classes of synthetic receptors (SNIPRs and CARs). The methods and evidence supporting the focused claims are very **solid**, although the small-scale nature of the investigation (number of modifications, number of minibinders, etc.) makes it difficult to determine how generalizable these results and potential design principles are. This work will be of interest to synthetic biologists and cell engineers as a starting point for systematic, larger-scale analysis and optimization of synthetic receptor designs for cellular therapy and other applications.

Introduction

Synthetic receptors enable engineered cells to detect specific signals in their environment and respond with therapeutic payloads to treat disease^{1–3}. Due to their therapeutic potential, synthetic biology groups have produced many different synthetic receptor systems. Perhaps the most widely exploited of these systems is the chimeric antigen receptor (CAR)⁴, which couples an antigen-sensing binder with signaling domains from the T cell receptor complex and associated co-stimulatory molecules. CAR T cells have proven tremendously valuable as anti-cancer therapeutics^{5,6}. More recently, synthetic receptors such as synNotch⁷, MESA⁸, GEMS⁹, SNIPRs¹⁰, and many others¹ were developed to use similar antigen-sensing domains coupled to customizable transcriptional programs. These synthetic receptor systems have shown great promise in enhancing existing cell-based therapies^{11–16} and bioengineering² applications ranging from building cell-based biosensors^{17–19} to understanding and engineering development^{20–26}.

Current receptors primarily rely on single chain variable fragment antibodies (scFvs) or less frequently variable heavy domain monomeric antibodies (nanobodies) to sense their targets. Monomeric antibodies that are generated for both research and therapeutic purposes are now mostly discovered through high-throughput screening using phage^{27–29} or yeast display³⁰ followed by sub-screening in an orthogonal assay and then affinity maturation. Although these strategies have proven useful for generating antibody therapeutics and research tools, scFvs and nanobodies do not always translate easily to antigen sensors for synthetic receptors. While it is unclear why not all scFvs and nanobodies function in cell engineering contexts, it seems likely that the physical properties of binding, including affinity, on- and off-rates, and the steric and conformational requirements for each antibody, all have a role to play^{31,32}. Further, although nanobodies and scFvs are relatively compact, even their modest size of >100 amino acids can be a limitation in the engineering of primary cells where genetic payload size is heavily constrained. For these reasons, small protein binders that can be engineered to bind specific targets hold promise as the next generation of antigen sensors.

De novo-designed binders that can be created using computational methods against almost any protein with known structure are a tantalizing new class of proteins that could serve as antigen sensors for synthetic receptors. Computational design of protein-protein interactions has a long history^{33,34} but recent advances have made this technology readily accessible for synthetic biology applications. Protein docking-based methods have enabled the design of binders against any protein structure³⁵. Deep learning-assisted methods improved the speed of binder design and have yielded nanomolar affinity binders or better directly from computational pipelines without the need for affinity maturation^{36–38}. Deep learning methods have also extended the design of binders to small helical peptides³⁹, dramatically expanding the scope of potential binder targets. Previously, we demonstrated that a *de novo*-designed binder created using a docking-based approach that targeted the SARS-CoV-2 Spike protein⁴⁰ could be combined with synNotch to drive customizable responses in a model of SARS-CoV-2 infection¹⁸. The SARS-CoV-2 minibinder used in those studies, LCB1, is a small three-helix bundle of 56 amino acids and binds directly to the interface it was designed for with picomolar affinity, overcoming many of the potential challenges for antigen sensor design. If minibinders like LCB1 generalize as antigen sensors across other synthetic receptors, these *de novo*-designed proteins could serve as powerful additions to the cell engineering toolbox.

In this study, we sought to test the generalizability of the LCB1 minibinder across different synthetic receptors. First, we demonstrate that LCB1 functions as an antigen sensor for a synthetic intermembrane proteolysis receptor (SNIPR)¹⁰ with distinct activation mechanics compared to synNotch. Unlike our previously described SARSNotch, the LCB1-SNIPR is capable of detecting both

cells expressing the SARS-CoV-2 Spike and live SARS-CoV-2 virus. We also show that LCB1 functions as an antigen sensor for a CAR, demonstrating that this minibinder's capabilities are not limited to proteolytic synthetic receptors but are also effective at modulating endogenous cellular pathways. Minibinder-coupled CARs were able to specifically activate T cell programs in primary human CD8⁺ T cells against SARS-CoV-2 Spike-expressing cells. Together, these findings suggest that minibinders can be easily deployed across synthetic receptors as antigen sensors.

Results

Minibinders adapt readily to next-generation synthetic proteolytic receptors

We previously demonstrated that the *de-novo* designed LCB1 and LCB3 minibinders for SARS-CoV-2 Spike protein could be used as functional antigen sensors when integrated with synNotch¹⁸. These anti-SARS Notch constructs successfully detected both purified Spike protein and Spike expressed on the surface of mammalian cells in a model of virally-infected cells to drive a synthetic transcriptional output.

We sought to extend these findings by testing the ability of the LCB1 and LCB3 minibinders to serve as antigen sensors for other synthetic receptors. Since the development of the original synNotch, a modular toolkit of parts for designing proteolysis-based receptors has been developed, allowing for tuning of receptor activation in various contexts¹⁰. These synthetic intramembrane proteolysis receptors (SNIPRs) use a collection of extracellular, transmembrane, juxtamembrane, and intracellular domains to tune baseline activity and receptor dynamic range for a variety of cellular and antigen contexts. Although these receptors function similarly to synNotch, the absence of the Notch regulatory repeats in the extracellular domain of SNIPRs suggests that their activation is mechanistically unique. Given the similar function but mechanistic differences in activation of SNIPRs, we hypothesized that these receptors would be a good test case for evaluating minibinder function across synthetic receptors.

We selected a SNIPR chassis using the CD8a hinge domain, the human Notch1 transmembrane domain, and the human Notch2 juxtamembrane domain that had displayed low baseline activation and high payload expression upon specific activation in Jurkat and primary human CD4⁺ and CD8⁺ T cells. We generated LCB1- and LCB3-equipped SNIPRs coupled to Gal4-VP64 transcription factor (**Figure 1A**). We transduced these constructs into a Jurkat line expressing an output circuit as described previously¹⁸ that expresses BFP downstream of Gal4-VP64 activation. SNIPR-expressing Jurkat T cells were sorted for surface staining of the expressed receptor, positive expression of a co-transduced mCherry marker, positive expression of mCitrine marking the integration of the output circuit, and the absence of BFP expression at baseline. See Supplemental Figure 1A for schema of output circuit and SNIPR expression constructs.

We incubated a polyclonal population of SNIPR-expressing cells with either untransduced K562 cells or K562s stably expressing a prefusion stabilized version of the SARS-CoV-2 Spike ectodomain⁴¹ displayed on a PDGFR transmembrane domain as previously described¹⁸ (**Figure 1B**). After 72 hours, we assessed our output circuit-only expressing Jurkat cells (No Notch) and LCB1- and LCB3-SNIPR expressing cells for BFP expression (**Figure 1C**). BFP expression was specifically increased for both SNIPR-expressing cell lines only in the presence of Spike-expressing K562s, demonstrating that minibinder-coupled SNIPRs specifically detected Spike protein on other cells.




Figure 1

Minibinders adapt readily to next-generation synthetic proteolytic receptors.

A) Schematic representation of a synthetic intramembrane proteolysis receptor (SNIPR) that enables customized genetic responses upon binding a target. LCB1 binder structure cartoon loosely based on PDB 7JZU. B) Assay for testing SNIPR function, whereby SNIPRs with various minibinder antigen sensors are expressed in Jurkat cells and cells are mixed with SARS- CoV-2 Spike-expressing K562 cells. SNIPR activation is readout as BFP expression. C) Histograms of BFP expression in SNIPR-expressing Jurkats after 72-hour incubation with K562s or K562s expressing Spike.

Minibinders perform similarly across different synthetic proteolytic receptors

Given the putatively different modes of activation between synNotch and SNIPRs, we wanted to compare the performance of minibinders between these different synthetic proteolytic receptors.

We sought to test performance in different antigen density regimes to better understand the dynamic range of the sensing capabilities of these minibinder-coupled receptors. We generated two Spike-expressing K562 populations expressing a moderate (Spike-K562 (M)) and high (Spike-K562 (H)) amount of Spike (**Figure 2A** [↗](#)). We also regenerated the LCB1-Notch and LCB3-Notch expressing Jurkat lines described in our previous work to match the expression level of LCB1- and LCB3-SNIPR in the lines generated above (**Figure 2B** [↗](#), Supplemental Figure 1A). We then compared polyclonal populations of these lines in the same 72 hour incubation assay described above.

All minibinder-coupled receptor lines expressed BFP specifically in response to both M and H Spike-expressing K562s (**Figure 2C** [↗](#)). To differentiate activated (BFP-expressing cells) from inactive cells, we fit a two component Gaussian mixture model to the BFP expression data across experimental populations. We compared the predicted mean of the activated component identified by the mixture model across each receptor to understand the amount of payload that each receptor could produce in response to Spike-K562s (**Figure 2D** [↗](#)). For both synNotch and SNIPR, LCB1-coupled receptors produced greater payload expression in response to Spike-expressing cells compared to LCB3-coupled receptors. SynNotch and SNIPR receptors coupled to the same minibinder produced similar levels of BFP expression. To determine the fraction of each population similarly, followed by LCB3-Notch ($84\% \pm 0.5\%$) and then LCB3-SNIPR ($71\% \pm 1.9\%$) with the lowest activation.

These findings suggest that despite distinct mechanisms of activation of synNotch and SNIPRs, minibinders perform similarly across these receptors. This is true both for the maximum efficacy of a specific minibinder, and also the relative performance of two different minibinders directed at the same target.

Novel synthetic proteolytic receptors enable minibinder-dependent detection of live SARS-CoV-2 virus

Although minibinders performed similarly between synNotch and SNIPRs in a cell-expressed antigen context, we hypothesized that the distinct activation mechanism of SNIPRs might provide advantages for detection in other contexts. Previously, we showed that LCB1-Notch was not capable of activating in response to successfully activated (expressing BFP), we fit our mixture model to each experiment and assessed the fraction of cells in the population with the higher mean expression (**Figure 2E** [↗](#)). All receptors showed an antigen dose-dependent increase in activation. Again, LCB1-coupled receptors showed the highest % activation, with LCB1-SNIPR ($90\% \pm 0.5\%$ activation against Spike-K562 (H) cells) and LCB1-Notch ($96\% \pm 0.3\%$) performing pseudotyped lentivirus decorated with the SARS-CoV-2 Spike. We sought to determine whether SNIPR's unique activation mechanism would enable it to better detect SARS-CoV-2 virus itself.

Using the Jurkat lines generated above, we took our best-performing receptors LCB1-Notch and LCB1-SNIPR and incubated them with live SARS-CoV-2 virus for 72 hours before fixing and assessing BFP expression (**Figure 3A** [↗](#)). We initially incubated receptor-expressing Jurkats with no virus, the WA1 viral isolate, or the B.1.617.2 (Delta) virus isolate, all at a multiplicity of infection of 1. When assessing median BFP expression (**Figure 3B** [↗](#)) we saw that LCB1-Notch produced no BFP in response to virus, LCB1-SNIPR had higher baseline activity, but importantly it also showed increased BFP expression in response to both viral isolates. We then conducted a dose-response




Figure 2

Minibinders perform similarly across different synthetic proteolytic receptors.

A) Expression of surface-displayed Spike protein (as assessed by anti-Myc labeling of an N-terminal extracellular Myc tag) for untransduced K562s and K562 lines sorted for moderate (M) and high (H) expression of Spike (median fluorescent counts for populations across 3 replicates). B) Surface expression levels of multiple minibinder-coupled synthetic receptors on Jurkat cells as assessed by anti-Myc labeling of an N-terminal extracellular Myc tag on each receptor (median fluorescent counts for populations across 3 replicates). C) Histograms describing BFP expression in minibinder receptor-expressing Jurkats after 72 hours of incubation with K562s with 3 different levels of Spike expression (pooled populations from 3 replicates). D) Mean log BFP expression in the activated populations of Jurkats after 72 hour incubation with Spike-K562 (H) cells as calculated by the estimated mean of the high BFP expression population after fitting a two component Gaussian mixture model on each of 3 replicates per condition. Data points are estimated mean of 'on' population in each replicate. E) The fraction of BFP-expressing Jurkats in the mixture model-defined activated population for each receptor as a function of Spike expression. All error bars (including E) are standard deviation.

study, incubating LCB1-SNIPR cells with a 2 log range of viral MOIs and assessing the fraction of cells that activated (**Figure 3C**). LCB1-SNIPR cells showed a modest dose-dependent increase in % cells activated for both viral isolates, with stronger activation for WA1.

These data suggest that although performance of these minibinders generalizes well across synthetic receptors, the minibinders can successfully enable unique capabilities for the different receptors they are coupled to.

Minibinders enable chimeric antigen receptor-mediated killing of SARS-CoV-2 Spike-expressing cells

After demonstrating minibinder generalization to multiple proteolytic receptors, we sought to extend these findings to a distinct receptor class to test minibinder viability as antigen sensors across a wider range of synthetic receptors. For this study, we selected the chimeric antigen receptor (CAR) because of its important therapeutic relevance.

We generated a minibinder-coupled second-generation CAR by combining LCB1 with the CD8⁺ hinge and transmembrane domains and the intracellular domains of 4-1BB and CD3 ζ (**Figure 4A**). This construct, dubbed LCB1-CAR, expressed well when lentivirally transduced into CD8⁺ primary human T cells (Supplemental Figure 2B).

We tested the ability of LCB1-CAR to induce an antigen-specific cytotoxic phenotype (**Figure 4B**). After transducing CD8⁺ T cells with LCB1-CAR, we co-cultured them with either parental K562 cells or Spike-K562 (H) cells. We co-cultured our effector and target cells at ratios ranging from 2 effectors for every target to 4 targets for every effector. We then evaluated the ability of LCB1-CAR to elicit cell proliferation, expression of the T cell activation marker CD25, and lysis of target cells.

LCB1-CAR promoted a specific cytotoxic reaction to Spike-expressing K562s (**Figure 4C-E**). LCB1-CAR-expressing T cells proliferated in the presence of Spike-K562s but not when cultured on their own or with parental K562s, as indicated by a dilution of the CellTrace labeling dye in cell populations only in the LCB1-CAR/ Spike-K562 condition (**Figure 4C**). The expression of the activation marker CD25 was upregulated for LCB1-CAR cells in the presence of Spike-K562s but not in other conditions (**Figure 4D**). Importantly, LCB1-CAR T cells were able to induce specific lysis of Spike-K562s at effector:target ratios of 2:1 (~40% lysis) and 1:1 (~20% lysis), with little off-target lysis of other cell types (**Figure 4E**). These data together demonstrate that the minibinder LCB1 is a functional antigen sensor for a CAR, albeit with moderate efficacy.

Short linker sequences optimize minibinder-coupled receptor function

We next sought to improve the efficacy of minibinder-coupled CARs. Although LCB1 proved the better of the two binders in proteolytic receptor function, we considered that LCB3 might have different efficacy when coupled to a CAR. LCB1 and LCB3 bind the same interface on the Spike protein, but in opposite orientations⁴⁰. We hypothesized that this binding difference, which was detrimental to LCB3 function on synNotch and SNIPR, might provide different activity for a CAR.

In comparing LCB1- and LCB3-CARs, we also added in a CD19-CAR as a benchmark, as this CAR is the gold standard in the field. Our CD19-CAR used the same 4-1BB and CD3 ζ intracellular domains as the LCB-CARs, swapping in the CD19 scFv FMC63⁴³ for the antigen sensor domain. All of the LCB1-, LCB3-, and CD19-CARs expressed at similar levels after lentiviral transduction into primary human CD8⁺ T cells (Supplemental Figure 2B).




Figure 3

Novel synthetic proteolytic receptors enable minibinder-dependent detection of live SARS-CoV-2 virus.

A) Schematic of live virus sensing experiment. Minibinder receptor-expressing Jurkats were incubated with live SARS-CoV-2 virus for 72 hours, and then fixed with paraformaldehyde and assessed for BFP expression as a sign of receptor activation. B) Median BFP expression in Jurkat cells exposed to no virus or WA1 or Delta isolates of SARS-CoV-2 at a multiplicity of infection of 1. C) Fraction of Jurkat population expressing BFP as a function of MOI of SARS-CoV-2 that cells were exposed to. All error bars are standard deviation.




Figure 4

Minibinders enable chimeric antigen receptor-mediated killing of SARS-CoV-2 Spike-expressing cells.

A) Schematic of a chimeric antigen receptor with 4-1BB costimulatory domain using the minibinder LCB1 as its antigen sensor. B) Assay schematic indicating that primary CD8⁺ human T cells expressing the LCB1-CAR were mixed with Spike-expressing K562s for 72 hours and then were assessed for proliferation, T cell activation, and killing of the target cells. C) Proliferation of T cells as a function of which target cells they were incubated with. Histogram showing Cell Trace dye fluorescence across a population of T cells, with homogenous high staining indicating no proliferation and multiple populations indicating proliferation. D) T cell activation as assessed by CD25 antibody-labeling as a function of which targets T cells were exposed to (median fluorescent for population in each of 3 replicates). E) The fraction of target cell populations that were lysed after T cell incubation as a function of the ratio of effector T cells to K562 targets that were incubated together. All error bars are standard deviation.

We evaluated cytotoxicity of each of these CARs against K562s, CD19-K562s (Supplemental Figure 2A), and Spike-K562 (H) cells (**Figure 5A** [↗](#)). As expected, the CD19-CAR performed well and lysed $96\% \pm 0.5\%$ of target cells. Both LCB-CARs performed similarly to each other and at about half the efficacy of the CD19-CAR, with LCB1-CAR lysing $43\% \pm 6.4\%$ of targets and LCB3-CAR lysing $38\% \pm 6.9\%$ of targets. Efficacy varied between T cells from different donors, with all 3 CARs showing similar specific cytotoxicity when expressed in donor cells that had higher levels of off-target cytotoxicity (Supplemental Figure 2C). Together, these results suggest that unlike with synNotch and SNIPR, the binding orientation of LCB1 and LCB3 do not affect CAR function, and that these minibinders are less efficacious antigen sensors compared to the gold-standard CD19 scFv.

We next sought to test optimizations in other aspects of the binders. Antigen sensor affinity^{44,45} [↗](#) and the composition of the linker region between the antigen sensor and the CD8a hinge^{44,46} [↗](#) have both been demonstrated to increase CAR-dependent lysis of target cells. Both LCB minibinders have picomolar affinity for their targets, similar to or better than the $\sim 300\text{pM}$ affinity described for the CD19 scFv we used⁴⁷ [↗](#), which suggests increasing affinity is unlikely to improve performance.

Instead, we targeted the linker region (**Figure 5B** [↗](#)). Our initial construct featured a minimal linker (GS, “simple”) between the minibinder and the CD8a hinge domain used in the CAR. We generated constructs including a short flexible linker (2x repeats of GGGGS, “short”), a longer flexible linker (4x repeats of GGGGS, “long”), and a synthetic stabilized immunoglobulin fold-like domain⁴⁸ [↗](#) predicted to be highly stable as a rigid alternative to the flexible linkers above (Fibcon, “rigid”). We constructed LCB1-based CARs with each of these linkers, and they all expressed at similar levels (Supplemental Figure 2B). We then evaluated these minibinder and linker variants in the same killing assay described above (**Figure 5C** [↗](#)). At 24 hours after co-culture with target cells, mild differences could be seen between linker variants, with modestly lower cytotoxicity from the long and rigid variants compared to the simple and short variants. By 72 hours post co-culture, these differences disappeared and all variants behaved similarly (Supplemental Figure 2D). Altogether, the simple GS linker appears to be optimal for CAR-dependent cytotoxicity.

With these linker variants in hand, we also explored whether these linkers would affect minibinder-coupled synNotch or SNIPR function. All LCB1 linker variants successfully expressed when coupled to synNotch or SNIPR, but lower expression was seen for the short, long, and rigid variants when coupled to SNIPR (Supplemental Figure 3A). We then tested Jurkats expressing these variants in a co-culture experiment with Spike-K562 (M) cells (**Figure 5D** [↗](#)). Expression differences did not correlate with differences in functionality. LCB1-Simple and -Short versions of synNotch functioned best, while the LCB1-Long and -Rigid activated a much smaller percentage of cells in response to Spike-K562s ($\sim 75\%$ of cells activated for LCB1-Simple and -Short Notch vs $\sim 45\%$ activated for LCB1-Long and -Rigid Notch). SNIPR was similarly unable to tolerate the long linker, but LCB1-Rigid SNIPR performed similarly to the variants with the simple and short linkers.

These data suggest that while minibinders function well across a variety of synthetic receptors, there is still more exploration needed to determine how to create minibinders with the best efficacy at chimeric antigen receptors. Overall, our data suggest that minibinders work best with a short flexible linker coupling them to the receptor they are serving as antigen sensors for.

Discussion

In this study, we demonstrated that *de novo*-designed protein minibinders generalize as antigen sensors across multiple synthetic receptors. The LCB1 and LCB3 binders targeting SARS-CoV-2 Spike successfully induced activation of two proteolytic receptors in the presence of cells expressing Spike, and LCB1 enabled SNIPR-based detection of live SARS-CoV-2 virus. LCB1 and




Figure 5

Short linker sequences optimize minibinder-coupled receptor function.

A) The fraction of K562s expressing no ligand, CD19, or Spike that were lysed after 72 hour incubation with LCB1-CAR, LCB3-CAR, or the benchmark CD19-CAR at a 1:1 effector:target ratio. B) Schematic describing 4 different linkers that were evaluated to explore the effect of linker length and rigidity on minibinder receptor function. Fibron linker structure cartoon loosely based on PDB 3TEU. C) The fraction of K562 of Spike-K562 targets lysed after 24 hour incubation with CD8+ T cells expressing 4 variants of the LCB1-CAR at a 1:1 effector:target ratio. D) The percent of proteolytic receptor-expressing cells that were expressing BFP after 72 hour incubation with K562 or Spike-K562 (M) target cells for 4 different linker variations. All error bars are standard deviation.

LCB3 similarly enabled chimeric antigen receptors expressed in human CD8⁺ T cells to recognize and drive killing of Spike-expressing cells in culture. Finally, we evaluated several potential optimizations to improve the performance of minibinder-coupled receptors and found that for the receptors tested short flexible linkers produced the optimal functionality. Together, these results identify *de novo*-designed minibinders as an important part of the burgeoning cell engineering toolbox.

The two minibinders LCB1 and LCB3 showed differential efficacy across receptors. For both of the proteolytic receptors, LCB1 was a dramatically better antigen sensor and resulted in as much as 10-fold higher payload expression and twice as much population activity compared to the LCB3-coupled receptors. We previously hypothesized that this difference was the result of the binding orientation of these two proteins, which bind to the same surface on Spike but in opposite orientations⁴⁰. Curiously, this hierarchy in function is not dependent on the activation mode of the proteolytic receptor – synNotch activation is thought to depend on its Notch-derive extracellular domain⁷ while the SNIPRs are hypothesized to function through dimerization and an as-yet unelucidated mechanism that is dependent on the CD8a extracellular hinge domain¹⁰. The shared functional hierarchy of LCB1 and LCB3 between these receptors suggests that despite their distinct mechanisms of activation, they share some common requirements of their antigen sensors. This hierarchy was not seen for the same binders on a CAR, where both LCB1 and LCB3 produced similar T cell activation profiles. Our results suggest that a systematic exploration into the role of binding orientation might aid in creating optimal *de novo*-designed antigen sensors in the future.

Our data suggest that the optimal linker sequence for connecting minibinders to a target receptor is a simple glycine-serine (GS) linker. All receptors tested tolerated both a GS linker and 2xGGGGS linker well, but the performance of a longer flexible linker was poor across all receptors, and only SNIPR tolerated a more rigid linker. The linkers tested in this study are not exhaustive – future studies incorporating minibinders into synthetic receptors could also explore distance from the plasma membrane as a key variable of linker length.

Computational approaches for creating *de novo*-designed binders against practically any protein^{34–38} or peptide³⁹ with known structure position minibinders as attractive new antigen sensors. The ability of these methods to target specific interfaces of proteins of interest will further enable novel synthetic receptors that can distinguish between different faces and perhaps even conformations of membrane-displayed proteins. Further, novel methods for designing high-affinity binders for small proteins like neuropeptides and peptide hormones³⁹ have the potential to build synthetic receptors capable of robust activation in the presence of secreted factors, a sorely lacking piece of the current cell engineering toolkit. Further, although binders have only been designed against proteins with known structure, structure prediction methods^{49,50} might enable the design of binders against proteins without experimentally elucidated structures. We anticipate that as nanobodies have enabled intracellular sensing of proteins^{51,52} and their conformations^{53–56}, *de novo*-designed binders will enable a new generation of intracellular tools that can control the stability^{57–59} and function^{60–62} of proteins intracellularly.

Together, we hope the minibinder-coupled receptors described here prompt the rapid adoption of novel minibinders against a diversity of targets to dramatically expand the perceptual repertoire of cell engineering tools.

Materials and Methods

DNA Constructs

The synthetic receptors and minibinders used in this study (described in Supplemental Table 1 and schematized in Supplemental Figure 1) were constructed using the Mammalian Toolkit (MTK)⁶³, a hierarchical DNA assembly method. The SNIPR used in this study was a generous gift from Dr. Kole Roybal and contains the CD8a hinge domain, the human Notch1 transmembrane domain, and the human Notch2 juxtamembrane domain. The SNIPR was domesticated as an MTK part 3B with the Gal4-VP64 transcriptional actuator domain. The chimeric antigen receptor used in this study is comprised of the CD8a hinge and transmembrane domains, the 4-1BB costimulatory domain, and the CD3ζ chain and was also domesticated as an MTK part 3B. LCB1 linker variants were generated *de novo* as gene blocks ordered from IDT. The rigid linker is the Fibcon domain that has been previously described⁴⁸ with a sequence of
MLDAPTDLQVTNVTDSITVSWTPPSATITGYRITYTPSNG
PGEPKELTVPPSSTSVTITGLTPGVEYVVSVALKDNQESP PLVGTQTG.

For lentiviral transduction, a new set of MTK plasmids were generated as part of this study. The lentiviral plasmids previously used for SARSNotch¹⁸ were mutated to remove BsaI and BsmBI restriction sites. In the process of this study, we discovered that these plasmids were suboptimal for transfecting primary T cells. A novel MTK lentiviral backbone, pMK1101, was generated based on the pHR series of lentiviral plasmids that used the PacCI restriction sites for assembly. A new series of transcriptional unit vectors (pMK1213, pMK1129, pMK1130, pMK1131, pMK1132) were generated that allow transcriptional unit assembly via the standard BsaI MTK Golden Gate reaction that can then be assembled into pMK1101 via a PacCI reaction following manufacturer instructions (New England Biolabs #R0745S).

All plasmids were propagated in Stbl3 *E. coli* (QB3 MacroLab). Domestication was verified via sequencing and transcriptional unit assembly was verified via restriction digest and sequencing. All plasmids will be available on Addgene, and the sequence for all constructs used in this study are included as supplemental data.

Cell Culture

K562 and Jurkat cell lines were cultured in RPMI-1640 media (Gibco #11875-093) supplemented with 10% fetal bovine serum (FBS) (Hyclone #SH30071.03), and 1% Anti-Anti (Gibco #15240-096).

Primary human CD8+ T cells were isolated from anonymous donor blood after apheresis by negative selection. T cells were cryopreserved in RPMI-1640 (Corning #10-040-CV) with 20% human AB serum (Valley Biomedical, #HP1022) and 5% DMSO (Sigma-Aldrich #472301). After thawing, T cells were cultured in human T cell medium consisting of X-VIVO 15 (Lonza #04-418Q), 5% Human AB serum and 10 mM neutralized N-acetyl L-Cysteine (Sigma-Aldrich #A9165) supplemented with 30 units/mL IL-2 (NCI BRB Pre-clinical Repository).

Lenti-X 293T packaging cells (Clontech #11131D) were cultured in medium consisting of Dulbecco's Modified Eagle Medium (DMEM) (Gibco #10569-010) and 10% fetal bovine serum (FBS) (University of California, San Francisco [UCSF] Cell Culture Facility). Lenti-X 293T cells were cultured in 150 or 225 flasks (Corning #430825 and #431082) and passaged upon reaching 80% confluency. To passage, cells were treated with TrypLE express (Gloco #12605010) at 37°C for 5 minutes. Then 10 mL of media was used to quench the reaction and cells were collected into a 50 mL conical tube and pelleted by centrifugation (400xg for 5 minutes). Cells were cultured until passage 30 whereupon fresh Lenti-X 293T cells were thawed.

Vero-TMPRSS2 obtained from ATCC were cultured in DMEM (Corning) supplemented with 10% fetal bovine serum (FBS) (GeminiBio), 1% glutamine (Corning), and 1% penicillin-streptomycin (Corning).

All cells were maintained at 37°C with 5% CO₂, and cell counting was conducted using a Trypan Blue (Invitrogen #T10282) exclusion stain and an automated Countess II FL (Invitrogen #AMQAF1000) cell counter.

Lentiviral Transduction

Lenti-X 293T cells (Takara Bio #632180) were seeded at approximately 7e5 cells/well in a 6-well plate to yield ~80% confluency the following day. The following day cells were transfected with 1.5µg of transfer vector containing the desired expression cassette, and the lentiviral packaging plasmids pMD2.G (170ng) and pCMV-dR8.91(1.33µg) using 10µl of Eugene HD (Promega #E2312) according to manufacturer protocols. At 48 hours the viral supernatant was filtered through a 0.45µm PVDF filter and added to Jurkat or K562 cells seeded at approximately 1e5 cells/well in a 12-well plate.

After 24 hours, the viral media was then exchanged for fresh media. Polyclonal cell populations were selected via fluorescence activated cell sorting (FACS) as described below.

For each construct tested, 1e6 Primary T cells were thawed and activated the next day using CD3/CD28 Dynabeads (Life Technologies #11131D) at a 1:3 cell: bead ratio. Lenti-X 293T cells were transfected to generate virus the same day that T cells were activated. At 48 hours post-transfection, the viral supernatant was filtered through a 0.45µm PVDF filter and concentrated using the Lenti-X concentrator (Takara #631231) according to the manufacturer's instructions. For each construct, 2 wells of a 6-well plate of virus were concentrated together and then incubated in one well of a 24-well plate with 1e6 T cells. 24 hours after the addition of virus to T cell culture, viral media was removed and fresh media was added. At day 5 post T cell stimulation, Dynabeads were removed and the T cells were sorted for construct expression via FACS. T cells were then expanded for at least eight days until assays.

Antibodies

Surface-expressed proteins were assayed using Alexa Fluor 647 Anti-Myc tag antibody (Cell Signaling Technologies #2233S). T cell activation was assessed using Alexa Fluor 647 anti-CD25 (Biolegend #302618). CD19 expression was assessed using FITC anti-CD19 (Biolegend #302206). All antibodies were diluted 1:100 in DPBS (UCSF Cell Culture Facility) for staining.

Fluorescence Activated Cell Sorting (FACS)

Cell lines were bulk sorted for high expression using the UCSF Laboratory for Cell Analysis Core Facility FACSARIAII (BD Biosciences) or a Sony SH800.

Fluorescence-negative controls were used to set detector power so that negative cells appeared to have mean fluorescence ~100 counts, and then transduced cells were sorted for cells with expression outside of the negative control expression level. Jurkat cells transduced with synNotch and SNIPR output circuit were sorted for low BFP and high mCitrine expression. SynNotch and SNIPR transduced Jurkats cells were stained for surface expression and then sorted for low BFP, high mCherry, high surface expression, and high mCitrine expression.

Spike-expressing K562s were stained for surface expression and sorted for high surface expression and high mCherry. T cells transduced with CAR variants were stained for surface expression and then sorted for high surface expression and high mCherry.

Flow Cytometry

Flow cytometry was performed using a LSR-Fortessa (BD Biosciences). Prior to the flow cytometry, cells were seeded at densities described below in 96 well U bottom plates (Falcon #877217) and incubated for 24-72 hours as specified by the experiment. Plates were then spun at 400xg for 5 minutes to settle all cells. Cells were then resuspended and stained for 45 minutes with appropriate antibodies. After incubation, plates were spun down, rinsed with 200µl DPBS, and resuspended in flow buffer (DPBS with 2% FBS). The plates were then run on the flow cytometer using a four laser configuration (405nm, 488nm, 561nm, 640nm), collecting fluorescence for TagBFP (405nm ex, 450/50 em), mCitrine (488nm ex, 530/30 em, 505lp dichroic), mCherry (561nm ex, 610/20 em, 600lp dichroic), and Alexa Fluor 647 (640nm ex, 670/14 em). At least 10,000 events were recorded for all single cell line assays, and 30,000 events for experiments involving two cell lines. All data were collected using FACSDiva (BD Biosciences)

synNotch and SNIPR Activation Assays

Jurkats expressing synNotch and SNIPR constructs were seeded at a density of 2.5e4 cells/well in a U bottom 96 well plate. Cells were plated alone or with an equal number of target K562 cells in a total of 200µl of growth medium. Plates were spun briefly (400xg for 1 minute) to increase likelihood of cell-cell interaction. Cells were co-cultured for between 24 and 72 hours depending on the assay, and the BFP expression and surface expression of receptors and Spike were conducted using flow cytometry. SynNotch and SNIPR activation was assessed as described previously by fitting a two-component Gaussian mixture model to BFP expression data and estimating the fraction of the population in the ‘off’ and ‘on’ components¹⁸[\[link\]](#).

CAR Evaluation Assays

Target K562 cells were spun down, resuspended in T cell growth medium, and were plated in U bottom 96 well dishes at different densities using serial dilution with the 2.5e4 cells plated in wells for the 1:1 effector:target condition. CAR-expressing T cells were stained 1:1000 with CellTrace CFSE (ThermoFisher #C34570) for 20 minutes at 37C and then washed with DPBS. Stained cells were then plated at a density of 2.5e4 cells per well with and without target cells. Cells were co-cultured between 24 and 72 hours depending on experimental conditions. At the selected endpoint, cells were pelleted, stained for either CD25 expression in co-culture conditions or surface markers (anti-Myc or anti-CD19) in mono-culture wells. Cells were then assessed via flow cytometry. T cells and K562s were separated by gating on SSC and CellTrace fluorescence, with K562s having low CellTrace fluorescence and moderate to high SSC values and T cells having low SSC values and moderate to high CellTrace fluorescence. To calculate % target lysis, the number of K562 cells was averaged for each K562 target line across the three replicate wells where those targets were plated with untransduced T cells and then the number of K562s in all experimental wells for that cell line was divided by that number.

SARS-CoV-2 culture

SARS-CoV-2 isolates USA-WA1/2020 (BEI NR-52281), California (B.1.429), Beta (B.1.351, California Department of Health), Delta (B.1.617.2, California Department of Health), and UK (B.1.1.7, California Department of Health) was used for infection studies. All virus experiments were performed in a Biosafety Level 3 laboratory. SARS-CoV-2 stocks were propagated in Vero-TMPRSS2 cells and their sequence verified by next-generation sequencing. Viral stock titer was calculated using plaque-forming assays.

Viral Infection Studies

Jurkat cells were seeded into 96-well plates at 2.5×10^4 cells per well. Cells were rested for at least 24h before infection. At the time of infection, medium containing viral inoculum (MOI of 0.1 and 1) was added on the cells. At 72 hours post infection the cells were washed once with PBS/- and then incubated for 30 minutes at 4C with 1:1000 Bioscience Fixable Viability Dye eFluor 780 (Invitrogen #65-0865-14) in PBS/- before fixing with 4% PFA for 30 minutes, washed twice with PBS/-, and stored at 4C for further processing. Cells were prepared for flow cytometry by pelleting, resuspending in flow buffer, and then were assessed for BFP expression in cells without uptake of the viability dye. Percent population activation was assessed as described above in synNotch and SNIPR assays.

Plaque Forming assay

Cell supernatants were analyzed for viral particle formation for viral stocks. In brief, Vero-TMPRSS2 cells were seeded and incubated overnight. Cells were inoculated with 10^{-1} to 10^{-6} dilutions of the respective homogenates or supernatant in serum-free DMEM. After the 1-h absorption period, the media in the wells were overlaid with 2.5% Avicel (RC-591, Dupont). After 72 h, the overlay was removed, and the cells were fixed in 10% formalin for 1 h and stained with crystal violet for visualization.

Data Presentation, Analysis, and Availability

All experiments were performed in at least biological triplicate. All data bars are the mean of 3 replicates, with error presented as standard deviation of replicates, and individual replicate values presented as circles where possible. For flow cytometry experiments, fluorescence is reported as median fluorescence value for all cells within a single replicate, and the presented means and error are calculated between replicates. Events collected from flow cytometry were filtered to remove small events and then gated on FSC and SSC to capture singlet populations.

Where presented, histogram density estimates represent all events for all 3 replicates in a flow cytometry experiment, and are calculated via seaborn's `kdeplot` function with bandwidth adjustment of 0.2. All data analysis was conducted using custom Python scripts, available on [github](#)⁶⁴. Analysis was conducted in Jupyter⁶⁵ and relied on [numpy](#)⁶⁶, [matplotlib](#)⁶⁷, [seaborn](#)⁶⁹, [pandas](#)⁷⁰, [SciPy](#)⁷², [scikit-learn](#)⁷³ and [fcsparser](#). Protein schematics were inspired by their respective structures as indicated in figure legends and assembled using pieces from a proposed protein emoticon⁷⁴. All primary data from flow cytometry experiments are available on [Zenodo](#)⁷⁵. All figures assembled with Affinity Designer 2.

Acknowledgements

We thank the members of the Hana El-Samad, Wendell Lim, and Jeremy Reiter laboratories for their comments and feedback throughout this project, especially Hersh Bhargava and Devan Shah. We thank Manny De Vera, Pilar Lopez, and Sim Sidhu for essential assistance. Human T cells used in this study were generously isolated and donated by the Lim lab. We thank the Whelan laboratory for providing the Vero cells overexpressing human TMPRSS2 (Vero-TMPRSS2). ZYW thanks Bad Moves, Martha, Stephanie E. Crilly, and Kayleigh S. Stevenson for essential support.

Funding

ZYW was supported in part by NIH 5K12GM081266 and in part by NIH 1K99GM147825. IPC was supported by NIH F31 grant AI164671-01. We gratefully acknowledge support from the Roddenberry Foundation (M.O.), P. and E. Taft (M.O.) and the Pendleton Foundation (M.O.). M.O. is a Chan Zuckerberg Biohub – San Francisco Investigator.

Author Contributions

ZYW: Conceptualization, formal analysis, funding acquisition, investigation, methodology, resources, writing - original draft, writing - review & editing SS: Investigation, methodology, resources, formal analysis, writing - original draft MK: Investigation, methodology, resources, writing - review & editing IPC: Investigation, methodology, funding acquisition, writing - original draft, writing - review & editing MO: Supervision, funding acquisition, resources HE-S: Conceptualization, supervision, funding acquisition, writing - review & editing

References

1. Manhas J., Edelstein H. I., Leonard J. N., Morsut L. (2022) **The evolution of synthetic receptor systems** *Nat. Chem. Biol* **18**:244–255
2. Lim W. A. (2022) **The emerging era of cell engineering: Harnessing the modularity of cells to program complex biological function** *Science* **378**:848–852
3. Weber E. W., Maus M. V., Mackall C. L. (2020) **The Emerging Landscape of Immune Cell Therapies** *Cell* **181**:46–62
4. Eshhar Z., Waks T., Gross G., Schindler D. G. (1993) **Specific activation and targeting of cytotoxic lymphocytes through chimeric single chains consisting of antibody-binding domains and the gamma or zeta subunits of the immunoglobulin and T-cell receptors** *Proc. Natl. Acad. Sci* **90**:720–724
5. June C. H., Sadelain M. (2018) **Chimeric Antigen Receptor Therapy** *N. Engl. J. Med* **379**:64–73
6. Majzner R. G., Mackall C. L. (2019) **Clinical lessons learned from the first leg of the CAR T cell journey** *Nat. Med* **25**:1341–1355
7. Morsut L., Roybal K. T., Xiong X., Gordley R. M., Coyle S. M., Thomson M., Lim W. A. (2016) **Engineering Customized Cell Sensing and Response Behaviors Using Synthetic Notch Receptors** *Cell* **164**:780–791
8. Daringer N. M., Dudek R. M., Schwarz K. A., Leonard J. N. (2014) **Modular Extracellular Sensor Architecture for Engineering Mammalian Cell-based Devices** *ACS Synth. Biol* **3**:892–902
9. Scheller L., Strittmatter T., Fuchs D., Bojar D., Fussenegger M. (2018) **Generalized extracellular molecule sensor platform for programming cellular behavior** *Nat. Chem. Biol* **14**:723–729

10. Zhu I., Liu R., Garcia J. M., Hyrenius-Wittsten A., Piraner D. I., Alavi J., Israni D. V., Liu B., Khalil A. S., Roybal K. T. (2022) **Modular design of synthetic receptors for programmed gene regulation in cell therapies** *Cell* **185**:1431–1443
11. Roybal K. T., Williams J. Z., Morsut L., Rupp L. J., Kolinko I., Choe J. H., Walker W. J., McNally K. A., Lim W. A. (2016) **Engineering T Cells with Customized Therapeutic Response Programs Using Synthetic Notch Receptors** *Cell* **167**:419–432
12. Allen G. M. *et al.* (2022) **Synthetic cytokine circuits that drive T cells into immuneexcluded tumors** *Science* **378**
13. Choe J. H., Williams J. Z., Lim W. A. (2020) **Engineering T Cells to Treat Cancer: The Convergence of Immuno-Oncology and Synthetic Biology** *Annu. Rev. Cancer Biol* **4**:121–139
14. Hyrenius-Wittsten A., Su Y., Park M., Garcia J. M., Alavi J., Perry N., Montgomery G., Liu B., Roybal K. T. (2021) **SynNotch CAR circuits enhance solid tumor recognition and promote persistent antitumor activity in mouse models** *Sci. Transl. Med* **13**
15. Cho J. H., Okuma A., Sofjan K., Lee S., Collins J. J., Wong W. W. (2021) **Engineering advanced logic and distributed computing in human CAR immune cells** *Nat. Commun* **12**
16. Cho J. H., Collins J. J., Wong W. W. (2018) **Universal Chimeric Antigen Receptors for Multiplexed and Logical Control of T Cell Responses** *Cell* **173**:1426–1438
17. Dolberg T. B., Gunnels T. F., Ling T., Sarnese K. A., Crispino J. D., Leonard J. N. (2023) **Dolberg, T. B., Gunnels, T. F., Ling, T., Sarnese, K. A., Crispino, J. D. & Leonard, J. N. Building synthetic biosensors using red blood cell proteins. 2023.12.16.571988 Preprint at 10.1101/2023.12.16.571988 (2023) <https://doi.org/10.1101/2023.12.16.571988>**
18. Weinberg Z. Y. *et al.* (2021) **Sentinel cells enable genetic detection of SARS-CoV-2 Spike protein** *bioRxiv* <https://doi.org/10.1101/2021.04.20.440678>
19. Matsunaga S. *et al.* (2020) **Engineering Cellular Biosensors with Customizable Antiviral Responses Targeting Hepatitis B Virus** *iScience* **23**
20. Lee J. C., Brien H. J., Walton B. L., Eidman Z. M., Toda S., Lim W. A., Brunger J. M. (2023) **Instructional materials that control cellular activity through synthetic Notch receptors** *Biomaterials* **297**
21. Malaguti M., Portero Migueles R., Annoh J., Sadurska D., Blin G., Lowell S. (2022) **SynNPL: Synthetic Notch pluripotent cell lines to monitor and manipulate cell interactions in vitro and in vivo** *Development* **149**
22. Garibyan M. *et al.* (2023) **Garibyan, M., Hoffman, T., Makaske, T., Do, S., March, A. R., Cho, N., Pedroncelli, N., Lima, R. E., Soto, J., Jackson, B., Khademosseini, A., Li, S., McCain, M. & Morsut, L. Engineering Programmable Material-To-Cell Pathways Via Synthetic Notch Receptors To Spatially Control Cellular Phenotypes In Multi-Cellular Constructs. 2023.05.19.541497 Preprint at 10.1101/2023.05.19.541497 (2023) <https://doi.org/10.1101/2023.05.19.541497>**
23. Huang T.-H., Velho T., Lois C. (2016) **Monitoring cell-cell contacts in vivo in transgenic animals** *Development* **143**:4073–4084

24. Huang T., Niesman P., Arasu D., Lee D., De La Cruz A. L., Callejas A., Hong E. J., Lois C. (2017) **Tracing neuronal circuits in transgenic animals by transneuronal control of transcription (TRACT)** *eLife* **6**
25. Zhang S., Zhao H., Liu Z., Liu K., Zhu H., Pu W., He L., Wang R. A., Zhou B. (2022) **Monitoring of cell-cell communication and contact history in mammals** *Science* **378**
26. Toda S., McKeithan W. L., Hakkinen T. J., Lopez P., Klein O. D., Lim W. A. (2020) **Engineering synthetic morphogen systems that can program multicellular patterning** *Science* **370**
27. Miersch S., Sidhu S. S. (2012) **Synthetic antibodies: Concepts, potential and practical considerations** *Methods* **57**:486–498
28. Kuhn P., Fühner V., Unkauf T., Moreira G. M. S. G., Frenzel A., Miethe S., Hust M. (2016) **Recombinant antibodies for diagnostics and therapy against pathogens and toxins generated by phage display** *PROTEOMICS – Clin. Appl* **10**:922–948
29. Beasley M. D., Niven K. P., Winnall W. R., Kiefel B. R. (2015) **Bacterial cytoplasmic display platform Retained Display (ReD) identifies stable human germline antibody frameworks** *Biotechnol. J* **10**:783–789
30. McMahon C. *et al.* (2018) **Yeast surface display platform for rapid discovery of conformationally selective nanobodies** *Nat. Struct. Mol. Biol* **25**:289–296
31. Saper C. B. (2009) **A Guide to the Perplexed on the Specificity of Antibodies** *J. Histochem. Cytochem* **57**:1–5
32. Lee J. H., Yin R., Ofek G., Pierce B. G. (2022) **Structural Features of Antibody-Peptide Recognition** *Front. Immunol* **13**
33. Kortemme T., Baker D. (2004) **Computational design of protein–protein interactions** *Curr. Opin. Chem. Biol* **8**:91–97
34. Marchand A., Van Hall-Beauvais A. K., Correia B. E. (2022) **Computational design of novel protein–protein interactions – An overview on methodological approaches and applications** *Curr. Opin. Struct. Biol* **74**
35. Cao L. *et al.* (2022) **Design of protein-binding proteins from the target structure alone** *Nature* **605**:551–560
36. Gainza P. *et al.* (2023) **De novo design of protein interactions with learned surface fingerprints** *Nature* **617**:176–184
37. Bennett N. R. *et al.* (2023) **Improving de novo protein binder design with deep learning** *Nat. Commun* **14**
38. Watson J. L. *et al.* (2023) **De novo design of protein structure and function with RFdiffusion** *Nature* **620**:1089–1100
39. Torres S. V. *et al.* (2023) **De novo design of high-affinity binders of bioactive helical peptides** *Nature* :1–3 <https://doi.org/10.1038/s41586-023-06953-1>
40. Cao L. *et al.* (2020) **De novo design of picomolar SARS-CoV-2 miniprotein inhibitors** *Science* **370**:426–431

41. Hsieh C.-L. *et al.* (2020) **Structure-based design of prefusion-stabilized SARS-CoV-2 spikes** *Science* **369**:1501–1505
42. Imai C., Mihara K., Andreansky M., Nicholson I. C., Pui C.-H., Geiger T. L., Campana D. (2004) **Chimeric receptors with 4-1BB signaling capacity provoke potent cytotoxicity against acute lymphoblastic leukemia** *Leukemia* **18**:676–684
43. Zola H., Macardle P. J., Bradford T., Weedon H., Yasu H., Kurosawa Y. (1991) **Preparation and characterization of a chimeric CD 19 monoclonal antibody** *Immunol. Cell Biol* **69**:411–422
44. Hudecek M., Lupo-Stanghellini M.-T., Kosasih P. L., Sommermeyer D., Jensen M. C., Rader C., Riddell S. R. (2013) **Receptor Affinity and Extracellular Domain Modifications Affect Tumor Recognition by ROR1-Specific Chimeric Antigen Receptor T Cells** *Clin. Cancer Res* **19**:3153–3164
45. Liu X. *et al.* (2015) **Affinity-Tuned ErbB2 or EGFR Chimeric Antigen Receptor T Cells Exhibit an Increased Therapeutic Index against Tumors in Mice** *Cancer Res* **75**:3596–3607
46. Watanabe N., Bajgain P., Sukumaran S., Ansari S., Heslop H. E., Rooney C. M., Brenner M. K., Leen A. M., Vera J. F. (2016) **Fine-tuning the CAR spacer improves T-cell potency** *OncoImmunology* **5**
47. Ghorashian S. *et al.* (2019) **Enhanced CAR T cell expansion and prolonged persistence in pediatric patients with ALL treated with a low-affinity CD19 CAR** *Nat. Med* **25**:1408–1414
48. Jacobs S. A., Diem M. D., Luo J., Teplyakov A., Obmolova G., Malia T., Gilliland G. L., O’Neil K. T. (2012) **Design of novel FN3 domains with high stability by a consensus sequence approach** *Protein Eng. Des. Sel* **25**:107–117
49. Jumper J. *et al.* (2021) **Highly accurate protein structure prediction with AlphaFold** *Nature* **596**:583–589
50. Baek M. *et al.* (2021) **Accurate prediction of protein structures and interactions using a three-track neural network** *Science* **373**:871–876
51. Rothbauer U. *et al.* (2006) **Targeting and tracing antigens in live cells with fluorescent nanobodies** *Nat. Methods* **3**:887–889
52. Kirchhofer A. *et al.* (2010) **Modulation of protein properties in living cells using nanobodies** *Nat. Struct. Mol. Biol* **17**:133–138
53. Irannejad R. *et al.* (2013) **Conformational biosensors reveal GPCR signalling from endosomes** *Nature* **495**:534–538
54. Livingston K. E., Mahoney J. P., Manglik A., Sunahara R. K., Traynor J. R. (2018) **Measuring ligand efficacy at the mu-opioid receptor using a conformational biosensor** *eLife* **7**
55. Stoeber M., Jullié D., Lobingier B. T., Laeremans T., Steyaert J., Schiller P. W., Manglik A., von Zastrow M. (2018) **A Genetically Encoded Biosensor Reveals Location Bias of Opioid Drug Action** *Neuron* **98**:963–976
56. Crilly S. E., Ko W., Weinberg Z. Y., Puthenveedu M. A. (2021) **Conformational specificity of opioid receptors is determined by subcellular location irrespective of agonist** *eLife* **10**

57. Kim M. (2023) **A Degron-Based bioPROTAC for Controlling Signaling in CAR T Cells**
58. Lim S. *et al.* (2020) **bioPROTACs as versatile modulators of intracellular therapeutic targets including proliferating cell nuclear antigen (PCNA)** *Proc. Natl. Acad. Sci* **117**:5791–5800
59. Caussinus E., Kanca O., Affolter M. (2012) **Fluorescent fusion protein knockout mediated by anti-GFP nanobody** *Nat. Struct. Mol. Biol* **19**:117–121
60. Cheloha R. W., Harmand T. J., W?ne C., Schwartz T. U., Ploegh H. L. (2020) **Exploring cellular biochemistry with nanobodies** *J. Biol. Chem* **295**:15307–15327
61. Helma J., Cardoso M. C., Muyldermans S., Leonhardt H. (2015) **Nanobodies and recombinant binders in cell biology** *J. Cell Biol* **209**:633–644
62. Gil A. A., Carrasco-López C., Zhu L., Zhao E. M., Ravindran P. T., Wilson M. Z., Goglia A. G., Avalos J. L., Toettcher J. E. (2020) **Optogenetic control of protein binding using light-switchable nanobodies** *Nat. Commun* **11**
63. Fonseca J. P. *et al.* (2019) **A Toolkit for Rapid Modular Construction of Biological Circuits in Mammalian Cells** *ACS Synth. Biol* **8**:2593–2606
64. Weinberg Z. (2024) **Weinberg, Z. weinberg/synbinders: Initial Release. (2024).** doi:10.5281/zenodo.10480824 <https://doi.org/10.5281/zenodo.10480824>
65. Kluyver T. *et al.* (2016) **Jupyter Notebooks -a publishing format for reproducible computational workflows** *Position. Power Acad. Publ. Play. Agents Agendas* :87–90
66. Harris C. R. *et al.* (2020) **Array programming with NumPy** *Nature* **585**:357–362
67. Caswell T. A. *et al.* (2023) **Caswell, T. A., Andrade, E. S. de Lee, A., Droettboom, M., Hoffmann, T., Klymak, J., Hunter, J., Firing, E., Stansby, D., Varoquaux, N., Nielsen, J. H., Gustafsson, O., Sunden, K., Root, B., May, R., Elson, P., Seppänen, J. K., hannah Lee, J.-J., Dale, D., McDougall, D., Straw, A., Hobson, P., Lucas, G., Comer, R., Gohlke, C., Vincent, A. F., Yu, T. S., Ma, E. & Silvester, S. matplotlib/matplotlib: REL: v3.7.4. (2023).** doi:10.5281/zenodo.10152802 <https://doi.org/10.5281/zenodo.10152802>
68. Hunter J. D. (2007) **Matplotlib: A 2D Graphics Environment** *Comput. Sci. Eng* **9**:90–95
69. Waskom M. L. (2021) **seaborn: statistical data visualization** *J. Open Source Softw* **6**
70. The pandas development team. pandas-dev/pandas: Pandas (2023) **The pandas development team. pandas-dev/pandas: Pandas. (2023).** doi:10.5281/zenodo.10426137 <https://doi.org/10.5281/zenodo.10426137>
71. McKinney W., Walt S. van der, Millman J (2010) **Data Structures for Statistical Computing in Python** *Proc. 9th Python Sci. Conf* :56–61 <https://doi.org/10.25080/Majora-92bf1922-00a>
72. Virtanen P. *et al.* (2020) **SciPy 1.0: fundamental algorithms for scientific computing in Python** *Nat. Methods* **17**:261–272
73. Pedregosa F. *et al.* (2011) **Scikit-learn: Machine Learning in Python** *J. Mach. Learn. Res* **12**:2825–2830

74. White A. (2023) **White, A. whitead/protein-emoji. (2023).**
<<https://github.com/whitead/protein-emoji>>
75. Weinberg Z., Soliman S., Kim M., Chen I., Ott M., El-Samad H. (2024) **Weinberg, Z., Soliman, S., Kim, M., Chen, I., Ott, M. & El-Samad, H. Data supporting 'De novo-designed minibinders expand the synthetic biology sensing repertoire'. (2024).** doi:10.5281/zenodo.10480660
<https://doi.org/10.5281/zenodo.10480660>

Article and author information

Zara Y. Weinberg

Department of Biochemistry & Biophysics, UCSF, San Francisco CA

For correspondence: weinberz@gmail.com

ORCID iD: [0000-0001-7176-038X](https://orcid.org/0000-0001-7176-038X)

Sarah S. Soliman

Department of Biochemistry & Biophysics, UCSF, San Francisco CA

ORCID iD: [0009-0003-8918-3332](https://orcid.org/0009-0003-8918-3332)

Matthew S. Kim

Tetrad Graduate Program, UCSF, San Francisco CA, Cell Design Institute, San Francisco CA

ORCID iD: [0000-0002-5836-8874](https://orcid.org/0000-0002-5836-8874)

Irene P. Chen

Gladstone Institutes, San Francisco CA, Department of Medicine, UCSF, San Francisco CA

ORCID iD: [0000-0002-5766-9253](https://orcid.org/0000-0002-5766-9253)

Melanie Ott

Gladstone Institutes, San Francisco CA, Department of Medicine, UCSF, San Francisco CA, Chan Zuckerberg Biohub–San Francisco, San Francisco CA

ORCID iD: [0000-0002-5697-1274](https://orcid.org/0000-0002-5697-1274)

Hana El-Samad

Department of Biochemistry & Biophysics, UCSF, San Francisco CA, Cell Design Institute, San Francisco CA, Chan Zuckerberg Biohub–San Francisco, San Francisco CA, Altos Labs, San Francisco CA

For correspondence: helsamad@altoslabs.com

ORCID iD: [0000-0001-6239-9916](https://orcid.org/0000-0001-6239-9916)

Copyright

© 2024, Weinberg et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Ahmad Khalil

Boston University, Boston, United States of America

Senior Editor

Satyajit Rath

Indian Institute of Science Education and Research (IISER), Pune, India

Reviewer #1 (Public Review):

Summary:

The authors want to explore how much two known minibinder protein domains against the Spike protein of SARS-CoV-2 can function as a binding domain of 2 sets of synthetic receptors (SNIPR and CAR); the authors also want to know how some modifications of the linkers of these new receptors affect their activation profile.

Major strengths and weaknesses of the methods and results:

- Strengths include: analysis of synthetic receptor function for 2 classes of synthetic receptors, with robust and appropriate assays for both kinds of receptors. The modifications of the linkers are also interesting and the types of modifications that are often used in the field.

- Weaknesses include: none of the data analysis provides statistical interpretation of the results (that I could find). One dataset is confusing: Figures 5A and C, are said to be the same assay with the same constructs, but the results are 30% in A, and 70% in C.

An appraisal of whether the authors achieved their aims, and whether the results support their conclusions:

Given the open-ended nature of the goal (implicit in it being an exploration), it is hard to say if the authors have reached their aims; they have done an exploration for sure; is it big enough an exploration? This reviewer is not sure.

The results are extremely clearly presented, both in the figures and in the text, both for the methods and the results. The claims put forward (with limited exceptions see below) are very solidly supported by the presented data.

A discussion of the likely impact of the work on the field, and the utility of the methods and data to the community:

The work may stimulate others to consider minibinders as potential binding domains for synthetic receptors. The modifications that are presented although not novel, do provide a starting point for larger-scale analysis.

It is not clear how much this is generalizable to other binders (the authors don't make such claims though). The claims are very focused on the tested modifications, and the 2 receptors and minibinder used, a scope that I would define as narrow; the take-home message if one wants to try it with other minibinders or other receptors seems to be: test a few things, and your results may surprise you.

Any additional context you think would help readers interpret or understand the significance of the work:

We are at the infancy stage of synthetic receptors optimization and next-generation derivation; there is a dearth of systematic studies, as most focus is on developing a few ones that work. This work is an interesting attempt to catalyze more research with these new minibinders. Will it be picked up based on this? Not sure.

<https://doi.org/10.7554/eLife.96154.1.sa1>

Reviewer #2 (Public Review):

Summary:

Weinberg et al. show that spike LCB minibinders can be used as the extracellular domain for SynNotch, SNIPR, and CAR. They evaluated their designs against cells expressing the target proteins and live virus.

Strengths:

This is a good fundamental demonstration of alternative use of the minibinder. The results are unsurprising but robust and solid in most cases.

Weaknesses:

The manuscript would benefit from better descriptions of the study's novelty. Given that LCB previously worked in SynNotch, what unexpected finding was uncovered by this study? It is well known that the extracellular domain of CAR is amenable to different types of binding domains (e.g., scFv, nanobody, DARPIn, natural ligands). So, it is not surprising that a minibinder also works with CAR. We don't know if the minibinders are more or less likely to be compatible with CAR or SNIPR.

The demonstrations are all done using just 1 minibinder. It is hard to conclude that minibinders, as a unique class of protein binders, are generalizable in different contexts. All it can conclude is that this specific Spike minibinder can be used in synNotch, SNIPR, and CAR. The LCB3 minibinder seems to be much weaker.

The sensing of live viruses is interesting, but the output is very weak. It is difficult to imagine a utility for such a weak response.

<https://doi.org/10.7554/eLife.96154.1.sa0>

Author response:

We thank the reviewers for their attention to our study and for their fair and reasonable assessment of the strengths and weaknesses of our work. We believe the reviewers adequately captured both the potential implications of our work as well as its major current limitations. As both reviewers noted, we believe the work presented in this manuscript is an exciting first step in adapting minibinders as antigen sensors for synthetic receptors but many questions remain before these new tools can be widely adopted. We hope that this work will catalyze others to try minibinders as potential antigen sensors when developing novel synthetic receptors, and we hope that future work will more thoroughly test a wide range of linkers to better optimize antigen sensor function across synthetic receptors.

In our future work, we intend to evaluate a greater diversity of minibinders across different relevant therapeutic targets. We are working to test both existing minibinders as well as generate novel minibinders using deep-learning-based de novo protein design methods. We further hope to explore additional linker modifications, especially focusing on modifications

that will allow minibinder coupled-synthetic receptors to escape the glycocalyx of engineered cells. We hope to share findings on these topics in either an update to this manuscript or in future manuscripts, depending on the results of our studies in progress.

Finally, reviewers noted a mismatch in the data displayed in Figure 5A and 5C, whereby LCB-CAR-expressing cells induced higher lysis in Figure 5C than in Figure 5A. This is due to figure 5C showing only 24 hours of incubation between effector and target cells, as opposed to the 72 hours of incubation that is quantitated in 5A. These mismatched timepoints were selected because linker-dependent differences in lysis were most readily apparent at 24 hours and were negligible at 72 hours. The full-time course of lysis for this experiment can be seen in Supplemental Figure 2D.