

Neurons enhance blood-brain barrier function via upregulating claudin-5 and VE-cadherin expression due to GDNF secretion

Lu Yang, Zijin Lin, Ruijing Mu, Wenhan Wu, Hao Zhi, Xiaodong Liu , Hanyu Yang , Li Liu 

Department of Pharmacology, School of Pharmacy, China Pharmaceutical University, Nanjing 210009, China

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Blood-brain barrier (BBB) prevents neurotoxins from entering central nervous system. We aimed to establish and characterize an *in vitro* triple co-culture BBB model consisting of brain endothelial cells hCMEC/D3, astrocytoma U251 cells, and neuroblastoma SH-SY5Y cells. Co-culture of SH-SY5Y and U251 cells markedly enhanced claudin-5 and VE-cadherin expression in hCMEC/D3 cells, accompanied by increased transendothelial electrical resistance and decreased permeability. Conditioned medium (CM) from SH-SY5Y cells (S-CM), U251 cells (U-CM), and co-culture of SH-SY5Y and U251 cells (US-CM) also promoted claudin-5 and VE-cadherin expression. Glial cell line-derived neurotrophic factor (GDNF) levels in S-CM and US-CM were significantly higher than CMs from hCMEC/D3 and U-CM. Both GDNF and US-CM upregulated claudin-5 and VE-cadherin expression, which were attenuated by anti-GDNF antibody and GDNF signaling inhibitors. GDNF increased claudin-5 expression via the PI3K/AKT/FOXO1 and MAPK/ERK pathways. Meanwhile, GDNF promoted VE-cadherin expression by activating PI3K/AKT/ETS1 and MAPK/ERK/ETS1 signaling. The roles of GDNF in BBB integrity were validated using brain-specific *Gdnf* silencing mice. The developed triple co-culture BBB model was successfully applied to predict BBB permeability. In conclusion, neurons enhance BBB integrity by upregulating claudin-5 and VE-cadherin expression through GDNF secretion and established triple co-culture BBB model may be used to predict drugs' BBB permeability.

eLife assessment

The manuscript by Yang and coworkers presents **valuable** evidence that an *in vitro* brain blood barrier composed of endothelial cells, astrocytes, and neuroblastoma cells of human origin, would resemble better the *in vivo* condition. The presented results constitute **solid** evidence that GDNF induces the expression of VE-Cadherin and Claudin-5. Further, silencing of GDNF in the brain of mice altered brain blood barrier properties. This provides a new perspective on the interaction between neurons and endothelial cells and this model can be used to screen the permeability of the brain blood barrier to different drugs.

<https://doi.org/10.7554/eLife.96161.1.sa2>

Introduction

As a dynamic interface between the blood circulatory system and the central nervous system (CNS), the blood-brain barrier (BBB) maintains homeostasis and normal function of CNS by strictly regulating material exchange between the blood and brain (Palmiotti et al., 2014 [↗](#)). The maintenance of BBB is mainly attributed to the expression of tight junctions (TJ) and adherent junctions (AJ) between adjacent brain endothelial cells as well as a variety of drug transporters. However, as a double-edged sword that protects CNS function, BBB also restricts the transport of some drugs from blood to brain, leading to poor CNS therapeutic effects and even CNS treatment failure (Banks, 2016 [↗](#)).

Several *in silico*, *in vitro*, *in situ*, and *in vivo* methods have been developed to assess the permeability of drugs across BBB, but each method has its limitations (Hanafy et al., 2021 [↗](#)). The *in situ* brain perfusion (ISBP) is considered the “gold standard” for assessing BBB permeability but there exist limits related to animal ethics. Moreover, ISBP is not suitable for human or high-throughput workflows. Thus, a suitable, accurate, and high-throughput *in vitro* BBB model is required to predict the permeability of drug candidates through BBB.

BBB is formed by neurovascular units (NVU), composed of neural (neurons, microglia, and astrocytes) and vascular components (vascular endothelial cells, pericytes, and vascular smooth muscle cells) (Potjewyd et al., 2018 [↗](#)). The constant crosstalk and interactions among these cells contribute to the structural and signaling-based regulation of transcellular and paracellular transport, control of BBB permeability, and regulation of cerebral circulation (Arvanitis et al., 2020 [↗](#); Muoio et al., 2014 [↗](#); Potjewyd et al., 2018 [↗](#)). Although brain microvascular endothelial cells (BMECs) are commonly utilized as an *in vitro* BBB model to assess drug permeability across BBB, *in vitro* mono-culture of BMECs tends to lose some unique characteristics without the support of other cell types. To overcome the drawbacks of mono-culture BBB models, a range of multicellular co-culture BBB models co-cultured with other NVU elements (such as astrocytes or pericytes) have been developed. Astrocytes are the most abundant glial cell type in brain (Clasadonte et al., 2017). Their terminal feet cover >80% of the surface of capillaries to form interdigitating coverage without slits (Mathiisen et al., 2010 [↗](#)). In addition, astrocytes also contain several proteins related to the tight binding of the basement membrane. The *in vitro* multicellular co-culture BBB models composed of astrocytes, pericytes, and endothelial cells are considered to mimic the vascular structure and offer paracellular tightness in BBB (Ito et al., 2019 [↗](#); Nakagawa et al., 2009 [↗](#); Watanabe et al., 2021 [↗](#)).

Neurons also play a crucial role in the NVU and may be involved in the regulation of BBB function. The maturation of BBB in mice considerably overlaps with the establishment of neuronal activity (Biswas et al., 2020 [\[1\]](#)). A study revealed that the conditioned medium collected from primary rat neurons also attenuated cell death caused by glucose-oxygen-serum deprivation (Lin et al., 2016 [\[2\]](#)). Furthermore, the primary rat neurons were reported to affect the differentiation and formation of BMECs (Savettieri et al., 2000 [\[3\]](#); Schiera G. et al., 2003; Xue et al., 2013 [\[4\]](#)). Recent research reported that, compared with the double co-culture of hCMEC/D3 and 1321N1 (human astrocytoma cells), the triple co-culture of hCMEC/D3, 1321N1, and human neuroblastoma SH-SY5Y cells exhibited a higher transendothelial electrical resistance (TEER) (Barberio et al., 2022 [\[5\]](#)). A similar report showed that co-cultivation of RBE4.B (rat brain capillary endothelial cells) and neurons resulted in the lower permeability of [³H] sucrose than RBE4.B cells grown alone (Schiera Gabriella. et al., 2005). All these studies indicate that neurons, as the elements of the NVU, may be tightly connected to the formation and maintenance of BBB functions.

The aims of the study were: 1) to establish and characterize an *in vitro* triple co-culture BBB model consisting of human brain endothelial cells (hCMEC/D3), human astrocytoma cells (U251), and human neuroblastoma cells (SH-SY5Y); 2) to investigate whether neurons were involved in the formation and maintenance of BBB integrity and explore their underlying mechanisms. The integrity of BBB was evaluated by quantifying the leakage of both fluorescein and FITC-Dextran 3–5 kDa (FITC-Dex); 3) to validate the *in vitro* results through *in vivo* experiments. Finally, the *in vivo/in vitro* correlation assay (IVIVC) was analyzed to prove that the triple co-culture BBB model could better predict the BBB penetration of CNS drugs compared with the mono-culture BBB model.

Result

Establishment and characterization of the *in vitro* triple co-culture BBB model

TEER values were measured during the co-culture (Figure 1C [\[6\]](#)). TEER values of the four *in vitro* BBB models gradually increased until day 6. On day 7, the TEER values showed a decreasing trend. Thus, a 6-day co-culture was used for subsequent experiments. The highest TEER values were observed in the triple co-culture BBB model, followed by double co-culture with U251 cells and double co-culture with SH-SY5Y cells BBB models, hCMEC/D3 cells mono-culture BBB model showed the lowest TEER values. We also measured the TEER values of U251 monolayer cells, and the results showed that U251 cells themselves also contributed to the physical barrier of the model (Figure 1D [\[7\]](#)). The apparent permeability coefficients (P_{app}) values of fluorescein and FITC-Dex were measured to characterize the integrity of the four BBB models (Figure 1E [\[8\]](#), 1F). Consistent with the TEER values, the triple co-culture BBB model showed the lowest permeability of fluorescein and FITC-Dex, followed by double co-culture with U251 cells and double co-culture with SH-SY5Y cells BBB models. It was noticed that the monolayer of U251 cells itself also worked as a barrier, preventing the leakage of permeability markers, which may explain why the permeability of FITC-Dex in the double co-culture model with U251 cells is lower than that in the double co-culture model with SH-SY5Y cells (Figure 1G [\[9\]](#)). Moreover, co-culture with SH-SY5Y, U251, and U251 + SH-SY5Y cells significantly enhanced the proliferation of hCMEC/D3 cells by 5.80%, 4.61%, and 10.41%, respectively (Figure 1H [\[10\]](#)).

The paracellular barrier of BBB is associated with TJs, AJs, and transporters (Abbott, 2013 [\[11\]](#)). The mRNA levels of TJs (claudin-5, ZO-1, and occludin), AJs (VE-cadherin and β -catenin), and transporters (P-gp, BCRP, OCT-1, OCT-2, OAT-3, and OATP1A1) in hCMEC/D3 cells from the four BBB models were analyzed using qPCR (Figure 1I [\[12\]](#) and 1J [\[13\]](#)). Compared with hCMEC/D3 cell mono-culture model, double co-culture with SH-SY5Y, double co-culture with U251, and triple co-culture BBB models showed markedly increases in claudin-5, VE-cadherin, and BCRP mRNA expression.




Figure 1.

The effects of co-culture with U251 and/or SH-SY5Y cells on the integrity of hCMEC/D3 and BBB function.

(A): Schematic diagram of the establishment process of the triple co-culture BBB model. (B): Four types of BBB models: h: hCMEC/D3 cells mono-culture model; h+S: double co-culture with SH-SY5Y cells BBB model; h+U: double co-culture with U251 cells BBB model; h+S+U: triple co-culture model. (C): The transendothelial electrical resistance (TEER) of four models, and the TEER values in day 6 were compared. Blank: the inserts without seeding cells. $n=4$. (D): The TEER of hCMEC/D3 and U251 cells monolayer. U: U251 cells mono-culture model. $n=4$. (E and F): The apparent permeability coefficient (P_{app} , $\times 10^{-6}$ cm/s) of fluorescein (NaF) and FITC-Dextran 3–5 kDa (FITC-Dex) of four BBB models. $n=4$. (G): The P_{app} ($\times 10^{-6}$ cm/s) of NaF and FITC-Dex across the blank inserts, and hCMEC/D3 or U251 mono-culture models. $n=4$. (H): The cell density of hCMEC/D3 cells after mono/co-culturing. $n=3$. (I and J): The mRNA levels of tight junction proteins, adherent junction proteins, and transporters. $n=4$. (K and L): The protein expression levels of claudin-5 (CLDN-5), ZO-1, occludin in hCMEC/D3 cells. $n=4$. (M and N): The protein expression levels of VE-cadherin (VE-Cad), β -catenin, and BCRP in hCMEC/D3 cells. $n=4$. (O and P): The correlations between the P_{app} ($\times 10^{-5}$ cm/s) of NaF and claudin-5 expression (O), or VE-cadherin expression (P). (Q and R): The correlation between P_{app} ($\times 10^{-5}$ cm/s) of FITC-Dex and claudin-5 expression (Q), or VE-cadherin expression (R). The above data are shown as the mean $\pm$ S.E.M. Statistical significance was determined using one-way ANOVA with Fisher's LSD test or Welch's ANOVA test. The simple linear regression analysis was used to examine the presence of a linear relationship between two variables.

Figure 1—source data1 (Raw and annotated data files for western blot)

Figure 1—source data2 (Summary of original data and analysis)

Expression of corresponding proteins was measured using Western blot (**Figure 1K-1N** [↗](#)). Notably increased claudin-5 expression was detected in double co-culture with SH-SY5Y cells and triple co-culture BBB models, while VE-cadherin expression was markedly increased in double co-culture with U251 cells and triple co-culture BBB models. Expression levels of other TJ proteins (ZO-1 and occludin) and AJ protein (β -catenin) were unaltered. The expression of BCRP was slightly affected by co-cultivation without statistical differences. Significant negative correlations were found between P_{app} values of fluorescein and the expression of claudin-5 or VE-cadherin. P_{app} values of FITC-Dex were also negatively correlated to the expression levels of claudin-5 or VE-cadherin (**Figure 1O-1R** [↗](#)). These results indicate that the decreased permeability of fluorescein and FITC-Dex mainly results from the upregulated expression of both claudin-5 and VE-cadherin.

Neurons and astrocytes upregulated the expression of claudin-5 and VE-cadherin by GDNF secretion

The hCMEC/D3 cells did not direct contact with U251 or SH-SY5Y cells in the double co-culture and triple co-culture BBB models, indicating that cell-cell interaction between U251, SH-SY5Y, and hCMEC/D3 cells relied on secreted active factors. To test this hypothesis, the effects of conditioned medium (CM) from SH-SY5Y cells (S-CM), U251 cells (U-CM), and co-culture of SH-SY5Y and U251 cells (US-CM) on the expression of claudin-5 and VE-cadherin in hCMEC/D3 cells were analyzed (**Figure 2A** [↗](#)). Both S-CM, U-CM, and US-CM markedly increased the expression of claudin-5 and VE-cadherin. US-CM showed the strongest induction effects on claudin-5 and VE-cadherin.

To investigate which cytokines were involved in the promotion of hCMEC/D3 cell integrity by U251 and SH-SY5Y cells, the mRNA expression levels of various cytokines in these three types of cells were compared. The results showed that U251 or SH-SY5Y cells exhibited significantly higher expression levels of GDNF, nerve growth factor (NGF), insulin-like growth factor-1 (IGF-1), vascular endothelial growth factor (VEGF), basic fibroblast growth factor (bFGF), and transforming growth factor- β (TGF- β) compared to hCMEC/D3 cells (**Figure 2B** [↗](#)). Furthermore, the mRNA expression of GDNF, IGF-1, TGF- β , and bFGF in SH-SY5Y cells was higher than those in U251 cells. In these cytokines, GDNF (Dong et al., 2018; Igarashi et al., 1999 [↗](#); Shimizu et al., 2012 [↗](#)), bFGF (Shimizu et al., 2011 [↗](#); Wang et al., 2016 [↗](#)), IGF-1 (Ko et al., 2009; Nowranghi et al., 2019 [↗](#)), and TGF- β (Fu et al., 2021 [↗](#)) have been reported to promote BBB integrity. Thus, the concentrations of GDNF, bFGF, IGF-1, and TGF- β in the CMs were measured (**Figure 2C** [↗](#)). The results showed that levels of GDNF, bFGF, and IGF-1 in S-CM and US-CM were significantly higher than CMs from hCMEC/D3 cells (H-CM) and U-CM, but levels of TGF- β in S-CM and U-CM were lower than those in H-CM. Interestingly, the level of IGF-1 in US-CM was remarkably lower than that in S-CM, indicating that U251 cells suppressed IGF-1 secretion from SH-SY5Y. The effects of GDNF, bFGF, IGF-1, and TGF- β on the expression of claudin-5 and VE-cadherin were investigated (**Figure 2D-2G** [↗](#)). Among the four tested neurotrophic factors, only GDNF induced the expression of claudin-5 and VE-cadherin in a concentration-dependent manner (**Figure 2D** [↗](#)). In contrast, a high level (5 ng/mL) of TGF- β slightly downregulated claudin-5 and VE-cadherin expression (**Figure 2G** [↗](#)). These results demonstrate that upregulation of claudin-5 and VE-cadherin expression by US-CM are attributed to secreted GDNF.

To provide additional verification of the deduction, anti-GDNF antibody was used to neutralize exogenous and endogenous GDNF in culture medium. Consistent with our expectation, the anti-GDNF antibody concentration-dependent reversed the US-CM-induced claudin-5 and VE-cadherin expression (**Figure 2H** [↗](#)). Furthermore, GDNF-induced upregulation of claudin-5 and VE-cadherin expression was also reversed by the anti-GDNF antibody (**Figure 2I** [↗](#)). The induction effects of US-CM and GDNF on the claudin-5 and VE-cadherin expression in hCMEC/D3 cells were also confirmed in primary rat brain microvascular endothelial cells (**Figure 2J** [↗](#)).




Figure 2.

Neurons and astrocytes upregulated claudin-5 and VE-cadherin expression in hCMEC/D3 cells due to GDNF secretion.

(A): Effects of conditioned medium (CM) on claudin-5 and VE-cadherin expression. Con: the normal medium; S-CM: the CM from SH-SY5Y cells; U-CM: the CM from U251 cells; US-CM: the CM from SH-SY5Y cells co-culture with U251 cells. (B): The mRNA expression levels of neurotrophic factors in hCMEC/D3, U251, and SH-SY5Y cells. (C): Concentrations of GDNF, bFGF, IGF-1, and TGF- β in the CMs. H-CM: the CM from hCMEC/D3 cells. (D-G): Effects of GDNF (D), IGF-1 (E), bFGF (F), and TGF- β (G) on the expression of claudin-5 and VE-cadherin. The dosages have been marked in the figure. (H and I): Effects of anti-GDNF antibody on the upregulation of claudin-5 and VE-cadherin expression induced by US-CM (H) or 200 pg/mL GDNF (I). (J): Effects of 200 pg/mL GDNF and US-CM on claudin-5 and VE-cadherin expression in primary rat brain microvascular endothelial cells. (K and L): Effects of 3 μ M RET tyrosine kinase inhibitor SSP-86 (SPP), and 5 μ M Src family kinases inhibitor PP2 on the upregulation of claudin-5 and VE-cadherin induced by 200 pg/mL GDNF (K) and US-CM (L). (M and N): Effects of SPP on the TEER (M), the permeability of NaF, and FITC-Dex (N) of the hCMEC/D3 mono-culture BBB model treating 200 pg/mL GDNF. The TEER values of day 6 were compared. (O and P): Effects of PP2 on the TEER (O), the permeability of NaF, and FITC-Dex (P) of the hCMEC/D3 mono-culture BBB model treating 200 pg/mL GDNF. The TEER values of day 6 were compared. (Q and R): Effects of SPP on the TEER (Q), the permeability of NaF, and FITC-Dex (R) of the triple co-culture BBB model. The TEER values of day 6 were compared. (S and T): Effects of PP2 on the TEER (S), the permeability of NaF, and FITC-Dex (T) of the triple co-culture BBB model. The TEER values of day 6 were compared. The above data are shown as the mean $\pm$ S.E.M.; n=4. Statistical significance was determined using one-way ANOVA with Fisher's LSD test or Welch's ANOVA test.

Figure 2—source data1 (Raw and annotated data files for western blot)

Figure 2—source data2 (Summary of original data and analysis)

GDNF forms a heterohexameric complex with two GFR α 1 molecules and two RET receptors to activate the GDNF-GFR α 1-RET signaling (Fielder et al., 2018 [\[3\]](#)). The RET receptor tyrosine kinase inhibitor SPP-86 (Bhallamudi et al., 2021 [\[4\]](#)) and Src-type kinase inhibitor PP2 (Morita et al., 2006 [\[5\]](#)) were used to further investigate whether GDNF upregulated the expression of claudin-5 and VE-cadherin in hCMEC/D3 cells by activating the GDNF-GFR α 1-RET signaling pathway. Both SPP-86 and PP2 markedly attenuated claudin-5 and VE-cadherin expression induced by GDNF (Figure 2K [\[3\]](#)) and US-CM (Figure 2L [\[3\]](#)).

The contributions of GDNF-induced claudin-5 and VE-cadherin expression to TEER and permeability were investigated using hCMEC/D3 cells mono-culture BBB model. GDNF significantly increased TEER values (Figure 2M [\[3\]](#), 2O) and decreased the permeability of fluorescein and FITC-Dex (Figure 2N [\[3\]](#), 2P), which were almost abolished by SPP-86 or PP2. Furthermore, treatment with SPP-86 or PP2 completely reversed the increased TEER values (Figure 2Q [\[3\]](#), 2S) and decreased permeability of fluorescein and FITC-Dex (Figure. 2R, 2T) in the triple co-culture BBB model. These results indicate that neurons but also astrocytes upregulate claudin-5 and VE-cadherin expression in hCMEC/D3 cells by secreting GDNF. Subsequently, GDNF induces claudin-5 and VE-cadherin expression by activating GDNF-GFR α 1-RET signaling.

GDNF induced claudin-5 and VE-cadherin expression of hCMEC/D3 by activating the PI3K/AKT and MAPK/ERK pathways

GDNF exerts its biological activities by activating several signaling pathways, including the phosphatidylinositol-3-kinase (PI3K)/ protein kinase B (AKT), mitogen-activated protein kinase (MAPK)/ extracellular regulated kinase (ERK), MAPK/ c-Jun N-terminal kinase (JNK), and MAPK/ p38 pathways (Fielder et al., 2018 [\[3\]](#)). The effects of the PI3K/AKT, MAPK/ERK, MAPK/JNK, and MAPK/p38 pathway inhibitors LY294002 (Figure 3A [\[3\]](#)), U0126 (Figure 3B [\[3\]](#)), SP600125 (Figure 3C [\[3\]](#)), and SB203580 (Figure 3D [\[3\]](#)), respectively, on GDNF-induced claudin-5 and VE-cadherin expression in hCMEC/D3 cells were investigated. GDNF increased claudin-5 and VE-cadherin expression, accompanied by the phosphorylation of AKT (p-AKT) and ERK (p-ERK). However, it did not stimulate the phosphorylation of JNK and p38. SPP-86, LY29002, and U0126 significantly suppressed GDNF-induced claudin-5 and VE-cadherin expression, while SP600125 and SB203580 had almost no effect on GDNF-induced claudin-5 and VE-cadherin expression. GDNF-induced phosphorylation of AKT and ERK was also markedly attenuated by the anti-GDNF antibody (Figure 3E [\[3\]](#)). Similarly, US-CM remarkably upregulated the expression of claudin-5, VE-cadherin, p-AKT, and p-ERK, which were also markedly reversed by SPP-86, LY29002, U0126, or anti-GDNF antibody (Figure 3F-3J [\[3\]](#)). These findings indicate that GDNF induces the expression of claudin-5 and VE-cadherin in hCMEC/D3 cells by activating both the PI3K/AKT and MAPK/ERK pathways.

GDNF upregulated the claudin-5 expression in hCMEC/D3 cells by activating the PI3K/AKT/FOXO1 pathway

Claudin-5 is negatively regulated by the transcriptional repressor forkhead box O1 (FOXO1) (Beard et al., 2020 [\[6\]](#)). FOXO1 is also an important target of PI3K/AKT signaling. FOXO1 phosphorylation results in FOXO1 accumulation in the cytoplasm (Zhang et al., 2011 [\[7\]](#)) and lowers its level in the nucleus. Here, we investigated whether GDNF-induced claudin-5 expression is involved in FOXO1 nuclear exclusion. As shown in Figure 4A [\[3\]](#), both GDNF and US-CM significantly enhanced FOXO1 phosphorylation (p-FOXO1). Similarly, GDNF and US-CM increased the levels of phosphorylated and unphosphorylated FOXO1 in the cytoplasm and decreased the levels of nuclear FOXO1 (Figure 4B [\[3\]](#)). Whether FOXO1 was involved in the GDNF-induced regulation of claudin-5 and VE-cadherin expression was investigated in hCMEC/D3 cells transfected with FOXO1 siRNA. Silencing FOXO1 significantly decreased FOXO1 levels in both whole-cell lysates and nucleus of hCMEC/D3 cells (Figure 4C [\[3\]](#)), demonstrating FOXO1 silencing efficacy. Consistent with our expectation, silencing FOXO1 upregulated the expression of claudin-5 rather than VE-cadherin expression




Figure 3.

GDNF induced claudin-5 and VE-cadherin expression in hCMEC/D3 cells by activating the PI3K/AKT and MAPK/ERK signaling.

(A): Effects of 3 μ M LY294002 (LY) on the levels of claudin-5, VE-cadherin, and p-AKT/AKT in hCMEC/D3 cells stimulated by 200 pg/mL GDNF. (B): Effects of 2 μ M U0126 (U0) on the levels of claudin-5, VE-cadherin, and p-ERK/ERK in hCMEC/D3 cells stimulated by 200 pg/mL GDNF. (C): Effects of 5 μ M SP600125 (SP) on the levels of claudin-5, VE-cadherin, and p-JNK/JNK in hCMEC/D3 cells stimulated by 200 pg/mL GDNF. (D): Effects of 2 μ M SB203580 (SB) on the levels of claudin-5, VE-cadherin, and p-p38/p38 in hCMEC/D3 cells stimulated by 200 pg/mL GDNF. (E): Effects of anti-GDNF antibody on the GDNF-induced p-AKT/AKT and p-ERK/ERK ratios. (F): Effects of 3 μ M LY on the levels of claudin-5, VE-cadherin, and p-AKT/AKT in hCMEC/D3 cells stimulated by US-CM. (G): Effects of 2 μ M U0 on the levels of claudin-5, VE-cadherin, and p-ERK/ERK in hCMEC/D3 cells stimulated by US-CM. (H): Effects of 5 μ M SP on the levels of claudin-5, VE-cadherin, and p-JNK/JNK in hCMEC/D3 cells stimulated by US-CM. (I): Effects of 2 μ M SB on the levels of claudin-5, VE-cadherin, and p-p38/p38 in hCMEC/D3 cells stimulated by 2 US-CM. (J): Effects of anti-GDNF antibody on the US-CM-induced p-AKT/AKT and p-ERK/ERK ratios. The above data are shown as the mean $\pm$ S.E.M.; $n=4$. Statistical significance was determined using one-way ANOVA with Fisher's LSD test or Welch's ANOVA test.

Figure 3—source data1 (Raw and annotated data files for western blot)

Figure 3—source data2 (Summary of original data and analysis)

(Figure 4D [↗](#)). In contrast, high levels of FOXO1 were observed in both whole-cell lysates and nucleus of hCMEC/D3 cells that were transfected with plasmids containing *FOXO1*. Meanwhile, FOXO1 overexpression resulted in a decrease in both basal and GDNF-induced claudin-5 expression (Figure 4E [↗](#)), consistent with the known role of FOXO1 on claudin-5 expression.

Several reports have demonstrated that nuclear localization of FOXO1 is modulated by multiple pathways, including the PI3K/AKT (Tang et al., 1999 [↗](#)) and MAPK/ERK (Asada et al., 2007 [↗](#)) pathways. The effects of LY294002 and U0126 on GDNF-induced FOXO1 phosphorylation were measured in hCMEC/D3 cells. LY294002, but not U0126, significantly reversed the GDNF-induced alterations in total p-FOXO1, cytoplasmic p-FOXO1, cytoplasmic FOXO1, and nuclear FOXO1 (Figure 4F [↗](#)). Furthermore, LY294002 reversed the upregulation of claudin-5 expression induced by *FOXO1* siRNA and GDNF (Figure 4G [↗](#)).

GDNF upregulated VE-cadherin expression in hCMEC/D3 cells by activating the PI3K/AKT/ETS1 and MAPK/ERK/ETS1 signaling pathways

E26 oncogene homolog 1 (ETS1) is a transcription factor that binds to the ETS-binding site located in the proximal region of the VE-cadherin promoter, hence regulating the expression of VE-cadherin (Lelièvre et al., 2000 [↗](#); Luo et al., 2022 [↗](#)). Activation of the PI3K/AKT (He et al., 2023 [↗](#); Hui et al., 2018 [↗](#)) and MAPK/ERK (Watanabe et al., 2004 [↗](#)) pathways was reported to upregulate the expression of ETS1. The previous results showed that GDNF induced VE-cadherin and claudin-5 expression in hCMEC/D3 cells by activating the PI3K/AKT and MAPK/ERK pathways. Therefore, we hypothesized that GDNF modulated ETS1 levels to promote VE-Cadherin and claudin-5 expression through PI3K/AKT and MAPK/ERK signaling pathways. Both US-CM and GDNF significantly increased total (Figure 5A [↗](#)) and nuclear ETS1 expression (Figure 5B [↗](#)). LY294002 and U0126 markedly attenuated GDNF-induced total (Figure 5C [↗](#)) and nuclear ETS1 expression (Figure 5D [↗](#)). To further confirm the involvement of the PI3K/AKT/ETS1 and MAPK/ERK/ETS1 pathways in GDNF-induced VE-cadherin expression, ETS1 in hCMEC/D3 cells was knocked down using *ETS1* siRNA. *ETS1* silencing saliently declined the expression levels of total (Figure 5E [↗](#)) and nuclear (Figure 5F [↗](#)) ETS1 in hCMEC/D3 cells, demonstrating silencing efficacy. In *ETS1* silencing hCMEC/D3 cells, GDNF no longer induced the expression of total (Figure 5E [↗](#)) and nuclear (Figure 5F [↗](#)) ETS1. Moreover, *ETS1* silencing substantially downregulated VE-cadherin expression and attenuated GDNF-induced VE-cadherin expression (Figure 5G [↗](#) and 5H), while having minimal impact on both basal and GDNF-induced claudin-5 expression (Figure 5G [↗](#) and 5I [↗](#)).

Brain GDNF deficiency increased BBB permeability partly due to the impairment of claudin-5 and VE-cadherin expression

To further demonstrate the positive effects of GDNF on BBB maintenance, GDNF in mice brains was knocked down via intracerebroventricular injection of AAV-PHP.eB packaged with *Gdnf* short hairpin RNA (shRNA) (Figure 6A [↗](#)). Knockdown efficiency was confirmed through Western blotting (Figure 6B [↗](#)). Consistent with *in vitro* results, GDNF knockdown greatly downregulated the expression of claudin-5 and VE-cadherin in the brains of mice (Figure 6B [↗](#)). The integrity of BBB was assessed by examining the brain distributions of fluorescein and FITC-Dex. The results showed that specifically knocking down brain GDNF little affected plasma levels of fluorescein (Figure 6C [↗](#)) and FITC-Dex (Figure 6F [↗](#)), but significantly elevated the concentrations of fluorescein (Figure 6D [↗](#)) and FITC-Dex (Figure 6G [↗](#)) in the brains, leading to notable increases in the brain-to-plasma concentration ratios of two probes (Figure 6E [↗](#) and 6H). These alterations were consistent with the decline in claudin-5 and VE-cadherin expression. In addition, significant reductions in the levels of p-AKT (Figure 6I [↗](#)), p-ERK (Figure 6J [↗](#)), p-FOXO1 (Figure 6K [↗](#)), and ETS1 expression (Figure 6L [↗](#)) were observed in the brains of *Gdnf* knockdown mice.




Figure 4.

GDNF induced the claudin-5 expression in hCMEC/D3 cells by activating the PI3K/AKT/FOXO1 pathway.

(A and B): Effects of US-CM and GDNF on the phosphorylated FOXO1 (p-FOXO1)/FOXO1 ratio, total FOXO1 expression (A), cytoplasmic p-FOXO1, cytoplasmic FOXO1, and nuclear FOXO1 expression (B). (C and D): The expression levels of total and nuclear FOXO1 (C), claudin-5, and VE-cadherin (D) in hCMEC/D3 cells transfected with *FOXO1* siRNA (siFOXO1). NC: negative control. (E): Effects of *FOXO1* overexpression (FOXO1-OE) and GDNF on the expression levels of claudin-5, total FOXO1, and nuclear FOXO1. FOXO1-NC: negative control plasmids. (F): Effects of LY and U0 on GDNF-induced alterations of total p-FOXO1/FOXO1 ratio, cytoplasmic p-FOXO1, cytoplasmic FOXO1, and nuclear FOXO1 expression. (G): Effects of LY on the claudin-5 expression upregulated by siFOXO1. The above data are shown as the mean $\pm$ S.E.M.; n=4. Statistical significance was determined using one-way ANOVA with Fisher's LSD test.

Figure 4—source data1 (Raw and annotated data files for western blot)

Figure 4—source data2 (Summary of original data and analysis)




Figure 5.

GDNF induced VE-cadherin expression in hCMEC/D3 cells by activating the PI3K/AKT/ETS1 and MAPK/ERK/ETS1 pathways.

(**A** and **B**): Effects of US-CM and GDNF on total (**A**) and nuclear (**B**) ETS1 expression. (**C** and **D**): Effects of LY and UO on 200 pg/mL GDNF-induced total (**C**) and nuclear (**D**) ETS1 expression. (**E** and **F**): Expression levels of total (**E**) and the nuclear ETS1 (**F**) in hCMEC/D3 cells after knocking down *ETS1* with siRNA (siETS1). (**G-I**): Effects of GDNF and siETS1 on the expression of VE-cadherin and claudin-5. The above data are shown as the mean $\pm$ S.E.M.; $n=4$. Statistical significance was determined using one-way ANOVA with Fisher's LSD test.

Figure 5—source data1 (Raw and annotated data files for western blot)

Figure 5—source data2 (Summary of original data and analysis)




Figure 6.

The deficiency of brain GDNF in mice increased the permeability of BBB and reduced claudin-5 and VE-cadherin expression in mice brains.

(A): Experimental configuration of AAV-GFP (shNC) or AAV-shGdnf (shGdnf) intracerebroventricular injection. (B): Effects of brain-specific *Gdnf* silencing on the expression levels of GDNF, claudin-5, and VE-cadherin in the brains. (C-E): Effects of brain-specific *Gdnf* silencing on NaF levels in plasma (C), brain (D), and the ratio of brain to plasma (E). (F-H): Effects of brain-specific *Gdnf* silencing on FITC-Dex levels in plasma (F), brain (G), and the ratio of brain to plasma (H). (I-K): The expression ratios of p-AKT/AKT (I), p-ERK/ERK (J), and p-FOXO1/FOXO1 (K) in the brains of *Gdnf* silencing mice. (L): The expression level of ETS1 in the brains of *Gdnf* silencing mice. The above data are shown as the mean $\pm$ S.E.M.; n=6. Statistical significance was determined using the unpaired t-test or unpaired t-test with Welch's correction.

Figure 6—source data1 (Raw and annotated data files for western blot)

Figure 6—source data2 (Summary of original data and analysis)

The triple co-culture BBB model better predicted the permeabilities of drugs across BBB

This study demonstrated that the developed triple co-culture BBB model displays better BBB characteristics compared with the hCMEC/D3 mono-culture BBB model. In this study, 18 drugs were utilized to further investigate the superiority of the developed triple co-culture BBB model over the hCMEC/D3 mono-culture BBB model. The P_{app} of the 18 drugs from the apical to the basolateral side based on the hCMEC/D3 mono-culture ($P_{app, Mono}$) and triple co-culture ($P_{app, Triple}$) BBB models are measured and listed in [Table 5](#). The results showed that $P_{app, Triple}$ values of all tested drugs were lower than the $P_{app, Mono}$ values. Significant differences were observed in 14 out of 18 drugs. The predicted permeability coefficient-surface area product values (PS) of the tested drugs were respectively calculated based on their $P_{app, Mono}$ values ($PS_{Pre, Mono}$), and $P_{app, Triple}$ values ($PS_{Pre, Triple}$). The predicted PS values were further compared to their corresponding observations (PS_{Obs}). The results showed that the predictive accuracy of $PS_{Pre, Triple}$ was superior to $PS_{Pre, Mono}$. Except for verapamil, amitriptyline, fluoxetine, and clozapine, the predicted $PS_{Pre, Triple}$ values of the other 14 drugs in the triple co-culture BBB models were within the 0.5–2 folds of PS_{Obs} ([Figure 7B](#)). However, in the hCMEC/D3 mono-culture BBB model, only 7 predicted $PS_{Pre, Mono}$ values were within the 0.5–2 folds range of their observations ([Figure 7A](#)).




Figure 7.

In vitro/in vivo correlation (IVIVC) assay of BBB permeability.

(A): The comparison of the estimated permeability coefficient-surface area product (PS_{Mono}) recalculated from $P_{app, Mono}$ with the observed *in vivo* PS values (PS_{Obs}). (B): The comparison of the estimated permeability coefficient-surface area product (PS_{Triple}) recalculated from $P_{app, Triple}$ with the observed *in vivo* PS values (PS_{Obs}). The solid line represents a perfect prediction, and the dashed lines represent the 0.5–2 folds of their observations. The PS_{Obs} values were determined by *in situ* brain perfusion in rodents, which were collected from the literature.

Discussion

The main findings of the study were to successfully develop an *in vitro* triple co-culture BBB model consisting of hCMEC/D3, U251, and SH-SY5Y cells and to confirm the involvement of neurons in BBB maintenance as well as the possible mechanisms. Co-culture with U251 and/or SH-SY5Y cells

markedly promoted the TEER of hCMEC/D3 cells and reduced the leakage of fluorescein and FITC-Dex due to the upregulation of claudin-5 and VE-cadherin expression.

The roles of claudin-5 and VE-cadherin in the maintenance of BBB function have been demonstrated (Dejana et al., 2008 [↗](#); Hashimoto et al., 2023 [↗](#); Li W. et al., 2018; Ohtsuki et al., 2007 [↗](#)). It was reported that *Cld-5*-deficient mice exhibited BBB impairment, allowing the transport of small molecules (<800 D) across BBB (Nitta et al., 2003 [↗](#)). In contrast, claudin-5 overexpression significantly restricted the permeability of inulin across the conditionally immortalized rat brain capillary endothelial cell monolayer (Ohtsuki et al., 2007 [↗](#)). Moreover, the claudin-5 expression in hCMEC/D3 is much lower than in human brain microvessels (Eigenmann et al., 2013 [↗](#); Ohtsuki et al., 2013 [↗](#); Weksler et al., 2013 [↗](#)), which may cause the low TEER values. VE-cadherin, a major member of the cadherin family in endothelial cells, is also required for BBB integrity (Dejana et al., 2008 [↗](#); Li W. et al., 2018). The absence of VE-cadherin resulted in faulty cell-to-cell junctions and disrupted distribution of ZO-1 (Sauteur et al., 2017 [↗](#); Tunggal et al., 2005 [↗](#)). Strongly negative correlations between P_{app} values of fluorescein or FITC-Dex and the expression of claudin-5 or VE-cadherin further demonstrated the importance of claudin-5 and VE-cadherin in BBB integrity.

Neurons and astrocytes, as important components of the NVU, may be involved in the formation and maintenance of BBB function. Several reports showed that co-culture with astrocytes CC-2565 or SC-1810 enhanced the TEER of hCMEC/D3 cells (Hatherell et al., 2011 [↗](#)), whereas co-culture of RBE4.B cells with primary rat neurons and astrocytes showed a lower permeability of [³H] sucrose than mono-culture of RBE4.B cells (Schiera Gabriella. et al., 2005). Similarly, co-cultured with hCMEC/D3 cells and 1321N1 (astrocytes) or 1321N1+SH-SY5Y cells possess higher TEER values and lower permeability to Lucifer yellow than hCMEC/D3 alone. Compared with double co-culture of hCMEC/D3 and 321N1 cells, the triple co-culture of hCMEC/D3, 1321N1, and SH-SY5Y cells showed higher TEER values (Barberio et al., 2022 [↗](#)). Our study also demonstrated that co-culture with U251 and/or SH-SY5Y cells significantly lowered BBB permeability and upregulated VE-cadherin or claudin-5 expression in hCMEC/D3 cells.

Next, we focused on the molecular mechanisms by which neurons and possibly astrocytes upregulated the expression of VE-cadherin and claudin-5 in BMECs. Co-culture with SH-SY5Y cells significantly upregulated claudin-5 and VE-cadherin expression in hCMEC/D3 cells. In the double co-culture with SH-SY5Y cells or triple co-culture BBB models, hCMEC/D3 cells were not in direct contact with SH-SY5Y cells, indicating that the interaction between SH-SY5Y and hCMEC/D3 cells depended on the release of some active compounds. It was consistent with the above deduction that the S-CM also markedly induced claudin-5 and VE-cadherin expression. Different from S-CM, U-CM mainly upregulated the expression of VE-cadherin and just had a slight impact on claudin-5 expression. In general, neurons but also astrocytes secrete some neurotrophic factors (Lonka-Nevalaita et al., 2010 [↗](#); Sweeney et al., 2019 [↗](#)) that could contribute to the maintenance of structural stability of BBB. High levels of bFGF, GDNF, IGF-1, and TGF- β were detected in U-CM, S-CM, and US-CM. Further research showed that only GDNF induced VE-cadherin and claudin-5 expression in hCMEC/D3 cells in a concentration-dependent manner, and the anti-GDNF antibody attenuated claudin-5 and VE-cadherin expression induced by US-CM or GDNF. These results indicated that neurons upregulated claudin-5 and VE-cadherin expression through GDNF secretion. Levels of GDNF in S-CM were higher than those in U-CM, which seemed to partly explain why S-CM has a stronger promoting effect on claudin-5 than U-CM. The roles of GDNF in BBB maintenance and the regulation of claudin-5 and VE-cadherin expression were further confirmed using brain-specific *Gdnf* knockdown C57BL/6J mice. Consistent with our *in vitro* results, brain-specific *Gdnf* silencing greatly increased BBB penetration of fluorescein and FITC-Dex, accompanied by the downregulation of claudin-5 and VE-cadherin expression.

Generally, GDNF activates several signal transduction pathways, such as the PI3K/AKT and MAPK signaling pathways (Fielder et al., 2018 [↗](#)) by forming a heterohexameric complex with two GFR α molecules and RET receptors. It was also reported that GDNF improved BBB barrier function due to the activation of MAPK/ERK1 (Dong et al., 2018) and PI3K/AKT (Liu D. et al., 2022) signaling. Consistent with previous reports, we found that signaling inhibitors SPP-86, PP2, LY294002, and U0126 markedly attenuated US-CM- and GDNF-induced claudin-5 and VE-cadherin expression in hCMEC/D3 cells, inferring that GDNF promoted claudin-5 and VE-cadherin expression via activating both the PI3K/AKT and MAPK/ERK pathways.

Signal transduction pathways control gene expression by modifying the function of nuclear transcription factors. The nuclear accumulation of FOXO1 negatively regulates claudin-5 expression (Beard et al., 2020 [↗](#); Taddei et al., 2008 [↗](#)). FOXO1 is an important target of the PI3K/AKT signaling axis (Zhang et al., 2011 [↗](#)), and AKT-induced phosphorylation of FOXO1 results in cytoplasmic FOXO1 accumulation and decreases nuclear FOXO1 accumulation (Zhang et al., 2011 [↗](#)). From these results, we inferred that GDNF-induced claudin-5 expression in hCMEC/D3 cells may be involved in the activation of PI3K/AKT/FOXO1 pathway. Similarly, both US-CM and GDNF increased cytoplasmic p-FOXO1 and decreased nuclear FOXO1 in hCMEC/D3 cells, which was reversed by LY29002 rather than U0126. Roles of FOXO1 in GDNF-induced claudin-5 were verified through silencing and overexpressing *FOXO1* in hCMEC/D3 cells. *FOXO1* silencing enhanced claudin-5 but not VE-cadherin expression, accompanied by a decline in total and nuclear FOXO1. In contrast, *FOXO1* overexpression significantly decreased claudin-5 expression. In hCMEC/D3 cells overexpressing *FOXO1*, GDNF lost its promotion effect on claudin-5 expression. It was noticed that U0126 attenuated the GDNF-induced upregulation of claudin-5 but had minimal impact on GDNF-mediated FOXO1 phosphorylation and the decline of nuclear FOXO1. In Sertoli cells, it was found that testosterone stimulated claudin-5 expression by activating the RAS/RAF/ERK/CREB pathway (Bulldan et al., 2016 [↗](#)). The transcriptional regulation of claudin-5 by CREB was confirmed in bEnd.3 (mouse brain endothelial cell). CREB overexpression significantly increased both gene and protein expression of claudin 5. In contrast, depletion of CREB decreased claudin-5 expression in gene and protein levels (Li Y. et al., 2022). However, another report showed that in human lung microvascular endothelial cells, U0126 attenuated phosphorylation of ERK and lipopolysaccharide-stimulated claudin-5 damage, indicating activation of MAPK/ERK pathway impaired rather than promoted claudin-5 expression (Liu Y. et al., 2019). Thus, the real mechanisms that GDNF-induced activation of the RET/MAPK/ERK pathway promotes claudin-5 expression need further investigation.

ETS1 is a member of the ETS family that plays an important role in cell adhesion, migration, and blood vessel information. ETS1 binds to an ETS-binding site located in the proximal region of the VE-cadherin promoter, controlling VE-cadherin expression (Lelièvre et al., 2001 [↗](#)). Several studies have demonstrated the role of ETS1 in the regulation of VE-cadherin expression. For example, IFN- γ and TNF- α impaired BBB integrity by decreasing ETS1-induced VE-cadherin expression (Luo et al., 2022 [↗](#)). *ETS1* silencing reduced VE-cadherin expression in umbilical vein endothelial cells (Colas-Algora et al., 2020 [↗](#)). In contrast, *ETS1* overexpression induced VE-cadherin expression in mouse brain capillary endothelial cells and fibroblasts (Lelièvre et al., 2000 [↗](#)). As expected, *ETS1* silencing resulted in a decrease in the expression of VE-cadherin in hCMEC/D3 cells, but claudin-5 expression remained unaffected. Additionally, *ETS1* silencing removed the inductive effect of GDNF on VE-cadherin expression while unaffected the upregulation of claudin-5 induced by GDNF. The activation of the PI3K/AKT (He et al., 2023 [↗](#); Hui et al., 2018 [↗](#)) and MAPK/ERK (Watanabe et al., 2004 [↗](#)) pathways promotes ETS1 expression. Our findings also demonstrated that US-CM and GDNF significantly increased total and nuclear ETS1 levels, which were eliminated by signaling inhibitors LY294002 and U0126. Both our results and previous research provide evidence that GDNF upregulates ETS1 expression via the activation of PI3K/AKT and MAPK/ERK signaling to promote VE-cadherin expression.

The present study showed that the characteristics of the *in vitro* triple co-culture BBB model were superior to those of the hCMEC/D3 mono-culture BBB model. The hCMEC/D3 mono-culture and triple co-culture BBB models were used to try to predict the *PS* values of 18 drugs by comparing them with their observations. The prediction success rate (14/18) of triple co-culture BBB model was greater than that of hCMEC/D3 mono-culture BBB model (7/18). However, poor predictions were observed for verapamil, amitriptyline, fluoxetine, and clozapine, which may partly be due to inaccuracies in their $f_{u, brain}$ values. These four drugs are high protein-binding drugs. Due to the methodological discordance and the limitations of the historic devices for these drugs, the $f_{u, brain} \leq 0.01$ maybe with low confidence and accuracy (Bowman et al., 2018; Di et al., 2017). For example, $f_{u, brain}$ values of amitriptyline from different reports display large differences (0.002 (Sanchez-Dengra et al., 2021), 0.0071 (Weber et al., 2013)). The same large differences of $f_{u, brain}$ values were also found in fluoxetine (0.0023 (Maurer et al., 2005), 0.004 (Summerfield et al., 2007), 0.00094 (Liu X. et al., 2005)), and clozapine (0.0056 (Bhyrapuneni et al., 2018), 0.014 (Cremers et al., 2012), 0.0094 (Summerfield et al., 2007)).

In summary, the developed triple co-culture BBB model outperformed the mono-culture or double co-culture BBB models, mainly attributing to the fact that neurons and possibly astrocytes upregulated claudin-5 and VE-cadherin expression by secreting GDNF. GDNF upregulates claudin-5 expression by activating the PI3K/AKT/FOXO1 and MAPK/ERK pathways, and the activation of PI3K/AKT and MAPK/ERK signaling also promotes ETS1 to modify VE-cadherin expression. Additionally, the developed *in vitro* triple co-culture BBB model accurately predicted the *in vivo* BBB permeability of CNS drugs. This suggests the potential of our triple co-culture BBB model for utilization in CNS candidates screening during the drug development process (Figure 8).




Figure 8.

The mechanism of neurons and astrocytes induced the integrity of brain endothelial cells.

Neurons but also astrocytes trigger the activation of PI3K/AKT and MAPK/ERK pathways in brain endothelial cells by GDNF secretion, which in turn regulates transcription factors of claudin-5 (FOXO1) and VE-cadherin (ETS1) to promote claudin-5 and VE-cadherin expression and leads to the enhancement of BBB integrity. Meanwhile, with the increase in barrier integrity, the *in vitro* BBB model also obtained a stronger *in vivo* correlation.

Materials and Methods

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Antibody	β -actin (mouse monoclonal)	Proteintech	66009 RRID:AB_2883475	1:10000
Antibody	GAPDH (mouse monoclonal)	Absin	Abs830030ss RRID:AB_2811228	1:50000
Antibody	β -Tubulin (mouse monoclonal)	Fdbio Science	FD0064 RRID: AB_3076327	1:10000
Antibody	Lamin B (mouse monoclonal)	Proteintech	66095 RRID:AB_2721256	1:10000
Antibody	Claudin-5 (rabbit polyclonal)	Wanleibio	WL03731 RRID: AB_3076320	1:1000
Antibody	Occludin (rabbit polyclonal)	Wanleibio	WL01996 RRID:AB_3076325	1:500
Antibody	ZO-1 (mouse polyclonal)	Proteintech	21773-1-AP RRID:AB_10733242	1:5000
Antibody	VE-cadherin (rabbit polyclonal)	Wanleibio	WL02033 RRID:AB_3076321	1:1000
Antibody	β -catenin (rabbit polyclonal)	Wanleibio	WL0962a RRID: AB_3076323	1:5000
Antibody	BCPR (rabbit polyclonal)	CST	4477S RRID:AB_10544928	1:1000
Antibody	P-gp (rabbit	CST	13978S RRID:AB_2798357	1:1500

	monoclonal)			
Antibody	p-AKT (mouse monoclonal)	Huaan Biotechnology	ET1607 RRID:AB_2940863	1:2000
Antibody	AKT (mouse monoclonal)	Huaan Biotechnology	ET1609 RRID:AB_3069857	1:2000
Antibody	p-ERK (rabbit polyclonal)	Proteintech	28733-1-AP RRID:AB_2881202	1:1000
Antibody	ERK (rabbit polyclonal)	Proteintech	11257-1-AP RRID:AB_2139822	1:1000
Antibody	p-p38 (rabbit monoclonal)	CST	4511S RRID:AB_10890701	1:250
Antibody	p38 (rabbit monoclonal)	CST	8690S RRID:AB_10999090	1:250
Antibody	p-JNK (rabbit polyclonal)	Wanleibio	WL01813 RRID:AB_2910628	1:1000
Antibody	JNK (rabbit polyclonal)	Wanleibio	WL01295 RRID:AB_3064853	1:1000
Antibody	FOXO1 (rabbit polyclonal)	Proteintech	18592 RRID:AB_2934932	1:1000
Antibody	p-FOXO1 (rabbit polyclonal)	Wanleibio	WL03634 RRID: AB_3076326	1:1000
Antibody	ETS1 (mouse monoclonal)	Santa Cruz	sc-55581 RRID:AB_831289	1:500
Antibody	ETS1 (mouse monoclonal)	Proteintech	66598 RRID:AB_2881958	1:3000
cell line (<i>Homo sapiens</i>)	hCMEC/D3 cells	JENNIO Biological Technology	Cat#JNO-H0520	Authenticated (STR profiling)

cell line (<i>Homo sapiens</i>)	U251 cells	Cellcook Biological Technology	Cat#CC1701	Authenticated (STR profiling)
cell line (<i>Homo sapiens</i>)	SH-SY5Y cells	Cellcook Biological Technology	Cat#CC2101	Authenticated (STR profiling)
Software, algorithm	GraphPad Prism	Version 8.0.2	RRID:SCR_002798	
Software, algorithm	BioTek Cytation 5 Cell Imaging Multi-Mode Reader	BioTek Cytation 5	RRID:SCR_019732	
Software, algorithm	QuantStudio 3 Real Time PCR System	QuantStudio 3	RRID:SCR_018712	
commercial assay or kit	GDNF-Elisa kit	R&D system	Cat#212-GD	
commercial assay or kit	bFGF-Elisa kit	Elabscience	Cat#E-EL-H6042	
commercial assay or kit	IGF-1-Elisa kit	Elabscience	Cat#E-EL-H0086	
commercial assay or kit	TGF- β -Elisa kit	Elabscience	Cat#E-EL-0162	
peptide, recombinant protein	GDNF	R&D system	Cat#212-GD	
peptide, recombinant protein	bFGF	MedChemExpress	Cat#HY-P7331	
peptide, recombinant protein	IGF-1	MedChemExpress	Cat#HY-P70783	
peptide, recombinant	TGF- β	MedChemExpress	Cat#HY-P70543	

protein				
chemical compound, drug	SPP-86	MedChemExpress	Cat#HY-110193	
chemical compound, drug	PP2	MedChemExpress	Cat#HY-13805	
chemical compound, drug	LY294002	MedChemExpress	Cat#HY-10108	
chemical compound, drug	U0126	MedChemExpress	Cat#S1102	
chemical compound, drug	SP600125	Selleck	Cat#HY-12041	
chemical compound, drug	SB203580	MedChemExpress	Cat#HY-10256	

Key Resources Table

Cell culture

Rat brain microvascular endothelial cells were isolated from Sprague-Dawley rats (male, 7–10 days old, Sino-British Sippr/BK Laboratory Animal Ltd, Shanghai, China) as the described method (Ji et al., 2013; Li M. S. et al., 2013) and cultured in Dulbecco's Modified Eagle Media (DMEM)/F12 (#12500-039, Gibco, Carlsbad, CA, USA) containing 10% fetal bovine serum (FBS) (#10100147C, Gibco, Carlsbad, CA, USA) and 62.5 µg/mL penicillin and 100 µg/mL streptomycin (SunShine Biotechnology Co., Ltd. Nanjing, China). Then, hCMEC/D3, U251, and SH-SY5Y cells were cultured in DMEM/F12 containing 10% FBS, 62.5 µg/mL penicillin and 100 µg/mL streptomycin.

Establishment of the triple co-culture model

The establishment process of the triple co-culture model is illustrated in **Figure 1A**. SH-SY5Y cells were seeded at a density of 4.5×10^4 cells/cm² in plates and differentiated with 10 µM retinoic acid (Sigma-Aldrich, St. Louis, MO, USA) for 72 h. The differentiated SH-SY5Y cells were cultured in the fresh DMEM/F12 medium containing 10% FBS. U251 cells were seeded at 2×10^4 cells/cm² on the bottom of Transwell inserts (0.4 µm pore size, SPL Life Sciences, Pocheon, Korea) coated with rat-tail collagen (Corning Inc., Corning, NY, USA). Next, the inserts were suspended in plate wells containing the culture medium after 5 h of incubation. After 24 h of incubation, hCMEC/D3 cells were seeded on the apical side of the inserts at 3×10^4 cells/cm² and cultured for another 48 h. Then, the inserts seeded with U251 and hCMEC/D3 cells were suspended in plates seeded with differentiated SH-SY5Y cells and co-cultured for another 6 days. The culture medium was replaced every 24 h.

Three other types of BBB *in vitro* models, namely the hCMEC/D3 cell mono-culture model, the double co-culture with SH-SY5Y cells BBB model, and the double co-culture with U251 cells BBB model were also established (**Figure 1B**). TEER was periodically measured using a Millicell-ERS (MERS00002) instrument (Millipore, Billerica MA, USA) to monitor cell confluence and development of tight junctions.

***In vitro* BBB permeability study**

The culture medium was removed from the apical and basolateral sides of the inserts and washed twice with preheated Hank's balanced salt solution (HBSS). Fresh HBSS was then added to both the apical and basolateral chambers. After 15 min of preincubation, HBSS in the apical and basolateral chambers was replaced with HBSS containing FITC-dextran 3–5 kDa (FITC-Dex) (Sigma-Aldrich, St. Louis, MO, USA), fluorescein sodium (Sigma-Aldrich, St. Louis, MO, USA), or other tested agents and blank HBSS, respectively. Next, 200 µL aliquots were collected from the basolateral chamber after 30 min of incubation at 37 °C. The concentrations of the tested agents in the basolateral chamber were measured.

The apparent permeability coefficients (P_{app} , cm/s) of the tested agents across the *in vitro* BBB model were calculated using the equation (Tavelin et al., 2002):

$$P_{app} = (Q/1800)/(S \times C_0) \quad (1)$$

where S is the surface area of the insert membrane (0.33 cm² for 6.5 mm inserts, 4.46 cm² for 24 mm inserts), Q is the transported amount of the tested agents transported from the donor chamber to the receiver chamber for 30 min (1800 s), and C_0 is the initial concentration of the tested agents in the donor chamber.

The quantification methods of prazosin, verapamil, lamotrigine

Prazosin, verapamil, and lamotrigine (Aladdin, Shanghai, China) were analyzed by high-performance liquid chromatography (Shimadzu, Kyoto, Japan) with YMC-Triart C18 column (5 µm, 150 × 4.6 mm, YMC America Inc., Allentown, PA, USA). Prazosin and verapamil were detected using the RF-20A fluorescence detector. Lamotrigine was detected using the SPD-20A ultraviolet detector. Samples were centrifugated at 12000 rpm for 10 mins, then 150 µL supernatant was taken and used for analysis. The run temperature was set at 40 °C, the injection volume was 20 µL and the flow rate was 1 mL/min. Initial concentrations in donor chamber and other chromatographic conditions of drugs are summarized in **Table 1**.

The quantification methods of clozapine, venlafaxine, bupropion, amantadine, carbamazepine, fluoxetine, amitriptyline, gabapentin, midazolam, risperidone, olanzapine, mirtazapine, metoclopramide, doxepin, donepezil

All compounds were purchased from Aladdin (Shanghai, China). Except for prazosin, verapamil, and lamotrigine, the other compounds were analyzed by using liquid chromatography-mass spectrometry (Shimadzu, Kyoto, Japan) with YMC-Triart C18 column (5 µm, 150 × 2.0 mm, YMC America Inc., Allentown, PA, USA). Each sample was mixed with 10 µL internal standard. Then 1 mL extraction was added to each sample. The samples were vortex vibrated on the oscillator for 10 min, and then centrifuged at 4 °C and 12000 rpm for 10 mins. The supernatant solvent was evaporated with nitrogen flow, then redissolved with 100 µL 40 % (v/v) acetonitrile, and centrifuged at 4 °C and 15000 rpm for 10 min. The supernatants were injected into LC-MS for analysis. The injection volume of each sample was 5 µL. The mass charge ratio, extraction, and initial concentrations in donor chamber of drugs were summarized in **Table 2**.

Table 1

Initial concentrations in donor chamber and chromatographic conditions of prazosin, verapamil, and lamotrigine

Compound	Concentration	Wavelength
Prazosin	5 μ M	Ex: 250 nm
		Em: 390 nm
Verapamil	5 μ M	Ex: 280 nm
		Em: 310 nm
Lamotrigine	6 μ M	220nm

Table 2

The summary of mass charge ratio, extraction, initial concentrations in donor chamber

Compound	Concentration	Mass charge ratio [M+H] ⁺	Extraction
Amantadine	3 μ M	181	Water-saturated N-butanol
Amitriptyline	1.5 μ M	278	Ethyl acetate
Bupropion	3 μ M	240	Ethyl acetate
Carbamazepine	3 μ M	237	Ethyl acetate
Clozapine	4 μ M	327	Ethyl acetate
Donepezil	3 μ M	380	Methyl tert-butyl ether
Doxepin	4 μ M	317	Methyl tert-butyl ether
Fluoxetine	3 μ M	310	Ethyl acetate
Gabapentin	10 μ M	172	Ethyl acetate
Metoclopramide	4 μ M	301	Ethyl acetate
Midazolam	3 μ M	327	Ethyl acetate
Mirtazapine	3 μ M	266	Methyl tert-butyl ether
Olanzapine	3 μ M	313	Methyl tert-butyl ether
Risperidone	4 μ M	427	Methyl tert-butyl ether
Venlafaxine	10 μ M	278	Ethyl acetate

Cell proliferation analysis

On day 6 after co-culture, hCMEC/D3 cells were fixed with 4% paraformaldehyde for 15 min and washed with phosphate buffered saline (PBS) for three times. Next, the fixed cells were blocked with 5% goat serum for 2 h and washed with PBS for four times. The blocked cells were incubated with DAPI (Invitrogen, Carlsbad, CA, USA) and washed with PBS for four times. Cell numbers were counted using Cytation5 (BioTek, Winooski, VT, USA).

Reverse transcription and Quantitative real-time PCR (qPCR)

Total RNA of cells was extracted using RNAiso Plus reagent (Takara Bio Inc. Otsu, Shiga, Japan) and reverse transcribed using HiScript III RT SuperMix (Vazyme, Shanghai, China) as the described method (Yang et al., 2023 [\[1\]](#)). Paired primers were synthesized by Tsingke Biotech Co., Ltd (Beijing, China), and their sequences were listed in **Table 3** [\[1\]](#). The SYBR Master Mix was purchased from Yeasen (Shanghai, China). Then, qPCR was performed on the Applied Biosystems QuantStudio 3 real-time PCR system (Thermo Fisher Scientific, Waltham, MA, USA). The mRNA levels of related genes were normalized to *ACTB* or *GAPDH* using the comparative cycle threshold method.

Western blotting analysis

Whole-cell and tissue lysates, nucleoprotein, and cytoplasmic protein were prepared using RIPA Lysis Buffer (Beyotime, Shanghai, China) as the described method (Wu et al., 2021 [\[1\]](#)). Proteins were separated through sodium dodecyl sulfate-polyacrylamide gel electrophoresis and transferred onto nitrocellulose or polyvinylidene difluoride membranes. The membranes were blocked with 5% skim milk and incubated with corresponding primary antibodies at 4 °C overnight. After being washed with Tris-buffered saline Tween buffer, the membranes were incubated with secondary antibodies (Cell Signaling Technology, MA, USA) at 1:3000 dilution: Anti-mouse IgG, HRP-linked Antibody (#7076), Anti-rabbit IgG, HRP-linked Antibody (#7074). Protein levels were visualized using a highly sensitive ECL western blotting substrate and a gel imaging system (Tanon Science & Technology, Shanghai, China).

Preparation of conditioned medium

Conditioned medium (CM) of U251 cells (U-CM), SH-SY5Y cells (S-CM), or co-culture of U251 and SH-SY5Y cells (US-CM) were prepared. U251 cells were seeded at the top of the insert membrane and suspended on 6-well plates seeded with differentiated SH-SY5Y cells to co-culture U251 cells with SH-SY5Y cells. The medium was collected every 24 h. The CMs were subsequently used for hCMEC/D3 cell culture after filtrating with 0.2 µm filters for 144 h. The levels of glia-derived neurotrophic factor (GDNF), basic fibroblast growth factor (bFGF), insulin-like growth factor-1 (IGF-1), and transforming growth factor-β (TGF-β) in CMs were measured using corresponding ELISA kits according to the manufacturers' instructions.

Neutralization of GDNF with Anti-GDNF Antibody

Exogenous and endogenous GDNF in the medium was neutralized with anti-GDNF antibody (#AF-212-NA, R&D system, Minneapolis, MN, USA). 0.25, 0.5, 1.0 µg/mL anti-GDNF antibody was added into the US-CM or medium containing 200 pg/mL GDNF, and then the medium was preincubated at 4 °C for 1 h. The hCMEC/D3 cells were incubated with 200 pg/mL GDNF or US-CM containing anti-GDNF antibody or not for 6 days, and then cell lysate was collected for western blot. The medium was replaced every 24 h.

Gene (protein)	Forwards Primer, 5'→3'	Reverse Primer, 3'→5'
<i>ACTB</i> (β -actin)	GGACTTCGAGCAAGAGATGG	AGCACTGTGTTGGCGTACAG
<i>GAPDH</i> (GAPDH)	TGTGGGCATCAATGGATTTGG	ACACCATGTATTCCGGGTCAAT
<i>CLDN5</i> (claudin-5)	CTCTGCTGGTTCGCCAACAT	CAGCTCGTACTTCTGCGACA
<i>OCN</i> (occludin)	ACAAGCGGTTTTATCCAGAGTC	GTCATCCACAGGCGAAGTTAAT
<i>TJP1</i> (ZO-1)	ACCAGTAAGTCGTCCTGATCC	TCGGCCAAATCTTCTCACTCC
<i>CDH5</i> (VE-cadherin)	AAGCGTGAGTCGCAAGAATG	TCTCCAGGTTTTCGCCAGTG
<i>ABCB1</i> (P-gp)	TTGCTGCTTACATTCAGGTTTCA	AGCCTATCTCCTGTTCGCATTA
<i>ABCG2</i> (BCRP)	ACGAACGGATTAACAGGGTCA	CTCCAGACACACCACGGAT
<i>SLC22A1</i> (OCT1)	ACGGTGGCGATCATGTACC	CCCATTCTTTTGAGCGATGTGG
<i>SLC22A2</i> (OCT2)	CATCGTCACCGAGTTTAACTG	AGCCGATACTCATAGAGCCAAT
<i>SLC22A8</i> (OAT3)	ATGGCCCAGTCTATCTTCATGG	GACGGTGCTCAGGGTAATGC
<i>SLCO1A1</i> (OATP1A1)	TAATGTGGGTGTACGTCCTAGT	GCTCCTGTTTCTACAAGCCCAA
<i>GDNF</i> (GDNF)	GCAGACCCATCGCCTTTGAT	CCACACCTTTTAGCGGAATGC
<i>BDNF</i> (BDNF)	CTACGAGACCAAGTGCAATCC	AATCGCCAGCCAATTCTCTTT
<i>NGF</i> (NGF)	TGTGGGTTGGGGATAAGACCA	GCTGTCAACGGGATTTGGGT
<i>IGF1</i> (IGF-1)	GCTCTTCAGTTCGTGTGTGGA	GGTCATGGATGGACCTTACTGT
<i>VEGFA</i> (VEGF)	CCCACTGAGGAGTCCAACAT	AAATGCTTTCTCCGCTCTGA
<i>FGF2</i> (bFGF)	AGAAGAGCGACCCTCACATCA	CGGTTAGCACACACTCCTTTG
<i>TGFBI</i> (TGF- β)	GGCCAGATCCTGTCCAAGC	GTGGGTTTCCACCATTAGCAC

Table 3

Primer sequences for qPCR for indicted genes

Transfection of hCMEC/D3

Here, hCMEC/D3 cells were plated in the plates or culture dishes at 6×10^4 cells/cm² and transfected with 10 nM of negative control or human *FOXO1* and *ETS1* small interfering RNA (siRNA) (Tsingke Biotechnology, Beijing, China) for 12 h using Lipofectamine™ 3000 (Invitrogen, Carlsbad, CA, USA) reagent according to the manufacturer's instructions. Cells were then incubated with a medium containing GDNF for 72 h. The siRNA sequences of human *FOXO1* and *ETS1* were summarized in [Table 4](#).

FOXO1 Overexpression by plasmids

The plasmids encoding FOXO1 (EX-Z7404-M02) were constructed by GeneCopoeia (Rockville, MD, USA). The hCMEC/D3 cells were plated in plates or dishes at 6×10^4 cells/cm². They were subsequently transfected with 1 µg of negative control or plasmids encoding FOXO1 6 h using Lipofectamine™ 3000 reagent according to the manufacturer's instructions. Transfected cells were then incubated with the medium containing GDNF for 72 h.

Animals

C57BL/6J mice (male, 4–5 weeks old, 16–18 g, 12 mice) were obtained from Sino-British Sippr/BKLaboratory Animal Ltd (Shanghai, China). Mice were maintained in groups under standard conditions with free access to food and water. Animal studies were performed in accordance with the Guide for the Care and Use of Laboratory Animals (National Institutes of Health) and approved by the Animal Ethics Committee of China Pharmaceutical University (Approval Number: 202307003).

Brain-specific *Gdnf* knockdown and evaluation of BBB permeability

Mice were randomly divided into control (shNC) and *Gdnf* silencing (sh*Gdnf*) groups (6 mice each group). The shRNA sequence of mice *Gdnf* was listed in [Table 4](#). The 2×10^9 viral genome each of pAAV-U6-shRNA (NC2)-CMV-EGFP or pAAV-U6-shRNA (*Gdnf*)-CMV-EGFP (OBio Technology, Beijing, China) were injected into the bilateral lateral ventricle area (relative to the bregma: anterior-posterior −0.3 mm; medial-lateral ±1.0 mm; dorsal-ventral −3.0 mm) through intracerebroventricular (*i.c.v*) infusion. Three weeks following *i.c.v* injection, BBB permeability and expression of corresponding targeted proteins were measured in the mice.

A mixture of FITC-Dex (50 mg/kg) and fluorescein sodium (10 mg/kg) was intravenously administered to experimental mice. Thirty minutes after the injection, the mice were euthanized under isoflurane anesthesia, and brain tissue and plasma samples were obtained quickly. The concentrations of FITC-Dex and fluorescein in the plasma and brain were measured as previously described (Li P. et al., 2022; Zhou Y. et al., 2019). No blinding was performed in animal studies.

The prediction of drug permeability across BBB using the developed in vitro BBB model

The P_{app} values of 18 drugs – prazosin, verapamil, lamotrigine, clozapine, venlafaxine, bupropion, amantadine, carbamazepine, fluoxetine, amitriptyline, gabapentin, midazolam, risperidone, olanzapine, mirtazapine, metoclopramide, doxepin, and donepezil – across the hCMEC/D3 cells mono-culture and triple co-culture models were measured. The predicted *in vivo* permeability-surface area product (PS , µL/min/g brain) values across BBB were calculated using the following equation:

$$PS_{Pre} = \frac{P_{app} \times 60 \times VSA \times 1000}{f_{u,brain}} \quad (2)$$

Gene	Target sequence
<i>ETS1</i>	CGCTATACCTCGGATTACT
<i>FOXO1</i>	AATCTCCTAGGAGAAGAGCTG
<i>Gdnf</i>	GCCAGTGTTTATCTGATAC

Table 4

The target sequences for siRNA or shRNA

where VSA is the luminal area of the vascular space of brain, which was set to 150 cm²/g (Fenstermacher et al., 1988), and $f_{u, brain}$ is the unbound fraction of brain.

The published *in vivo* brain permeability values were unified to observed PS (PS_{Obs}) by multiplying by VSA equal to 150 cm²/g. The PS_{Obs} and $f_{u, brain}$ of these 18 drugs were summarized in **Table 5**. If PS_{Pre} values were within 0.5–2.0 folds of observations, the prediction was considered successful.

Statistical analyses

All results are expressed as mean ± S.E.M. Data were analyzed using GraphPad Prism software version 8.0.2 (GraphPad Software, La Jolla, CA, USA). $P < 0.05$ was considered statistically significant. The unpaired Student’s t-test or unpaired t-test with Welch’s correction for two-group comparisons, and one-way ANOVA followed by Fisher’s LSD test or Welch’s ANOVA test was used for multiple group comparisons. The simple linear regression analysis was used to examine the presence of a linear relationship between two variables.

Acknowledgements

The authors would like to give special thanks to the support of the China Pharmaceutical University Pharmaceutical Animal Laboratory Center.

Funding information

Funder	Grant reference number	Author
National Natural Science Foundation of China	82373943	Xiaodong Liu
National Natural Science Foundation of China	82173884	Li Liu
National Natural Science Foundation of China	82204511	Hanyu Yang
“Double First-Class” university project	No. CPU2022QZ21	Li Liu
Jiangsu Funding Program for Excellent Postdoctoral Talent	No. 2022ZB305	Hanyu Yang
China Postdoctoral Science Foundation	2023M733897	Hanyu Yang
The funding sources were involved in study design, data collection, and interpretation, or the decision to submit the work for publication.		

Author Contributions

Lu Yang: Conceptualization, Methodology, Investigation, Data curation, and Writing—original draft preparation; **Zijin Lin:** Data curation, Formal analysis; **Ruijing Mu:** Visualisation, Software; **Wenhan Wu:** Formal analysis; **Hao Zhi:** Data curation; **Xiaodong Liu:** Supervision, Conceptualization, Project administration, Funding acquisition; **Hanyu Yang:** Supervision,

Compounds	$f_{u, \text{ brain }}$	PS_{Obs} μl/min/g	$P_{\text{app, Mono}}$ cm/s×10 ⁻⁶	$PS_{\text{Pre, Mono}}$ μl/min/g	$P_{\text{app, Triple}}$ cm/s×10 ⁻⁶	$PS_{\text{Pre, Triple}}$ μl/min/g
Amantadine	0.1985 ^(a)	11.61 ^(b)	6.84±1.90	31.02	3.64±0.53	16.52
Amitriptyline	0.01 ^(c)	459.00 ^(k)	15.24±1.28	1371.60	14.61±0.54	1314.90
Bupropion	0.12 ^(d)	151.92 ^(b)	15.19±0.40	113.96	11.34±0.88	85.06
Carbamazepine	0.116 ^(e)	95.94 ^(b)	34.37±2.51	266.69	11.71±0.30	90.87
Clozapine	0.014 ^(f)	226.08 ^(b)	38.97±1.08	2505.26	12.87±4.11	827.27
Donepezil	0.07 ^(g)	158.13 ^(b)	20.91±1.49	268.84	14.47±1.69	186.04
Doxepin	0.025 ^(b)	219.24 ^(b)	16.88±2.17	607.68	10.66±1.84	383.76
Fluoxetine	0.004 ^(b)	269.82 ^(b)	11.48±1.69	2583.00	9.97±2.05	2243.25
Gabapentin	0.782 ^(b)	16.29 ^(b)	16.75±3.23	19.28	8.78±0.46	10.10
Lamotrigine	0.273 ^(b)	12.60 ^(b)	14.26±0.74	47.01	5.97±0.22	19.69
Metoclopramide	0.365 ^(b)	12.51 ^(b)	14.14±2.87	34.87	6.63±0.84	16.34
Midazolam	0.045 ^(h)	272.70 ^(b)	25.30±1.99	506.00	19.09±0.48	381.80
Mirtazapine	0.08 ^(b)	191.25 ^(b)	22.96±0.89	258.30	17.86±0.43	200.93
Olanzapine	0.034 ^(b)	227.97 ^(b)	24.33±7.60	644.03	12.49±1.06	330.62
Prazosin	0.09 ⁽ⁱ⁾	16.92 ⁽ⁱ⁾	5.61±0.76	56.09	2.81±1.04	28.10
Risperidone	0.099 ^(b)	84.96 ^(b)	16.10±5.74	146.36	11.70±0.50	106.36
Venlafaxine	0.205 ^(a)	58.41 ^(b)	9.58±0.56	42.06	8.25±0.71	36.20
Verapamil	0.033 ^(c)	33.57 ^(c)	7.21±0.81	196.52	5.56±0.12	151.77

^(a)(Esaki et al., 2019); ^(b)(Summerfield et al., 2007); ^(c)(Friden et al., 2011); ^(d)(Bhattacharya et al., 2021); ^(e)(Maurer et al., 2005); ^(f)(Cremers et al., 2012); ^(g)(Di et al., 2011); ^(h)(Kodaira et al., 2011); ⁽ⁱ⁾(Zhou L. et al., 2009); ^(j)(Di Marco et al., 2019); ^(k)(Avdeef et al., 2010).

Table 5—source data (Summary of original data and analysis)

Table 5.

The unbound fraction in brain ($f_{u, \text{ brain }}$), the observed PS_{Obs} , and the predicted PS (PS_{Pre}), P_{app} across the hCMEC/D3 mono-culture model ($P_{\text{app, Mono}}$) and triple co-culture model ($P_{\text{app, Triple}}$) of the tested drugs. n=4.

Conceptualization, Funding acquisition; **Li Liu**: Supervision, Conceptualization, Project administration, Funding acquisition, Resources. All authors reviewed and approved the final version of this manuscript.

Declaration of interests

The authors declare no competing interests.

Data availability

All data are available in the manuscript and supporting files; source data files for western blots have been provided for all figures.

References

1. Abbott NJ (2013) **Blood-brain barrier structure and function and the challenges for CNS drug delivery** *J Inherit Metab Dis* **36**:437–449 <https://doi.org/10.1007/s10545-013-9608-0>
2. Arvanitis C D, Ferraro G B, Jain R K (2020) **The blood-brain barrier and blood-tumour barrier in brain tumours and metastases** *Nat Rev Cancer* **20**:26–41 <https://doi.org/10.1038/s41568-019-0205-x>
3. Asada S *et al.* (2007) **Mitogen-activated protein kinases, Erk and p38, phosphorylate and regulate Foxo1** *Cell Signal* **19**:519–527 <https://doi.org/10.1016/j.cellsig.2006.08.015>
4. Avdeef A, Sun N (2010) **A New In Situ Brain Perfusion Flow Correction Method for Lipophilic Drugs Based on the pH-Dependent Crone-Renkin Equation** *Pharmaceutical Research* **28**:517–530 <https://doi.org/10.1007/s11095-010-0298-0>
5. Banks W A (2016) **From blood-brain barrier to blood-brain interface: new opportunities for CNS drug delivery** *Nat Rev Drug Discov* **15**:275–292 <https://doi.org/10.1038/nrd.2015.21>
6. Barberio C, Withers A, Mishra Y, Couraud P-O, Romero I A, Weksler B, Owens R M (2022) **A human-derived neurovascular unit in vitro model to study the effects of cellular cross-talk and soluble factors on barrier integrity** *Frontiers in Cellular Neuroscience* **16** <https://doi.org/10.3389/fncel.2022.1065193>
7. Beard R S, Hoettels B A, Meegan J E, Wertz T S, Cha B J, Yang X, Oxford J T, Wu M H, Yuan S Y (2020) **AKT2 maintains brain endothelial claudin-5 expression and selective activation of IR/AKT2/FOXO1-signaling reverses barrier dysfunction** *J Cereb Blood Flow Metab* **40**:374–391 <https://doi.org/10.1177/0271678X18817512>
8. Bhallamudi S, Roos B B, Teske J J, Wicher S A, McConico A C M P, Sathish V, Prakash Y S (2021) **Glial-derived neurotrophic factor in human airway smooth muscle** *J Cell Physiol* **236**:8184–8196 <https://doi.org/10.1002/jcp.30489>

9. Bhattacharya C, Masters A R, Bach C, Stratford R E (2021) **Population model analysis of chiral inversion and degradation of bupropion enantiomers, and application to enantiomer specific fraction unbound determination in rat plasma and brain** *J Pharm Biomed Anal* **195** <https://doi.org/10.1016/j.jpba.2020.113872>
10. Bhyrapuneni G, Thentu J B, Palacharla V R C, Muddana N, Aleti R R, Ajjala D R, Nirogi R (2018) **A definite measure of occupancy exposures, seeking with non-radiolabeled in vivo 5-HT_{2A} receptor occupancy and in vitro free fractions** *J Recept Signal Transduct Res* **38**:359–366 <https://doi.org/10.1080/10799893.2018.1531888>
11. Biswas S, Cottarelli A, Agalliu D (2020) **Neuronal and glial regulation of CNS angiogenesis and barrierogenesis** *Development* **147** <https://doi.org/10.1242/dev.182279>
12. Bowman C M, Benet L Z (2018) **An examination of protein binding and protein-facilitated uptake relating to in vitro-in vivo extrapolation** *Eur J Pharm Sci* **123**:502–514 <https://doi.org/10.1016/j.ejps.2018.08.008>
13. Buldan A, Dietze R, Shihan M, Scheiner-Bobis G (2016) **Non-classical testosterone signaling mediated through ZIP9 stimulates claudin expression and tight junction formation in Sertoli cells** *Cellular Signalling* **28**:1075–1085 <https://doi.org/10.1016/j.cellsig.2016.04.015>
14. Clasadonte J, Prevot V (2017) **The special relationship: glia–neuron interactions in the neuroendocrine hypothalamus** *Nature Reviews Endocrinology* **14**:25–44 <https://doi.org/10.1038/nrendo.2017.124>
15. Colas-Algora N, Garcia-Weber D, Cacho-Navas C, Barroso S, Caballero A, Ribas C, Correias I, Millan J (2020) **Compensatory increase of VE-cadherin expression through ETS1 regulates endothelial barrier function in response to TNF α** *Cell Mol Life Sci* **77**:2125–2140 <https://doi.org/10.1007/s00018-019-03260-9>
16. Cremers T I, Flik G, Hofland C, Stratford R E (2012) **Microdialysis evaluation of clozapine and N-desmethylozapine pharmacokinetics in rat brain** *Drug Metab Dispos* **40**:1909–1916 <https://doi.org/10.1124/dmd.112.045682>
17. Dejana E, Orsenigo F, Lampugnani M G (2008) **The role of adherens junctions and VE-cadherin in the control of vascular permeability** *J Cell Sci* **121**:2115–2122 <https://doi.org/10.1242/jcs.017897>
18. Di L, Umland J P, Chang G, Huang Y, Lin Z, Scott D O, Troutman M D, Liston T E. (2011) **Species independence in brain tissue binding using brain homogenates** *Drug Metab Dispos* **39**:1270–1277 <https://doi.org/10.1124/dmd.111.038778>
19. Di L *et al.* (2017) **Industry Perspective on Contemporary Protein-Binding Methodologies: Considerations for Regulatory Drug-Drug Interaction and Related Guidelines on Highly Bound Drugs** *J Pharm Sci* **106**:3442–3452 <https://doi.org/10.1016/j.xphs.2017.09.005>
20. Di Marco A *et al.* (2019) **Application of an in Vitro Blood-Brain Barrier Model in the Selection of Experimental Drug Candidates for the Treatment of Huntington’s Disease** *Mol Pharm* **16**:2069–2082 <https://doi.org/10.1021/acs.molpharmaceut.9b00042>
21. Dong C, Ubogu E E (2018) **GDNF enhances human blood-nerve barrier function in vitro via MAPK signaling pathways** *Tissue Barriers* **6**:1–22 <https://doi.org/10.1080/21688370.2018.1546537>

22. Eigenmann D E, Xue G, Kim K S, Moses A V, Hamburger M, Oufir M (2013) **Comparative study of four immortalized human brain capillary endothelial cell lines, hCMEC/D3, hBMEC, TY10, and BB19, and optimization of culture conditions, for an in vitro blood–brain barrier model for drug permeability studies** *Fluids Barriers CNS* **10** <https://doi.org/10.1186/2045-8118-10-33>
23. Esaki T, Ohashi R, Watanabe R, Natsume-Kitatani Y, Kawashima H, Nagao C, Mizuguchi K (2019) **Computational Model To Predict the Fraction of Unbound Drug in the Brain** *J Chem Inf Model* **59**:3251–3261 <https://doi.org/10.1021/acs.jcim.9b00180>
24. Fenstermacher J, Gross P, Sposito N, Acuff V, Pettersen S, Gruber K (1988) **Structural and Functional Variations in Capillary Systems within the Brain** *Ann N Y Acad Sci* **529**:21–30 <https://doi.org/10.1111/j.1749-6632.1988.tb51416.x>
25. Fielder G C, Yang T W, Razdan M, Li Y, Lu J, Perry J K, Lobie P E, Liu D X (2018) **The GDNF Family: A Role in Cancer?** *Neoplasia* **20**:99–117 <https://doi.org/10.1016/j.neo.2017.10.010>
26. Friden M, Bergstrom F, Wan H, Rehngren M, Ahlin G, Hammarlund-Udenaes M, Bredberg U (2011) **Measurement of unbound drug exposure in brain: modeling of pH partitioning explains diverging results between the brain slice and brain homogenate methods** *Drug Metab Dispos* **39**:353–362 <https://doi.org/10.1124/dmd.110.035998>
27. Fu J *et al.* (2021) **Astrocyte-Derived TGFβ1 Facilitates Blood–Brain Barrier Function via Non-Canonical Hedgehog Signaling in Brain Microvascular Endothelial Cells** *Brain Sciences* **11** <https://doi.org/10.3390/brainsci11010077>
28. Hanafy A S, Dietrich D, Fricker G, Lamprecht A (2021) **Blood-brain barrier models: Rationale for selection** *Adv Drug Deliv Rev* **176** <https://doi.org/10.1016/j.addr.2021.113859>
29. Hashimoto Y, Greene C, Munnich A, Campbell M (2023) **The CLDN5 gene at the blood-brain barrier in health and disease** *Fluids and Barriers of the CNS* **20** <https://doi.org/10.1186/s12987-023-00424-5>
30. Hatherell K, Couraud P O, Romero I A, Weksler B, Pilkington G J (2011) **Development of a three-dimensional, all-human in vitro model of the blood-brain barrier using mono-, co-, and tri-cultivation Transwell models** *J Neurosci Methods* **199**:223–229 <https://doi.org/10.1016/j.jneumeth.2011.05.012>
31. He F, Wang Q F, Li L, Yu C, Liu C Z, Wei W C, Chen L P, Li H Y (2023) **Melatonin Protects Against Hyperoxia-Induced Apoptosis in Alveolar Epithelial type II Cells by Activating the MT2/PI3K/AKT/ETS1 Signaling Pathway** *Lung* **201**:225–234 <https://doi.org/10.1007/s00408-023-00610-0>
32. Hui K, Wu S, Yue Y, Gu Y, Guan B, Wang X, Hsieh J T, Chang L S, He D, Wu K (2018) **RASAL2 inhibits tumor angiogenesis via p-AKT/ETS1 signaling in bladder cancer** *Cell Signal* **48**:38–44 <https://doi.org/10.1016/j.cellsig.2018.04.006>
33. Igarashi Y *et al.* (1999) **Glial cell line-derived neurotrophic factor induces barrier function of endothelial cells forming the blood-brain barrier** *Biochem Biophys Res Commun* **261**:108–112 <https://doi.org/10.1006/bbrc.1999.0992>
34. Ito R *et al.* (2019) **A Human Immortalized Cell-Based Blood-Brain Barrier Triculture Model: Development and Characterization as a Promising Tool for Drug-Brain Permeability Studies** *Mol Pharm* **16**:4461–4471 <https://doi.org/10.1021/acs.molpharmaceut.9b00519>

35. Ji B S, Cen J, He L, Liu M, Liu Y Q, Liu L (2013) **Modulation of P-glycoprotein in rat brain microvessel endothelial cells under oxygen glucose deprivation** *J Pharm Pharmacol* **65**:1508–1517 <https://doi.org/10.1111/jphp.12122>
36. Ko J A, Yanai R, Nishida T (2009) **IGF-1 released by corneal epithelial cells induces up-regulation of N-cadherin in corneal fibroblasts** *J Cell Physiol* **221**:254–261 <https://doi.org/10.1002/jcp.21850>
37. Kodaira H, Kusuhara H, Fujita T, Ushiki J, Fuse E, Sugiyama Y (2011) **Quantitative evaluation of the impact of active efflux by p-glycoprotein and breast cancer resistance protein at the blood-brain barrier on the predictability of the unbound concentrations of drugs in the brain using cerebrospinal fluid concentration as a surrogate** *J Pharmacol Exp Ther* **339**:935–944 <https://doi.org/10.1124/jpet.111.180398>
38. Lelièvre E, Mattot V, Huber P, Vandenbunde B, Soncin F (2000) **ETS1 lowers capillary endothelial cell density at confluence and induces the expression of VE-cadherin** *Oncogene* **19**:2438–2446 <https://doi.org/10.1038/sj.onc.1203563>
39. Lelièvre E, Lionneton F, Soncin F, Vandenbunder B (2001) **The Ets family contains transcriptional activators and repressors involved in angiogenesis** *Int J Biochem Cell Biol* **33**:391–407 [https://doi.org/10.1016/s1357-2725\(01\)00025-5](https://doi.org/10.1016/s1357-2725(01)00025-5)
40. Li M S, Cen J, He L, Liu L, Ji B S (2013) **CJY, an isoflavone, interacts with ATPase of P-glycoprotein in the rat brain microvessel endothelial cells (RBMECs)** *J Chemother* **25**:347–354 <https://doi.org/10.1179/1973947813Y.0000000094>
41. Li P, Yang Y, Lin Z, Hong S, Jiang L, Zhou H, Yang L, Zhu L, Liu X, Liu L (2022) **Bile Duct Ligation Impairs Function and Expression of Mrp1 at Rat Blood-Retinal Barrier via Bilirubin-Induced P38 MAPK Pathway Activations** *Int J Mol Sci* **23** <https://doi.org/10.3390/ijms23147666>
42. Li W, Chen Z, Chin I, Chen Z, Dai H (2018) **The Role of VE-cadherin in Blood-brain Barrier Integrity Under Central Nervous System Pathological Conditions** *Current Neuropharmacology* **16**:1375–1384 <https://doi.org/10.2174/1570159x16666180222164809>
43. Li Y, Li Y, Zhang Y, Zhao Q, Zhang P, Sun M, Liu B, Yang H, Li P (2022) **Muscone and (+)-Borneol Cooperatively Strengthen CREB Induction of Claudin 5 in IL-1 β -Induced Endothelium Injury** *Antioxidants* **11** <https://doi.org/10.3390/antiox11081455>
44. Lin C H, Wang C H, Hsu S L, Liao L Y, Lin T A, Hsueh C M (2016) **Molecular Mechanisms Responsible for Neuron-Derived Conditioned Medium (NCM)-Mediated Protection of Ischemic Brain** *PLoS One* **11** <https://doi.org/10.1371/journal.pone.0146692>
45. Liu D, Yang L, Liu P, Ji X, Qi X, Wang Z, Chi T, Zou L (2022) **Sigma-1 receptor activation alleviates blood-brain barrier disruption post cerebral ischemia stroke by stimulating the GDNF-GFRalpha1-RET pathway** *Exp Neurol* **347** <https://doi.org/10.1016/j.expneurol.2021.113867>
46. Liu X *et al.* (2005) **Use of a physiologically based pharmacokinetic model to study the time to reach brain equilibrium: an experimental analysis of the role of blood-brain barrier permeability, plasma protein binding, and brain tissue binding** *J Pharmacol Exp Ther* **313**:1254–1262 <https://doi.org/10.1124/jpet.104.079319>

47. Liu Y, Mu S, Li X, Liang Y, Wang L, Ma X (2019) **Unfractionated Heparin Alleviates Sepsis-Induced Acute Lung Injury by Protecting Tight Junctions** *Journal of Surgical Research* **238**:175–185 <https://doi.org/10.1016/j.jss.2019.01.020>
48. Lonka-Nevalaita L, Lume M, Leppänen S, Jokitalo E, Peränen J, Saarma M (2010) **Characterization of the Intracellular Localization, Processing, and Secretion of Two Glial Cell Line-Derived Neurotrophic Factor Splice Isoforms** *The Journal of Neuroscience* **30**:11403–11413 <https://doi.org/10.1523/jneurosci.5888-09.2010>
49. Luo Y *et al.* (2022) **Endothelial ETS1 inhibition exacerbate blood-brain barrier dysfunction in multiple sclerosis through inducing endothelial-to-mesenchymal transition** *Cell Death Dis* **13** <https://doi.org/10.1038/s41419-022-04888-5>
50. Mathiisen T M, Lehre K P, Danbolt N C, Ottersen O P (2010) **The perivascular astroglial sheath provides a complete covering of the brain microvessels: an electron microscopic 3D reconstruction** *Glia* **58**:1094–1103 <https://doi.org/10.1002/glia.20990>
51. Maurer T S, DeBartolo D B, Tess D A, Scott D O (2005) **Relationship between exposure and nonspecific binding of thirty-three central nervous system drugs in mice** *Drug Metab Dispos* **33**:175–181 <https://doi.org/10.1124/dmd.104.001222>
52. Morita A *et al.* (2006) **Regulation of Dendritic Branching and Spine Maturation by Semaphorin3A-Fyn Signaling** *The Journal of Neuroscience* **26**:2971–2980 <https://doi.org/10.1523/jneurosci.5453-05.2006>
53. Muoio V, Persson P B, Sendeski M M (2014) **The neurovascular unit – concept review** *Acta Physiol* **210**:790–798 <https://doi.org/10.1111/apha.12250>
54. Nakagawa S, Deli M A, Kawaguchi H, Shimizudani T, Shimono T, Kittel A, Tanaka K, Niwa M (2009) **A new blood-brain barrier model using primary rat brain endothelial cells, pericytes and astrocytes** *Neurochem Int* **54**:253–263 <https://doi.org/10.1016/j.neuint.2008.12.002>
55. Nitta T, Hata M, Gotoh S, Seo Y, Sasaki H, Hashimoto N, Furuse M, Tsukita S (2003) **Size-selective loosening of the blood-brain barrier in claudin-5-deficient mice** *J Cell Biol* **161**:653–660 <https://doi.org/10.1083/jcb.200302070>
56. Nowrangi D S, McBride D, Manaenko A, Dixon B, Tang J, Zhang J H (2019) **rhIGF-1 reduces the permeability of the blood-brain barrier following intracerebral hemorrhage in mice** *Exp Neurol* **312**:72–81 <https://doi.org/10.1016/j.expneurol.2018.11.009>
57. Ohtsuki S, Sato S, Yamaguchi H, Kamoi M, Asashima T, Terasaki T (2007) **Exogenous expression of claudin-5 induces barrier properties in cultured rat brain capillary endothelial cells** *J Cell Physiol* **210**:81–86 <https://doi.org/10.1002/jcp.20823>
58. Ohtsuki S *et al.* (2013) **Quantitative targeted absolute proteomic analysis of transporters, receptors and junction proteins for validation of human cerebral microvascular endothelial cell line hCMEC/D3 as a human blood-brain barrier model** *Mol Pharm* **10**:289–296 <https://doi.org/10.1021/mp3004308>
59. Palmiotti C A, Prasad S, Naik P, Abul K M, Sajja R K, Achyuta A H, Cucullo L (2014) **In vitro cerebrovascular modeling in the 21st century: current and prospective technologies** *Pharm Res* **31**:3229–3250 <https://doi.org/10.1007/s11095-014-1464-6>

60. Potjewyd G, Moxon S, Wang T, Domingos M, Hooper N M (2018) **Tissue Engineering 3D Neurovascular Units: A Biomaterials and Bioprinting Perspective** *Trends Biotechnol* **36**:457–472 <https://doi.org/10.1016/j.tibtech.2018.01.003>
61. Sanchez-Dengra B, Gonzalez-Alvarez I, Sousa F, Bermejo M, Gonzalez-Alvarez M, Sarmiento B (2021) **In vitro model for predicting the access and distribution of drugs in the brain using hCMEC/D3 cells** *Eur J Pharm Biopharm* **163**:120–126 <https://doi.org/10.1016/j.ejpb.2021.04.002>
62. Sauteur L, Affolter M, Belting H-G (2017) **Distinct and redundant functions of Esam and VE-cadherin during vascular morphogenesis** *Development* <https://doi.org/10.1242/dev.140038>
63. Savettieri G *et al.* (2000) **Neurons and ECM regulate occludin localization in brain endothelial cells** *Neuroreport* **11**:1081–1084 <https://doi.org/10.1097/00001756-200004070-00035>
64. Schiera G, Bono E, Raffa M P, Gallo A, Pitarresi G L, Di Liegro I, Savettieri G. (2003) **Synergistic effects of neurons and astrocytes on the differentiation of brain capillary endothelial cells in culture** *J Cell Mol Med* **7**:165–170 <https://doi.org/10.1111/j.1582-4934.2003.tb00215.x>
65. Schiera G, Sala S, Gallo A, Raffa M P, Pitarresi G L, Savettieri G, Di Liegro I. (2005) **Permeability properties of a three-cell type in vitro model of blood-brain barrier** *Journal of Cellular and Molecular Medicine* **9**:373–379 <https://doi.org/10.1111/j.1582-4934.2005.tb00362.x>
66. Shimizu F, Sano Y, Abe M A, Maeda T, Ohtsuki S, Terasaki T, Kanda T (2011) **Peripheral nerve pericytes modify the blood-nerve barrier function and tight junctional molecules through the secretion of various soluble factors** *J Cell Physiol* **226**:255–266 <https://doi.org/10.1002/jcp.22337>
67. Shimizu F, Sano Y, Saito K, Abe M A, Maeda T, Haruki H, Kanda T (2012) **Pericyte-derived glial cell line-derived neurotrophic factor increase the expression of claudin-5 in the blood-brain barrier and the blood-nerve barrier** *Neurochem Res* **37**:401–409 <https://doi.org/10.1007/s11064-011-0626-8>
68. Summerfield S G, Read K, Begley D J, Obradovic T, Hidalgo I J, Coggon S, Lewis A V, Porter R A, Jeffrey P (2007) **Central nervous system drug disposition: the relationship between in situ brain permeability and brain free fraction** *J Pharmacol Exp Ther* **322**:205–213 <https://doi.org/10.1124/jpet.107.121525>
69. Sweeney M D, Zhao Z, Montagne A, Nelson A R, Zlokovic B V (2019) **Blood-Brain Barrier: From Physiology to Disease and Back** *Physiol Rev* **99**:21–78 <https://doi.org/10.1152/physrev.00050.2017>
70. Taddei A, Giampietro C, Conti A, Orsenigo F, Breviario F, Pirazzoli V, Potente M, Daly C, Dimmeler S, Dejana E (2008) **Endothelial adherens junctions control tight junctions by VE-cadherin-mediated upregulation of claudin-5** *Nat Cell Biol* **10**:923–934 <https://doi.org/10.1038/ncb1752>
71. Tang E D, Nuñez G, Barr F G, Guan K-L (1999) **Negative Regulation of the Forkhead Transcription Factor FKHR by Akt** *Journal of Biological Chemistry* **274**:16741–16746 <https://doi.org/10.1074/jbc.274.24.16741>
72. Tavelin S, Gråsjö J, Taipalensuu J, Ocklind G, Artursson P, Wise C. (2002) **Applications of Epithelial Cell Culture in Studies of Drug Transport** *Epithelial Cell Culture Protocols* :233–272 <https://doi.org/10.1385/1-59259-185-x:233>

73. Tunggal J A, Helfrich I, Schmitz A, Schwarz H, Günzel D, Fromm M, Kemler R, Krieg T, Niessen C M (2005) **E-cadherin is essential for in vivo epidermal barrier function by regulating tight junctions** *The EMBO Journal* **24**:1146–1156 <https://doi.org/10.1038/sj.emboj.7600605>
74. Wang Z G *et al.* (2016) **bFGF Protects Against Blood-Brain Barrier Damage Through Junction Protein Regulation via PI3K-Akt-Rac1 Pathway Following Traumatic Brain Injury** *Mol Neurobiol* **53**:7298–7311 <https://doi.org/10.1007/s12035-015-9583-6>
75. Watanabe D *et al.* (2004) **Transcription factor Ets-1 mediates ischemia- and vascular endothelial growth factor-dependent retinal neovascularization** *Am J Pathol* **164**:1827–1835 [https://doi.org/10.1016/S0002-9440\(10\)63741-8](https://doi.org/10.1016/S0002-9440(10)63741-8)
76. Watanabe D, Nakagawa S, Morofuji Y, Toth A E, Vastag M, Aruga J, Niwa M, Deli M A (2021) **Characterization of a Primate Blood-Brain Barrier Co-Culture Model Prepared from Primary Brain Endothelial Cells, Pericytes and Astrocytes** *Pharmaceutics* **13** <https://doi.org/10.3390/pharmaceutics13091484>
77. Weber M L, Hofland C M, Shaffer C L, Flik G, Cremers T, Hurst R S, Rollema H (2013) **Therapeutic doses of antidepressants are projected not to inhibit human alpha4beta2 nicotinic acetylcholine receptors** *Neuropharmacology* **72**:88–95 <https://doi.org/10.1016/j.neuropharm.2013.04.027>
78. Weksler B, Romero I A, Couraud P-O (2013) **The hCMEC/D3 cell line as a model of the human blood brain barrier** *Fluids and Barriers of the CNS* **10** <https://doi.org/10.1186/2045-8118-10-16>
79. Wu T *et al.* (2021) **Bile duct ligation causes opposite impacts on the expression and function of BCRP and P-gp in rat brain partly via affecting membrane expression of ezrin/radixin/moesin proteins** *Acta Pharmacol Sin* **42**:1942–1950 <https://doi.org/10.1038/s41401-020-00602-3>
80. Xue Q, Liu Y, Qi H, Ma Q, Xu L, Chen W, Chen G, Xu X (2013) **A Novel Brain Neurovascular Unit Model with Neurons, Astrocytes and Microvascular Endothelial Cells of Rat** *International Journal of Biological Sciences* **9**:174–189 <https://doi.org/10.7150/ijbs.5115>
81. Yang H, Su M, Liu M, Sheng Y, Zhu L, Yang L, Mu R, Zou J, Liu X, Liu L (2023) **Hepatic retinaldehyde deficiency is involved in diabetes deterioration by enhancing PCK1- and G6PC-mediated gluconeogenesis** *Acta Pharm Sin B* **13**:3728–3743 <https://doi.org/10.1016/j.apsb.2023.06.014>
82. Zhang X, Tang N, Hadden T J, Rishi A K (2011) **Akt, FoxO and regulation of apoptosis** *Biochimica et Biophysica Acta* **1813**:1978–1986 <https://doi.org/10.1016/j.bbamcr.2011.03.010>
83. Zhou L, Schmidt K, Nelson F R, Zelesky V, Troutman M D, Feng B (2009) **The effect of breast cancer resistance protein and P-glycoprotein on the brain penetration of flavopiridol, imatinib mesylate (Gleevec), prazosin, and 2-methoxy-3-(4-(2-(5-methyl-2-phenyloxazol-4-yl)ethoxy)phenyl)propanoic acid (PF-407288) in mice** *Drug Metab Dispos* **37**:946–955 <https://doi.org/10.1124/dmd.108.024489>
84. Zhou Y *et al.* (2019) **Increase in P-glycoprotein levels in the blood-brain barrier of partial portal vein ligation /chronic hyperammonemia rats is mediated by ammonia/reactive oxygen species/ERK1/2 activation: In vitro and in vivo studies** *Eur J Pharmacol* **846**:119–127 <https://doi.org/10.1016/j.ejphar.2019.01.005>

Editors

Reviewing Editor

Ivan Velasco

Universidad Nacional Autónoma de México, Mexico City, Mexico

Senior Editor

Jonathan Cooper

Fred Hutchinson Cancer Research Center, Seattle, United States of America

Reviewer #1 (Public Review):

Summary:

In this manuscript, the authors established an in vitro triple co-culture BBB model and demonstrated its advantages compared with the mono or double co-culture BBB model. Further, the authors used their established in vitro BBB model and combined it with other methodologies to investigate the specific mechanism that co-culture with astrocytes but also neurons enhanced the integrity of endothelial cells.

Strengths:

The results persuasively showed the established triple co-culture BBB model well mimicked several important characteristics of BBB compared with the mono-culture BBB model, including better barrier function and in vivo/in vitro correlation. The human-derived immortalized cells used made the model construction process faster and more efficient, and have a better in vivo correlation without species differences. This model is expected to be a useful high-throughput evaluation tool in the development of CNS drugs.

Based on the previous experimental results, detailed studies investigated how co-culture with neurons and astrocytes promoted claudin-5 and VE-cadherin in endothelial cells, and the specific signaling mechanisms were also studied. Interestingly, the authors found that neurons also released GDNF to promote barrier properties of brain endothelial cells, as most current research has focused on the promoting effect of astrocytes-derived GDNF on BBB. Meanwhile, the author also validated the functions of GDNF for BBB integrity in vivo by silencing GDNF in mouse brains. Overall, the experiments and data presented support their claim that, in addition to astrocytes, neurons also have a promoting effect on the barrier function of endothelial cells through GDNF secretion.

Weaknesses:

Although the authors demonstrated a highly usable for predicting the BBB permeability, recorded TEER measurements are still far from the human BBB in vivo reported measurements of TEER, and expression of transporters was not promoted by co-culture, which may lead to the model being unsuitable for studying drug transport mediated by transporters on BBB.

<https://doi.org/10.7554/eLife.96161.1.sa1>

Reviewer #2 (Public Review):

Summary:

Yang and colleagues developed a new in vitro blood-brain barrier model that is relatively simple yet outperforms previous models. By incorporating a neuroblastoma cell line, they

demonstrated increased electrical resistance and decreased permeability to small molecules.

Strengths:

The authors initially elucidated the soluble mediator responsible for enhancing endothelial functionality, namely GDNF. Subsequently, they elucidated the mechanisms by which GDNF upregulates the expression of VE-cadherin and Claudin-5. They further validated these findings *in vivo*, and demonstrated predictive value for molecular permeability as well. The study is meticulously conducted and easily comprehensible. The conclusions are firmly supported by the data, and the objectives are successfully achieved. This research is poised to advance future investigations in BBB permeability, leakage, dysfunction, disease modeling, and drug delivery, particularly in high-throughput experiments. I anticipate an enthusiastic reception from the community interested in this area. While other studies have produced similar results with tri-cultures (PMID: 25630899), this study notably enhances electrical resistance compared to previous attempts.

Weaknesses:

Considerable effort has been directed towards developing *in vitro* models that more closely resemble their *in vivo* counterparts, utilizing stem cell-derived NVU cells. Although these examples are currently rudimentary, they offer better BBB mimicry than Yang's study.

Additionally, some instances might benefit from more robust statistical tests; nonetheless, I do not think this would significantly alter the experimental conclusions.

Similar experiments with tri-cultures yielding analogous results have been reported by other authors (PMID: 25630899). TEER values are a bit higher than the aforementioned experiments; however, this study has values at least one order of magnitude lower than physiological levels.

<https://doi.org/10.7554/eLife.96161.1.sa0>