

Cytoneme-mediated intercellular signaling in keratinocytes essential for epidermal remodeling

Reviewed Preprint

Published from the original preprint after peer review and assessment by eLife.

About eLife's process

Reviewed preprint version 1

May 16, 2024 (this version)

Sent for peer review

March 12, 2024

Posted to preprint server

February 28, 2024

Yi Wang, Qingan He, Oliver Has, Kiarash Forouzes, Dae Seok Eom 

Department of Developmental and Cell Biology, University of California, Irvine, CA 92617, USA • Center for Complex Biological Systems, University of California, Irvine, CA 92697, USA • UC Irvine Skin Biology Resource Center, University of California, Irvine, CA 92697, USA

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

Summary

The skin, the largest organ, functions as a primary defense mechanism. Epidermal stem cells supply undifferentiated keratinocytes that differentiate as they migrate toward the outermost skin layer. Although such a replenishment process is disrupted in various human skin diseases, its underlying mechanisms remain elusive. With high-resolution live imaging and in vivo manipulations, we revealed that Notch signaling between keratinocytes is mediated by signaling filopodia called cytonemes and is essential for proper keratinocyte differentiation and proliferation. Inhibiting keratinocyte cytonemes reduced Notch expression within undifferentiated keratinocytes, leading to abnormal differentiation and hyperproliferation, resembling human skin disease phenotypes. Overproduction of Interleukin (IL)-17 signal, associated with skin diseases like *psoriasis*, induces psoriatic phenotypes via cytonemes in zebrafish. Our study suggests that intercellular signaling between keratinocytes through cytonemes is critical for epidermal maintenance, and its misregulation could be an origin of human skin diseases.

eLife assessment

This study provides **valuable** evidence that differentiated cells of the zebrafish skin form membrane protrusions called cytonemes, that contact and potentially transmit Notch signals to cells of the intermediate layer below. Evidence that periderm cells send out cytoneme-like protrusions is **solid**, and perturbations that affect cytoneme number clearly affect periderm structure and gene expression. However, evidence that these effects are directly due to cytoneme mediated-Notch signaling is **incomplete**.

Introduction

The vertebrate skin is a multilayered structure comprised of three main layers: the epidermis, the dermis, and the hypodermis (Akat et al., 2022 [↗](#)). Each layer has distinct anatomical characteristics and serves specific functions. Of these layers, the epidermis, as the outermost layer, plays a pivotal role as the body's first line of defense against environmental threats such as pathogens. It also plays a crucial role in regulating body temperature and preventing water loss into the surrounding environment. The epidermis consists of various layers, including the stratum basale, which houses a single layer of stem cells, and several layers of differentiated cells, including the stratum spinosum, stratum granulosum, and stratum lucidum. The outermost layer, the stratum corneum, primarily comprises dead keratinocytes that are continuously shed and replenished by underlying cells differentiated from epidermal stem cells in mammals (Segre, 2006 [↗](#)). In zebrafish, the epidermis shares a similar composition to that of mammals. However, unlike mammals, aquatic animals lack a cornified layer and instead rely on mucous layer to protect the epidermis in the aquatic environment. The basal layer of zebrafish, equivalent to mammalian stratum basale, consists of stem cells that provide keratinocytes for replenishment during epidermal remodeling or wound healing. The periderm, comparable to the mammalian stratum granulosum and lucidum, is where fully differentiated keratinocytes are found. Similar to mammalian stratum granulosum, an intermediate layer is present between the basal layer and the periderm, housing undifferentiated keratinocytes (Chang and Hwang, 2011 [↗](#)). This intermediate layer is formed during metamorphosis and persists throughout adulthood in zebrafish (Lee et al., 2014 [↗](#)).

The maintenance of a healthy epidermal barrier requires on the intricately controlled proliferation and differentiation of keratinocytes, which are supplied from underlying basal stem cells (Blanpain and Fuchs, 2009 [↗](#)). As keratinocytes migrate apically from the stem cell layer, they undergo differentiation while concurrently exhibit proliferation within the intermediate layer. Ultimately, fully differentiated keratinocytes replenish the periderm. This replenishment mechanism is well-conserved in both zebrafish and humans, and its mis-regulation constitutes a pivotal factor in the etiology of various human skin diseases, including *psoriasis*, *atopic dermatitis*, and others (Chang and Hwang, 2011 [↗](#); Lowes et al., 2014 [↗](#); Nowell and Radtke, 2013 [↗](#); Schempp et al., 2009 [↗](#); Zhou et al., 2022 [↗](#)). Studies have also revealed that maintaining a healthy epidermis requires complex intercellular communications between keratinocytes and the immune system, adding complexity to our efforts to comprehensively understand the mechanism behind this process (Blanpain and Fuchs, 2009 [↗](#); Park et al., 2021 [↗](#)).

One of the major signaling pathways critical for keratinocyte differentiation and proliferation is the Notch pathway. Studies have demonstrated that Notch receptors and ligands exhibit spatially restricted expression patterns in layers in epidermis (Nowell and Radtke, 2013 [↗](#)). Notch 1, 2, and 3 are notably prevalent in the undifferentiated keratinocytes in mice (Blanpain et al., 2006 [↗](#); Nickoloff et al., 2002 [↗](#); Thelu et al., 2002 [↗](#)). Moreover, dysregulated Notch signaling has been linked to abnormal keratinocyte differentiation and hyperproliferation, which are one of the hallmarks of various skin diseases including *psoriasis* and *atopic dermatitis* (Gratton et al., 2020 [↗](#)), highlighting the importance of proper Notch signaling in epidermal maintenance to retain the balance between keratinocyte differentiation and proliferation (Ota et al., 2014 [↗](#); Rangarajan et al., 2001 [↗](#)). Nonetheless, the precise mechanisms underlying the initiation of Notch signaling in undifferentiated keratinocytes and its dynamic regulations are not yet understood.

Furthermore, various combinations of cytokines produced by different immune cells under inflammatory conditions can significantly contribute to the onset or progression of these skin diseases. Among these, interleukin-17 (IL-17) is a well-characterized cytokine closely associated with the symptoms of these conditions. The overproduction of IL-17 by T helper (Th)17 cells and

other immune cells is a primary driver of certain skin diseases, such as *psoriasis*. Biologic treatments that inhibit IL-17 have proven to effectively alleviating the symptoms of these conditions (Mosca et al., 2021 [DOI](#)).

Unfortunately, discontinuing these treatments often lead to symptom recurrence. Also known side effects associated with these inhibitors include headaches, nasopharyngitis, and infections (Campa et al., 2016 [DOI](#)). Consequently, a definitive cure remains elusive currently (Fragoulis et al., 2016 [DOI](#); Lowes et al., 2014 [DOI](#)).

Thus, IL-17 is one of the key molecules that contribute to the development of *psoriasis* and other skin diseases. The IL-17 pathway exerts its effects on keratinocytes, endothelial cells, and immune cells resulting in the production of additional cytokines that promote positive feedback loops, sustaining the inflammatory state (Mosca et al., 2021 [DOI](#)). However, the underlying cellular and molecular mechanisms of this process are not yet fully understood. Therefore, gaining a comprehensive understanding of the interplay between keratinocytes, the Notch pathway, IL-17, and other players is essential for advancing the treatment of these diseases.

Intercellular signaling plays a critical role in skin development and maintenance. In recent years, a growing body of evidence has demonstrated that cells can establish communication through long, thin cellular protrusions extended by cells involved in either sending or receiving signals. These protrusions can be categorized based on their cytoskeletal composition, mode of signal delivery, morphology, and other characteristics. Examples include cytonemes, airinemes, tunneling nanotubes, intercellular bridges, migrasomes, exophers, and more (Eom, 2020 [DOI](#); Kornberg and Roy, 2014b [DOI](#); Ma et al., 2015 [DOI](#); Melentijevic et al., 2017 [DOI](#); Zhang and Scholpp, 2019 [DOI](#)). These specialized cellular protrusions have been observed in diverse cell types and species, ranging from fruit flies and sea urchins to zebrafish and mice, with their signaling roles experimentally confirmed *in vivo* (Daly et al., 2022 [DOI](#); Eom, 2020 [DOI](#); Hall et al., 2023 [DOI](#); Kornberg and Roy, 2014a [DOI](#); Zhang and Scholpp, 2019 [DOI](#)). These specialized protrusions are significantly longer than typical filopodia and establish direct contact with target cells and serve as highways for transporting signaling molecules for major signaling pathways such as Hedgehog, Wnt, TGF β , FGF, Notch in many *in vivo* and *in vitro* contexts (Daly et al., 2022 [DOI](#)). These signaling cellular protrusions temporally exist and have thin actin or actin/tubulin-based filaments. Consequently, live-imaging is considered one of the most effective methods for observing these structure. For that reason, zebrafish has emerged as an excellent model system for studying these cellular protrusions. The transparent nature of their early developing embryos makes zebrafish particularly well-suited for this purpose. Moreover, zebrafish are vertebrates that share common mechanisms with mammals in terms of epidermal remodeling and maintenance (Eisenhoffer et al., 2017 [DOI](#); Li et al., 2011 [DOI](#); Martinez-Navarro et al., 2019 [DOI](#)).

In this study, we show evidence that cytonemes extended by fully differentiated keratinocytes play a crucial role in activating Notch signaling in undifferentiated keratinocytes, thus contributing to the maintenance of epidermal homeostasis and remodeling in zebrafish. Furthermore, we demonstrate that IL-17, a key player in the pathogenesis of human skin diseases, and *clint1*, a gene required for epidermal homeostasis, regulate keratinocyte cytonemes (Dodd et al., 2009 [DOI](#); Sahlen et al., 2021 [DOI](#)). Collectively, our findings shed light on the mechanisms that govern the regulation of skin replenishment through keratinocyte cytonemes, providing a novel perspective on understanding how skin diseases can originate from keratinocytes autonomously through the mediation of cellular protrusions.

Results

Differentiated keratinocytes extend cytoneme-like cellular protrusions

In previous studies, we have reported a unique type of signaling cellular protrusions known as ‘airinemes’ and have investigated their signaling roles in pigment cells of zebrafish (Bowman et al., 2023 [↗](#); Eom, 2020 [↗](#); Eom et al., 2015 [↗](#); Eom and Parichy, 2017 [↗](#); Park et al., 2022 [↗](#)). To extend our understanding of airinemes in other cell types, we employed random cell labelling and identified several cell types that extend airineme-like protrusions featured by highly curved filaments and large vesicles at their tips (Eom, 2020 [↗](#); Eom et al., 2015 [↗](#); Lee et al., 2014 [↗](#)). Fully differentiated keratinocytes, marked by *krt4*, were among the cell types displaying these airineme-like protrusions (Lee et al., 2014 [↗](#)) (Figure 1A [↗](#), arrowhead). These keratinocytes intriguingly also extend cytoneme-like protrusions, characterized by relatively straight filaments and a lack of vesicle at the tips (Eom, 2020 [↗](#); Kornberg and Roy, 2014b [↗](#); Zhang and Scholpp, 2019 [↗](#)) (Figure 1A, B [↗](#), arrowheads). Nevertheless, we observed a low incidence of airineme-like protrusions, contrasting with the more frequent occurrence of cytoneme-like structures in these keratinocytes (Figure 1C [↗](#)). Consequently, we conducted a more in-depth characterization of these cytoneme-like protrusions. It is noted that cytonemes primarily consist of actin filaments, whereas airinemes are both actin and tubulin-based structures (Eom, 2020 [↗](#)). To affirm their nature, we utilized an actin marker, *LifeAct-mRuby*, which clearly labeled these cytoneme-like protrusions while showing no co-localization with *tubulin-mCherry* (Figure 1D, E [↗](#)). Additionally, cytoneme-like protrusions were significantly reduced with actin inhibitor treatment (Figure S1). Therefore, based on their cytoskeletal composition and morphology, we classified them as cytonemes.

We observed that keratinocytes marked by *krt4* expression extend cytonemes with the highest frequency during post-embryonic stages, specifically between 6.5 to 7.5 SSL (Standardized Standard Length (Parichy et al., 2009 [↗](#))) (Figure 1C [↗](#)). It is worth noting that embryonic keratinocytes are replaced by post-embryonic cells around these stages, suggesting the potential role of cytonemes in epidermal remodeling and maintenance (Lee et al., 2014 [↗](#); Parichy et al., 2009 [↗](#)). These keratinocyte cytonemes exhibit an approximate speed of 2.98µm/min during extension and retract at 1.9µm/min, with an average length of 18.21µm (Figure 1F, G [↗](#)).

Keratinocyte cytonemes establish physical contact with underlying undifferentiated keratinocytes

Similar to mammals, the zebrafish epidermis consists of three layers (Chang and Hwang, 2011 [↗](#)). We specifically visualized the periderm, which is the outermost layer and consists of *krt4*⁺ fully differentiated keratinocytes. The intermediate layer, where undifferentiated keratinocytes reside, was marked with *krtt1c19e* promoter driving tdTomato expression, pseudo-colored in magenta (Chang and Hwang, 2011 [↗](#); Lee et al., 2014 [↗](#)). The basal layer was also labelled with *krtt1c19e* but exhibited a typical polygonal morphology (Figure 2A-A'' [↗](#)). This allowed us to distinguish all three epidermal layers using the *Tg(krt4:lyn-EGFP; krtt1c19e:lyn-tdTomato)*.

In addition to *krt4*⁺ keratinocytes, we explored whether undifferentiated keratinocytes (*krtt1c19e*⁺) also extend cytonemes or other types of cellular protrusions during metamorphic stages. However, we observed that cytoneme-like protrusions from undifferentiated keratinocytes were significantly fewer in numbers compared to those from fully differentiated keratinocytes, and we did not observe cytoneme-like protrusions from basal stem cells (Figure 2B [↗](#)). In this study, our focus was primarily on cytonemes from *krt4*⁺ differentiated keratinocytes.




Figure 1.

Differentiated keratinocytes extend cytoneme-like cellular protrusions

(A) Airinome-like protrusion (arrowhead), and (B) cytoneme-like protrusions (arrowheads) labelled in *krt4:palmeGFP* injected keratinocytes. (C) Keratinocytes expressing *krt4* extend cytoneme-like protrusions most frequently during metamorphic stages [6.5-7.5 SSL; N=457 cells, 19 larvae total]. A cytoneme-like protrusion (arrowhead) labeled for (D) F-actin, revealed by *LifeAct-mRuby* but did not colocalize (E) with tubulin, revealed by *Tuba-mCherry*. (F) Keratinocyte cytonemes extend faster than they retract (N=125 cells, 4 larvae total), with an average length (G) of 18.21μm (N=190 cells, 4 larvae total). Scale bars: 20μm (A, B, D, E). Error bars indicate mean ± SEM.




Figure 2.

Keratinocyte cytonemes establish physical contact with underlying undifferentiated keratinocytes

(A-A'') Postembryonic zebrafish epidermis of *Tg(krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato)* comprises three distinct layers. (A) Expression of *lyn-EGFP* in *krt4*⁺ cells in the periderm layer. (A'-A'') Expression of *lyn-tdTomato* (pseudo-colored magenta) in *krtt1c19e*⁺ cells in (A') the intermediate layer and (A'') the basal layer. Note that the green circular labelling in (A', arrowheads) represents mucous-secreting goblet cells. (Chang and Hwang, 2011) (B) Undifferentiated keratinocytes (KC) extend significantly fewer cytoneme-like protrusions ($P < 0.0001$, $N = 49$, 3 larvae; $N = 91$ cells, 6 larvae total). (C) Cytonemes from fully differentiated keratinocytes make physical contact with undifferentiated keratinocytes in the intermediate layer (red dotted circles). Note that dotted lines indicate the layers in the cross-sectional views. Statistical significances were assessed by Student *t* test. Scale bars: 20 μ m (A-A'', C). Error bars indicate mean $\pm$ SEM.

To investigate the role of keratinocyte cytonemes, we first asked what the target cells of those cytonemes are. High resolution confocal imaging revealed that cytonemes originating from peridermal keratinocytes (*krt4+*) establish physical contact with underlying undifferentiated keratinocytes (*krtt1c19e+*) in the intermediate layer. Cross-sectional views showed that the tips of these cytonemes stop at the intermediate layer, with no observed instances reaching the basal layer (**Figure 2C** [↗](#), red dashed circles).

Cytoneme-mediated signaling regulates keratinocyte differentiation and proliferation

The observations that cytonemes most often observed during epidermal remodeling and their target cells are undifferentiated keratinocytes led us to hypothesize that cytoneme-mediated signaling play a pivotal role in regulating keratinocyte differentiation and proliferation.

To test, we generated a transgenic line capable of temporally expressing a dominant negative form of *cdc42* (*cdc42DN*) in peridermal keratinocytes, a strategy frequently used to inhibit cytoneme extension in various contexts ([Daly et al., 2022](#) [↗](#); [Zhang and Scholpp, 2019](#) [↗](#)). We implemented a dually inducible TetON system in the transgene, denoted as *Tg(krt4:TetGBDTRE-v2a-cdc42DN)*, and immersed the transgenic and control fish in water containing inducer drugs, doxycycline and dexamethasone, for temporal *cdc42DN* transgene induction ([Eom et al., 2015](#) [↗](#); [Knopf et al., 2010](#) [↗](#)). To minimize potential off-target effects of *cdc42* manipulation, we determined the drug concentrations at which normal filopodial or lamellipodial activities were maintained while inhibiting cytoneme extensions (Video S1). We employed two control groups in this experiment: one consisting of a transgenic line with no drugs and the other comprising non-transgenic fish treated with the same drug concentrations as the experimental group. Under these conditions, we observed a significant reduction in the frequency of cytoneme extension in the *cdc42DN*-expressing transgenic line within the drug treated group (**Figure 3A** [↗](#)). In contrast to the two control groups, the peridermal layer in the cytoneme-inhibited group displayed severe disorganization, with areas devoid of *krt4+* keratinocytes (**Figure 3B'** [↗](#), arrowhead). Furthermore, the expression of the undifferentiated keratinocyte marker, *krtt1c19e* (magenta), was dramatically increased in the periderm when cytoneme extension was inhibited (**Figure 3B-B', C** [↗](#)). Additionally, we found a substantial increase in the number of keratinocytes within the peridermal layer in the cytoneme-inhibited group (**Figure 3D** [↗](#)). However, it is important to note that cytoneme inhibition did not affect the cell death rate or the number of goblet cells (**Figure 3E** [↗](#), **2A'** [↗](#), Figure S2A). These findings strongly suggest that cytoneme-mediated signaling controls keratinocyte differentiation and proliferation.

Cytonemes activate Notch in undifferentiated keratinocytes

Previous studies have established the role of Notch signaling in controlling keratinocyte differentiation and proliferation. However, the mechanisms triggering Notch signaling in keratinocytes have remained obscure ([Moriyama et al., 2008](#) [↗](#); [Nguyen et al., 2006](#) [↗](#); [Rangarajan et al., 2001](#) [↗](#)). To investigate whether the target cells of cytonemes, specifically undifferentiated keratinocytes in the intermediate layer, are responsive to Notch signaling, we crossed Notch signal reporter line, *Tg(Tp1:H2B-mCherry)*, with *Tg(krt4:lyn-EGFP; krtt1c19e:lyn-Tdtomato)* to label all three epidermal layers. In this Notch reporter line, the promoter from the Epstein Barr Virus terminal protein 1 (*Tp1*) gene was used as a Notch responsive element, containing two Rbp-Jk binding sites that drive expression of Histone 2B-Cherry ([Parsons et al., 2009](#) [↗](#)). We observed that only undifferentiated keratinocytes in the intermediate layer exhibit Notch responsiveness, whereas keratinocytes in the periderm or basal layer do not (**Figure 4A** [↗](#), arrowheads, Figure S3).

Subsequently, we tested whether Notch activation in undifferentiated keratinocytes is dependent on cytonemes. First, we validated the *Tp1* Notch reporter line, *Tg(Tp1:EGFP)* by measuring cytoplasmic EGFP signal before and after Notch inhibitor, LY411575 treatment, which




Figure 3.

Cytoneme-mediated signaling regulates keratinocyte differentiation and proliferation

(A) Expression of *cdc42DN* effectively inhibits cytoneme extension ($F_{2, 231} = 25.38$, $P < 0.0001$, $N = 234$ cells, 10 larvae total). (B) Periderm layer of Tg(*krt4:TetGBDTRE-v2a-Cdc42DN*; *krt4:lyn-EGFP*; *krtt1c19e:lyn-tdTomato*) without drug treatment. (B') Periderm layer of Tg(*krt4:lyn-EGFP*; *krtt1c19e:lyn-tdTomato*) with drug treatment. (B'') Periderm layer of Tg(*krt4:TetGBDTRE-v2a-cdc42DN*; *krt4:lyn-EGFP*; *krtt1c19e:lyn-tdTomato*) with drug treatment exhibits disorganization and regions lacking *krt4*+ keratinocytes (arrowhead). Cytoneme inhibition increases *krtt1c19e* expression in the periderm (C) ($F_{2, 387} = 226.7$, $P < 0.0001$, $N = 39$ larvae total) and leads to an increased number of keratinocytes within the periderm (D) ($F_{2, 18} = 27.36$, $P < 0.0001$, $N = 21$ larvae total). (E) The cell death rate in the periderm is not affected by cytoneme inhibition, tested with acridine orange incorporation ($F_{2, 17} = 3.192$, $P = 0.0666$, $N = 20$ larvae total). Statistical significances were assessed by One-way ANOVA followed by Tukey's HSD post hoc test. Scale bars: 20 μ m (B-B''). Error bars indicate mean $\pm$ SEM.




Figure 4.

Cytonemes activate Notch in undifferentiated keratinocytes

(A) Postembryonic zebrafish epidermis of *Tg(krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato;TP1:H2BmCherry)* shows Notch responsiveness in undifferentiated keratinocytes within the intermediate layer but not in the periderm or basal layer. (B) Inhibition of cytonemes reduces Notch responsiveness in undifferentiated keratinocytes. ($F_{2,33} = 48.27$, $P < 0.0001$, $N=36$ larvae total). (C) Expression of DeltaC-mCherry fusion protein along the cytoneme (yellow arrowheads). (D-D'') Notch inhibition by expressing SuHDN in undifferentiated keratinocytes results in the disorganization of periderm (D''). (C) Properly organized periderm layer of *Tg(krtt1c19e:tetGBDTRV2a-SuHDN;krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato)* without drug treatment and (C') *Tg(krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato)* with drug treatment. (E-G) Notch inhibition (E) increases *krtt1c19e* signal in the periderm ($F_{2,397} = 89.47$, $P < 0.0001$, $N=40$ larvae total), (F) reduces Notch responsiveness in undifferentiated keratinocytes ($F_{2,32} = 13.18$, $P < 0.0001$, $N=35$ larvae total), and leads to (G) an increased number of keratinocytes in the periderm ($F_{2,19} = 6.620$, $P = 0.0066$, $N=22$ larvae total). Statistical significances were assessed by One-way ANOVA followed by Tukey's HSD post hoc test. Scale bars, 20 μ m (A, C, D). Error bars indicate mean $\pm$ SEM.

demonstrated a significant decrease in Notch signaling, as expected (Figure S4A, B). Next, we measured the difference in *Tp1* intensity before and after cytoneme inhibition. Notably, we observed a significant reduction in Notch signal under conditions where cytonemes were compromised (Figure 4B, Figure S4C). We also found more abundant Notch activated keratinocytes during metamorphic stages (Figure S4D, Figure 1C).

Next, we predicted that if cytonemes contribute to transmitting a Notch-Delta signal from differentiated keratinocytes to undifferentiated keratinocytes, then cytonemes should contain Delta ligands. Upon detecting DeltaC mRNA expression in the cytoneme extending keratinocytes (Fig. S3), we investigated the localization of DeltaC protein within the cytonemes of the *krt4+* differentiated keratinocytes. For this experiment, we utilized a construct where a 109kb BAC containing zebrafish *dlc* coding sequence regulatory elements that were recombineered to generate a mCherry fusion, resulting *dlc:Dlc-mCherry* (Eom et al., 2015). We confirmed the presence of Dlc-mCherry along the cytonemes (Fig. 4C, yellow arrowheads). These findings suggest that cytonemes are responsible for Notch activation in undifferentiated keratinocytes.

We then asked why Notch activation is dependent on cytonemes since peridermal keratinocytes and underlying intermediate keratinocytes are in constant contact with each other. We revealed that intermediate keratinocytes exclusively express *lfn* (lunatic fringe), which potentially act as a default inhibitor of Notch activation in these cells (Dale et al., 2003; Panin et al., 1997) (Figure S5A, S3). However, this needs further studies to understand how cytonemes overcome this Notch signaling barrier or potentiate the signal.

Next, we hypothesized that if Notch activation by cytonemes is essential for keratinocyte differentiation and proliferation, then direct inhibition of Notch signal in undifferentiated keratinocytes should yield phenotypes similar to those observed when cytonemes are inhibited (Gratton et al., 2020). To investigate this, we generated a transgenic line capable of temporally expressing dominant negative form of Suppressor of Hairless (SuHDN) in the undifferentiated keratinocytes, *Tg(krtt1c19e:TetGBDTRE-v2a-SuHDN)* (Eom et al., 2015). We observed that Notch inhibition in the undifferentiated keratinocytes resulted in a disorganized periderm, abnormally increased expression of undifferentiated keratinocyte marker, and hyperproliferation of keratinocytes in the periderm (Figure 4D-D", E, F, G, Figure S4C). These results closely mirrored the phenotypes observed when we inhibited cytoneme extension in peridermal keratinocytes (Figure 3B-B", C, D, Figure S4C). These observations suggest that Notch signaling, activated by cytonemes in undifferentiated keratinocytes, plays a critical role in epidermal remodeling and maintenance.

It has been documented that both hypo- or hyper-activation of Notch signaling can lead to the onset of skin diseases (Abdou et al., 2012; Gratton et al., 2020). Thus, we further investigated whether gain-of-function experiments for Notch also disrupt epidermal homeostasis. We overexpressed the Notch IntraCellular Domain (NICD) in undifferentiated keratinocytes, using the *Tg(krtt1c19e:nVenus-v2a-NICD)*. Consistent with previous studies in mammals and the cytoneme or Notch loss-of-function experiments, we observed disorganized periderm, abnormal keratinocyte differentiation and hyper-proliferation (Figure S5B-B', C, D). These results imply that uncontrolled Notch signaling in keratinocytes could lead to disruption of epidermal remodeling and maintenance in both zebrafish and mammals.

Interleukin-17 regulates keratinocyte cytonemes

The aberrant differentiation and hyperproliferation of keratinocytes observed when cytoneme extension was inhibited in peridermal keratinocytes (*krt4+*) or Notch signaling was inhibited in undifferentiated keratinocytes (*krtt1c19e+*) are hallmark features of many human skin diseases such as *psoriasis*, *atopic dermatitis* and more (Armstrong and Read, 2020; Gratton et al., 2020) (Figure 3, 4). Also, it has been well established that impaired Notch signaling is associated

with some of these human skin disorders (Gratton et al., 2020 [↗](#); Nowell and Radtke, 2013 [↗](#)). Thus, we asked which key molecules are critical for the onset or progression of these skin diseases and whether they have an impact on keratinocyte cytoneme signaling.

One of them is interleukin-17 (IL-17) and we hypothesized IL-17 signaling could influence cytoneme extension in peridermal keratinocytes. Given that IL-17 is a cytokine released into the extracellular space, and to test its role in peridermal keratinocytes, we decided to overexpress the IL-17 receptors. It is noted that interleukin-17 receptors and ligands are evolutionarily conserved across chordates (Gonzalez-Fernandez et al., 2020 [↗](#); Wu et al., 2011 [↗](#)).

To test, we generated transgenic lines that cell-autonomously overexpressed il17 receptor D (*il17rd*) or il17 receptor A1a (*il17ra1a*) in fully differentiated peridermal keratinocytes (*krt4+*), *Tg(krt4:nVenus-v2a-il17rd)* and *Tg(krt4:il17ra1a-v2a-mCherry)*. Both of these IL-17 receptor-overexpressing transgenic lines exhibited a significant reduction in cytoneme extension, resulting in a decreased Notch activation in undifferentiated keratinocytes (**Figure 5A, B** [↗](#)). Moreover, similar to the two previous manipulations, we observed a severely disorganized periderm, an upregulation of the undifferentiated keratinocyte marker expression, and hyperproliferation of keratinocytes within the periderm (**Figure 5C-C', D, E** [↗](#), Figure S3).

To assess whether IL-17 signaling regulates keratinocyte differentiation and proliferation through cytonemes, first we overexpressed the il17a ligand in the epidermis and examined its effect on cytoneme extension. Given the abundance of tissue-resident macrophages in the epidermis, we generated a transgenic line that overexpressed *il17a* under a macrophage promoter, *Tg(mpeg1:il17a-v2a-mCherry)* (Bowman et al., 2023 [↗](#)). Consistent with the experiments that overexpressed il17 receptors, we confirmed that il17a ligand overexpression in the epidermal microenvironment led to a significant reduction in cytoneme extension in the differentiated keratinocytes (**Fig. 5A** [↗](#), **6A** [↗](#)). Next, we generated an *il17a* loss-of-function mutant using CRISPR/Cas9 approach. This mutant harbor an 11bp deletion in the second exon of the *il17a* gene, resulting in a premature stop codon (Fig. S6). We observed a significant increase in cytoneme extension frequency in *il17a* heterozygote mutants as compared to their wild-type siblings (**Fig. 6B** [↗](#)). Together, our findings suggest that IL-17 signaling controls cytoneme extension.

Subsequently, we sought to understand how the IL-17 signal impacts cytoneme extension. Studies have shown that IL-17 signaling can regulate actin cytoskeleton in keratinocytes (Borowczyk et al., 2020 [↗](#); Das et al., 2019 [↗](#)). To evaluate whether *il17rd* overexpression alters the actin cytoskeleton, thus cytonemes, we compared keratinocyte microridges, which are actin cytoskeleton-rich structures, between control and *il17rd*-overexpressed peridermal keratinocytes (Pinto et al., 2019 [↗](#); van Loon et al., 2020 [↗](#)). Our observation revealed severe disorganization of microridges in the *il17rd* overexpressed keratinocytes, coupled with a significant down-regulation of Cdc42 GTPase expression in these cells (**Figure 6C, D** [↗](#)). These data suggest that IL-17 signaling negatively regulates cytoneme extension by down-regulating Cdc42 that are essential in control of actin cytoskeleton (Daly et al., 2022 [↗](#); Hall, 1998 [↗](#); Jaffe and Hall, 2005 [↗](#); Stanganello et al., 2015 [↗](#)).

We then crossed two transgenic lines: *Tg(krt4:TetGBDTRE-v2a-cdc42DN)* and either *Tg(krt4:nVenus-v2a-il17rd)* or *Tg(krt4:il17ra1a-v2a-mCherry)*. Given that il17 receptor overexpression reduced Cdc42 expression in differentiated keratinocytes, we anticipated observing a more severe inhibition of cytoneme extension when these manipulations are combined compared to individual manipulations (**Fig. 6C** [↗](#)). Indeed, when inhibited cytoneme extension with the cdc42DN along with *il17rd* or *il17ra1a* overexpression, we observed a more significant inhibition of cytoneme extension than when manipulated individually (**Figure 6E** [↗](#)).




Figure 5.

Disruption of epidermal maintenance by Interleukin-17 receptor overexpression

Overexpression of *il17rd* or *il17ra1a* in *krt4+* keratinocytes results in (A) a significant reduction in cytoneme extension ($F_{2, 325} = 38.48$, $P < 0.0001$, $N=328$ cells, 10 larvae total), and (B) decreased Notch responsiveness ($F_{2, 34} = 23.16$, $P < 0.0001$, $N=37$ larvae total). (C-C') The periderm is disrupted in (C') *Tg(krt4:nVenus-v2a-il17rd;krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato)* compared to the (C) properly arranged periderm of *Tg(krt4:lyn-EGFP;krtt1c19e:lyn-tdTomato)*, resulting in (D) an increased *krtt1c19e* signal in the periderm (D) ($F_{2, 287} = 109.7$, $P < 0.0001$, $N=29$ larvae total) and (E) an increased number of keratinocytes in the periderm ($F_{2, 15} = 27.79$, $P < 0.0001$, $N=18$ larvae total). Statistical significances were assessed by One-way ANOVA followed by Tukey's HSD post hoc test. Scale bars: 20 μ m (C). Error bars indicate mean $\pm$ SEM.




Figure 6.

Interleukin-17 regulates keratinocyte cytonemes

(A) Significant reduction in cytoneme extension in response to overexpressed *il17a* ligands in the epidermis ($P < 0.0001$, $N = 3$ larvae per group). (B) Cytoneme extension frequency increased significantly in *il17a*^{+/-} mutants as compared to their wild-type siblings ($P = 0.0009$, $N = 4$ larvae for wild-type, $N = 3$ for mutants). (C) FACS sorted *krt4*⁺ peridermal cells show significantly reduced *Cdc42* expression ($P = 0.001$, $N = 5$ independent experiments). (D) *il17rd*-overexpressing *krt4*⁺ keratinocytes exhibit disorganization of microridges (arrowheads). (E) Simultaneous overexpression of *cdc42DN* and *il17rd* or *il17ra1a* in *krt4*⁺ keratinocytes further reduces cytoneme extension (*il17rd*: $P < 0.0001$, $N = 164$ cells, 7 larvae total, *il17ra1a*: $P < 0.0001$, $N = 115$ cells, 6 larvae total). Statistical significances were assessed by Student t test. Scale bars: 20 μ m (C, G). Error bars indicate mean $\pm$ SEM.

Our data collectively support the idea that cytoneme-mediated intercellular signaling between peridermal and underlying undifferentiated keratinocytes is essential for epidermal remodeling and maintenance. IL-17 functions as a regulator capable of affecting cytoneme extension by modulating the actin cytoskeleton in keratinocytes. Consequently, our data suggest that the overproduction of IL-17 ligands by immune cells in conditions such as psoriatic or other skin diseases leads to epidermal imbalance through altered cytoneme signaling in keratinocytes.

In addition, recent human Genome-Wide Association Studies (GWAS) have identified a close association between *clint1* and individuals with *psoriasis* and *atopic dermatitis* (Sahlen et al., 2021 [↗](#)). It has also been reported that *clint1* is important for epidermal homeostasis and development in zebrafish (Dodd et al., 2009 [↗](#)). Given these reported findings, we investigated to determine whether *clint1* plays a role in epidermal maintenance through its influence on keratinocyte cytonemes. Indeed, we found that *clint1* mutants exhibited a significant decrease in cytoneme extension in peridermal keratinocytes, accompanied by reduced Notch signaling in undifferentiated keratinocytes. As a result, these mutant fish displayed abnormal keratinocyte differentiation and hyperproliferation of keratinocytes (Figure S7). These findings suggest that *clint1* is yet another regulator of cytoneme-mediated intercellular signaling in keratinocytes.

In conclusion, the intercellular signaling facilitated by cytonemes between fully differentiated and undifferentiated keratinocytes is crucial for the maintenance and remodeling of the epidermis. Our study implies that the patients with human skin diseases may have defective cytoneme signaling between keratinocytes.

Discussion

It is intriguing to observe that keratinocytes have the ability to extend both cytoneme and airneme-like protrusions. While these protrusions differ in their extension frequency and developmental context, this observation implies that a single cell type can employ two distinct types of cellular protrusions for signaling, each capable of transferring unique signaling molecules to potentially different targets. Previous research in *Drosophila* has demonstrated that the air sac primordium (ASP) can extend multiple cytonemes that appear morphologically identical but convey different signaling molecules (Roy et al., 2011 [↗](#)). However, it remains unknown whether the same cell type can extend two morphologically (and functionally) distinct protrusions that might also serve distinct functions in separate contexts. Consequently, future studies aimed at unraveling the regulation of these two types of protrusions could provide insights into how cells choose their signaling modalities based on their specific signaling requirements and environmental cues.

We have asked why cytoneme-mediated signaling is necessary between fully differentiated- and undifferentiated keratinocytes, even though they are separated by opposing cell membranes. Notably, we have identified the Notch signaling pathway as being involved, which operates through membrane receptor and ligand interactions between adjacent cells. This led us to hypothesize that Notch signaling between cells in these layers is suppressed by Fringe proteins, which modulate Notch signaling. In our research, we have discovered that *lunatic fringe* (*lfng*) is exclusively expressed in the intermediate layer in zebrafish ((Panin et al., 1997 [↗](#)), Figure S3). Early studies in mice have also reported that three fringe genes exhibit differential expression in different epidermal layers (Thelu et al., 1998 [↗](#)). Hence, it appears that Notch signaling is tightly regulated or inhibited by default between epidermal layers in both zebrafish and mice.

While it is well documented that Notch signaling is essential for keratinocyte differentiation and proliferation, the mechanism by which Notch signaling is initiated remains unclear. Our study suggests that cytonemes from fully differentiated keratinocytes in the periderm can activate Notch

signaling in undifferentiated keratinocytes in intermediate layer. However, further investigations are necessary to fully comprehend how cytoneme-mediated signaling coordinate with Lunatic fringe or other Fringes to trigger the Notch signal.

It is widely believed that the overproduction of IL-17 is a major contributing factor to *psoriasis*, and many studies have demonstrated that IL-17 signaling triggers inflammation (Liu et al., 2020 [↗](#)). IL-17 affects various cellular targets, such as keratinocytes, neutrophils, endothelial cells, promoting tissue inflammation (Blauvelt and Chiricozzi, 2018 [↗](#)). Thus, further studies are essential to determine whether uncontrolled cytoneme-mediated signaling in keratinocytes correlates with the onset or progression of psoriatic or other skin diseases. Nevertheless, our data strongly indicate that cell-autonomous overexpression of *il17rd* in peridermal keratinocytes (*krt4+*) can induce psoriasis-like phenotypes by inhibiting cytoneme extensions, which subsequently leads to a reduction in Notch signaling in undifferentiated keratinocytes. A diminished Notch signal has been observed in psoriatic lesions (Ota et al., 2014 [↗](#)). Our study suggests that cytoneme-mediated intercellular communication between keratinocytes plays a crucial role in epidermal maintenance, and its dysregulation may contribute to the development of human skin diseases. This idea has yet to be fully appreciated, and it could open up new avenues for understanding the onset or progression of human skin diseases.

Moreover, we have demonstrated that IL-17 can influence cytoneme extension by regulating Cdc42 GTPases, ultimately affecting actin polymerization. Consequently, it would be intriguing to investigate whether genes associated with *psoriasis* or other human skin diseases also function as regulators of cytonemes. For instance, the epidermal-specific deletion of inhibitor of NF- κ B kinase 2 (IKK2) or the double knock out of the c-Jun and JunB genes in mice resulted in psoriasis-like phenotypes in a T cell-independent manner (Pasparakis et al., 2002 [↗](#); Zenz et al., 2005 [↗](#)). Among the various roles of IKK and JunB, they are also known to regulate the actin cytoskeleton (Dubin-Bar et al., 2008 [↗](#); Lennikov et al., 2018 [↗](#); Otani et al., 2016 [↗](#)). Thus, it is conceivable that dysregulated cytoneme-mediated signaling may contribute to psoriatic conditions in these contexts.

Similarly, we have shown that *clint1* mutants exhibited compromised cytoneme signaling and keratinocyte hyperproliferation (Figure S7) (Dodd et al., 2009 [↗](#)). Clint1 (Clathrin Interactor 1) plays an important role in vesicle trafficking, and it is well established that endocytic pathways are critical for multiple steps in cytoneme-mediated morphogen delivery (Kalthoff et al., 2002 [↗](#)). Therefore, it is interesting to investigate whether clint1 functions as a cytoneme regulator could provide valuable insights.

Another important question requires an answer: the proper remodeling and maintenance of the epidermis seemingly depend on a well-controlled supply of keratinocytes from basal stem cells. Our study demonstrated the relay of cytoneme-mediated signals from the periderm to the underlying undifferentiated keratinocytes, leading to their differentiation and subsequent replacement of peridermal keratinocytes. However, the mechanism through which basal stem cells detect the need for an increased population of keratinocytes to maintain the intermediate layer remains unclear. This process does not necessarily involve cellular protrusion-mediated signaling. However, we discovered cytoneme-like protrusions originating from undifferentiated keratinocytes (**Figure 2B** [↗](#)). Our preliminary data, not presented in this manuscript, suggests that these cellular protrusions project downward, indicating their interaction with basal stem cells. Although further investigation is needed, it is conceivable that a replenishment signal from external sources or the periderm can be transmitted to stem cells through a multi-step cytoneme-mediated intercellular signaling pathways.

Acknowledgements

We thank Drs. Plikus, Andersen and Parsons for the invaluable discussions, and all the members of Eom lab for maintaining our fish lines. This study was supported by the National Institutes of Health grant R35GM142791 (to D.S.E.) and National Institutes of Health T32 training grant AR080622 (to Y.W.).

Author Contributions

Conceptualization, Y.W. and D.S.E.; Investigation, Y.W., Q.H., O.H., K.F. and D.S.E.; Writing – Original Draft, Y.W. and D.S.E.; Writing – Review & Editing, Y.W. and D.S.E.; Funding Acquisition; D.S.E. and Y.W.; Supervision, D.S.E.

Competing interests

The authors declare no competing interests.

Experimental model and subject details

Zebrafish

Adult zebrafish were maintained at a constant temperature of 28.5°C under a 16:8 (L:D) cycle. The zebrafish used in this study were wild-type AB^{WP} or its derivative WT(ABb). Additionally, transgenic lines used include *Tg(krt4:lyn-EGFP)^{sq18}*, *Tg(krtt1c19e:lyn-tdTomato)^{sq16}* provided by T. Carney; *Tg(Tp1:EGFP)^{um13}*, *Tg(Tp1:H2BmCherry)^{jh11}* provided by M. Parsons; *clint1a^{hi1520Tg/+}(AB)*, obtained from the Zebrafish International Resource Center (ZIRC). The sexes of individual fish could not be distinguished since experiments were performed before the development of secondary sexual characteristics. All zebrafish stocks used in this study were known to produce balanced sex ratios, ensuring that experiments sampled similar numbers of males and females. All animal work conducted in this study received approval from the University of California Irvine Institutional Animal Care and Use Committee (protocol #AUP-22-023) and adhere to institutional and federal guidelines for the ethical use of animals.

Method details

Transgenesis and transgenic line production

Transgenic fish lines with cell-type specific and temporally inducible expression of human *cdc42DN* and *SuHDN* were generated following the protocol described by Eom et al., (2015) <https://doi.org/10.1016/j.dev.2015.05.010>. These constructs were driven by a 2.2kb *krt4* promoter obtained from T. Carney for the expression in fully differentiated keratinocytes and a 3.9kb *krtt1c19e* promoter, also from T. Carney, for the expression in undifferentiated keratinocytes and epidermal stem cells. The efficiency of gene expression in both *Tg(krt4:TetGBDTRE-v2a-cdc42DN)^{ir.rt3}* and *Tg(krtt1c19e:TetGBDTRE-v2a-SuHDN)^{ir.rt18}*, induced by doxycycline (Dox; Sigma-Aldrich) and dexamethasone (Dex; Sigma-Aldrich), was performed in both F0 mosaic fish and non-mosaic stable lines, with consistent results. The coding sequence of *il17rd* (ZDB-GENE-020320-5) and *il17ra1a* (ZDB-GENE-070705-242)

were isolated from WT(ABb) cDNA and used to generate constructs driven by the *krt4* promoter, which were assembled using Gateway assembly into the pDestTol2CG2 destination vector to produce *Tg(krt4:nVenus-v2a-il17rd)^{ir.rt16}* and *Tg(krt4:il17ra1a-v2a-mCherry)^{ir.rt17}*. Similarly, *Tg(mpeg1:il17a-v2a-mCherry)^{ir.rt21}* was produced by isolating the coding sequence of il17a (ZDB-GENE-061031-3) from WT(ABb) cDNA and using Gateway assembly to assemble the construct driven by the mpeg1 promoter into the pDestTol2-exorh:mCherry destination vector [Addgene #195989]. Constitutive Notch activation used the intracellular domain of notch1a (NICD) linked by 2a sequence to nVenus and driven by the *krtt1c19e* promoter to produce *Tg(krtt1c19e:nVenus-v2a-NICD)^{ir.rt19}*. The lunatic fringe reporter line, *Tg(lfng:mCherry)^{ir.rt20}*, was generated by first amplifying the *lfng* promoter sequence (ZDB-GENE-980605-16) using WT gDNA and then used Gateway assembly into the pDestTol2-exorh:EGFP destination vector [Addgene #195983]. These constructs were then injected into WT(ABb) fish to generate mosaic F0 fish, which were then bred in pairs to find germline carriers to produce non-mosaic transgenic lines. Actin was examined with *LifeAct-mRuby* [Addgene #166977] (Barros-Becker et al., 2017 [DOI](#)) driven by *krt4* promoter. Microtubules in *krt4+* keratinocytes and cytonemes were labelled using subclones from Tuba1-mCherry [(Addgene #49149) (Friedman et al., 2010 [DOI](#))].

Generation of *il17a* heterozygous mutants by the CRISPR/Cas9 System

Disruption of *il17a/f1* (GenBank accession number NM_001020787) was performed using CRISPR/Cas9 technology. The target site in exon 2 was selected using SMART and the corresponding sequence was 5'-AGCAGATCATTCATACCGGC-3'. Single guide RNA (sgRNA) was synthesized using MEGashortscriptTM T7 High Yield Transcription Kit (Invitrogen). Then, 1μg/μL TrueCutTM Cas9 Protein v2 (InvitrogenTM) and 300ng/μL sgRNA were co-injected into the one-cell stage zebrafish embryos to knockout the *il17a* gene. Founder fish (F0) were outcrossed with WT(ABb) to generate *il17a* heterozygous mutants, which were identified upon adulthood and used for experiments. The PCR primers for genotyping were as follows: Forward 5'-CCTCCGCTTCTTATGGTGAGTATAGC -3' and Reverse 5'-GGAACCACTGAATGCCAATATAGCAG -3'.

Drug treatments

Zebrafish were kept in E3 medium with or without drugs (Dox/Dex) during both the light and the dark cycles for long-term drug treatment experiments. For overnight time-lapse images, fish subjected to drug-treatment were transferred to fresh E3 medium 30min before explant preparation to minimize background fluorescence from the drugs.

Zebrafish with *krt4:palmEGFP* injected TetGBD-containing transgenics and non-transgenic siblings were treated with 37.5μM Dox and 50μM Dex one day prior to time-lapse imaging, while fish for long-term drug treatment received 27.5 μM Dox and 50μM Dex starting from SSL7.0. Notch inhibitor, LY411575 (10mM; Thomas Scientific) was prepared in DMSO (Sigma-Aldrich), and fish were treated with drug [LY411575 (3μM)] or vehicle from SSL7.0. For acute drug administration, Cdc42 inhibitor ML141 (Sigma-Aldrich), prepared in DMSO, was added to fish medium [ML141 (2μM)] 30min prior to explant preparation for time-lapse imaging.

Acridine Orange Cell Death Assay

Acridine Orange (5mg/mL; Sigma-Aldrich) was prepared in Invitrogen UltraPureTM Distilled Water. Fish were washed in 1x E3 medium 30min before being transferred to 1x E3 medium containing AO (2μg/mL). The fish were incubated in AO for 30min and then rinsed in regular 1x E3 medium three times for 10min each to minimize background fluorescence during imaging. Dead cells, pseudo-colored in magenta, were counted for all groups.

Time-lapse and still imaging

Ex vivo time-lapse imaging of zebrafish epidermal cells within their native tissue environment was acquired at 3-minute intervals for 10 hours. This was achieved using a 40x water-immersion objective on a Leica TCS SP8 confocal microscope equipped with a resonant scanner and two HyD detectors. Larvae were imaged at SSL7.5, unless otherwise specified. For still images, the region directly underneath the dorsal fin of the zebrafish was captured, with a total of 10 tiles per fish.

RT-PCR, qRT-PCR and FACS

Zebrafish at SSL7.5 were skinned (N=15 per group), and the dissected tissues were placed in cold 1x PBS. Tissues were washed by pipetting up and down and then centrifuged to remove the supernatant. Next, 1mL Gibco Stem[®] Pro[®] Accutase Cell Dissociation Reagent was added to resuspend tissues, and the tube was incubated for at least 10 min at 37°C to dissociate the cells. After incubation, the cells were passed through 40µm cell strainers to remove large tissue chunks before subjecting them to FACS (Fluorescence-activated cell sorting). FACS enabled the separation and collection of desired cell populations based on different cell membrane markers. EGFP⁺ and tdtomato⁺ cells were collected for downstream non-quantitative RT-PCR and quantitative RT-PCR. For cDNA synthesis, the Invitrogen Cells Direct[™] One-Step qRT-PCR Kit was employed. Quantitative PCR was conducted on an Applied Biosystems QuantStudio[™] 3 Real-Time PCR Instrument using custom TaqMan probes for *cdc42* (Dr03139234_m1) and *actin* (Dr03432610_m1). Quantitative PCRs were run with at least three triplicates for each sample. Non-quantitative RT-PCR amplifications were performed with 40 cycles (*actb1*, *dlc*, *lfng*, *krt4*, *krt19*, *il17rd*, *notch2*, *notch3*, *il17ra1a*) using Takara PrimeStar GXL DNA Polymerase. *actb1*: 5'-CATCCGTAAGGACCTGTATGCCAAC- 3', 5'- AGGTTGGTCGTTTCGTTTGAATCTC- 3' (Hu et al., 2016 [\[4\]](#)); *krt4*: 5'- TCAACCAGGTCTATCTCTTACTCC- 3', 5'-AACAGGCTCTGGTTGACAGTTAC- 3'; *krt19*: 5'-GCTACCACCTTCTCCAGCGGAAG-3', 5'- TGCAGCACCAATCCTCCACCAG- 3'; *lfng*: 5'-CATATCGCGGAAAAGTCGTGGTC- 3', 5'- GCTTATCCGCTTCCCTCCTGCTG- 3'; *dlc*:5'-CGGGAATCGTCTCTTTGATAAT- 3', 5'- CTCACCGATAGCGAGTCTTCTT- 3'; *il17rd*: 5'-CTGGGTACCGGAGACTTGCGC- 3', 5'- TGGCGTGTTCCTCCAGCGGAC-3'; *il17ra1a*: 5'-TCACCTTTATACCTGTAGTGCTG- 3'; 5'- CATGTTTCTCTGACATCAAAAC-3'; *notch2*: 5'-GAGTGTGTGGACCCGTTAGTATG- 3', 5'- GCAGGCATCATCAATGTGACAC- 3'; *notch3*: 5'-TCAGGATTGTTCTCTCGTTGATG- 3', 5'- GTGTAAAGCATGTACCACCATTG- 3'.

Cell Counts

For all experiments, 6 out of 10 tiles of still images, which represent the central trunk area directly beneath the dorsal fin, were selected for cell counts. To ensure consistency across all samples and minimize discrepancies resulting from variations in fish trunk widths, a region of interest (ROI), 290µm x 200µm per tile, was created using the rectangle selection tool in ImageJ. Z-stack imaging was confined to the superficial/periderm layer based on the cross-sectional view, and the final count included both *krt4*⁺ and *krtr1c19e*⁺ cells at the periderm layer.

Intensity analyses

To analyze Notch expression level, tile images were merged separately for each fish. Merged images were then sectioned in ImageJ using the polygon selection tool to encompass only the fish trunk, and maximal projection was utilized to obtain the mean gray value. For *krtr1c19e*⁺ expression in the periderm layer, the z-stack was set as previously described, and the mean gray value of *krtr1c19e*⁺ expression was obtained.

Quantification and statistical analysis

GraphPad Prism software version 9.0.0 for Windows (GraphPad Software, San Diego, CA, USA) was used to perform statistical analyses. Continuous data were evaluated by Student t-test and Post hoc means were compared by Tukey-Kramer HSD.

References

- Abdou A.G., Maraee A.H., Sharaf A., Elnaidany N.F (2012) **Up-regulation of Notch-1 in psoriasis: an immunohistochemical study** *Ann Diagn Pathol* **16**:177–184
- Akat E., Yenmis M., Pombal M.A., Molist P., Megias M., Arman S., Vesely M., Anderson R., Ayaz D (2022) **Comparison of vertebrate skin structure at class level: A review** *Anat Rec (Hoboken)* **305**:3543–3608
- Armstrong A.W., Read C (2020) **Pathophysiology, Clinical Presentation, and Treatment of Psoriasis: A Review** *JAMA* **323**:1945–1960
- Barros-Becker F., Lam P.Y., Fisher R., Huttenlocher A (2017) **Live imaging reveals distinct modes of neutrophil and macrophage migration within interstitial tissues** *J Cell Sci* **130**:3801–3808
- Blanpain C., Fuchs E (2009) **Epidermal homeostasis: a balancing act of stem cells in the skin** *Nat Rev Mol Cell Biol* **10**:207–217
- Blanpain C., Lowry W.E., Pasolli H.A., Fuchs E (2006) **Canonical notch signaling functions as a commitment switch in the epidermal lineage** *Genes Dev* **20**:3022–3035
- Blauvelt A., Chiricozzi A (2018) **The Immunologic Role of IL-17 in Psoriasis and Psoriatic Arthritis Pathogenesis** *Clin Rev Allergy Immunol* **55**:379–390
- Borowczyk J., Buerger C., Tadjrischi N., Drukala J., Wolnicki M., Wnuk D., Modarressi A., Boehncke W.H., Brembilla N.C (2020) **IL-17E (IL-25) and IL-17A Differentially Affect the Functions of Human Keratinocytes** *J Invest Dermatol* **140**:1379–1389
- Bowman R.L., Wang D., Eom D.S (2023) **A macrophage subpopulation promotes airineme-mediated intercellular communication in a matrix metalloproteinase-9 dependent manner** *Cell Rep* **42**
- Campa M., Mansouri B., Warren R., Menter A (2016) **A Review of Biologic Therapies Targeting IL-23 and IL-17 for Use in Moderate-to-Severe Plaque Psoriasis** *Dermatol Ther (Heidelb)* **6**:1–12
- Chang W.J., Hwang P.P (2011) **Development of zebrafish epidermis** *Birth Defects Res C Embryo Today* **93**:205–214
- Dale J.K., Maroto M., Dequeant M.L., Malapert P., McGrew M., Pourquie O (2003) **Periodic notch inhibition by lunatic fringe underlies the chick segmentation clock** *Nature* **421**:275–278
- Daly C.A., Hall E.T., Ogden S.K (2022) **Regulatory mechanisms of cytoneme-based morphogen transport** *Cell Mol Life Sci* **79**
- Das S. et al. (2019) **Differential Influence of IL-9 and IL-17 on Actin Cytoskeleton Regulates the Migration Potential of Human Keratinocytes** *J Immunol* **202**:1949–1961

- Dodd M.E., Hatzold J., Mathias J.R., Walters K.B., Bennin D.A., Rhodes J., Kanki J.P., Look A.T., Hammerschmidt M., Huttenlocher A (2009) **The ENTH domain protein Clint1 is required for epidermal homeostasis in zebrafish** *Development* **136**:2591–2600
- Dubin-Bar D., Bitan A., Bakhrat A., Kaiden-Hasson R., Etzion S., Shaanan B., Abdu U (2008) **The Drosophila IKK-related kinase (Ik2) and Spindle-F proteins are part of a complex that regulates cytoskeleton organization during oogenesis** *BMC Cell Biol* **9**
- Eisenhoffer G.T. *et al.* (2017) **A toolbox to study epidermal cell types in zebrafish** *J Cell Sci* **130**:269–277
- Eom D.S (2020) **Airinemes: thin cellular protrusions mediate long-distance signalling guided by macrophages** *Open Biol* **10**
- Eom D.S., Bain E.J., Patterson L.B., Grout M.E., Parichy D.M (2015) **Long-distance communication by specialized cellular projections during pigment pattern development and evolution** *Elife* **4**
- Eom D.S., Parichy D.M (2017) **A macrophage relay for long-distance signaling during postembryonic tissue remodeling** *Science* **355**:1317–1320
- Fragoulis G.E., Siebert S., McInnes I.B (2016) **Therapeutic Targeting of IL-17 and IL-23 Cytokines in Immune-Mediated Diseases** *Annu Rev Med* **67**:337–353
- Friedman J.R., Webster B.M., Mastronarde D.N., Verhey K.J., Voeltz G.K (2010) **ER sliding dynamics and ER-mitochondrial contacts occur on acetylated microtubules** *J Cell Biol* **190**:363–375
- Gonzalez-Fernandez C., Chaves-Pozo E., Cuesta A (2020) **Identification and Regulation of Interleukin-17 (IL-17) Family Ligands in the Teleost Fish European Sea Bass** *Int J Mol Sci* **21**
- Gratton R., Tricarico P.M., Moltrasio C., de Oliveira Lima Estevao, Brandao A.S., Marzano L., Zupin A.V., and Crovella L. (2020) **Pleiotropic Role of Notch Signaling in Human Skin Diseases** *Int J Mol Sci* **21**
- Hall A (1998) **Rho GTPases and the actin cytoskeleton** *Science* **279**:509–514
- Hall E.T. *et al.* (2023) **Cytoneme signaling provides essential contributions to mammalian tissue patterning** *Cell*
- Hu Y., Xie S., Yao J (2016) **Identification of Novel Reference Genes Suitable for qRT-PCR Normalization with Respect to the Zebrafish Developmental Stage** *PLoS One* **11**
- Jaffe A.B., Hall A (2005) **Rho GTPases: biochemistry and biology** *Annu Rev Cell Dev Biol* **21**:247–269
- Kalthoff C., Groos S., Kohl R., Mahrhold S., Ungewickell E.J (2002) **Clint: a novel clathrin-binding ENTH-domain protein at the Golgi** *Mol Biol Cell* **13**:4060–4073
- Knopf F., Schnabel K., Haase C., Pfeifer K., Anastassiadis K., Weidinger G (2010) **Dually inducible TetON systems for tissue-specific conditional gene expression in zebrafish** *Proc Natl Acad Sci U S A* **107**:19933–19938

- Kornberg T.B., Roy S (2014) **Communicating by touch--neurons are not alone** *Trends Cell Biol* **24**:370–376
- Kornberg T.B., Roy S (2014) **Cytonemes as specialized signaling filopodia** *Development* **141**:729–736
- Lee R.T., Asharani P.V., Carney T.J (2014) **Basal keratinocytes contribute to all strata of the adult zebrafish epidermis** *PLoS One* **9**
- Lennikov A., Mirabelli P., Mukwaya A., Schaupper M., Thangavelu M., Lachota M., Ali Z., Jensen L., Lagali N. (2018) **Selective IKK2 inhibitor IMD0354 disrupts NF-kappaB signaling to suppress corneal inflammation and angiogenesis** *Angiogenesis* **21**:267–285
- Li Q., Frank M., Thisse C.I., Thisse B.V., Uitto J (2011) **Zebrafish: a model system to study heritable skin diseases** *J Invest Dermatol* **131**:565–571
- Liu T., Li S., Ying S., Tang S., Ding Y., Li Y., Qiao J., Fang H (2020) **The IL-23/IL-17 Pathway in Inflammatory Skin Diseases: From Bench to Bedside** *Front Immunol* **11**
- Lowes M.A., Suarez-Farinas M., Krueger J.G (2014) **Immunology of psoriasis** *Annu Rev Immunol* **32**:227–255
- Ma L., Li Y., Peng J., Wu D., Zhao X., Cui Y., Chen L., Yan X., Du Y., Yu L (2015) **Discovery of the migrasome, an organelle mediating release of cytoplasmic contents during cell migration** *Cell Res* **25**:24–38
- Martinez-Navarro F.J., Martinez-Menchon T., Mulero V., Galindo-Villegas J (2019) **Models of human psoriasis: Zebrafish the newly appointed player** *Dev Comp Immunol* **97**:76–87
- Melentijevic I. *et al.* (2017) **C. elegans neurons jettison protein aggregates and mitochondria under neurotoxic stress** *Nature* **542**:367–371
- Moriyama M., Durham A.D., Moriyama H., Hasegawa K., Nishikawa S., Radtke F., Osawa M (2008) **Multiple roles of Notch signaling in the regulation of epidermal development** *Dev Cell* **14**:594–604
- Mosca M., Hong J., Haderl E., Hakimi M., Liao W., Bhutani T (2021) **The Role of IL-17 Cytokines in Psoriasis** *Immunotargets Ther* **10**:409–418
- Nguyen B.C. *et al.* (2006) **Cross-regulation between Notch and p63 in keratinocyte commitment to differentiation** *Genes Dev* **20**:1028–1042
- Nickoloff B.J., Qin J.Z., Chaturvedi V., Denning M.F., Bonish B., Miele L (2002) **Jagged-1 mediated activation of notch signaling induces complete maturation of human keratinocytes through NF-kappaB and PPARgamma** *Cell Death Differ* **9**:842–855
- Nowell C., Radtke F (2013) **Cutaneous Notch signaling in health and disease** *Cold Spring Harb Perspect Med* **3**
- Ota T., Takekoshi S., Takagi T., Kitatani K., Toriumi K., Kojima T., Kato M., Ikoma N., Mabuchi T., Ozawa A (2014) **Notch signaling may be involved in the abnormal differentiation of epidermal keratinocytes in psoriasis** *Acta Histochem Cytochem* **47**:175–183

Otani T., Ogura Y., Misaki K., Maeda T., Kimpara A., Yonemura S., Hayashi S (2016) **IKKepsilon inhibits PKC to promote Fascin-dependent actin bundling** *Development* **143**:3806–3816

Panin V.M., Papayannopoulos V., Wilson R., Irvine K.D (1997) **Fringe modulates Notch-ligand interactions** *Nature* **387**:908–912

Parichy D.M., Elizondo M.R., Mills M.G., Gordon T.N., Engeszer R.E (2009) **Normal table of postembryonic zebrafish development: staging by externally visible anatomy of the living fish** *Dev Dyn* **238**:2975–3015

Park S., Kim H., Wang Y., Eom D.S., Allard J (2022) **Zebrafish airinemes optimize their shape between ballistic and diffusive search** *Elife* **11**

Park S. *et al.* (2021) **Skin-resident immune cells actively coordinate their distribution with epidermal cells during homeostasis** *Nat Cell Biol* **23**:476–484

Parsons M.J., Pisharath H., Yusuff S., Moore J.C., Siekmann A.F., Lawson N., Leach S.D (2009) **Notch-responsive cells initiate the secondary transition in larval zebrafish pancreas** *Mech Dev* **126**:898–912

Pasparakis M. *et al.* (2002) **TNF-mediated inflammatory skin disease in mice with epidermis-specific deletion of IKK2** *Nature* **417**:861–866

Pinto C.S., Khandekar A., Bhavna R., Kiesel P., Pigino G., Sonawane M (2019) **Microridges are apical epithelial projections formed of F-actin networks that organize the glycan layer** *Sci Rep* **9**

Rangarajan A. *et al.* (2001) **Notch signaling is a direct determinant of keratinocyte growth arrest and entry into differentiation** *EMBO J* **20**:3427–3436

Roy S., Hsiung F., Kornberg T.B (2011) **Specificity of Drosophila cytonemes for distinct signaling pathways** *Science* **332**:354–358

Sahlen P. *et al.* (2021) **Chromatin interactions in differentiating keratinocytes reveal novel atopic dermatitis- and psoriasis-associated genes** *J Allergy Clin Immunol* **147**:1742–1752

Schempp C., Emde M., Wolfle U (2009) **Dermatology in the Darwin anniversary. Part 1: Evolution of the integument** *J Dtsch Dermatol Ges* **7**:750–757

Segre J.A (2006) **Epidermal barrier formation and recovery in skin disorders** *J Clin Invest* **116**:1150–1158

Stanganello E., Hagemann A.I., Mattes B., Sinner C., Meyen D., Weber S., Schug A., Raz E., Scholpp S (2015) **Filopodia-based Wnt transport during vertebrate tissue patterning** *Nat Commun* **6**

Thelu J., Rossio P., Favier B (2002) **Notch signalling is linked to epidermal cell differentiation level in basal cell carcinoma, psoriasis and wound healing** *BMC Dermatol* **2**

Thelu J., Viallet J.P., Dhouailly D (1998) **Differential expression pattern of the three Fringe genes is associated with epidermal differentiation** *J Invest Dermatol* **111**:903–906

van Loon A.P., Erofeev I.S., Maryshev I.V., Goryachev A.B., Sagasti A. (2020) **Cortical contraction drives the 3D patterning of epithelial cell surfaces** *J Cell Biol* **219**

Wu B., Jin M., Zhang Y., Wei T., Bai Z. (2011) **Evolution of the IL17 receptor family in chordates: a new subfamily IL17REL** *Immunogenetics* **63**:835–845

Zenz R., Eferl R., Kenner L., Florin L., Hummerich L., Mehic D., Scheuch H., Angel P., Tschachler E., Wagner E.F (2005) **Psoriasis-like skin disease and arthritis caused by inducible epidermal deletion of Jun proteins** *Nature* **437**:369–375

Zhang C., Scholpp S (2019) **Cytonemes in development** *Curr Opin Genet Dev* **57**:25–30

Zhou X., Chen Y., Cui L., Shi Y., Guo C (2022) **Advances in the pathogenesis of psoriasis: from keratinocyte perspective** *Cell Death Dis* **13**

Article and author information

Yi Wang

Department of Developmental and Cell Biology, University of California, Irvine, CA 92617, USA

Qingan He

Department of Developmental and Cell Biology, University of California, Irvine, CA 92617, USA

Oliver Has

Department of Developmental and Cell Biology, University of California, Irvine, CA 92617, USA

Kiarash Forouzesh

Department of Developmental and Cell Biology, University of California, Irvine, CA 92617, USA

Dae Seok Eom

Department of Developmental and Cell Biology, University of California, Irvine, CA 92617, USA,
Center for Complex Biological Systems, University of California, Irvine, CA 92697, USA, UC Irvine
Skin Biology Resource Center, University of California, Irvine, CA 92697, USA

For correspondence: dseom@uci.edu

ORCID iD: [0000-0002-0617-8788](https://orcid.org/0000-0002-0617-8788)

Copyright

© 2024, Wang et al.

This article is distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editors

Reviewing Editor

Valerie Horsley

Yale University, New Haven, United States of America

Senior Editor

Jonathan Cooper

Fred Hutchinson Cancer Research Center, Seattle, United States of America

Reviewer #1 (Public Review):**Summary:**

In this paper, Wang et al show that differentiated peridermal cells of the zebrafish epidermis extend cytoneme-like protrusions toward the less differentiated, intermediate layer below. They present evidence that expression of a dominant-negative cdc42, inhibits cytoneme formation and leads to elevated expression of a marker of undifferentiated keratinocytes, *krtt1c19e*, in the periderm layer. Data is presented suggesting the involvement of Delta-Notch signaling in keratinocyte differentiation. Finally, changes in expression of the inflammatory cytokine IL-17 and its receptors is shown to affect cytoneme number and periderm structure in a manner similar to Notch and cdc42 perturbations.

Strengths:

Overall, the idea that differentiated cells signal to underlying undifferentiated cells via membrane protrusions in skin keratinocytes is interesting and novel, and it is clear that periderm cells send out thin membrane protrusions that contain a Notch ligand. Further, perturbations that affect cytoneme number, Notch signaling, and IL-17 expression clearly lead to changes in periderm structure and gene expression.

Weaknesses:

More work is needed to determine whether the effects on keratinocyte differentiation are due to a loss of cytonemes themselves, or to broader effects of inhibiting cdc42. Moreover, more evidence is needed to support the claim that periderm cytonemes deliver Delta ligands to induce Notch signaling below. Without these aspects of the study being solidified, understanding how IL-17 affects these processes seems premature.

<https://doi.org/10.7554/eLife.97400.1.sa2>

Reviewer #2 (Public Review):**Summary:**

The aim of the study was to understand how cells of the skin communicate across dermal layers. The research group has previously demonstrated that cellular connections called airinemes contribute to this communication. The current work builds upon this knowledge by showing that differentiated keratinocytes also use cytonemes, specialized signaling filopodia, to communicate with undifferentiated keratinocytes. They show that cytonemes are the more abundant type of cellular extension used for communication between the differentiated keratinocyte layer and the undifferentiated keratinocytes. Disruption of cytoneme formation led to the expansion of the undifferentiated keratinocytes into the periderm, mimicking skin diseases like psoriasis. The authors go on to show that disruption of cytonemes results in perturbations in Notch signaling between the differentiated keratinocytes of the periderm and the underlying proliferating undifferentiated keratinocytes. Further, the authors show that Interleukin-17, also known to drive psoriasis, can restrict the formation of periderm cytonemes, possibly through the inhibition of Cdc42 expression. This work suggests that cytoneme-mediated Notch signaling plays a central role in normal epidermal regulation. The authors propose that disruption of cytoneme function may be an underlying cause of various human skin diseases.

Strengths:

The authors provide strong evidence that periderm keratinocytes cytonemes contain the notch ligand DeltaC to promote Notch activation in the underlying intermediate layer to

regulate accurate epidermal maintenance.

Weaknesses:

The impact of the study would be increased if the mechanism by which Interleukin-17 and Cdc42 collaborate to regulate cytonemes was defined. Experiments measuring Cdc42 activity, rather than just measuring expression, would strengthen the conclusions.

<https://doi.org/10.7554/eLife.97400.1.sa1>

Reviewer #3 (Public Review):

Summary:

Leveraging zebra fish as a research model, Wang et al identified "cytoneme-like structures" as a mechanism for mediating cell-cell communications among skin epidermal cells. The authors further demonstrated that the "cytoneme-like structures" can mediate Notch signaling, and the "cytoneme-like structures" are influenced by IL17 signaling.

Strengths:

Elegant zebrafish genetics, reporters, and live imaging.

Weaknesses: (minor)

This paper focused on characterizing the "cytoneme-like structures" between different layers and the NOTCH signaling. However, these "cytoneme-like structures" observed in undifferentiated KC (Figure 2B), although at a slightly lower frequency, were not interpreted. In addition, it is unclear if these "cytoneme-like structures" can mediate other signaling pathways than NOTCH.

Overall, this is a solid paper with convincing data reporting the "cytoneme-like structures" in vivo, and with compelling data demonstrating the roles in NOTCH signaling and the regulation by IL17.

These findings provide a foundation for future work exploring the "cytoneme-like structures" in the mammalian system and other epithelial tissue types. This paper also suggests a potential connection between the "cytoneme-like structures" and psoriasis, which needs to be further explored in clinical samples.

<https://doi.org/10.7554/eLife.97400.1.sa0>