

Interneuron FGF13 regulates seizure susceptibility via a sodium channel-independent mechanism

Reviewed Preprint

v2 • November 19, 2024

Revised by authors

Reviewed Preprint

v1 • June 21, 2024

Susan Lin, Aravind R Gade, Hong-Gang Wang, James E Niemeyer, Allison Galante, Isabella DiStefano, Patrick Towers, Jorge Nunez, Maiko Matsui, Theodore H Schwartz, Anjali M Rajadhyaksha, Geoffrey S Pitt 

Cardiovascular Research Institute, Weill Cornell Medicine, New York, United States • Department of Neurological Surgery and Brain and Mind Research Institute, Weill Cornell Medicine of Cornell University, New York Presbyterian Hospital, New York, United States • Department of Pediatrics, Division of Pediatric Neurology, Weill Cornell Medicine, New York, United States • Brain and Mind Research Institute, New York, United States

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

eLife Assessment

This **important** study advances our understanding of how FGF13 variants confer seizure susceptibility. By acting in a set of inhibitory interneurons, FGF13 regulates synaptic transmission and excitability. The data presented here are **convincing** and combine cell type-specific knockouts and electrophysiology, complemented by histology/RNA studies. Collectively, this research will be of interest to a wide audience, particularly those involved in the study of epilepsy, inhibitory neurons, and ion channels.

<https://doi.org/10.7554/eLife.98661.2.sa3>

Abstract

Developmental and Epileptic Encephalopathies (DEEs), a class of devastating neurological disorders characterized by recurrent seizures and exacerbated by disruptions to excitatory/inhibitory balance in the brain, are commonly caused by mutations in ion channels. Disruption of, or variants in, *FGF13* were implicated as causal for a set of DEEs, but the underlying mechanisms were clouded because *FGF13* is expressed in both excitatory and inhibitory neurons, *FGF13* undergoes extensive alternative splicing producing multiple isoforms with distinct functions, and the overall roles of FGF13 in neurons are incompletely cataloged. To overcome these challenges, we generated a set of novel cell type-specific conditional knockout mice. Interneuron-targeted deletion of *Fgf13* led to perinatal mortality associated with extensive seizures and impaired the hippocampal inhibitory/excitatory balance while excitatory neuron-targeted deletion of *Fgf13* caused no detectable seizures and no survival deficits. While best studied as a voltage-gated sodium channel (Na_v) regulator, we observed no effect of *Fgf13* ablation in interneurons on Na_vs but rather a marked reduction in K⁺ channel currents. Re-expressing different *Fgf13* splice isoforms could partially rescue deficits in interneuron excitability and restore K⁺ channel current amplitude. These results enhance our understanding of the molecular mechanisms that drive the pathogenesis of

Introduction

Developmental and Epileptic Encephalopathies (DEEs) are a devastating group of rare epilepsies and neurodevelopmental disorders (Scheffer, French et al. 2016 [\[1\]](#), Scheffer, Berkovic et al. 2017 [\[2\]](#), Scheffer and Liao 2020 [\[3\]](#)) that affect 4 in 10,000 infants per year (Covanis 2012 [\[4\]](#), Ware, Huskins et al. 2019 [\[5\]](#)). DEEs are characterized by repeated seizures, severe developmental delay, and are often drug resistant. Febrile seizures, which are more common in young children, can contribute to the progression of disease (Skotte, Fadista et al. 2022 [\[6\]](#)). DEEs are lethal when the seizures lead to Sudden Unexpected Death in Epilepsy (SUDEP) (Scheffer and Nabbout 2019 [\[7\]](#), Chilcott, Diaz et al. 2022 [\[8\]](#), Donnan, Schneider et al. 2023 [\[9\]](#)). Although not all DEEs have been linked to a genetic basis, variants in ion channels and their auxiliary proteins are among the best-characterized genetic causes for DEEs, and sodium channelopathies have been identified as a top cause (Meisler, Hill et al. 2021 [\[10\]](#)).

Epilepsy-associated variants in voltage-gated sodium channels (Na_v s), necessary for action potential initiation, exert gain-of-function effects in excitatory neurons or loss of function effects in inhibitory neurons, leading to an overall pro-excitatory state. Gain-of-function variants in *SCN8A* coding for the neuronal sodium channel $\text{Na}_v1.6$ (Adam, Feldman et al. 1993 [\[11\]](#), Bunton-Stasyshyn, Wagnon et al. 2019 [\[12\]](#), Meisler 2019 [\[13\]](#), Talwar and Hammer 2021 [\[14\]](#)) or *SCN2A* coding for $\text{Na}_v1.2$ (Kim, Yang et al. 2020 [\[15\]](#), Li, Jancovski et al. 2021 [\[16\]](#)) (Miao, Tang et al. 2020 [\[17\]](#)) affect excitatory neurons of the forebrain, and lead to network hyperexcitability that drives seizures. In contrast, loss-of-function variants in *SCN1A* coding for $\text{Na}_v1.1$, the most abundant Na_v expressed in forebrain inhibitory neurons, reduce inhibitory drive to excitatory neurons and thus drive network hyperexcitability as the mechanism for seizures (Vormstein-Schneider, Lin et al. 2020 [\[18\]](#), Ding, Li et al. 2021 [\[19\]](#)). Variants in $\text{Na}_v1.1$ underlie the most common genetic subgroup in the prototypical DEE, Dravet Syndrome (Scheffer and Nabbout 2019 [\[7\]](#), Han, Chen et al. 2020 [\[20\]](#), Brunklaus, Pérez-Palma et al. 2022 [\[21\]](#), He, Li et al. 2022 [\[22\]](#)). DEEs also result from variants in Na_v auxiliary subunits, such as in the Na_v beta subunit *SCN1B* (Patino, Claes et al. 2009 [\[23\]](#), Ogiwara, Nakayama et al. 2012 [\[24\]](#), Kim, Dibbens et al. 2013 [\[25\]](#), Ramadan, Patel et al. 2017 [\[26\]](#), Bouza and Isom 2018 [\[27\]](#), Hull, O'Malley et al. 2020 [\[28\]](#)) as well as *FGF12* (Siekierska, Isrie et al. 2016 [\[29\]](#), Oda, Uchiyama et al. 2019 [\[30\]](#), Saleem, Chenchieri et al. 2024 [\[31\]](#)) and *FGF13* (Puranam, He et al. 2015 [\[32\]](#), Fry, Marra et al. 2021 [\[33\]](#), Velišková, Marra et al. 2021 [\[34\]](#), Narayanan, Majethia et al. 2022 [\[35\]](#)), both fibroblast growth factor homologous factors (FHF) (Goldfarb 2005 [\[36\]](#)).

Here, we focus on *FGF13*, which is expressed in both excitatory and inhibitory neurons (Puranam, He et al. 2015 [\[32\]](#), Joglekar, Prijibelski et al. 2021 [\[37\]](#)). Like other members of the FHF family (FGF11–14), *FGF13* is a non-canonical fibroblast growth factor that is not secreted and does not bind to FGF receptors (Schoorlemmer and Goldfarb 2001 [\[38\]](#), Olsen, Garbi et al. 2003 [\[39\]](#)). FHFs reside in the cytoplasm where they can bind to the intracellular C-terminus of various Na_v s and modulate their function (Goldfarb 2005 [\[36\]](#), Pablo, Wang et al. 2016 [\[40\]](#)). *FGF13* has been shown to be upregulated in the hippocampus of a temporal lobe epilepsy mouse model, whereas knockdown of *Fgf13* attenuated hyperexcitability in hippocampal cells (Shen, Yue et al. 2022 [\[41\]](#)). Other *FGF13* functions, such as the regulation of microtubules (Wu, Yang et al. 2012 [\[42\]](#)), control of mitogen-activated protein kinases (Lu, Shi et al. 2015 [\[43\]](#)), and effects on neurodevelopment (Nishimoto and Nishida 2007 [\[44\]](#)) including development of chandelier cells (Favuzzi, Deogracias et al. 2019 [\[45\]](#)) have been ascribed, but less well characterized. Moreover, the full complement of *FGF13* functions has yet to be defined, obscuring insight into how *FGF13* variants lead to DEEs.

While DEE associated variants in *FGF13* are thought to affect Na_v function, this has not been demonstrated in neurons and the contribution of other FGF13 functions has not been investigated, thus preventing an understanding of disease pathology and the development of precision therapies. Lack of mechanistic clarity also arises because DEE associated variants in *FGF13* have been proposed to cause gain-of-function effects in excitatory neurons (Fry, Marra et al. 2021 [↗](#)) or via an apparently contradictory loss-of-function mechanism in inhibitory neurons (Puranam, He et al. 2015 [↗](#)). Further complicating mechanistic understanding, *FGF13* can generate multiple isoforms, driven by alternative promoter usage and consequent alternative splicing, with different but overlapping functions (Munoz-Sanjuan, Smallwood et al. 2000 [↗](#), Puranam, He et al. 2015 [↗](#)). We recently discovered that alternatively spliced *Fgf13* mRNAs exhibit distinct cell type-specific expression in the mouse cortex. In the developing mouse brain excitatory neurons express almost exclusively *Fgf13-S* while inhibitory neurons express both *Fgf13-S* and *Fgf13-VY*, a notable example of differential expression of an alternatively spliced neuronal gene in excitatory vs. inhibitory neurons (Furlanis, Traunmuller et al. 2019 [↗](#), Joglekar, Prijbelski et al. 2021 [↗](#)). This cell type-specific alternative splicing pattern is similar to other genes which are implicated in the control of synaptic interactions in the hippocampus (Furlanis, Traunmuller et al. 2019 [↗](#)), yet the differential splicing complicates the understanding of current models.

DEE associated variants in *FGF13* have been identified in *FGF13-S* and modeled as causing pro-excitatory Na_v channel dysfunction in excitatory neurons. Conversely, a maternally transmitted chromosomal translocation that disrupts the X-linked *FGF13* locus and eliminates *FGF13-VY* expression while preserving expression of *FGF13-S* causes a seizure disorder in two brothers (Puranam, He et al. 2015 [↗](#)). Thus, how *FGF13* variants confer seizure susceptibility is unclear. Finally, whether *FGF13* variants even exert their effects through Na_v dysfunction or through alternative mechanisms has not been defined.

We generated a series of conditional knockout mouse lines to probe FGF13 function in excitatory and inhibitory hippocampal neurons and investigated the isoform-specific and cell type mechanisms by which variants in *Fgf13* cause DEEs. We define a critical role for FGF13 in interneurons but not excitatory neurons underlying a pro-excitatory state associated with seizures. Moreover, as we observe limited effects on Na_v currents underlying the changes associated with a pro-excitatory state, our data add to growing observations that FGF13 can function independently of Na_vs.

Methods

Animals

Mice were handled in accordance with the ethical guidelines of the National Institutes of Health Guide for the Care and Use of Laboratory Animals. This study was approved by the Weill Cornell Medical Center Institutional Animal Care and Use Committee. Genetically modified mice were maintained on C57BL6/J background (000664; The Jackson Laboratory, Bar Harbor, ME, USA). All mice were maintained on a standard rodent chow diet (PicoLab Rodent Diet 20; 5053; LabDiet, St. Louis, MO, USA) with a 12-h light/dark cycle. Mice were weaned at post-natal day 21 (P21) and group housed in cages holding between two and five mice. To generate neuron-specific mutant mice, female *Fgf13^{fl/fl}* mice (Wang, Tang et al. 2017 [↗](#)) were crossed with male *Nestin-Cre^{+/+}* (*B6.Cg-Tg(Nes-cre)1Kln/J*, JAX #003771) to produce male hemizygous knockouts (*Nestin^{Fgf13} KO*), female heterozygous mutants (*Nestin^{Fgf13} HET*), and wild type littermates. To generate excitatory neuron-targeted mutant mice, female *Fgf13^{fl/fl}* mice were crossed with male *Emx1-Cre^{+/+}* (*B6.129S2-Emx1tm1(cre)Krj/J*, #005628) to produce male hemizygous knockouts (*Emx1^{Fgf13} KO*), female heterozygous mutants (*Emx1^{Fgf13} HET*), and wild type littermates. Female *Emx1* knockouts were generated by crossing male hemizygous knockouts (*Emx1^{Fgf13} KO*) with female *Fgf13^{fl/fl}* mice. To generate interneuron-specific mutant mice, male *Gad2-Cre^{+/+}* (*Gad2tm2(cre)Zjh/J*, #010802) were

crossed with female *Fgf13^{fl/fl}* mice to produce male hemizygous knockouts (*Gad2^{Fgf13 KO}*), female heterozygous mutants (*Gad2^{Fgf13 HET}*), and wild type littermates. To generate interneuron-GFP reporter mice, male *Gad2-Cre^{+/+}* (*Gad2tm2(cre)Zjh/J*, #010802) were crossed with *Cre*-dependent *Ai6^{+/+}* mice (B6.Cg-Gt(ROSA)26Sortm6(CAG-ZsGreen1)Hze/J), and the resulting *Gad2^{GFP}* (*Gad2-Cre^{+/+}*; *Ai6^{+/+}*) males were crossed with female *Fgf13^{fl/fl}* mice to generate *Fgf13* knockout interneuron-GFP reporter mice (*Gad2-Cre^{+/+}*; *Fgf13^{FL/X}*; *Ai6^{+/+}* or *Gad2^{GFP-Fgf13 KO}*). To generate medial ganglionic eminence (MGE) derived interneuron-targeted mutant mice, female *Fgf13^{fl/fl}* mice were crossed with male *Nkx2.1-Cre^{+/+}* (*C57BL/6J-Tg(Nkx2-1-cre)2Sand/J*, # 008661) to produce male hemizygous knockouts (*Nkx2.1^{Fgf13 KO}*), female heterozygous mutants (*Nkx2.1^{Fgf13 HET}*), and wild type littermates.

Hyperthermia induced seizures

Mice at P12 were tested using a modified protocol (Cheah, Yu et al. 2012 [↗](#), Puranam, He et al. 2015 [↗](#)). Mouse core body temperature was recorded with a rodent rectal temperature probe placed at a depth of 1 cm and connected to a rodent temperature controller (TCAT-2DF, Physitemp). Behavior was monitored by a video camera (Logitech). Command temperature $\pm 0.3^\circ\text{C}$ was maintained using an infrared heat lamp positioned directly over the recording chamber (2L glass beaker). Mice were acclimated to the chamber for 5 minutes, and any mice with a lower core body temperature were warmed with the heat lamp up to 36.5°C . For baseline monitoring, temperature was adjusted to an initial set-point of 36.5°C for 5 minutes. Body temperature was elevated in 0.5°C increments, for 2 minutes at each increment. Temperature was increased until a seizure or 42.0°C was reached. At the end of hyperthermia induction, mice were placed on a room temperature surface to recover and monitored for 10 minutes.

Electroencephalogram recording

Animal preparation

Mice (P12) were anesthetized under isoflurane (3.5% induction, 1-2% maintenance), provided pre-surgical analgesia (Meloxicam, 2mg/kg SC; Bupivacaine 0.15 ml under scalp), and head-fixed in a stereotaxic frame (KOPF Instruments) with ear bars and an anesthesia-passing nose cone. A heatpad was placed under the animal and the eyes were protected with ointment (Paralube Vet). Hair was removed with fine scissors and an oval-shaped incision ($\sim 4 \times 6$ mm) was made on the scalp to expose the skull. Fine forceps were used to scrape the membrane, remove any remaining hair, and expose bregma and skull sutures. The hippocampal EEG site was located at -2.5 mm AP, 1.5 mm ML, -1.25 mm DV. A dental drill was then used to open small craniotomies at the (AP, ML) sites for EEG contacts with a 0.5 mm bur (Meisinger). Reference and ground contact craniotomies were made on the contralateral skull. The EEG recording system (Pinnacle Technology Inc) was attached to headmounts (Pinnacle product #8201-ss) with conductive metallic holes and fastened to the skull by small conductive screws (Pinnacle product #8209). An insulated tungsten wire (0.002", California Fine Wire Company) was soldered to the EEG mount and trimmed to 2 mm to target the hippocampus (Klorig, Alberto et al. 2019 [↗](#)), using epoxy to insulate the soldered portion of the electrode. The mount, with hippocampus electrode attached, was slowly lowered onto the skull with a micromanipulator such that the electrode was inserted directly down into the craniotomy over hippocampus. The headmount was then attached with a small amount of dental cement (C&B Metabond, Parkell) to hold the device steady as skull screws were then implanted into the remaining craniotomies. Skull screws were then reinforced to the headmount by conductive silver paint (SPI supplies). Dental cement (Metabond) was then used to cover all parts of the device and attach it firmly to the skull.

Recording and analysis

After recovery from surgery the headmount was attached to a preamplifier and animals placed in a bedded recording chamber with ad libitum food and water. Pinnacle EEG software recorded electrographic data at 400 Hz from the tethered preamplifier. Spontaneous activity was recorded for up to 5 hours, while heat-lamp experiments were recorded as described above. After recording, data were transferred to a PC and analyzed in MATLAB (Mathworks). Raw data were bandpass filtered (1-60 Hz, infinite impulse response filter with bandpass.m) and events of interest were automatically flagged by using line length of the signal (difference of successive voltage values with a moving 5-sec window), which permitted detection of large deviations (MATLAB findpeaks.m). Interictal and seizure events were then visually examined. In heat-lamp experiments, the times of temperature steps were recorded for alignment with the simultaneous EEG recording.

Hippocampal Neuronal Cultures

Hippocampi were dissected from P0-1 newborn pups and dissociated through enzymatic treatment with 0.25% trypsin and subsequent trituration. The cells were plated on glass coverslips previously coated with poly-D-lysine and laminin in 24-well cell culture plates. The hippocampal cells were grown in neurobasal A medium (ThermoFisher Scientific) supplemented with 2% B-27, 2mM glutamine, 10% heat-inactivated fetal bovine serum, and 1% penicillin/streptomycin in a 5% CO₂ incubator at 37°C overnight. After 24 hours this medium was replaced by a culture medium containing 2% B-27, 0.5mM glutamine, 1% heat-inactivated fetal bovine serum, 70 µM uridine and 25 µM 5-fluorodeoxyuridine. Cultured neurons were used for electrophysiology and immunocytochemistry.

Immunocytochemistry

Cells were fixed in 4% PFA 20 minutes, rinsed three times with phosphate buffered saline (PBS), and blocked in 2.5% bovine serum albumin with 0.2% Triton-X in PBS for 30 minutes. Primary anti-FGF13 antibody (Wang, Hennessey et al. 2011 [DOI](#)) was diluted in 2.5% bovine serum albumin and incubated in 4°C overnight. Coverslips were rinsed three times in PBS, then incubated in Alexa-568 secondary antibody (1:500 in 2.5% bovine serum albumin) for 1 hour at room temperature followed by DAPI for 5 minutes. Coverslips were mounted onto glass slides with mounting media (Vector Laboratories).

Acute Slice Preparation

Acute hippocampal slices were prepared from P12 to P14 mice. Briefly, after the mouse was anesthetized with isoflurane and then decapitated, the brain was quickly extracted and transferred into ice-cold cutting solution bubbled with 95% O₂ and 5% CO₂. The cutting solution contained (in mM): NaCl 77, sucrose 75, KCl 2.5, NaH₂PO₄ 1.4, NaHCO₃ 25, MgSO₄ 7, CaCl₂ 0.5, glucose 25, and sodium pyruvate 3, pH 7.4. Coronal brain slices (300 µm) including hippocampus were prepared using a Leica VT 1200S vibratome (Leica, Inc), and were incubated in a BSK-2 brain slice keeper (Automate Scientific, CA) containing oxygenated artificial cerebrospinal fluid (aCSF) at 35 °C for 30 min to recover. Afterwards, the slices were maintained at room temperature at least 30 min before use.

Electrophysiology

For spontaneous excitatory and inhibitory synaptic currents (sEPSCs and sIPSCs) in brain slices, the slice was placed in a recording chamber on the stage of an upright, infrared-differential interference contrast microscope (BX51WI, Olympus Optical) equipped with an ORCA-Flash2.8 C11440 Digital CMOS Camera (Hamamatsu Photonics), and was continuously perfused at a rate of 2 ml/min with aCSF bubbled with 95% O₂ and 5% CO₂ at 35°C. Pyramidal neurons in hippocampal

CA1 stratum pyramidale were visualized with a 40X water-immersion objective. Spontaneous EPSCs and IPSCs were recorded in the whole-cell voltage clamp configuration at a holding potential of -70 mV and sampled at 10 kHz and filtered at 2 kHz using an Axopatch 200B amplifier and Digidata 1322A digitizer (Molecular Devices). The pipette internal solution for sEPSC contained (in mM): potassium gluconate (KGluc) 125, KCl 10, MgCl₂ 5, EGTA 0.6, HEPES 5, CaCl₂ 0.06, phosphocreatine disodium 10, Mg-ATP 2, Na₂-GTP 0.2, creatine phosphokinase 50 U/ml, and N-Ethyl lidocaine bromide 5, osmolarity 289 mOsm/L, pH 7.2 adjusted with KOH. The pipette internal solution for sIPSC contained (in mM): KCl 125, MgCl₂ 5, EGTA 0.6, HEPES 5, CaCl₂ 0.06, phosphocreatine disodium 10, Mg-ATP 2, Na₂-GTP 0.2, creatine phosphokinase 50 U/ml and N-Ethyl lidocaine bromide 5, osmolarity 291 mOsm/L, pH 7.2 adjusted with KOH. The external solution aCSF contained (in mM): NaCl 126, KCl 2.5, NaH₂PO₄ 1.25, NaHCO₃ 26, CaCl₂ 2, MgSO₄ 2, glucose 10. Bicuculline methiodide 20 μM was added to aCSF for sEPSC, and 2-amino-5-phosphonovaleic acid (APV) 50 μM and 6,7-dinitroquinoxaline-2,3-dione (DNQX) 20 μM was added to aCSF for sIPSC.

Whole-cell sodium Na⁺ currents and whole-cell K⁺ currents, or action potentials (APs) from cultured neurons were recorded with a HEKA EPC10 amplifier in the voltage-clamp or current-clamp configuration, respectively.

For Na⁺ current recordings in cultured neurons, the internal pipette solution contained (in mM): CsF 125, NaCl 10, HEPES 10, TEA-Cl 15, EGTA 1.1, Na-GTP 0.5, pH 7.4 with NaOH; the external solution contained NaCl 125, KCl 5, CaCl₂ 2, MgCl₂ 1, TEA-Cl 20, HEPES 5, Glucose 10, pH 7.4 with NaOH. For K⁺ current recordings and AP initiation in cultured neurons, the internal pipette solution contained (in mM): potassium gluconate 130, KCl 10, MgCl₂ 5, ethylene glycol-bis(β-aminoethyl ether)-N,N,N',N'-tetraacetic acid 0.6, HEPES 5, CaCl₂ 0.06, phosphocreatine disodium 10, Mg-ATP 2, Na₂-GTP 0.2, and creatine phosphokinase 50 U/ml, pH 7.2 adjusted with KOH; the external solution contained (in mM): NaCl 119, KCl 5, HEPES 20, CaCl₂ 2, MgCl₂ 2, glucose 30, APV 0.05, DNQX 0.02, bicuculline 0.02, pH 7.3 adjusted with NaOH. Additionally for K⁺ current recordings, the external solution contained 100 nM tetrodotoxin (TTX) and 5 μM nifedipine to block Na⁺ and Ca²⁺ channels.

Recording pipettes were pulled from borosilicate glass with Sutter P-97 Micropipette Puller (Sutter Instrument Co.). For Na⁺ and K⁺ current recordings, pipette resistance ranged from 1.5 to 2.5 MΩ and the series resistance was compensated by at least 70%. For AP initiation recordings, pipette resistance ranged from 2.0 to 3.0 MΩ. For recordings of sEPSCs and sIPSCs in brain slices, pipette resistance ranged from 2.6 to 6.1 MΩ and series resistance was 10.5 ± 0.5 MΩ for WT and 10.8 ± 0.4 MΩ for *Gad2^{Fgf13} KO* without compensation.

Analysis of Electrophysiological Data

Action potentials were elicited by holding the cell at 0 pA in current clamp mode and then stimulating with current increments for 1 s. The numbers of action potentials elicited for each current injection were quantified using Fitmaster software. Action potential parameters were analyzed using Clampfit 10.4 software. Specifically, action potential amplitude and action potential duration measured at 50% of the amplitude (APD₅₀) values for the first three action potentials were determined using the statistics function. Additionally, dV/dt values were computed using the differential function. The threshold potential was determined by analyzing the phase plot of the initial action potential at rheobase by identifying the voltage value on the x-axis where the y-axis reached a rate of 1 mV/ms.

Na⁺ and K⁺ channel currents were activated by holding the neuron at -90mV and applying incremental voltages for a duration of 150 ms. Fitmaster software was used for the quantification of various channel parameters. The non-inactivating portion of the K⁺ channel current was assessed by analyzing the peak current at the conclusion of 150 ms depolarizing pulses. To generate steady-state inactivation curves, currents were elicited at a holding potential of -120 mV followed by a 500-ms prepulse to voltages ranging from -120 to +20 mV. Subsequently, currents

were measured at -20 mV for 20 ms. Normalized current values obtained at -20 mV were utilized to construct the steady-state inactivation curves. A Boltzmann equation, $[I/I_{\text{max}} = (1 + \exp((V - V_{1/2}) / k))^{-1}]$, was employed to fit the data, where I_{max} denotes the maximum current, $V_{1/2}$ represents the half-inactivation voltage, and k signifies the slope. This equation facilitated the calculation of $V_{1/2}$ of inactivation.

Data for Na^+ and K^+ current or APs were analyzed with Axon Clampfit (Molecular Devices), and data for sEPSC and sIPSC were analyzed with Mini Analysis (Synaptosoft).

Immunohistochemistry

Mice were sacrificed at postnatal day 13-16 (P13-16). Brains were extracted after transcardial perfusion of PBS followed by 4% paraformaldehyde (PFA). They were incubated overnight rocking at 4 °C in 4% paraformaldehyde, then equilibrated in 15% sucrose followed by 30% sucrose. Brains were embedded in Optimal Cutting Temperature (OCT) compound in cryomolds and cryosectioned at 50 μm into 0.1 M PBS. Floating sections were washed 3 times with PBS, then blocked for 1 hour in 2.5% bovine serum albumin and 0.2% Triton-X in PBS. Tissue was incubated with the anti-FGF13 antibody overnight at 4°C. After sections were washed three times with PBS, they were incubated with Alexa-568 secondary antibody for 2 hours at room temperature. Sections mounted onto glass slides with mounting media (Vector laboratories) and sealed with glass cover slips. Slides were stored at 4 °C. Hippocampal tissue was imaged using a confocal microscope (Zeiss LSM 880) and analyzed using Image-J software.

Western Blot

Tissue was immersed in ice cold RIPA buffer supplemented with Halt protease and phosphatase inhibitor cocktails (Thermo Fisher Scientific) and placed in a manual Dounce tissue homogenizer, in which they were homogenized for 1 minute on ice. The homogenate was then centrifuged at 21,000 $\times g$ at 4 °C. Supernatant was collected and protein concentration was determined by bicinchoninic acid assay. Protein was separated on an 8-16% gradient Tris-glycine gels and transferred to PVDF blotting membrane (GE Healthcare Life Sciences). The membrane was immunoblotted with 1:200 anti-FGF13 antibody or 1:200 anti-pan FHF-A (Neuromab clone N235/22) for two hours at RT to immunoblot for FGF13-S. Vinculin (Sigma V9131) was used as a loading control. The blots were visualized by chemiluminescence and images were captured using ChemiDoc™ Tough Imaging System (BioRad).

BaseScope *in situ* hybridization

Mice were sacrificed at P13-15. Brains were extracted after transcardial perfusion of PBS followed by 4% paraformaldehyde. The tissue was incubated for 24 hours rocking at 4°C in 4% paraformaldehyde, then equilibrated in 15% sucrose followed by 30% sucrose. Brains were embedded in OCT compound in cryomolds and sectioned coronally in 14 μm sections onto SuperFrost (ThermoFisher) slides, then stored at -80°C up to three months. Custom *Fgf13* isoform probes were generated as previously described (Joglekar, Prjibelski et al. 2021 [\[1\]](#)) and used according to the manufacturer's protocol for fixed frozen tissues. Hematoxylin followed by 0.02% ammonia water was used for nuclear counterstain.

Targeted metabolomics

Mice were sacrificed at P13 and brains were immediately extracted, homogenized, and placed into pre-chilled 80% methanol (-80L°C). The extract was dried with a Speedvac, and redissolved in HPLC grade water before it was applied to the hydrophilic interaction chromatography LC-MS. Metabolites were measured on a Q Exactive Orbitrap mass spectrometer (Thermo Scientific), which was coupled to a Vanquish UPLC system (Thermo Scientific) via an Ion Max ion source with a HESI II probe (Thermo Scientific). A Sequant ZIC-pHILIC column (2.1 mm i.d. $\times$ 150 mm, particle size of 5L μm , Millipore Sigma) was used for separation of metabolites. A 2.1L $\times$ 120Lmm guard

column with the same packing material was used for protection of the analytical column. Flow rate was set at 150 $\mu\text{L}/\text{min}$. Buffers consisted of 100% acetonitrile for mobile phase A, and 0.1% $\text{NH}_4\text{OH}/20\text{ mM CH}_3\text{COONH}_4$ in water for mobile phase B. The chromatographic gradient ran from 85% to 30% A in 20 min followed by a wash with 30% A and re-equilibration at 85% A. The Q Exactive was operated in full scan, polarity-switching mode with the following parameters: the spray voltage 3.0 kV, the heated capillary temperature 300°C, the HESI probe temperature 350°C, the sheath gas flow 40 units, the auxiliary gas flow 15 units. MS data acquisition was performed in the m/z range of 70–1,000, with 70,000 resolution (at 200 m/z). The AGC target was 1e6 and the maximum injection time was 250 ms. The MS data was processed using Xcalibur 4.1 (Thermo Scientific) to obtain the metabolite signal intensities. Identification required exact mass (within 5 ppm) and standard retention times. Statistical analysis was performed as follows: The intensity values were subjected to a Log2 transformation to normalize the data distribution. Student's *t*-test was conducted to calculate the significance of observed differences between groups. Enrichment analyses were performed using Metaboanalyst online tools (<https://www.metaboanalyst.ca/> .

Reverse transcriptase quantitative polymerase chain reaction

From hippocampal neuron cultures prepared for electrophysiology, RNA was isolated using RNeasy Mini Kit (QIAGEN) according to the manufacturer's instructions. Reverse transcription to generate cDNA was performed using SuperScriptTM IV VLOTM Master Mix with ezDNase (Invitrogen). qPCR was performed in duplicate for each sample with QantStudio 3 (Applied Biosystems) using SYBR green-based detection chemistries (Bio-Rad). For primer sequences used for qPCR target genes, see Supplemental Table 1.

Statistics

Statistical analyses were performed in GraphPad Prism v10 and specific analyses are described in the accompanying figure legends.

Results

Whole brain knockout of *Fgf13* results in spontaneous seizures and premature death

To determine how *FGF13* affects neuronal excitability and provide insight into how *FGF13* variants contributes to seizure disorders, we developed genetic mouse models that eliminate *Fgf13* in specific neuronal cell types. We started by eliminating the X-linked *Fgf13* in all neurons, crossing a male *Nestin*-Cre driver with a *Fgf13*^{*fl/fl*} female mouse from a previously validated line (Wang, Tang et al. 2017 ) (**Figure 1A** ) to generate hemizygous male *Fgf13* knockouts (*Nes-cre*;*Fgf13*^{*fl/Y*} or "*Nestin*^{*Fgf13*} *KO*") and heterozygous females (*Fgf13*^{*fl/+*} or "*Nestin*^{*Fgf13*} *HET*") alongside wild type (WT) littermates. We validated our mouse lines by examining the hippocampus, where *Fgf13* is highly expressed (Pablo, Wang et al. 2016 ) and which has previously been implicated in *FGF13*-related seizure pathology (Puranam, He et al. 2015 , Shen, Yue et al. 2022 ) and cognition (Wu, Yang et al. 2012 ). In hippocampus from WT mice, western blotting with a pan-FGF13 antibody that recognizes all FGF13 isoforms detected four distinct bands that we were able to assign to specific FGF13 isoforms by running standards (isoform specific cDNAs expressed in HEK293 cells) (**Figure 1B** ) and Supplemental Figure 1A-B). In hippocampi from *Nestin*^{*Fgf13*} *KO* brains we observed near complete absence of FGF13 (**Figure 1B** ) by western blot and by immunohistochemistry (**Figure 1C** ). In WT mice, FGF13 signal is prominent in the hippocampal pyramidal cell layer as well as in sparse presumed interneuron somata (see below) throughout all layers. Offspring were born at expected Mendelian ratios (**Table 1** ) , but all *Nestin*^{*Fgf13*} *KO* male mice died before weaning with a median survival age of 17 days postnatal (P17; **Figure 1D** ) . *Nestin*^{*Fgf13*} *HET* females also exhibited postnatal death, with only 60% surviving past one month

(Figure 1D [↗](#)), implying a gene dosage effect of *Fgf13* loss. *Nestin^{Fgf13 KO}* male mice suffered spontaneous seizures in their home cage (Supplemental video 1) as early as P12, followed by premature death (Figure 1D [↗](#)), consistent with mouse models of DEE that exhibit SUDEP (Bunton-Stasyshyn, Wagnon et al. 2019 [↗](#), Han, Chen et al. 2020 [↗](#)). *Nestin^{Fgf13 KO}* males were smaller than their wild type littermates by P14 (Figure 1E [↗](#)), suggesting developmental delay, likely a consequence of recurrent seizures and consistent with DEE mouse models (Yu, Mantegazza et al. 2006 [↗](#), Kalume, Westenbroek et al. 2013 [↗](#), Teran, Sainju et al. 2023 [↗](#)).

As febrile seizures are a hallmark of early developmental onset epilepsy and were observed in subjects with a gene translocation that disrupted the *FGF13* locus (Puranam, He et al. 2015 [↗](#)), we tested hyperthermia-induced seizure susceptibility in the *Nestin^{Fgf13 KO}* male mice, following a modified protocol (Cheah, Yu et al. 2012 [↗](#), Puranam, He et al. 2015 [↗](#)). Mice were tested at P12, the age at which we first observed spontaneous home cage seizures. Briefly, mice were acclimated to the testing chamber and a heat lamp was used to increase core body temperature at a steady rate of 0.5 °C every 2 minutes until 42.0 °C to simulate fever onset. All *Nestin^{Fgf13 KO}* males exhibited tonic-clonic seizures during the hyperthermic seizure protocol while wild type mice remained seizure free (Figure 1F [↗](#)).

Excitatory neuronal knockout of *Fgf13* does not result in spontaneous seizures

Excitatory neurons have been the focus of most studies investigating *Fgf13* dysfunction. *Fgf13* deficiency in excitatory forebrain neurons is linked to neurodevelopmental delay and cognitive impairment (Wu, Yang et al. 2012 [↗](#)), and excitatory neurons are hypothesized to be the relevant cell type for seizures in a DEE associated with *FGF13* variants (Fry, Marra et al. 2021 [↗](#)). To test the contribution of excitatory neurons to the seizure and SUDEP phenotype identified in *Nestin^{Fgf13 KO}* mice, we generated an excitatory neuron-targeted knockout mouse using an *Emx1*-Cre driver (Gorski, Talley et al. 2002 [↗](#)). We assessed efficacy of knockout by western blot using the pan-FGF13 antibody. In *Emx1^{Fgf13 KO}* mice, western blots showed the loss of two FGF13 isoforms (Figure 2A [↗](#)), FGF13-S and FGF13-U, consistent with single-cell long-read RNA sequencing that showed that excitatory neurons predominantly express FGF13-S (Joglekar, Prjibelski et al. 2021 [↗](#)), which we confirmed with a FGF13-S specific antibody (Supplemental Figure 1C). FGF13-VY and FGF13-V, the major isoforms expressed in inhibitory interneurons (Joglekar, Prjibelski et al. 2021 [↗](#)), were preserved. Thus, these data confirm at the protein level the differential expression of specific *Fgf13* transcripts in excitatory vs. inhibitory neurons observed by single cell RNA sequencing. Fluorescent immunohistochemistry of hippocampi from *Emx1^{Fgf13 KO}* showed loss of FGF13 from the CA1 pyramidal cell layer but maintenance of FGF13 signal from sparse putative interneuron somata (Figure 2B [↗](#)). *Emx1^{Fgf13 KO}* mice were born at expected Mendelian ratios (Table 1 [↗](#)) and all survived to adulthood (Figure 2C [↗](#)), similar to a previous report (Wu, Yang et al. 2012 [↗](#)). Average body size and mass at P14 revealed no differences between wild type and *Emx1^{Fgf13 KO}* genotypes (Figure 2D [↗](#)). None of the *Emx1^{Fgf13 KO}* succumbed to seizures during the hyperthermia protocol (Figure 2E [↗](#)), and EEG recordings during the hyperthermia protocol confirmed no seizure activity in *Emx1^{Fgf13 KO}* or wild type mice (Figure 2F [↗](#)). These data show that loss of *Fgf13* from excitatory forebrain neurons is not sufficient to induce seizure susceptibility.

The mechanism by which DEE-associated variants in *FGF13*-S contribute to a pro-excitatory state and seizures was hypothesized to be loss of a FGF13-S dependent long-term inactivation, a process that describes a successive reduction in peak Na_v current following repetitive depolarizations and is thought to derive from accreting channel pore blockade by the alternatively spliced N-terminus of FGF13-S (Fry, Marra et al. 2021 [↗](#)). Whether FGF13-S confers long-term inactivation in neurons has not been demonstrated. We employed the knockout model to assess the effects of FGF13-S on Na_v currents in *Emx1⁺* excitatory neurons and determine if FGF13-S bestows long-term inactivation. We cultured hippocampal neurons from *Emx1^{Fgf13 KO}* or wild type littermates (*Emx*-

Figure 1.

Whole brain knockout of *Fgf13* results in premature death and seizure susceptibility.

A. Breeding scheme to generate *Nestin^{Fgf13 KO}*, *Nestin^{Fgf13 HET}*, and wildtype littermates. **B.** Western blot of whole brain *Nestin^{Fgf13 KO}* (KO) and wildtype (WT) littermates at P2 validates *Fgf13* knockout. **C.** Fluorescent immunohistochemistry of hippocampal tissue validates *Fgf13* knockout (scale bar, *Fgf13 KO* μm). **D.** Survival curve of *Nestin* mutant mice shows decreased survival at one month of age log-rank test, ****, $p < 0.0001$). **E.** Body mass at P14 shows *Nestin^{Fgf13 KO}* are smaller in size (t-test, **, $p < 0.01$). **F.** *Nestin^{Fgf13 KO}* are susceptible to hyperthermia induced seizures (log-rank test, ****, $p < 0.0001$), unlike wildtype littermates.




Table 1

Mutant mice are born in Mendelian ratios

Mouse Strain	Probability Density Function
Nestin-Fgf13	0.23
Gad2-Fgf13	0.22
Nkx2.1-Fgf13	0.24
Emx1-Fgf13	0.21

Mutant mice were born in expected Mendelian ratios. Despite embryonic targeted deletion of *Fgf13*, mutant suffered no prenatal mortality related to developmental deficits.




Figure 2.

Excitatory neuronal knockout of *Fgf13* does not result in premature death and seizure susceptibility.

A. Western blot shows partial loss of Fgf13 from *Emx1^{Fgf13} KO* hippocampal tissue, compared to full knockout in *Nestin^{Fgf13} KO* hippocampus. **B.** Fluorescent immunohistochemistry of hippocampal tissue validates *Fgf13* knockout (scale bar, 100 μ m). **C.** *Emx1^{Fgf13} KO* mutant mice survive past one month of age (log-rank test, $p=ns$). **D.** Body mass at postnatal day 14 (P14) shows that *Emx1^{Fgf13} KO* are not different in size (scale bar, 2 cm) (t-test, $p=ns$). **E.** *Emx1^{Fgf13} KO* are not susceptible to hyperthermia induced seizures (log-rank test, $p=ns$). **F.** EEG recordings during hyperthermia protocol show *Emx1^{Fgf13} KO* do not exhibit heat-induced seizures. **G.** *Emx1^{Fgf13} KO* neurons exhibit diminished long-term inactivation (two-way ANOVA, *, $p < 0.05$), though the deficit is not sufficient to cause seizures (WT, $N=2$, $n = 15$; KO, $N=2$, $n = 15$). Example traces for WT and *Emx1^{Fgf13} KO* neurons are shown on the left.

Cre) and infected them with a Cre-dependent AAV8-DIO-GFP virus to allow identification of the *Emx1*⁺ neurons by GFP (Supplemental Figure 2A). The major neuronal voltage gated sodium channels (Na_v1.1, Na_v1.2, Na_v1.3, and Na_v1.6) are expressed in these neuron cultures consistent with levels previously reported (Heighway, Sedo et al. 2022 [\[10\]](#)), as assessed by reverse transcriptase quantitative polymerase chain reaction (Supplemental Figure 2B). Current density and the I-V relationship were not different between WT and *Emx1*^{Fgf13 KO} neurons (Supplemental Figure 2C), and the V_{1/2} of steady-state inactivation was not significantly different in *Emx1*^{Fgf13 KO} neurons compared to WT controls (Supplemental Figure 2D). To assess for long-term inactivation, we applied successive depolarizations (Figure 2G [\[10\]](#)) and observed successive reductions in peak current in WT neurons, a process blunted in *Emx1*^{Fgf13 KO} neurons. Thus, although *Fgf13* ablation in *Emx1*⁺ neurons diminished long-term inactivation—similar to that observed for DEE associated variants in *FGF13-S* expressed in Neuro2A cells (Fry, Marra et al. 2021 [\[11\]](#))—these data suggest that FGF13-S dysfunction in excitatory neurons is an unlikely contributor to a pro-excitatory state and seizures.

Inhibitory neuron knockout of *Fgf13* recapitulates spontaneous seizures and premature death in complete neuronal knockout

Since *Fgf13* knockout in excitatory neurons did not recapitulate the seizure phenotype seen in the *Nestin*^{Fgf13 KO} animals, we suspected that absence of *Fgf13* specifically in inhibitory interneurons may be playing the critical role. With immunohistochemistry on brains from mice in which a *Gad2*-Cre driver (Taniguchi, He et al. 2011 [\[12\]](#)) expressed a Cre-dependent GFP (Ai6) reporter (Madisen, Zwingman et al. 2010 [\[13\]](#)) in inhibitory interneurons (Supplemental Figure 3A-B), one-third of the GFP⁺ interneurons expressed FGF13 at P13 (Figure 3A [\[10\]](#)). We eliminated *Fgf13* specifically in inhibitory neurons by crossing the *Gad2*-Cre driver and *Fgf13*^{fl/fl} mice. The FGF13 signal by immunohistochemistry was completely eliminated from the GFP⁺ cells in mouse brains also expressing GFP from the *Gad2*-Cre driver while the diffuse staining pattern in the excitatory pyramidal cell layer and overall hippocampal morphology were unaffected (Supplemental Figure 3A-B). Further, higher magnification images showed that in *Gad2*^{Fgf13 KO} mice the FGF13⁺ staining in sparse somata (presumed interneurons above) was completely eliminated (Figure 3B [\[10\]](#)). Western blots with the pan-FGF13 antibody showed partial depletion of FGF13 in *Gad2*^{Fgf13 KO} (Figure 3C [\[10\]](#)), with the most significant reduction in FGF13-VY and FGF13-Y while FGF13-S was partially spared, consistent with our analysis of splice-variant specific transcript data (Joglekar, Prjibelski et al. 2021 [\[14\]](#)).

In conjunction with the generation of the *Nestin*^{Fgf13 KO} and *Emx1*^{Fgf13 KO} mice, the *Gad2*^{Fgf13 KO} mice provided the opportunity to validate and extend the previously published transcript data (Joglekar, Prjibelski et al. 2021 [\[14\]](#)) and the protein (western blots in this study) data in cell type-specific knockout models. Using BaseScope *in situ* hybridization we characterized *Fgf13-S* and *Fgf13-VY* expression in intact hippocampi on fixed frozen tissue. We detected diffuse *Fgf13-S* expression in excitatory cell layers while *Fgf13-VY* expression was concentrated in interneuron somata in wild type hippocampi (Supplemental Figure 4A-B). Quantification of the mRNA signal intensity revealed a decrease in *Fgf13-S* for genotypes in which we deleted *Fgf13* in excitatory cells (*Nestin*^{Fgf13 KO} and *Emx1*^{Fgf13 KO}); meanwhile quantification of the cell somata expressing *Fgf13-VY* revealed a decrease in the *Fgf13-VY*⁺ cell count for genotypes in which we deleted *Fgf13* in interneurons (*Nestin*^{Fgf13 KO} and *Gad2*^{Fgf13 KO}), as shown in Supplemental Figure 4C. *In situ* hybridization in these knockout models confirms the splice variant-specific transcript data previously obtained (Joglekar, Prjibelski et al. 2021 [\[14\]](#)), although we did not detect the expected decrease in the *Fgf13-S* signal in *Gad2*^{Fgf13 KO} mice, likely because we lacked the resolution due to the overwhelming *Fgf13-S* signal from excitatory neurons.

With this confirmation, we investigated the consequences of *Fgf13* loss from interneurons. None of the male *Gad2*^{Fgf13 KO} hemizygous knockout mice survived to adulthood (Figure 3D [\[10\]](#)), similar to the *Nestin*^{Fgf13 KO} male mice. The median survival in male *Gad2*^{Fgf13 KO} mice was 14.5 days, and




Figure 3.

Inhibitory neuronal knockout of *Fgf13* recapitulates premature death and seizure susceptibility.

A. Quantification of hippocampal interneuron histology reveals 31% of *Gad2*⁺ interneurons co-express FGF13. **B.** Fluorescent immunohistochemistry of hippocampal tissue validates *Fgf13* knockout in sparse inhibitory interneurons (scale bar, 50 μ m). **C.** Western blot validates partial loss of Fgf13 from *Emx1*^{Fgf13 KO} and *Gad2*^{Fgf13 KO} hippocampal tissue, and full knockout in *Nestin*^{Fgf13 KO} mice. **D.** *Gad2*^{Fgf13 KO} mutant mice have survival deficits around one month of age (log-rank test, ****, $p < 0.0001$). **E.** Body mass at P14 shows *Gad2*^{Fgf13 KO} are smaller in size than wildtype littermates (t-test, ****, $p < 0.001$; scale bar, 2 cm) and brains from *Gad2*^{Fgf13 KO} mice are smaller (t-test, ***, $p < 0.001$; scale bar, 0.5 cm). **F.** *Gad2*^{Fgf13 KO} mice are susceptible to hyperthermia induced seizures (log-rank test, ***, $p < 0.001$). **G.** Example EEG recording of wildtype and *Gad2*^{Fgf13 KO} mice during hyperthermia protocol. Insets show i.) example interictal spike, ii.) and iii.) tonic-clonic seizures.

survivors at P14 were smaller than their wild type littermates (**Figure 3E**). Female *Gad2^{Fgf13}^{KO}* mice also had a survival deficit (**Figure 3D**), though less severe than their male knockout littermates, suggesting a gene dosage effect. Brains of male *Gad2^{Fgf13}^{KO}* mice were smaller compared to wild type littermates, as was body size (**Figure 3E**).

Interneuron-specific knockout of *Fgf13* increased susceptibility to seizures in the hyperthermia-induced seizure protocol. All *Gad2^{Fgf13}^{KO}* males suffered tonic-clonic seizures while wild type littermates did not (**Figure 3F-G**). Furthermore, we observed some *Gad2^{Fgf13}^{KO}* males were susceptible to seizures during acclimation to the testing chamber, prior to the initiation of the hyperthermia protocol. With witnessed seizures immediately preceding death in some mice, we interpret that the previously noted unwitnessed premature deaths in the home cage were likely related to recurrent seizures. Together, these data suggest that FGF13 deficiency in inhibitory cells, rather than excitatory cells, is the major contributor to the seizures and premature mortality seen in the *Nestin^{Fgf13}^{KO}* mice.

Given the seizures and mortality in the *Gad2^{Fgf13}^{KO}* mice and because seizures can be associated with neuronal death (Henshall and Simon 2005), we considered if *Fgf13* ablation led to loss of hippocampal interneurons. We quantified *Gad2*-GFP⁺ interneurons in *Gad2^{GFP}* (*Gad2^{GFP}* vs. *Gad2^{GFP-Fgf13}^{KO}*) mice, in which *Gad2* drove both *Fgf13* ablation and expression of a GFP reporter. We found no difference in the number of *Gad2*-GFP⁺ hippocampal neurons between genotypes (Supplemental Figure 3C), indicating that *Gad2^{Fgf13}^{KO}* mice did not have a deficit of interneurons underlying their seizure phenotype. Further, an unbiased metabolomic analysis on brains from *Gad2^{Fgf13}^{KO}* and WT control mice revealed marked differences between genotypes (Supplemental Figure 5A and Supplemental Table 2) and several features consistent with severe seizures. For example, the metabolite most enriched in *Gad2^{Fgf13}^{KO}* brains was L-kynurenine (Log₂fold change = 2.58, *P* = 0.0009), a metabolite most enriched in patients with status epilepticus (Hanin, Chollet et al. 2024). Metabolite set enrichment analysis in *Gad2^{Fgf13}^{KO}* vs. WT (Supplemental Table 2) revealed that the most enriched Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway among upregulated metabolites was glycolysis (*P* = 4.29E-6), also consistent with data obtained in patients in status epilepticus (Hanin, Chollet et al. 2024). The most downregulated pathway was methionine metabolism (*P* = 9.88E-4), consistent with methionine synthase deficiency—an inborn error of metabolism—that present with seizures (Kripps, Sremba et al. 2022). Thus, this metabolomic analysis supported the severity of the seizure phenotype observed in *Gad2^{Fgf13}^{KO}* mice.

MGE-derived interneuron knockout of *Fgf13* partially recapitulates spontaneous seizures and premature death

As not all *Gad2*⁺ interneurons express *Fgf13* (Supplemental Figure 3A), we analyzed a single cell RNA-seq dataset from mouse CA1 interneurons (Harris, Hochgerner et al. 2018) to search for a subpopulation functionally relevant to the seizure pathology. We found the highest enrichment of *Fgf13* in subsets assigned as axo-axonic and medial ganglionic eminence (MGE) neurogliaform/Ivy cells (Supplemental Figure 6A-B) expressing the homeodomain transcription factor *Nkx2.1*. Further, *Fgf13* expression has previously been identified in *Nkx2.1*⁺ MGE-derived interneurons, including axo-axonic chandelier cells (Paul, Crow et al. 2017; Mahadevan, Mitra et al. 2021), which depend upon *Fgf13* expression for proper development (Favuzzi, Deogracias et al. 2019). Single-cell profiling of human postmortem cortical tissue has identified high *FGF13* expression in neurogliaform and parvalbumin interneurons throughout early development (Herring, Simmons et al. 2022). While *Fgf13* expression was not completely restricted to *Nkx2.1*⁺ MGE-derived interneurons, we thus chose an *Nkx2.1*-Cre driver (Xu, Tam et al. 2008) to target *Fgf13* deletion from interneurons generated in the MGE.

Fluorescent immunohistochemistry in brain slices from *Nkx2.1^{Fgf13} KO* mice showed partial depletion of the sparse somatic distribution of FGF13⁺ somata observed in the *Gad2^{Fgf13} KO* mice (Supplemental Figure 6C). Consistent with that result, BaseScope *in situ* hybridization in *Nkx2.1^{Fgf13} KO* mice showed a pattern similar to *Gad2^{Fgf13} KO* mice but with less complete loss of the interneuron somata (Supplemental Figure S4B). *Nkx2.1^{Fgf13} KO* mice were born at expected Mendelian ratios (Table 1) but only ~ 40% of male *Nkx2.1^{Fgf13} KO* mice survived past one month of age (Supplemental Figure 6D), consistent with a partial loss of FGF13 expression in hippocampal interneurons (Supplemental Figure 6C). The median survival age for male *Nkx2.1^{Fgf13} KO* mice was 22 days (P22), and surviving mutants were smaller than their wild type littermates at P14 (Supplemental Figure 6E). Female *Nkx2.1^{Fgf13} HET* survived to adulthood. We found that almost all male *Nkx2.1^{Fgf13} KO* suffered tonic-clonic seizures during the hyperthermia induction protocol, while wild type littermates did not (Supplemental Figure 6F). Because the lethality and seizure susceptibility in the *Nkx2.1^{Fgf13} KO* mice were reduced compared to the *Gad2^{Fgf13} KO* mice, we hypothesize that MGE-derived neurons are partially responsible for the phenotype observed in *Gad2^{Fgf13} KO* mice. The loss of FGF13 from other interneuron classes in which FGF13 is expressed—as evident from single cell RNA sequencing (Supplemental Figure 6A-B)—likely contributes to the more severe phenotype after broader inhibitory interneuron knockout by the *Gad2*-Cre driver.

Inhibitory neuronal knockout of *Fgf13* results in deficits of synaptic transmission

To identify if the seizure susceptibility of the *Gad2^{Fgf13} KO* mice is caused by loss of excitatory/inhibitory balance in the hippocampus, we recorded spontaneous excitatory and inhibitory inputs to the CA1 pyramidal cells of the dorsal hippocampus in slice recordings. Pyramidal cells in *Gad2^{Fgf13} KO* hippocampi, compared to wild type littermates, showed an increased frequency of sEPSCs (Figure 4A-B) and a decreased frequency of sIPSCs mice relative to wild type littermates (Figure 4C-D). Because there was no significant change in sEPSC or sIPSC amplitude (Figure 4B, D), the increased sEPSC and decreased sIPSC frequencies indicate a presynaptic and not postsynaptic mechanism. Together, these data suggest that ablation of *Fgf13* specifically in inhibitory neurons (in *Gad2^{Fgf13} KO* mice) decreased direct and downstream inhibitory drive onto pyramidal cells, thereby increasing pyramidal cell excitability.

Inhibitory neuronal knockout of *Fgf13* results in deficits of interneuron excitability

To assess the cellular mechanisms underlying decreased inhibitory drive from the inhibitory neurons in *Gad2^{Fgf13} KO* mice, we cultured hippocampal neurons from *Gad2^{Fgf13} KO* or wild type littermates (*Gad2*-Cre) and infected them with a Cre-dependent AAV8-DIO-GFP virus, thus allowing identification of the *Gad2*⁺ interneurons by GFP (Figure 5A). We quantified trains of action potentials in response to current injection in the GFP⁺ neurons to assess intrinsic excitability. Increasing current injection augmented the number of spikes in WT (*Gad2*-Cre) neurons until a plateau at ~260 pA (Figure 5B). At low levels (<100 pA) current injection elicited a similar number of spikes in neurons from *Gad2^{Fgf13} KO* mice (albeit spike morphology and frequency were different, see below) but the number of spikes then plateaued and was less than the number elicited in neurons from WT mice at higher amounts of current injection (Figure 5B), suggesting a decrease in intrinsic excitability in neurons from *Gad2^{Fgf13} KO* mice. Examination of individual traces revealed that more *Gad2^{Fgf13} KO* cells entered depolarization block, and at lower levels of current injection than wildtype interneurons (Figure 5C). The detected difference between genotypes is striking considering that only a third of the *Gad2*⁺ neurons from *Gad2^{Fgf13} KO* infected with the Cre virus would be expected to express FGF13 before viral Cre expression (Figure 3A).




Figure 4.

Inhibitory neuronal knockout of *Fgf13* results in deficits of synaptic transmission.

A. Example traces from spontaneous excitatory postsynaptic currents (sEPSCs). **B.** Cumulative fraction (Kolmogorov Smirnov test, ****, $p < 0.0001$) frequency, and current amplitude for sEPSCs (t-test, ****, $p < 0.0001$, WT $N=7$, $n=24$; KO $N=4$, $n=22$). **C.** Example traces from spontaneous inhibitory postsynaptic currents (sIPSCs). **D.** Cumulative fraction (Kolmogorov Smirnov test, ****, $p < 0.0001$), frequency, and current amplitude for sIPSCs (t-test, ****, $p < 0.0001$, WT $N=3$, $n=26$; KO $N=4$, $n=25$).




Figure 5.

Inhibitory neuronal knockout of *Fgf13* results in deficits of interneuron excitability.

A. Examples of FGF13-stained neurons from primary hippocampal neuron cultures generated from *Gad2*-Cre wildtype (top) and *Gad2^{Fgf13} KO* (bottom) mice (scale bar, 20 μ m). **B.** Example traces of AP spike trains from wildtype (left) and *Gad2^{Fgf13} KO* interneurons. Input-output curve shows decreased firing of evoked action potentials from *Gad2^{Fgf13} KO* interneurons (two-way ANOVA, **, $p < 0.01$, WT $N=4$, $n=28$; KO $N=4$, $n=26$). **C.** *Gad2^{Fgf13} KO* interneurons enter depolarization block at earlier current injections than wildtype interneurons (log-rank test, **, $p < 0.01$). **D.** Rheobase and threshold potential for cultured wildtype and *Gad2^{Fgf13} KO* interneurons. **E.** Action potential wave forms and phase plots (mean $\pm$ s.e.m.) for the first elicited action potential of the spike train for wildtype and *Gad2^{Fgf13} KO* interneurons. **F.** Action potential wave forms and phase plots for the second and third action potentials of the spike train of wildtype and *Gad2^{Fgf13} KO* interneurons. **G.** Analysis of the first three action potentials in the spike train show a decrease in the dV/dt_{max} , a decrease in action potential (AP) peak, a decrease in AP amplitude, and an increase in APD50 by the third AP of *Gad2^{Fgf13} KO* interneurons (two-way ANOVA, *, $p < 0.05$). **H.** Analysis of the first three action potentials in the spike train show an increase in membrane potential at the end of the AP for *Gad2^{Fgf13} KO* interneurons (two-way ANOVA, **, $p < 0.01$, ****, $p < 0.0001$). **I.** Example traces of K^+ currents from wildtype and *Gad2^{Fgf13} KO* interneurons (left). I-V curve for K^+ currents (right), (two-way ANOVA, *, $p < 0.05$, WT $N=3$, $n=26$; KO $N=5$, $n=29$).

To further investigate the mechanisms contributing to decreased excitability and depolarization block, we examined the first elicited action potential properties for each spike train at rheobase. Resting membrane potential, rheobase, and threshold potential were not different between genotypes, **Figure 5D** [↗](#). The single action potential wave forms and phase plots show little difference between *Gad2^{Fgf13 KO}* neurons infected with the Cre virus and wild type controls (**Figure 5E** [↗](#)), although action potential duration measured at 50% of the amplitude (APD50) trended longer in the *Gad2^{Fgf13 KO}* interneurons infected with the Cre virus (Supplemental Figure 7A). However, we observed marked genotype differences in the second and third elicited action potentials. Most notably, in *Gad2^{Fgf13 KO}* interneurons we observed a progressive reduction in the action potential peak amplitude and dV/dt max, an increase in APD50, and a greater increase in the interspike membrane voltage (**Figure 5F-G** [↗](#)), suggesting a growing repolarization deficit in the *Gad2^{Fgf13 KO}* neurons each subsequent spike, thus increasing the likelihood to undergo depolarization block.

Since the first spike action potential parameters were not significantly different between *Gad2^{Fgf13 KO}* neurons and wild type controls, we hypothesized that loss of FGF13 had minimal effects upon Na_V channel currents in these neurons. Indeed, we observed no differences in current amplitude nor steady-state inactivation between *Gad2^{Fgf13 KO}* neurons and wild type controls (Supplemental Figure 7B, C). Because we did not detect significant differences in Na_V current properties typically associated with FGF13 in *Fgf13* knockout in *Gad2^{Fgf13 KO}* neurons, we hypothesized that FGF13 regulated other currents that led to the observed changes in action potentials and decrease in repolarization. Since in cardiomyocytes we showed that FGF13 can also affect various K_V channels that contribute to the repolarizing phase of the cardiac action potential (Wang, Tang et al. 2017 [↗](#)), here we recorded macroscopic potassium currents and observed a reduced K_V current density in *Gad2^{Fgf13 KO}* interneurons compared to WT neurons (**Figure 5I** [↗](#)).

Virally mediated re-expression of *Fgf13-S* and

Fgf13-VY restores output to *Gad2^{Fgf13 KO}* neurons

Since inhibitory neurons express both *Fgf13-S* and *Fgf13-VY*, we tested if re-expression of either of these isoforms in *Gad2^{Fgf13 KO}* neurons could rescue the observed deficit in the input/output relationship. We isolated primary hippocampal neurons from *Gad2^{Fgf13 KO}* or controls (*Gad2-Cre*) and infected them with the Cre-dependent AAV8-DIO-GFP virus along with either an AAV8-DIO-*Fgf13-S* or AAV8-DIO-*Fgf13-VY*. Immunocytochemistry of GFP⁺ neurons showed expression of endogenous FGF13 in *Gad2-Cre* cultures and the absence of FGF13 in *Gad2^{Fgf13 KO}* cultures (**Figure 6A** [↗](#)). Infection with *Fgf13-S* in a *Gad2^{Fgf13 KO}* neuron showed concentrated expression in the axon initial segment and membrane-enriched signal in the soma and the dendritic branches (**Figure 6A** [↗](#)) while infection with *Fgf13-VY* produced prominent signal throughout the soma and branches, demonstrating successful re-expression of these splice variants in the knockout background. After viral re-expression of *Fgf13-VY* or *Fgf13-S*, neither threshold potential nor rheobase were affected and were not different compared to values from either WT or *Gad2^{Fgf13 KO}* neurons (**Figure 6B** [↗](#)). After rescue with either *Fgf13-VY* or *Fgf13-S*, current injection >100 pA elicited more action potentials than in *Gad2^{Fgf13 KO}* neurons, suggesting that excitability was at least partially restored (**Figure 6C** [↗](#)). Indeed, compared to *Gad2^{Fgf13 KO}* neurons a smaller fraction of *Gad2⁺* cells re-expressing *Fgf13-VY* or *Fgf13-S* reached depolarization block when injected with >100 pA (**Figure 6D** [↗](#)), although neither *Fgf13-VY* nor *Fgf13-S* alone reduced the probability of conduction block to what was observed in WT neurons. Further analysis showed that, even though *Fgf13-S* or *Fgf13-VY* partially restored spiking, the action potential wave forms and the phase plots were not restored by either *Fgf13-S* or *Fgf13-VY* alone (**Figure 6E** [↗](#)). Examination of specific parameters revealed that neither *Fgf13-S* nor *Fgf13-VY* rescue was able to prevent the prolongation of APD50 nor reduction in the action potential amplitude, dV/dt max or action potential peak amplitude observed in *Gad2^{Fgf13 KO}* neurons (**Figure 6F** [↗](#)), again suggesting that neither FGF13-S nor FGF13-VY is sufficient for maintaining excitability in *Gad2* neurons. On the other hand, infection with *Fgf13-S* partially hyperpolarized the interspike membrane potential

(compared to *Gad2^{Fgf13 KO}*, **Figure 6G** [↗](#)), suggesting that FGF13-S increased the repolarizing K⁺ currents. Indeed, infection with *Fgf13-S* but not *Fgf13-VY* increased K⁺ current density to WT levels (**Figure 6H** [↗](#)). These data suggest that FGF13-S and FGF13-VY, both expressed in inhibitory interneurons (Joglekar, Prjibelski et al. 2021 [↗](#)) contribute to inhibitory interneuron function and modulation of pyramidal neuron excitability.

Discussion

While variants in or disruption of *FGF13* have been associated with DEEs (Puranam, He et al. 2015 [↗](#), Fry, Marra et al. 2021 [↗](#), Velíšková, Marra et al. 2021 [↗](#), Narayanan, Majethia et al. 2022 [↗](#)), the pathogenesis was unknown. Employing multiple cell type specific mouse knockout models complemented by isoform specific rescue strategies, our data reveal how *FGF13* ablation contribute to seizure susceptibility and provide new insights into FGF13 functions in neurons. We found that *Fgf13* ablation and consequent loss of FGF13-S and FGF13-VY in a subset of interneurons reduced overall inhibitory drive from *Fgf13*-expressing interneurons onto hippocampal pyramidal neurons. Rescue experiments suggest that both FGF13-S and FGF13-VY isoforms are required for proper interneuron function as re-expression of FGF13-S or FGF13-VY only partially restored excitability (our experimental setup did not allow rescue with both isoforms simultaneously). Building upon our recent analysis of *Fgf13* splice variant expression (Joglekar, Prjibelski et al. 2021 [↗](#)), our data resolve apparently conflicting reports in which reduced inhibitory interneuron function, suggested as the main contributor in a global *Fgf13* female heterozygous knockout model, (Puranam, He et al. 2015 [↗](#)) contrasts with a model that proposes that *FGF13* variants within excitatory neurons drive increased excitability because of diminished long-term inactivation (Fry, Marra et al. 2021 [↗](#)).

The apparent requirement of both FGF13-S and FGF13-VY extends previous studies showing their distinct roles in neurons (Pablo, Wang et al. 2016 [↗](#)) and emphasizes that alternative splicing is an important source of protein diversity in the mammalian brain, an evolutionarily conserved feature thought to contribute to the nervous system's complexity (Raj and Blencowe 2015 [↗](#), Zhang, Chen et al. 2016 [↗](#), Karlsson and Linnarsson 2017 [↗](#), Porter, Jaamour et al. 2018 [↗](#), Furlanis, Traunmuller et al. 2019 [↗](#), Joglekar, Foord et al. 2023 [↗](#)). Cell-specific alternative splicing in the brain has been posited as a mechanism to efficiently adapt to the external environment by quickly modifying protein activity (Lipscombe and Lopez Soto 2019 [↗](#)). Not only does *FGF13* have the most extensive set of alternatively spliced isoforms among FHF's (Smallwood, Munoz-Sanjuan et al. 1996 [↗](#), Munoz-Sanjuan, Smallwood et al. 2000 [↗](#), Pablo, Wang et al. 2016 [↗](#)), but long-range single cell sequencing data show that *Fgf13* has a uniquely (among FHF's) segregated alternative splice variant expression pattern among distinct neuron cell type. Further, among all genes, has one of the highest differential isoform expression (DIE) values across multiple neuronal cell-types (Joglekar, Prjibelski et al. 2021 [↗](#)).

Our results are also consistent with the available patient data. The responsible splice variant(s) were proposed to be either *FGF13-V*, *FGF13-VY*, and *FGF13-Y* that were lost in two affected brothers who inherited a maternal balanced translocation that disrupts *FGF13* on the X-chromosome and preserves the *FGF13-S* and the *FGF13-U* splice variants (Puranam, He et al. 2015 [↗](#)) or, in contrast, *FGF13-S* in a series of unrelated individuals presenting with DEE symptoms (Fry, Marra et al. 2021 [↗](#)). As our data suggest that both *FGF13-S* and *FGF13-VY* contribute within inhibitory interneurons, we provide a unifying explanation for those reports. Here, we were able to assess FGF13-S and FGF13-VY, chosen because they are most abundantly expressed isoforms in the adult mouse brain, but the inability to rescue electrophysiological consequences completely with either isoform alone leaves open the possibility that other isoforms (e.g., FGF13-U, FGF13-V) also make critical contributions.




Figure 6.

AAV-mediated expression of FGF13 isoforms rescues excitability deficits in *Gad2^{Fgf13} KO* neurons.

A. Examples of FGF13-stained neurons from primary hippocampal neuron cultures generated from *Gad2^{Fgf13} KO* mice transduced with AAV8-DIO-GFP only, AAV8-DIO-GFP and AAV8-DIO-Fgf13-S, or AAV8-DIO-GFP and AAV8-DIO-Fgf13-VY (scale bar, 20 μ m). **B.** *Gad2^{Fgf13} KO* neurons expressing FGF13-VY or FGF13-S were not different in terms of threshold potential or rheobase, and were not different from wildtype (black line, from **Figure 5**) and *Gad2^{Fgf13} KO* neurons (red line, from **Figure 5**) (t-test, p =ns, KO+VY N =3, n =14; KO+S N =3, n =17). **C.** Evoked action potential traces from *Gad2^{Fgf13} KO* interneurons expressing FGF13-VY or FGF13-S. Input-output curve shows increased firing of evoked action potentials from the FGF13-VY or FGF13-S expressing interneurons, relative to *Gad2^{Fgf13} KO* interneurons (red line, from **Figure 5**; black line = wild type, from **Figure 5**) (two-way ANOVA, *, p <0.05). **D.** *Gad2^{Fgf13} KO* interneurons expressing FGF13-VY and FGF13-S do not enter depolarization block as early as *Gad2^{Fgf13} KO* interneurons (Red line [*Gad2^{Fgf13} KO*] and black line [wild type] are from **Figure 5**) (log-rank test, *, p <0.05). **E.** Action potential wave forms and phase plots for the initial three action potentials of the spike train for FGF13-VY and FGF13-S rescued *Gad2^{Fgf13} KO* interneurons. The black and red lines are from **Figure 5**. **F.** For the first three action potentials in the spike train, *Gad2^{Fgf13} KO* neurons re-expressed with FGF13-VY show no difference from *Gad2^{Fgf13} KO* neurons in terms of dV/dt max, AP peak, and AP amplitude, and AP50. *Gad2^{Fgf13} KO* neurons re-expressed with FGF13-S show difference from *Gad2^{Fgf13} KO* neurons only for the first action potential for AP peak and AP amplitude, but not dV/dt max or AP50. (two-way ANOVA, ***, p <0.001, KO+S vs. KO; **, p <.01, KO+S vs. KO). **G.** FGF13-S rescued neurons show a significant decrease in membrane voltage from *Gad2^{Fgf13} KO* neurons by the third action potential in the spike train (two-way ANOVA, *, p <0.05, KO+S vs. KO). **H.** Example traces of K^+ currents from *Gad2^{Fgf13} KO* neurons expressing FGF13-VY and FGF13-S. K^+ currents are rescued by expression of FGF13 in *Gad2^{Fgf13} KO* interneurons (two-way ANOVA, *, p <0.05, KO+VY N =5, n =21, KO+S N =5, n =21).

These results also expand our understanding of how *FGF13* contributes to neuronal functions, which have been challenging to define because of *FGF13*'s location on the X chromosome, the embryonic lethality of complete knockout models, absence of cell-type specific knockout models, *FGF13*'s complex alternative splicing, and an incomplete cataloging of specific functions for the various isoforms generated. Various roles for *FGF13* in inhibitory interneurons have previously been suggested. *Fgf13* silencing in *Nkx2.1-CreER* targeted neurons reduced the number of chandelier cell boutons (Favuzzi, Deogracias et al. 2019 [\[1\]](#)) and a more general role for *FGF13* in interneuron development was suggested by a human scRNA-seq dataset (Herring, Simmons et al. 2022 [\[2\]](#)). Our data here, in which seizure susceptibility was attributable solely to *Fgf13* knockout in inhibitory interneurons and our demonstration that *Fgf13* knockout restricted to *Nkx2.1*-expressing neurons increased seizure susceptibility—albeit to a lesser extent than complete knockout in inhibitory interneurons—support not only a possible role for *Fgf13* in chandelier cell development but also confirm more general inhibitory interneuron roles. Along with the reduced seizure susceptibility of female heterozygote knockout animals, these data also imply a gene dosage effect within interneurons. The differences in survival among *Gad2^{Fgf13 KO}*, *Gad2^{Fgf13 HET}*, *Nkx2.1^{Fgf13 KO}*, and *Nkx2.1^{Fgf13 HET}* also suggest a gene-dosage effect. Although we were unable to generate female homozygous *Gad2^{Fgf13 KO}* mice since the male hemizygous knockouts did not survive to adulthood, the reduced survival in male knockouts compared to female heterozygous knockouts further supports our hypothesis. The relative comprehensiveness of the specific Cre drivers also provides support for our hypothesis. In the hyperthermic seizure assay, mice from both the *Nestin^{Fgf13 KO}* and *Gad2^{Fgf13 KO}* lines, in which *Fgf13* was ablated in the entire interneuron population, suffered seizures at baseline (near 36 °C) as well as at elevated core body temperatures. In contrast, the majority of the *Nkx2.1^{Fgf13 KO}* mice, generated with a more restrictive Cre driver, mostly suffered seizures only at elevated core body temperatures. We also observed occasional seizures in the home cage for all genotypes in which *Fgf13* was ablated in interneurons (*Nestin^{Fgf13 KO}*, *Nkx2.1^{Fgf13 KO}*, and *Gad2^{Fgf13 KO}*) and often discovered dead mice, which we suspect was a SUDEP phenotype due to recurrent seizures, given that recurrent uncontrolled seizures are a major risk factor for sudden death (Devinsky, Hesdorffer et al. 2016 [\[3\]](#), Richerson 2023 [\[4\]](#)). Indeed, long term recording of *Gad2^{Fgf13 KO}* mice (data not shown) confirmed that multiple spontaneous seizures occur in the knockouts prior to death in the home cage.

While our data provide a clear connection between ablation of *Fgf13* in inhibitory neurons and seizure susceptibility, we find that *Fgf13* ablation in excitatory neurons does not increase seizure susceptibility. Indeed, complete neuronal knockout with *Nestin-Cre* conferred a relative increased survival compared to the interneuron specific knockout with *Gad2-Cre*. We speculate that this relative increased survival in the *Nestin^{Fgf13 KO}* mice could be technical, due to less efficient targeting of interneurons by *Nestin-Cre* (Liang, Hippenmeyer et al. 2012 [\[5\]](#)), but it may also be biological if *Nestin-Cre* targeting of both excitatory and inhibitory neurons is comparatively neuroprotective because of simultaneous loss of both excitation and inhibition and consequent compensation for the loss of inhibition. The previous observation that a global *Fgf13* heterozygous mouse line survived with a normal life span (Puranam, He et al. 2015 [\[6\]](#)) could support the interpretation that deletion from other cell types is comparatively neuroprotective. Moreover, our observations that our *Emx1^{Fgf13 KO}* mice do not suffer seizures is consistent with an earlier study of *Emx1^{Fgf13 KO}* mice, in which seizures were not reported (Wu, Yang et al. 2012 [\[7\]](#)). Previous studies suggested that FGF13-S, the isoform most highly expressed in excitatory neurons, confers long-term inactivation (Venkatesan, Liu et al. 2014 [\[8\]](#), Fry, Marra et al. 2021 [\[9\]](#)), and the reduction of long-term inactivation due to DEE-associated variants drives increased excitability as the mechanism for seizures (Fry, Marra et al. 2021 [\[9\]](#)). Our data confirm the ability of FGF13-S to impart long-term inactivation and that ablation of *Fgf13* in excitatory neurons eliminates long-term inactivation, yet this mechanism does not likely contribute to seizure susceptibility. Rather, knockout of *Fgf13* in central nervous system excitatory neurons appears to affect learning and memory (Wu, Yang et al. 2012 [\[7\]](#)), which we did not assess here. An important limitation of our knockout models, however, is that they cannot provide direct insight into how missense variants reported in patients contribute to seizures. Nevertheless, since our *Emx1^{Fgf13 KO}* mice show a loss

of long-term inactivation in excitatory neurons—the proposed mechanism by which the missense variants drive seizures—our model provides new boundaries on how those missense variants act in excitatory neurons.

Further, other than long-term inactivation, data herein show that *Fgf13* ablation led to relatively minor effects on neuronal voltage-gated Na^+ currents in *Emx1*⁺ excitatory neurons compared to previous results from others and us (Venkatesan, Liu et al. 2014 [\[1\]](#), Barbosa, Xiao et al. 2016 [\[2\]](#), Pablo, Wang et al. 2016 [\[3\]](#), Wang, Yang et al. 2021 [\[4\]](#), Shen, Yue et al. 2022 [\[5\]](#)). Those neuronal studies investigated DRG neurons, or CNS neurons via RNAi-mediated knockdown of infusion of antibodies targeting the FGF13-S N-terminus, and not a cell-type specific knockout as we studied here, thus suggesting differences in the models studied. As the *FGF13* DEEs result from a stable germ-line variants, the knockout model here (in contrast to knockdown approaches in cultured CNS neurons or acute infusion of antibodies) likely represents a more relevant disease model.

How does loss of FGF13 in inhibitory interneurons confer seizure susceptibility? Although knockout of *Fgf13* in *Gad2*⁺ neurons decreased excitability and reduced overall inhibitory drive to pyramidal cells, knockout surprisingly did not affect Na_V currents in inhibitory neurons. This conclusion fits well with our previous report that the cytoplasmic C-terminal in $\text{Na}_V1.1$, the major Na_V channel expressed in inhibitory neurons, has a substitution of an amino acid that reduces FHF binding (Wang, Chung et al. 2012 [\[6\]](#)), thus rendering these $\text{Na}_V1.1$ channels—and thus Na_V current in inhibitory interneurons—poor targets for FHF regulation. Rather, we found that perturbed excitability in interneurons lacking *Fgf13* results from reduced K^+ currents. This fits with our demonstration in cardiomyocytes that loss of FGF13 similarly reduces repolarizing K_V currents (Wang, Tang et al. 2017 [\[7\]](#)). Moreover, these data build upon the growing recognition that FHFs exert effects beyond Na_V channel regulation—the focus of most FHF studies since FHFs were identified as capable of directly interacting with Na_V channel C-termini (Liu, Dib-Hajj et al. 2001 [\[8\]](#)). Other identified roles for FGF13, such as regulating microtubule stability (Wu, Yang et al. 2012 [\[9\]](#)), affecting ribosome biogenesis (Bublik, Bursac et al. 2017 [\[10\]](#)), or generating axoaxonic synapses in chandelier cells (Favuzzi, Deogracias et al. 2019 [\[11\]](#)) may also contribute, but we did not investigate those roles here. The mechanism(s) by which FGF13 affects K_V currents is not known—we have not detected direct interaction between FHFs and K_V channels—but we speculate that the identified role for FGF13 in stabilizing microtubules may contribute, as microtubules regulate trafficking of ion channels to the plasma membrane during channel biogenesis (Vacher, Mohapatra et al. 2008 [\[12\]](#)).

Although we did not test it directly, our data are consistent with a possible role for FGF13 in chandelier cell function. Chandelier cells are fast-spiking GABAergic interneurons that uniquely target the axon initial segment of pyramidal cells. Chandelier cells are found sparsely in the brain, but are highly branched, enabling contact on hundreds of neighboring pyramidal cells (Compans and Burrone 2023 [\[13\]](#)). *Fgf13* has previously been identified as a key molecule in the development of chandelier cell axons (Favuzzi, Deogracias et al. 2019 [\[11\]](#)), but whether *Fgf13* subsequently contributes to their function has not been reported. Chandelier cells originate from the MGE, alongside parvalbumin- and somatostatin-expressing interneurons (Gallo, Paul et al. 2020 [\[14\]](#)). As our *Nkx2.1*^{*Fgf13* KO} mice target deletion of *Fgf13* from all the above subtypes, our data suggest that FGF13 may regulate chandelier cell function in seizure generation. Although chandelier cells have been shown to depolarize pyramidal cells and therefore exert a pro-excitatory role in the cortex (Szabadics, Varga et al. 2006 [\[15\]](#)), we observe otherwise in our mouse model in which decreased interneuron excitability results in a pro-excitatory state.

Understanding the molecular mechanisms that drive the pathogenesis of *Fgf13*-related seizures will enhance our understanding of DEEs and epilepsy at large. Future work will reveal the mechanisms of FGF13 interaction with potassium channels or other molecules that enable interneuron repolarization and excitability.

Acknowledgements

Supported by T32 DA03980 and F31 DA053796 (SL) and R01 HL160089 (GSP). We thank Andrew S. Lee and Gülcan Akgül for critical reading of the manuscript, as well as Mattia Malvezzi and Maiko Matsui for thoughtful discussions.

Conflicts of Interest

GSP is a scientific advisory board member for Tevard Biosciences. None of the other authors report any conflicts.

References

- Adam M. P., Feldman J., Mirzaa G. M., Pagon R. A., Wallace S. E., Bean L. J. H., Gripp K. W., Amemiya A. (1993) **GeneReviews** Seattle (WA): University of Washington, Seattle
- Barbosa C., Xiao Y., Johnson A. J., Xie W., Strong J. A., Zhang J.-M., Cummins T. R. (2016) **FHF2 isoforms differentially regulate Nav1.6-mediated resurgent sodium currents in dorsal root ganglion neurons** *Pflügers Archiv - European Journal of Physiology* :1–18
- Bouza A. A., Isom L. L. (2018) **Voltage-Gated Sodium Channel beta Subunits and Their Related Diseases** *Handb Exp Pharmacol* **246**:423–450
- Brunklaus A. *et al.* (2022) **Development and Validation of a Prediction Model for Early Diagnosis of Neurology** **98**:e1163–e1174
- Bublik D. R. *et al.* (2017) **Regulatory module involving FGF13, miR-504, and p53 regulates ribosomal biogenesis and supports cancer cell survival** *Proc Natl Acad Sci U S A* **114**:E496–E505
- Bunton-Stasyshyn R. K. A. *et al.* (2019) **Prominent role of forebrain excitatory neurons in SCN8A encephalopathy** *Brain* **142**:362–375
- Cheah C. S., Yu F. H., Westenbroek R. E., Kalume F. K., Oakley J. C., Potter G. B., Rubenstein J. L., Catterall W. A. (2012) **Specific deletion of Nav1.1 sodium channels in inhibitory interneurons causes seizures and premature death in a mouse model of Dravet syndrome** *Proc Natl Acad Sci U S A* **109**:14646–14651
- Chilcott E., Diaz J. A., Bertram C., Berti M., Karda R. (2022) **Genetic therapeutic advancements for Dravet Syndrome** *Epilepsy Behav* **132**
- Compans B., Burrone J. (2023) **Chandelier cells shine a light on the formation of GABAergic synapses** *Curr Opin Neurobiol* **80**
- Covanis A. (2012) **Epileptic encephalopathies (including severe epilepsy syndromes)** *Epilepsia* **53**:114–126
- Devinsky O., Hesdorffer D. C., Thurman D. J., Lhatoo S., Richerson G. (2016) **Sudden unexpected death in epilepsy: epidemiology, mechanisms, and prevention** *Lancet Neurol* **15**:1075–1088
- Ding J., Li X., Tian H., Wang L., Guo B., Wang Y., Li W., Wang F., Sun T. (2021) **Mutation-Beyond Dravet Syndrome: A Systematic Review and Narrative Synthesis** *Front Neurol* **12**
- Donnan A. M., Schneider A. L., Russ-Hall S., Churilov L., Scheffer I. E. (2023) **Rates of Status Epilepticus and Sudden Unexplained Death in Epilepsy in People With Genetic Developmental and Epileptic Encephalopathies** *Neurology* **100**:e1712–e1722
- Favuzzi E. *et al.* (2019) **Distinct molecular programs regulate synapse specificity in cortical inhibitory circuits** *Science* **363**:413–417

Fry A. E. *et al.* (2021) **Missense variants in the N-terminal domain of the A isoform of FHF2/FGF13 cause an X-linked developmental and epileptic encephalopathy** *Am J Hum Genet* **108**:176–185

Fry A. E. *et al.* (2021) **Missense variants in the N-terminal domain of the A isoform of FHF2/FGF13 cause an X-linked developmental and epileptic encephalopathy** *Am J Hum Genet* **108**:176–185

Furlanis E., Traunmuller L., Fucile G., Scheiffele P. (2019) **Landscape of ribosome-engaged transcript isoforms reveals extensive neuronal-cell-class-specific alternative splicing programs** *Nat Neurosci* **22**:1709–1717

Gallo N. B., Paul A., Van Aelst L. (2020) **Shedding Light on Chandelier Cell Development, Connectivity, and Contribution to Neural Disorders** *Trends Neurosci* **43**:565–580

Goldfarb M. (2005) **Fibroblast growth factor homologous factors: evolution, structure, and function** *Cytokine Growth Factor Rev* **16**:215–220

Gorski J. A., Talley T., Qiu M., Puelles L., Rubenstein J. L., Jones K. R. (2002) **Cortical excitatory neurons and glia, but not GABAergic neurons, are produced in the Emx1-expressing lineage** *J Neurosci* **22**:6309–6314

Han Z. *et al.* (2020) **Antisense oligonucleotides increase** *Sci Transl Med* **12**

Han Z. *et al.* (2020) **Antisense oligonucleotides increase Scn1a expression and reduce seizures and SUDEP incidence in a mouse model of Dravet syndrome** *Sci Transl Med* **12**

Hanin A., Chollet C., Demeret S., Di Meglio L., Castelli F., Navarro V. (2024) **Metabolomic changes in adults with status epilepticus: A human case-control study** *Epilepsia* **65**:929–943

Harris K. D., Hochgerner H., Skene N. G., Magno L., Katona L., Bengtsson Gonzales C., Somogyi P., Kessaris N., Linnarsson S., Hjerling-Leffler J. (2018) **Classes and continua of hippocampal CA1 inhibitory neurons revealed by single-cell transcriptomics** *PLoS Biol* **16**

He Z., Li Y., Zhao X., Li B. (2022) **Dravet syndrome: Advances in etiology, clinical presentation, and treatment** *Epilepsy Res* **188**

Heighway J., Sedo A., Garg A., Eldershaw L., Perreau V., Berecki G., Reid C. A., Petrou S., Maljevic S. (2022) **Sodium channel expression and transcript variation in the developing brain of human, Rhesus monkey, and mouse** *Neurobiol Dis* **164**

Henshall D. C., Simon R. P. (2005) **Epilepsy and apoptosis pathways** *J Cereb Blood Flow Metab* **25**:1557–1572

Herring C. A. *et al.* (2022) **Human prefrontal cortex gene regulatory dynamics from gestation to adulthood at single-cell resolution** *Cell* **185**:4428–4447

Hull J. M., O'Malley H. A., Chen C., Yuan Y., Denomme N., Bouza A. A., Anumonwo C., Lopez-Santiago L. F., Isom L. L. (2020) **Excitatory and inhibitory neuron defects in a mouse model of Scn1b-linked EIEE52** *Ann Clin Transl Neurol* **7**:2137–2149

Joglekar A., Foord C., Jarroux J., Pollard S., Tilgner H. U. (2023) **From words to complete phrases: insight into single-cell isoforms using short and long reads** *Transcription* **14**:92–104

- Joglekar A. *et al.* (2021) **A spatially resolved brain region- and cell type-specific isoform atlas of the postnatal mouse brain** *Nat Commun* **12**
- Kalume F., Westenbroek R. E., Cheah C. S., Yu F. H., Oakley J. C., Scheuer T., Catterall W. A. (2013) **Sudden unexpected death in a mouse model of Dravet syndrome** *J Clin Invest* **123**:1798–1808
- Karlsson K., Linnarsson S. (2017) **Single-cell mRNA isoform diversity in the mouse brain** *BMC Genomics* **18**
- Kim H. J., Yang D., Kim S. H., Kim B., Kim H. D., Lee J. S., Choi J. R., Lee S. T., Kang H. C. (2020) **The phenotype and treatment of SCN2A-related developmental and epileptic encephalopathy** *Epileptic Disord* **22**:563–570
- Kim Y. O. *et al.* (2013) **Do mutations in SCN1B cause Dravet syndrome?** *Epilepsy Res* **103**:97–100
- Klorig D. C., Alberto G. E., Smith T., Godwin D. W. (2019) **Optogenetically-Induced Population Discharge Threshold as a Sensitive Measure of Network Excitability** *eNeuro* **6**
- Kripps K. A. *et al.* (2022) **“Methionine synthase deficiency: Variable clinical presentation and benefit of early diagnosis and treatment.”** *J Inherit Metab Dis* **45**:157–168
- Li M. *et al.* (2021) **Antisense oligonucleotide therapy reduces seizures and extends life span in an SCN2A gain-of-function epilepsy model** *J Clin Invest* **131**
- Liang H., Hippenmeyer S., Ghashghaei H. T. (2012) **A Nestin-cre transgenic mouse is insufficient for recombination in early embryonic neural progenitors** *Biol Open* **1**:1200–1203
- Lipscombe D., Lopez Soto E. J. (2019) **Alternative splicing of neuronal genes: new mechanisms and new therapies** *Curr Opin Neurobiol* **57**:26–31
- Liu C., Dib-Hajj S. D., Waxman S. G. (2001) **Fibroblast growth factor homologous factor 1B binds to the C terminus of the tetrodotoxin-resistant sodium channel rNav1.9a (NaN)** *J Biol Chem* **276**:18925–18933
- Lu H., Shi X., Wu G., Zhu J., Song C., Zhang Q., Yang G. (2015) **FGF13 regulates proliferation and differentiation of skeletal muscle by down-regulating Spry1** *Cell Prolif* **48**:550–560
- Madisen L. *et al.* (2010) **A robust and high-throughput Cre reporting and characterization system for the whole mouse brain** *Nat Neurosci* **13**:133–140
- Mahadevan V. *et al.* (2021) **NMDARs Drive the Expression of Neuropsychiatric Disorder Risk Genes Within GABAergic Interneuron Subtypes in the Juvenile Brain** *Front Mol Neurosci* **14**
- Meisler M. H. (2019) **SCN8A encephalopathy: Mechanisms and models** *Epilepsia* **60**:S86–S91
- Meisler M. H., Hill S. F., Yu W. (2021) **Sodium channelopathies in neurodevelopmental disorders** *Nat Rev Neurosci* **22**:152–166
- Miao P., Tang S., Ye J., Wang J., Lou Y., Zhang B., Xu X., Chen X., Li Y., Feng J. (2020) **Electrophysiological features: The next precise step for SCN2A developmental epileptic encephalopathy** *Mol Genet Genomic Med* **8**

- Munoz-Sanjuan I., Smallwood P. M., Nathans J. (2000) **Isoform diversity among fibroblast growth factor homologous factors is generated by alternative promoter usage and differential splicing** *J Biol Chem* **275**:2589–2597
- Narayanan D. L., Majethia P., Shrikiran A., Siddiqui S., Dalal A., Shukla A. (2022) **Further evidence of affected females with a heterozygous variant in FGF13 causing X-linked developmental and epileptic encephalopathy 90** *Eur J Med Genet* **65**
- Nishimoto S., Nishida E. (2007) **Fibroblast growth factor 13 is essential for neural differentiation in *Xenopus* early embryonic development** *J Biol Chem* **282**:24255–24261
- Oda Y. *et al.* (2019) **Entire FGF12 duplication by complex chromosomal rearrangements associated with West syndrome** *J Hum Genet* **64**:1005–1014
- Ogiwara I., Nakayama T., Yamagata T., Ohtani H., Mazaki E., Tsuchiya S., Inoue Y., Yamakawa K. (2012) **A homozygous mutation of voltage-gated sodium channel β (I) gene SCN1B in a patient with Dravet syndrome** *Epilepsia* **53**:e200–203
- Olsen S. K., Garbi M., Zampieri N., Eliseenkova A. V., Ornitz D. M., Goldfarb M., Mohammadi M. (2003) **Fibroblast growth factor (FGF) homologous factors share structural but not functional homology with FGFs** *J Biol Chem* **278**:34226–34236
- Pablo J. L., Wang C., Presby M. M., Pitt G. S. (2016) **Polarized localization of voltage-gated Na⁺ channels is regulated by concerted FGF13 and FGF14 action** *Proc Natl Acad Sci U S A* **113**:E2665–2674
- Patino G. A. *et al.* (2009) **A functional null mutation of SCN1B in a patient with Dravet syndrome** *J Neurosci* **29**:10764–10778
- Paul A., Crow M., Raudales R., He M., Gillis J., Huang Z. J. (2017) **Transcriptional Architecture of Synaptic Communication Delineates GABAergic Neuron Identity** *Cell* **171**:522–539
- Porter R. S., Jaamour F., Iwase S. (2018) **Neuron-specific alternative splicing of transcriptional machineries: Implications for neurodevelopmental disorders** *Mol Cell Neurosci* **87**:35–45
- Puranam R. S., He X. P., Yao L., Le T., Jang W., Rehder C. W., Lewis D. V., McNamara J. O. (2015) **Disruption of Fgf13 causes synaptic excitatory-inhibitory imbalance and genetic epilepsy and febrile seizures plus** *J Neurosci* **35**:8866–8881
- Raj B., Blencowe B. J. (2015) **Alternative Splicing in the Mammalian Nervous System: Recent Insights into Mechanisms and Functional Roles** *Neuron* **87**:14–27
- Ramadan W. *et al.* (2017) **Confirming the recessive inheritance of SCN1B mutations in developmental epileptic encephalopathy** *Clin Genet* **92**:327–331
- Richerson G. B. (2023) **Divergent Causes and Convergent Mechanisms of SUDEP** *Ann Neurol* **94**:809–811
- Saleem N. M., Chencheri N., Thomas S., Alexander G., Madathil B. (2024) **Early-Onset Epileptic Encephalopathy Responsive to Phenytoin: A Diagnostic Clue for Fibroblast Growth Factor 12 Mutation** *Cureus* **16**

Scheffer I. E. *et al.* (2017) **ILAE classification of the epilepsies: Position paper of the ILAE Commission for Classification and Terminology** *Epilepsia* **58**:512–521

Scheffer I. E. *et al.* (2016) **Classification of the epilepsies: New concepts for discussion and debate-Special report of the ILAE Classification Task Force of the Commission for Classification and Terminology** *Epilepsia Open* **1**:37–44

Scheffer I. E., Liao J. (2020) **“Deciphering the concepts behind “Epileptic encephalopathy” and “Developmental and epileptic encephalopathy”.”** *Eur J Paediatr Neurol* **24**:11–14

Scheffer I. E., Nabbout R. (2019) **SCN1A-related phenotypes: Epilepsy and beyond** *Epilepsia* **60**:S17–S24

Schoorlemmer J., Goldfarb M. (2001) **Fibroblast growth factor homologous factors are intracellular signaling proteins** *Curr Biol* **11**:793–797

Shen K. F. *et al.* (2022) **Fibroblast growth factor 13 is involved in the pathogenesis of temporal lobe epilepsy** *Cereb Cortex*

Siekierska A., Isrie M., Liu Y., Scheldeman C., Vanthillo N., Lagae L., de Witte P. A., Van Esch H., Goldfarb M., Buyse G. M. (2016) **Gain-of-function FHF1 mutation causes early-onset epileptic encephalopathy with cerebellar atrophy** *Neurology* **86**:2162–2170

Skotte L. *et al.* (2022) **Genome-wide association study of febrile seizures implicates fever response and neuronal excitability genes** *Brain* **145**:555–568

Smallwood P. M., Munoz-Sanjuan I., Tong P., Macke J. P., Hendry S. H., Gilbert D. J., Copeland N. G., Jenkins N. A., Nathans J. (1996) **Fibroblast growth factor (FGF) homologous factors: new members of the FGF family implicated in nervous system development** *Proc Natl Acad Sci U S A* **93**:9850–9857

Szabadics J., Varga C., Molnár G., Oláh S., Barzó P., Tamás G. (2006) **Excitatory effect of GABAergic axo-axonic cells in cortical microcircuits** *Science* **311**:233–235

Talwar D., Hammer M. F. (2021) **SCN8A Epilepsy, Developmental Encephalopathy, and Related Disorders** *Pediatr Neurol* **122**:76–83

Taniguchi H. *et al.* (2011) **A resource of Cre driver lines for genetic targeting of GABAergic neurons in cerebral cortex** *Neuron* **71**:995–1013

Teran F. A., Sainju R. K., Bravo E., Wagnon J., Kim Y., Granner A., Gehlbach B. K., Richerson G. B. (2023) **Seizures Cause Prolonged Impairment of Ventilation, CO₂** *J Neurosci* **43**:4959–4971

Vacher H., Mohapatra D. P., Trimmer J. S. (2008) **Localization and targeting of voltage-dependent ion channels in mammalian central neurons** *Physiol Rev* **88**:1407–1447

Velíšková J., Marra C., Liu Y., Shekhar A., Park D. S., Iatckova V., Xie Y., Fishman G. I., Velíšek L., Goldfarb M. (2021) **Early onset epilepsy and sudden unexpected death in epilepsy with cardiac arrhythmia in mice carrying the early infantile epileptic encephalopathy 47 gain-of-function FHF1(FGF12) missense mutation** *Epilepsia* **62**:1546–1558

Venkatesan K., Liu Y., Goldfarb M. (2014) **Fast-onset long-term open-state block of sodium channels by A-type FGFs mediates classical spike accommodation in hippocampal pyramidal neurons** *J Neurosci* **34**:16126–16139

- Vormstein-Schneider D. *et al.* (2020) **Viral manipulation of functionally distinct interneurons in mice, non-human primates and humans** *Nat Neurosci* **23**:1629–1636
- Wang C., Chung B. C., Yan H., Lee S. Y., Pitt G. S. (2012) **Crystal structure of the ternary complex of a NaV C-terminal domain, a fibroblast growth factor homologous factor, and calmodulin** *Structure* **20**:1167–1176
- Wang C., Hennessey J. A., Kirkton R. D., Wang C., Graham V., Puranam R. S., Rosenberg P. B., Bursac N., Pitt G. S. (2011) **Fibroblast growth factor homologous factor 13 regulates Na⁺ channels and conduction velocity in murine hearts** *Circ Res* **109**:775–782
- Wang Q. *et al.* (2021) **Fibroblast growth factor 13 stabilizes microtubules to promote Na⁺ channel function in nociceptive DRG neurons and modulates inflammatory pain** *J Adv Res* **31**:97–111
- Wang X. *et al.* (2017) **Conditional knockout of Fgf13 in murine hearts increases arrhythmia susceptibility and reveals novel ion channel modulatory roles** *J Mol Cell Cardiol* **104**:63–74
- Ware T. L. *et al.* (2019) **Epidemiology and etiology of infantile developmental and epileptic encephalopathies in Tasmania** *Epilepsia Open* **4**:504–510
- Wu Q. F., Yang L., Li S., Wang Q., Yuan X. B., Gao X., Bao L., Zhang X. (2012) **Fibroblast growth factor 13 is a microtubule-stabilizing protein regulating neuronal polarization and migration** *Cell* **149**:1549–1564
- Xu Q., Tam M., Anderson S. A. (2008) **Fate mapping Nkx2.1-lineage cells in the mouse telencephalon** *J Comp Neurol* **506**:16–29
- Yu F. H., Mantegazza M., Westenbroek R. E., Robbins C. A., Kalume F., Burton K. A., Spain W. J., McKnight G. S., Scheuer T., Catterall W. A. (2006) **Reduced sodium current in GABAergic interneurons in a mouse model of severe myoclonic epilepsy in infancy** *Nat Neurosci* **9**:1142–1149
- Zhang X. *et al.* (2016) **Cell-Type-Specific Alternative Splicing Governs Cell Fate in the Developing Cerebral Cortex** *Cell* **166**:1147–1162

Author information

Susan Lin^{*}

Cardiovascular Research Institute, Weill Cornell Medicine, New York, United States
ORCID iD: [0000-0001-9344-6594](https://orcid.org/0000-0001-9344-6594)

^{*}These authors contributed equally

Aravind R Gade^{*}

Cardiovascular Research Institute, Weill Cornell Medicine, New York, United States

^{*}These authors contributed equally

Hong-Gang Wang

Cardiovascular Research Institute, Weill Cornell Medicine, New York, United States

James E Niemeyer

Department of Neurological Surgery and Brain and Mind Research Institute, Weill Cornell Medicine of Cornell University, New York Presbyterian Hospital, New York, United States

Allison Galante

Cardiovascular Research Institute, Weill Cornell Medicine, New York, United States

Isabella DiStefano

Cardiovascular Research Institute, Weill Cornell Medicine, New York, United States

Patrick Towers

Cardiovascular Research Institute, Weill Cornell Medicine, New York, United States

Jorge Nunez

Cardiovascular Research Institute, Weill Cornell Medicine, New York, United States

Maiko Matsui

Cardiovascular Research Institute, Weill Cornell Medicine, New York, United States

Theodore H Schwartz

Department of Neurological Surgery and Brain and Mind Research Institute, Weill Cornell Medicine of Cornell University, New York Presbyterian Hospital, New York, United States

Anjali M Rajadhyaksha

Department of Pediatrics, Division of Pediatric Neurology, Weill Cornell Medicine, New York, United States, Brain and Mind Research Institute, New York, United States

Geoffrey S Pitt

Cardiovascular Research Institute, Weill Cornell Medicine, New York, United States
ORCID iD: [0000-0003-2246-0289](https://orcid.org/0000-0003-2246-0289)

For correspondence: geoffrey.pitt@med.cornell.edu

Editors

Reviewing Editor

John Huguenard

Stanford University School of Medicine, Stanford, United States of America

Senior Editor

John Huguenard

Stanford University School of Medicine, Stanford, United States of America

Reviewer #2 (Public review):

Summary

The authors address three primary questions:

(1) how FGF13 variants confer seizure susceptibility,

- (2) the specific cell types involved, and
- (3) the underlying mechanisms, particularly regarding Nav dysfunction.

They use different Cre drivers to generate cell type-specific knockouts (KOs). First, using Nestin-Cre to create a whole-brain Fgf13 KO, they observed spontaneous seizures and premature death. While KO of Fgf13 in excitatory neurons does not lead to spontaneous seizures, KO in inhibitory neurons recapitulates the seizures and premature death observed in the Nestin-Cre KO. They further narrow down the critical cell type to MGE-derived interneurons (INs), demonstrating that MGE-neuron-specific KO partially reproduces the observed phenotypes. "All interneuron" KOs exhibit deficits in synaptic transmission and interneuron excitability, not seen in excitatory neuron-specific KOs. Finally, they rescue the defects in the interneuron-specific KO by expressing specific Fgf13 isoforms. This is an elegant and important study adding to our knowledge of mechanisms that contribute to seizures.

Strengths

- The study provides much-needed cell type-specific KO models.
- The authors use appropriate Cre lines and characterize the phenotypes of the different KOs.
- The metabolomic analysis complements the rest of the data effectively.
- The study confirms and extends previous research using improved approaches (KO lines vs. in vitro KD or antibody infusion).
- The methods and analyses are robust and well-executed.

Weaknesses

- One weakness lies in the use of the Nkx2.1 line (instead of Nkx2.1CreER) in the paper. As a result, some answers to key questions are incomplete. For instance, it remains unclear whether the observed effects are due to Chandelier cells or NGFCs, potentially both MGE and CGE derived, explaining why Nkx2.1 alone does not fully replicate the overall inhibitory KO. Using Nkx2.1CreER could have helped address the cell specificity. With the Nkx2.1 line used in the paper, the answer is partial.
- While the mechanism behind the reduced inhibitory drive in the IN-specific KO is suggested to be presynaptic, the chosen method does not allow them to exactly identify the mechanisms (spontaneous vs mEPSC/mIPSC), and whether it is a loss of inhibitory synapses (potentially axo-axonic) or release probability.

General Assessment

The general conclusions of this paper are supported by data. As it is, the claim that "these results enhance our understanding of the molecular mechanisms that drive the pathogenesis of Fgf13-related seizures" is partially supported. A more cautious term may be more appropriate, as the study shows the mechanism is not Nav-mediated and suggests alternative mechanisms without unambiguously identifying them. The conclusion that the findings "expand our understanding of FGF13 functions in different neuron subsets" is supported, although somewhat overstated, as the work is not conclusive about the exact neuron subtypes. However, it does indeed show differential functions for specific neuronal classes, which is a significant result.

Impact and Utility

This paper is undoubtedly valuable. Understanding that excitatory neurons are not the primary contributors to the observed phenotypes is crucial. The finding that the effects are not MGE-unique is also important. This work provides a solid foundation for further research and will be a useful resource for future studies.

<https://doi.org/10.7554/eLife.98661.2.sa2>

Reviewer #3 (Public review):**Summary:**

The authors aimed to determine the mechanism by which seizures emerge in Developmental and Epileptic Encephalopathies caused by variants in the gene FGF13. Loss of FGF13 in excitatory neurons had no effect on seizure phenotype as compared to loss of FGF13 in GABAergic interneurons, which in contrast caused a dramatic pro-seizure phenotype and early death in these animals. They were able to show that Fgf13 ablation and consequent loss of FGF13-S and FGF13-VY reduced overall inhibitory input from Fgf13-expressing interneurons onto hippocampal pyramidal neurons. This was shown to occur not via disruption to voltage-gated sodium channels but rather by reducing potassium currents and action potential repolarisation in these interneurons.

Strengths:

The authors employed multiple well validated, novel mouse lines with FGF13 knocked out in specific cell types including all neurons, all excitatory cells, all GABAergic interneurons, or a subset of MGE-derived interneurons, including axo-axonic chandelier cells. The phenotypes of each of these four mouse lines were carefully characterised to reveal clear differences with the most fundamental being that Interneuron-targeted deletion of FGF13 led to perinatal mortality associated with extensive seizures and impaired the hippocampal inhibitory/excitatory balance while deletion of FGF13 in excitatory neurons caused no detectable seizures and no survival deficits.

The authors made excellent use of western blotting and in situ hybridisation of the different FGF13 isoforms to determine which isoforms are expressed in which cell types, with FGF3-S predominantly in excitatory neurons and FGF13-VY and FGF13-V predominantly in GABAergic neurons.

The authors performed highly detailed electrophysiological analysis of excitatory neurons and GABAergic interneurons with FGF13 deficits using whole-cell patch clamp. This enabled them to show that FGF13 removal did not affect voltage-gated sodium channels in interneurons, but rather reduced the action of potassium channels, with the resultant effect of making it more likely that interneurons enter depolarisation block. These findings were strengthened by the demonstration that viral re-expression of different Fgf13 splice isoforms could partially rescue deficits in interneuron action potential output and restore K⁺ channel current size.

Additionally, the discussion was nuanced, and demonstrated how the current findings resolved previous apparent contradictions in the field involving the function of FGF13.

These findings will have a significant impact on our understanding of how FGF13 causes seizures and death in DEEs, and the action of different FGF13 isoforms within different neuronal cell types, particularly GABAergic interneurons.

Comments on revisions:

I appreciate the author's responses to the previous round of reviews. All my comments have been addressed. Congratulations on an excellent body of work.

<https://doi.org/10.7554/eLife.98661.2.sa1>

Author response:

The following is the authors' response to the original reviews.

Public Reviews:

Reviewer #1 (Public Review):

Summary:

A subset of fibroblast growth factor (FGF) proteins (FGF11-FGF14; often referred to as fibroblast growth factor homologous factors because they are not thought to be secreted and do not seem to act as growth factors) have been implicated in modulating neuronal excitability, however, the exact mechanisms are unclear. In part, this is because it is unclear how different FGF isoforms alter ion channel activity in different neuronal populations. In this study, the authors explore the role of FGF 13 in epilepsy using a variety of FGF13 knock-out mouse models, including several targeted cell-type specific conditional knockout mouse lines. The study is intriguing as it indicates that FGF13 plays an especially important role in inhibitory neurons. Furthermore, although FGF13 has been studied as a regulator of neuronal voltage-gated sodium channels, the authors present data indicating that FGF13 knockout in inhibitory neurons induces seizures not by altering sodium current properties but by reducing voltage-gated potassium currents in inhibitory neurons. While intriguing, the data are incomplete in several aspects and thus the mechanisms by which various FGF13 variants induce Developmental and Epileptic Encephalopathies are not resolved by the data presented.

Strengths:

A major strength is the array of techniques used to assess the mice and the electrical activity of the neurons.

The multiple mouse knock-out models utilized are a strength, clearly demonstrating that FGF13 expression in inhibitory neurons, and possibly specific sub-populations of inhibitory neurons, is critically important.

The data on the increased sensitivity to febrile seizures in KO mice are very nice, provide clear evidence for regulation of excitability in inhibitory neurons by FGF13.

The Gad2Fgf13-KO mice indicated that several Fgf13 splice variants may be expressed in inhibitory neurons and suggest that the Fgf13-VY splice variants may have previously unrecognized specific roles in regulating neuronal excitability.

The data on males and females from the various KO mice lines indicates a clear gene dosage effect for this X-linked gene.

The unbiased metabolomic analysis supports the assertion that Fgf13 expression in inhibitory neurons is important in regulating seizure susceptibility.

Weaknesses:

The knockout approach can be powerful but also has distinct limitations. Multiple missense mutations in FGF13-S have been identified. The knockout models employed here are not appropriate for understanding how these missense variants lead to altered neuronal excitability. While the data show that complete loss of Fgf13 from excitatory forebrain neurons is not sufficient to induce seizure susceptibility, it does not rule out that specific variants (e.g., R11C) might alter the excitability of forebrain neurons. The missense variants may alter excitatory and/or inhibitory neuron excitability in distinct ways from a full FGF13 knockout.

We agree with this overall interpretation of our data and have updated our language in the Discussion to make the distinction between mechanisms attributable to a knockout compared

to a missense variant. We note, however, that the proposed mechanism by which missense variants (e.g., R11C) drive seizures is through loss of long-term inactivation in excitatory neurons and our excitatory knockout model shows loss of long-term inactivation in excitatory neurons. Thus, our knockout model demonstrates that the mechanism(s) by which the missense variants alter neuronal excitability in excitatory neurons must exclude long-term inactivation, thereby providing some clarity regarding the proposed mechanism for those missense variants.

The electrophysiological experiments are intriguing but not comprehensive enough to support all of the conclusions regarding how FGF13 modulates neuronal excitability.

We agree and have updated the language in our Discussion to clarify speculation from conclusions that are directly supported by data.

Another concern is the use of different ages of neurons for different experiments. For example, sodium currents in Figures 2 and 5 (and Supplemental Figures 2 and 7) are recorded from cultured neurons, which may have very different properties (including changes in sodium channel complexes) from neurons in vivo that drive the development of seizure activity.

We agree and acknowledge the important differences between neurons examined in culture and in vivo, yet the in vitro vs in vivo preparations were necessitated by the specific experiments. While these differences are important, previous gene profiling studies comparing primary hippocampal neurons with developing mouse hippocampus have found that although gene expression is accelerated in vitro, gene expression profiles in vitro and in vivo are similar (PMID: 11438693). Moreover, the relative immaturity of the cultured neurons is balanced at least in part because the in vivo experiments were performed on very young animals (~P12), which also have relatively immature neurons. Thus, we predict that sodium channel complexes studied in vitro are informative for the in vivo aspects of this investigation.

Reviewer #2 (Public Review):

Summary:

The authors address three primary questions:

- (1) how FGF13 variants confer seizure susceptibility,*
- (2) the specific cell types involved, and*
- (3) the underlying mechanisms, particularly regarding Nav dysfunction.*

They use different Cre drivers to generate cell type-specific knockouts (KOs). First, using Nestin-Cre to create a whole-brain Fgf13 KO, they observed spontaneous seizures and premature death. While KO of Fgf13 in excitatory neurons does not lead to spontaneous seizures, KO in inhibitory neurons recapitulates the seizures and premature death observed in the Nestin-Cre KO. They further narrow down the critical cell type to MGE-derived interneurons (INs), demonstrating that MGE-neuron-specific KO partially reproduces the observed phenotypes. "All interneuron" KOs exhibit deficits in synaptic transmission and interneuron excitability, not seen in excitatory neuron-specific KOs. Finally, they rescue the defects in the interneuron-specific KO by expressing specific Fgf13 isoforms. This is an elegant and important study adding to our knowledge of mechanisms that contribute to seizures.

Strengths

- *The study provides much-needed cell type-specific KO models.*
- *The authors use appropriate Cre lines and characterize the phenotypes of the different KOs.*
- *The metabolomic analysis complements the rest of the data effectively.*
- *The study confirms and extends previous research using improved approaches (KO lines vs. in vitro KD or antibody infusion).*
- *The methods and analyses are robust and well-executed.*

Weaknesses

- *One weakness lies in the use of the Nkx2.1 line (instead of Nkx2.1CreER) in the paper. As a result, some answers to key questions are incomplete. For instance, it remains unclear whether the observed effects are due to Chandelier cells or NGFCs, potentially both MGE and CGE derived, explaining why Nkx2.1 alone does not fully replicate the overall inhibitory KO. Using Nkx2.1CreER could have helped address the cell specificity. With the Nkx2.1 line used in the paper, the answer is partial.*

We agree that while our data is consistent with the possibility of a role for Fgf13 in chandelier function, the current Cre driver does not provide sufficient direct evidence. We performed preliminary experiments (unpublished) using a Nkx2.1CreER driver, with late embryonic induction with a tamoxifen dosage validated for sparse labeling of chandelier cells (30846310). While we successfully replicated sparse labeling of neocortical chandelier cells (using a Cre-dependent Ai9 reporter), we were unable to determine if there was a significant loss of FGF13 as measured by immunohistochemistry since FGF13+ cells are only a small subset of the already sparse cells. Because multiple snRNA-seq studies identified Fgf13 as a marker for chandelier cells, we speculated—now more carefully circumspect—about the role of chandelier cells vs NGFCs.

- *While the mechanism behind the reduced inhibitory drive in the IN-specific KO is suggested to be presynaptic, the chosen method does not allow them to exactly identify the mechanisms (spontaneous vs mEPSC/mIPSC), and whether it is a loss of inhibitory synapses (potentially axo-axonic) or release probability.*

We agree that this is an important limitation of our work, and that we are unable to identify the exact mechanism behind the reduced inhibitory drive. We are continuing to explore this question in a follow-up study.

- *Some supporting data (e.g. Supplemental Figure 7 and 8) appear to come from only one (or two) WT and one (or two) KO mice. Supplementary data, like main data, should come from at least three mice in total to be considered complete/solid (even if the statistical analysis is done with cells).*

All panels in the manuscript, including supplementary data, except supplementary 7D and 8A, have N(mouse)≥3. Time limitations (graduating student) prevented us from obtaining a larger N. Because those supplementary data are not critical for supporting our conclusions, we removed them.

General Assessment

The general conclusions of this paper are supported by data. As it is, the claim that "these results enhance our understanding of the molecular mechanisms that drive the pathogenesis of Fgf13-related seizures" is partially supported. A more cautious term may

be more appropriate, as the study shows the mechanism is not Nav-mediated and suggests alternative mechanisms without unambiguously identifying them. The conclusion that the findings "expand our understanding of FGF13 functions in different neuron subsets" is supported, although somewhat overstated, as the work is not conclusive about the exact neuron subtypes. However, it does indeed show differential functions for specific neuronal classes, which is a significant result.

Impact and Utility

This paper is undoubtedly valuable. Understanding that excitatory neurons are not the primary contributors to the observed phenotypes is crucial. The finding that the effects are not MGE-unique is also important. This work provides a solid foundation for further research and will be a useful resource for future studies.

Reviewer #3 (Public Review):

Summary:

The authors aimed to determine the mechanism by which seizures emerge in Developmental and Epileptic Encephalopathies caused by variants in the gene FGF13. Loss of FGF13 in excitatory neurons had no effect on seizure phenotype as compared to the loss of FGF13 in GABAergic interneurons, which in contrast caused a dramatic pro-seizure phenotype and early death in these animals. They were able to show that Fgf13 ablation and consequent loss of FGF13-S and FGF13-VY reduced overall inhibitory input from Fgf13-expressing interneurons onto hippocampal pyramidal neurons. This was shown to occur not via disruption to voltage-gated sodium channels but rather by reducing potassium currents and action potential repolarisation in these interneurons.

Strengths:

The authors employed multiple well-validated, novel mouse lines with FGF13 knocked out in specific cell types including all neurons, all excitatory cells, all GABAergic interneurons, or a subset of MGE-derived interneurons, including axo-axonic chandelier cells. The phenotypes of each of these four mouse lines were carefully characterised to reveal clear differences with the most fundamental being that Interneuron-targeted deletion of FGF13 led to perinatal mortality associated with extensive seizures and impaired the hippocampal inhibitory/excitatory balance while deletion of FGF13 in excitatory neurons caused no detectable seizures and no survival deficits.

The authors made excellent use of western blotting and in situ hybridisation of the different FGF13 isoforms to determine which isoforms are expressed in which cell types, with FGF13-S predominantly in excitatory neurons and FGF13-VY and FGF13-V predominantly in GABAergic neurons.

The authors performed a highly detailed electrophysiological analysis of excitatory neurons and GABAergic interneurons with FGF13 deficits using whole-cell patch clamp. This enabled them to show that FGF13 removal did not affect voltage-gated sodium channels in interneurons, but rather reduced the action of potassium channels, with the resultant effect of making it more likely that interneurons enter depolarisation block. These findings were strengthened by the demonstration that viral re-expression of different Fgf13 splice isoforms could partially rescue deficits in interneuron action potential output and restore K⁺ channel current size.

Additionally, the discussion was nuanced and demonstrated how the current findings resolved previous apparent contradictions in the field involving the function of FGF13.

These findings will have a significant impact on our understanding of how FGF13 causes seizures and death in DEEs, and the action of different FGF13 isoforms within different neuronal cell types, particularly GABAergic interneurons.

Recommendations for the authors:

Reviewer #1 (Recommendations For The Authors):

The limitations of the KO model should be fully discussed in the discussion. It should be clear that knocking out FGF13 does not provide insight into how missense mutations such as R11C may alter excitatory and/or inhibitory neuron excitability.

We agree with this overall interpretation of our data and have updated our language in the Discussion to make the distinction between mechanisms attributable to a knockout compared to a missense variant. We note, however, that the proposed mechanism by which missense variants (e.g., R11C) drive seizures is through loss of long-term inactivation in excitatory neurons and our excitatory knockout model shows loss of long-term inactivation in excitatory neurons. Thus, our knockout model demonstrates that the mechanism(s) by which the missense variants alter neuronal excitability in excitatory neurons must exclude long-term inactivation, thereby providing some clarity regarding the proposed mechanism for those missense variants.

It is important to know what sodium channel isoforms are expressed in the cultured neurons used in the experiments for Figures 2 and 5. Are Nav1.1, Nav1.2, Nav1.3, and Nav1.6 expressed at appropriate levels in the cultures?

We agree it is important to know that the sodium channel isoforms expressed in our hippocampal neurons are expressed at physiologically relevant levels, for further validation of our primary culture system. We have added RT-qPCR data from our hippocampal neuron cultures (Supplemental Figure 2B) showing the relative levels of SCN1A, SCN2A, SCN3A, and SCN8A, which are similar to the relative levels of voltage-gated sodium channel isoforms found in rodent and human forebrain in early development (Figure 1 in PMID: 35031483).

The electrophysiological experiments are intriguing but limited. One, it would be helpful to report if there were any changes in resting membrane potential for the cells reported in Figure 5. It is also inappropriate to unequivocally state that "Nav currents were not significantly affected by Fgf13 knockout in Gad2Fghf13 KO neurons" as only a sampling of properties was investigated. Recovery from inactivation and persistent current amplitudes were not evaluated. Furthermore, while it looks like long-term inactivation is not altered, only one specific protocol was used and currents measured from cultured neurons may not be fully representative of neuronal properties in vivo.

We agree that we performed a selective analysis of Nav currents—selected because those are the major parameters that have been associated with FGF13 modulation. Because we did not observe significant differences in NaV currents, we therefore hypothesized that FGF13 affected other currents, as previously observed, and consequently assessed potassium currents, for which we did observe a difference. Further, we note that our sodium current and potassium current results are consistent with, and supportive of, our action potential data in which we find no deficit in AP initiation, but rather a deficit in AP repolarization. We revised the text to reflect the more limited analysis of Nav currents. Regarding long-term inactivation, we also agree that measurements in cultured neurons may not fully represent neuronal properties in vivo; however, we note that regulation of long-term inactivation by FGF13 has previously been assessed only in cultured cells (and not in neurons). Thus, our protocols were designed to query that modulation previously reported.

The first sentence of the results section is misleading: "To determine how FGF13 variants contribute to seizure disorders, we developed genetic mouse models that eliminate Fgf13 in specific neuronal cell types." The knockouts do not target specific splice isoforms and do not help determine how missense variants contribute to DEE. This should be modified to reflect better what is actually being tested.

We agree and have revised our text to state that our goal was to assess how FGF13 contributes to neuronal excitability and thereby accurately reflect the cell type-specific, but not isoform specific, targeting.

Reviewer #2 (Recommendations For The Authors):

• *The sentence in the introduction stating "an unusual example of differential expression of an alternatively spliced neuronal gene in excitatory vs. inhibitor neurons" is factually incorrect, especially for transcripts regulating intrinsic properties like FGF13. Refer to PMID: 31451803 for more details and consider rephrasing this statement.*

We updated our text to reflect the similarity of Fgf13's cell type-specific alternative splicing to other genes known to control synaptic interactions and neuronal architecture and added the suggested reference.

• *Consistency is needed in the manuscript regarding the term "BASEscope" or "basescope"; the correct version is "BaseScope."*

We corrected the text accordingly.

• *In the discussion, the term "reduced overall inhibitory drive" might be more appropriate than "input."*

We updated the text accordingly.

• *The authors should refer to the Fgf13 data in the database from Furlanis et al., which complements their findings: <https://scheiffele-splice.scicore.unibas.ch/>.*

We agree and now incorporate this reference.

• *The phrase "Fgf13 silencing in Nkx2.1 expressing neurons" should be clarified to include the use of CreER, which was crucial and effectively resulted in the labeling of a different subtype of interneurons, see PMID: 23180771.*

We agree and have updated our text accordingly.

• *Be more cautious when discussing the role of FGF13 in chandelier function; while it seems probable, the current Cre driver used provides no direct evidence.*

We agree (as noted above) that while our data are consistent with the possibility of a role for Fgf13 in chandelier function, the current Cre driver used is insufficient to offer direct evidence and therefore updated our text in the discussion.

• *The gene dosage effect is interesting, it would be interesting to explore it further in the future.*

We agree. Because our data suggest that seizures result from loss of inhibitory neuron input, we hypothesize that the gene dosage effect derives from further loss of inhibitory neuron input and thus more hyperexcitability.

• Another critical aspect not addressed here and of interest for the future is the distinction between the role of FGF13 in interneuron development versus general maintenance. Using Nkx2.1CreER could have helped address both cell specificity and developmental roles.

We agree that there may be an interesting distinction between the role of Fgf13 in development versus general maintenance. We have piloted an Nkx2.1-CreER targeted deletion of Fgf13 from cortical interneurons but have been unsuccessful with significant deletion of Fgf13, likely because the Nkx2.1-CreER strategy targets only a sparse subset of interneurons and FGF13 is expressed in only a subset of total interneurons. Thus, use of the Nkx2.1-CreER strategy is challenging. We are looking for ways to optimize.

Reviewer #3 (Recommendations For The Authors):

This was a truly fabulous paper, with an exceptional quantity of beautiful data. I would like to congratulate the authors on their superb work.

In the discussion, the authors correctly draw attention to the fact that the clear pro-seizure phenotype they see when FGF13 was knocked out more specifically in a subset of interneurons including chandelier cells, adds to our understanding of the role of FGF13 in chandelier cells. More than that though, given that FGF13 is reducing excitability in these cells AND this results in a strong pro-seizure phenotype, they may want to postulate that this lends further weight to the argument that chandelier cells are likely powerful regulators of network excitability despite suggestions in the field that they could potentially have a proexcitatory function (see Szabadics et al. Science 2006).

We agree this is interesting and have elaborated on our discussion of chandelier cells to include this point while also addressing the important caveats noted by reviewer 2.

A minor point:

On page 26 the sentence:

"Here, we were able to assess FGF13-S and FGF13-VY, chosen because they are most abundantly expressed isoforms in the adult mouse brain, but the inability to rescue electrophysiological consequences completely with either isoform alone leaves open the possibility that other isoforms (e.g., FGF13-U, FGF13-V, and FGF13-VY) also make critical contributions." Should the last "FGF13-VY" be removed?

We thank the reviewer for noticing the error and have updated the text accordingly.

<https://doi.org/10.7554/eLife.98661.2.sa0>