

Encoding of cerebellar dentate neuronal activity during visual attention in rhesus macaques

Reviewed Preprint

v1 • September 4, 2024

Not revised

Nico A Flierman, Sue Ann Koay, Willem S van Hoogstraten, Tom JH Ruigrok, Pieter R Roelfsema, Aleksandra Badura , Chris I De Zeeuw 

Netherlands Institute for Neuroscience, Amsterdam, The Netherlands • Department of Neuroscience, Erasmus MC, Rotterdam, The Netherlands • Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, USA • Department of Integrative Neurophysiology, VU University, Amsterdam, The Netherlands • Department of Psychiatry, Academic Medical Centre, Amsterdam, The Netherlands

 https://en.wikipedia.org/wiki/Open_access

 Copyright information

Abstract

The role of cerebellum in controlling eye movements is well established, but its contribution to more complex forms of visual behavior has remained elusive. To study cerebellar activity during visual attention we recorded extracellular activity of dentate nucleus (DN) neurons in two non-human primates (NHPs). NHPs were trained to read the direction indicated by a peripheral visual stimulus while maintaining fixation at the center, and report the direction of the cue by performing a saccadic eye movement into the same direction following a delay. We found that single unit DN neurons modulated spiking activity over the entire time-course of the task, and that their activity often bridged temporally separated intra-trial events, yet in a heterogeneous manner. To better understand the heterogeneous relationship between task structure, behavioral performance and neural dynamics, we constructed a behavioral, an encoding and a decoding model. Both NHPs showed different behavioral strategies, which influenced the performance. Activity of the DN neurons reflected the unique strategies, with the direction of the visual stimulus frequently being encoded long before an upcoming saccade. Retrograde labeling of the recording location indicated that these neurons receive predominantly inputs from Purkinje cells in the lateral cerebellum as well as neurons of the principal olive and medial pons, all regions known to connect with neurons in the prefrontal cortex contributing to planning of saccades. Together, our results highlight that DN neurons can dynamically modulate their activity during a visual attention task, comprising not only sensorimotor but also cognitive attentional components.

eLife assessment

This **important** study examined neuronal activity in the dentate nucleus of the cerebellum when monkeys performed a difficult perceptual decision-making task. The authors provide **convincing** evidence that the cerebellum represents sensory, motor, and behavioral outcome signals that are sent to the attentional system, but further analysis focusing on the disparity of performance between animals would improve the quality of the paper. This paper is of great general interest in that it shows the involvement of the cerebellum in cognitive processes at the neuronal level.

<https://doi.org/10.7554/eLife.99696.1.sa3>

Introduction

The crucial role of the cerebellum in sensorimotor processing, such as learning and execution of eye movements, is well established (Inoshita and Hirano, 2018 [↗](#); Ivry and Diener, 1991 [↗](#); Kunimatsu et al., 2016 [↗](#)). Yet, over the last few decades evidence is emerging that the cerebellum also participates in more complex cognitive visual functions (Baier et al., 2010 [↗](#); Brissenden et al., 2016 [↗](#); Courchesne et al., 1994 [↗](#); Nicolson et al., 2001 [↗](#); Voogd et al., 2012 [↗](#)). For example, while interacting with areas such as the frontal eye field (FEF), lateral intraparietal area, superior colliculus and basal ganglia, the cerebellum uses visual information to guide the planning of saccades (Kunimatsu et al., 2018 [↗](#); Scerra et al., 2019 [↗](#); Tanaka, 2006 [↗](#); Wardak et al., 2011 [↗](#)). An open question is to what extent the cerebellum also has access to information in the peripheral visual field to adjust the planning of saccades. If the cerebellum is prominently involved in such processing, one expects to find encodings of both visual cues and the related eye movements at its output stage in the cerebellar nuclei.

In daily life visual scenes often contain considerably more information than the visual system can process in short periods of time. Therefore, saccades are made to align the fovea, the region of the retina having the highest visual acuity and largest representation in the visual cortex, with the target of interest. Yet, how saccades are planned remains an open question. Neuroimaging studies suggest that network mechanisms in the parietal, frontal and temporal lobes that control the programming of saccades overlap with those underlying covert attention (Corbetta et al., 1998 [↗](#); Nobre et al., 2000 [↗](#)). Along the same avenue, the “premotor theory of attention” by Rizzolatti and colleagues (1987 [↗](#)) proposes that the circuit used for generating movements is also active during attention shifts. This theory is also supported by the observation that the working peripheral location of the covert cue stimulus is restricted by the ultimate potential outer position of the eye in that it is not possible to attend to a covert cue when the cue is outside of the range of the oculomotor system (Craighero et al., 2004 [↗](#); Hanning and Deubel, 2020 [↗](#)). At the same time, it should be noted that when it comes to neuronal responses at the single cell level, the cortical networks engaged in covert attention do not always perfectly overlap with those that control saccades. For example, findings in the frontal eye fields (FEF) show that at least part of the visual cells that respond to covert attention do not modulate in relation to the saccades, implying that there may well be cell type - specificity for responses to covert attention in this region (Gregoriou et al., 2012 [↗](#)).

In this work we aimed to examine if visual attention stimuli are processed in the dentate nucleus (DN) of the cerebellum, and to what extent the signaling during a visuomotor task controls the directional saccadic eye movements. Therefore we studied the activity of single units in the DN during a complex, peripheral Landolt C saccade task in two non-human primates (NHPs). The task

comprised the following consecutive components: 1) reading of C-direction-sensitive information in the peripheral visual field while holding gaze fixation on the center, 2) planning of a saccade into a direction that depends on that peripheral visual information, 3) executing the corresponding saccade following a delay, and 4) receiving a reward if the direction of the saccade is correct. We found that DN neurons could modulate their activity during multiple task epochs, often selectively for an individual C-gap direction or saccade direction. Further, animals employed different behavioral strategies during execution of the task. Finally, the location of the task-relevant neurons in the DN was in line with the known connections from and to the FEF, i.e., one of the main classical attention and saccade planning center(s) in the cerebral cortex (Gregoriou et al., 2012 [↗](#)).

Materials and Methods

Animals

All procedures complied with the NIH Guide for the Care and Use of Laboratory Animals (National Institutes of Health, Bethesda, Maryland), and were approved by the institutional animal care and use committee of the Royal Netherlands Academy of Arts and Sciences (AVD8010020184587). Two adult male non-human primates (NHPs, *Macaca mulatta*), referred to as Mi and Mo, were used in this study. Before being subjected to the peripheral Landolt C saccade task of the current study, both of them were used for two other studies, one on glissade control and another one on mechanisms underlying anti-saccades (Avila et al., 2022 [↗](#); Flierman et al., 2019 [↗](#)).

Experimental procedures

Surgery

Animals were prepared for eye movement recordings as well as extracellular single unit recordings in the cerebellum using a two-step surgical procedure (Chen et al., 2017 [↗](#)). First, under general anesthesia induced with ketamine (15 mg/kg, i.m.) and maintained under intubation by ventilating a mixture of 70% N₂O and 30% O₂, supplemented with 0.8% isoflurane, fentanyl (0.005 mg/kg, i.v.), and midazolam (0.5 mg/kg • h, i.v.), we implanted a titanium head holder, which allowed for immobilization of the NHP's head. Four months later, under the same anesthesia conditions, a custom-made 40 mm-wide chamber was implanted to gain access to the cerebellum. Animals recovered for at least 21 days before behavioral training and testing (for more details on surgical procedures, see (Avila et al., 2022 [↗](#); Flierman et al., 2019 [↗](#)).

Behavioral experiments

While they were head-restrained, Mi and Mo were trained to focus on a fixation dot at the center of the monitor, which was placed at a viewing point distance of 52 cm and operating at a frame rate of 100 Hz (1152 x 864 pixels). Eye movements of their left eye were recorded with an infrared video eye tracker scanning at a 1000 Hz (Eyelink 1000 plus, SR Research), (Figure 1 [↗](#)). Their eye movements were calibrated before every experiment (5 Standard Eyelink points for 10° eccentricity). We used the Landolt C task as described in (Ignashchenkova et al., 2004 [↗](#)) (Figure 1A, B [↗](#)) (Ignashchenkova et al., 2004 [↗](#)). At trial onset the animals were given a window of 500 ms to make a saccade to the central fixation point (red dot, 0.2° diameter) and required fixation to this point within a 4° diameter circular window for a period of at least 100 ms. If fixation was broken a new trial was started after a 1.5 s time-out. If not, the red C-stimulus was presented in 1 out of 4 positions (Figure 1C [↗](#)), randomly selected every trial. The gap direction of the C was always perpendicular to the position of the C relative to the fixation dot. The C-stimulus was presented for 250 ms at 5° eccentricity and had an outer and inner diameter of 1° and 0.6°, respectively. During

the presentation of the C-stimulus the animals had to maintain fixation on the fixation dot. After the 250 ms period, the C-stimulus was masked for 100 ms with a full red circle to prevent the animals from gaining extra information from the retinal after-images. After the offset of the mask presentation, the fixation dot was visible for another randomly drawn interval between 500-700 ms (50 ms steps) during which the animals were also not allowed to break fixation. After this delay period, two potential saccade targets were presented perpendicular from the target location; subsequently, the fixation dot turned gray (go-cue), indicating that the animal was allowed to make a saccade into the direction indicated by the initial cue, i.e., the gap of the C-stimulus. A trial was rewarded with a juice reward when the animals made a saccade to the correct target indicated by the gap of the C-stimulus within 500 ms after the central target turning gray. The correct target had to be fixated within a 4° diameter circular window around the target and fixation had to last at least 100 ms. Incorrect trials, in which the animal made a saccade to the wrong target, or if any of the before mentioned requirements were not met, triggered a 1.5 s time-out and the reward duration was reset to a minimum value of 100 ms (see next section for details on juice reward).

Motivation and controlled water intake

Animals had restricted water intake on the day before the experiment and they were water deprived on the day of the experiment. Controlled water intake was performed in consultation with the animal caretakers and in accordance with the rules of the Dutch law. During the task the animals were allowed to drink as much as they liked, yet always in the context of a reward for correctly executed trials. Rewards were in the form of strawberry or tropical lemonade, depending on the animal's preference. The duration and thereby amount of reward delivery started with a pulse of 100 ms. To motivate the animal to perform well, 50 ms was added to the reward pulse for every consecutively correct trial, up to a maximum 350 ms. If the animal made an incorrect response, no reward was given for that trial and the duration of the reward was reset to the initial 100 ms. After the experiment the animals were given water *ad libitum*.

Electrophysiological recordings

MRI-images were used as a reference frame for anatomical localization of the left DN (**Figure 2A**). Single-unit recordings were obtained using tungsten glass-coated electrodes (1.5 MΩ, Alpha Omega Engineering, Nazareth, Israel) through a 23-gauge guide tube, which was inserted only through the dura. A motorized microdriver (Alpha Omega Engineering, Nazareth, Israel) with a 1-mm spaced grid was used to introduce the electrode and map the recording sites. Extracellular recordings were digitized and sampled at 44 kHz and subsequently stored during the experiment using a Multi-Channel Processor (Alpha Omega Engineering, Nazareth, Israel). Single units were determined to be DN neurons by the x, y and z coordinates of the electrode tip in relation to those of the MRI images, as well as their waveform in combination with the absence of complex spikes (CS). The discovery of a task-related site was often guided by bursting or pausing of single unit action potentials in relation to task related variables, such as the C-stimulus, saccade and/or reward. The electrophysiological data, as detected with an online spike sorter (Multi-Spike Detector, Alpha Omega Engineering), were stored for offline analysis using custom-written MATLAB code.

Analyses of behavioral and electrophysiological data

Eye movement analysis

Saccade onset and offset were detected on the basis of an adaptive velocity threshold, which consisted of 3 standard deviations (std) of the noise during fixation (see also [Flierman et al., 2019](#)). A Savitzky-Golay filter was applied for smoothing the raw traces. Position traces were differentiated to find the eye velocity and acceleration signals.




Figure 1.

Task and behavioral performance.

A) Left, drawing of the animal performing the task. Right, schematic overview of directionality in the task, only one C-stimulus is presented per trial in one of 4 cardinal directions from the central fixation dot. This could be to the top/bottom of the central fixation, with possible gap directions to the left and right [L/R] task, or to the left/right of the central fixation, with possible gap directions to the top/bottom [U/D] task. **B)** Example trial with a C-stimulus presented to the right, with an "Up" gap direction, where the animal has to make an upward saccade. Delay period between the mask and the onset of the target varied between 500-700 ms (randomly drawn from: 500, 550, 600, 650 and 700 ms). The saccade had to be performed in the epoch between the appearance of the choice target and the reward. Beige arrow indicates the correct choice, not actually visible to the monkey. **C)** Overview of performance per animal per gap direction. Black bar at the middle of the violin plot represents the 25th and 75th percentile, the white dot at the center is the median. **D)** Average fraction of correct responses per retraining session, after the animal had had several weeks of 'holiday'. Each session represents the median performance of one day. **E)** Weight of the C-stimulus, i.e., the gap direction, in predicting the correct response through logistic regression. Each point is the average of one session, error bars indicate 95% confidence interval from 100 bootstrap permutations. Beige data points and fit represent the up/down task, the green data the left/right task.




Figure 2.

Neural activity in the DN was strongly modulated during the Landolt C task.

A) MRI images of Mi and Mo, showing the electrode position in the DN. **B**) Example trial of extracellularly recorded DN cell in Mi aligned to C-stimulus onset (blue shaded area shows when C is visible), while the animal maintained fixation until the potential targets were presented (in this case an upward saccade was correct). Top panel depicts eye position, bottom panel shows the single unit activity of the DN cell. Pink shaded area represents juice reward pulse duration. **C**) Raster plot of correct trials sorted per gap direction, represented per color, of the cell presented in B. Within each direction trials are sorted to saccade onset (black dot in each trial). **D**) Average activity per direction in 100 ms bins in spikes per second. Left panel displays activity aligned to C-stimulus onset, right panel displays activity aligned to saccade onset. **E**) Preferred directions of the population of neurons. Normalized polar histograms show the fraction cells that preferred the stimulus direction (left) or the action direction (right). **F**) same as E for direction preference during saccade epoch. **G**) Normalized activity of cells displaying facilitating activity deviating from baseline in either the window after C-stimulus onset (gray bar, 0 to 800 ms, left panel) or around the saccade (gray bar, -200 to 200 ms, right panel). **H**) Same as in G, but for suppressing cells. **I**) Timing of peak / trough responses from G and H, stars indicate significance at * $p=0.05$, ** $p=0.01$, *** $p=0.001$ level, according to Kolmogorov-Smirnov test; ns = not significant.

Neuronal modulation analysis

Task-related neuronal modulation was determined by the application of 200 ms sliding windows around the main events of interest (i.e., C-stimulus onset, saccade onset), ([Table 1](#)). Spike counts in the windows were compared with baseline activity (taken during -500 to 0 ms from fixation). If average activity over different trials in one of the sliding windows was significantly different from the average activity in the baseline window, the activity of that neuron was considered to be significantly modulated for that event of interest. In case of the stimulus event, if more than half of the windows were significantly different from the baseline window, the cell was considered to have sustained activity during the epoch surrounding the stimulus event.

Parametric statistics

If statistical testing involved two groups, and the data were normally distributed, a student's t-test was used. When more than two groups were involved, we used ANOVA tests with the Tukey-Kramer post-hoc test. To determine directional selectivity of neurons we used one-way ANOVA unless specified otherwise. When comparing fractions, the χ^2 for proportions was applied to determine significance. For circular statistics the circle statistics toolbox in MATLAB was used.

Directional selectivity

For each of the 4 possible stimulus directions (i.e., C-stimulus position, or gap direction, dependent on sorting trials) we determined the time point in the stimulus window and the saccade window where the variance in firing rate among the different directions was the largest (i.e., the point with the biggest difference between the direction with the lowest and the highest firing rate). In a 200 ms window around that time point, the spike rate per trial was determined. ANOVA was computed for the spike rate in the window with the different directions as a grouping variable. If the test was significant, the modulation was determined to be direction selective ($\alpha=0.05$). The direction that showed the biggest change in firing rate from the baseline was selected as the preferred direction. To determine if a cell preferably modulated for the stimulus or action direction, i.e., the preferred direction sorting, the ANOVA with the lowest p-value was selected. The stimulus direction and action direction are inherently correlated since they are always mutually perpendicular. Therefore, both stimulus and action direction being significant was quite common, but does not necessarily mean that the cell actually is direction selective for both sortings.

Behavioral model

To determine which task modality and behavioral variables (e.g., gap direction and saccade direction in trial n-1) were most predictive of outcome response at the end of a trial (e.g., left or rightward saccade in trial n) a logistic regression model was fitted. Because logistic regression models data with binary outcomes, the model was fitted separately for the L/R gap direction trials (i.e., L/R task) and U/D gap direction trials (i.e., U/D task; [Figure 1A](#)). This logistic regression model provides a description of the data in terms of weights of behavioral variables that best predict the outcome of the task. Confidence intervals were created by performing 100 bootstrap permutations and represent 5 to 95% percentiles.

Encoding model

To more fully understand the dynamics of neural responses over the time course of a trial, we fitted a Generalized Linear Model (GLM) to predict the spike train data of each neuron as temporally overlapping responses to behavioral events in the trial, using the method and code described by ([Park et al., 2014](#)). In brief, the GLM predicts the probability $P[y_j(t) | \vec{\tau}_j, \vec{w}]$ of a given neuron producing a number $y_j(t)$ of spikes in a 25 ms time window at time t in trial j . This probability is a function of the times $\vec{\tau}_j$ of various behavioral events in that trial, and has free

<i>Event</i>	<i>Description</i>
<i>Stimulus</i>	<i>0-800 ms after C-onset; 200 ms sliding window with 100 ms steps, centered around 100 to 700 ms</i>
<i>Saccade</i>	<i>from 200 ms before until 200 ms after saccade onset; 200 ms sliding window with 100 ms steps</i>

Table 1.

Description of the sliding windows used to determine significant modulation for each category.

parameters $\vec{w}$ that were fit to data as described below. The model assumes that the neuron's spike counts are Poisson distributed around the mean spike rate $\lambda_j(t) = \exp[\mu_j + \sum_k \sum_i w_{ik} b_i^k(t)]$, where μ_j is the baseline firing rate of the neuron in trial j (moving average firing rate using a 5-trial window). $b_i^k(t) \equiv b_i(t - \tau_k)$ are linear kernels (basis functions) that capture the time-varying response of the neuron to the behavioral event at time τ_k . We chose $b_i(\tau)$ (the basis functions aligned to the start of the event) to be a series of five ($i \in \{1, \dots, 5\}$) cosine-shaped bumps with progressively larger widths and spanning up to 1.5s from the start of the event. The weights w_{ik} for each of these basis functions allow the neuron's modeled response to have a flexible time course, including the possibilities of rising above baseline (facilitation) if $w_{ik} > 0$, falling below baseline (suppression) if $w_{ik} < 0$, or even switching in time between the two depending on the relative values of $\vec{w}_k \equiv \{w_{ik} | i = 1, \dots, 5\}$. Six behavioral events were included in the model: the start of fixation (beginning of the trial), onset of the C-stimulus, start of the delay period, time of appearance of the choice target, time at which the saccade begins, and (for correct trials only) the time at which the reward is delivered (see [Figure 1B](#)). To account for the possibility that the neuron responds to only a subset of behavioral events, we fitted the model using sparse group LASSO regularization penalty (Huang and Zhang, 2010) with software from (Mairal et al., 2014). The variable groups in the penalty terms correspond to $\vec{w}_k$, i.e., all the basis-function weights for the response to a given behavioral event k , so that the fitted model will have $\vec{w}_k = \vec{0}$ (all weights being exactly zero) if a dependency on event k does not significantly improve the model prediction. We selected the regularization strength by maximizing the 5-fold cross-validated model likelihood.

Because neural responses also differed across subsets of trials with different task conditions (e.g., the orientation of the C-stimulus gap direction), we additionally fitted a variant of the GLM where the model parameters depended on the category c (defined below) of the trial. This model thus predicts that the neuron has spike counts $P[y_j^{(c)}(t) | \tau_j, \vec{w}^{(c)}]$ in trials of category c , with potentially different parameters $\vec{w}^{(c)}$ for every category c . We call the previous model with no category dependence the “unspecialized” model, and the model with category dependence the “category-modulated” model. We considered all possible categories of trials as specified by four types of trial conditions: C direction, gap direction, saccade direction, and values of the same in the previous trial (past-gap and past-saccade). The gap direction (same for saccade direction) can take on one of four values {U,D,L,R}, which means that there are 14 different ways to construct categories of trials by gap direction (all partitions of the set). These 14 possibilities are:

	categories
1	{U}, {D}, {L}, {R}
2	{U, D}, {L, R}
3	{U, L}, {D, R}
4	{U, R}, {D, L}
5	{U}, {D, L, R}
6	{D}, {U, L, R}
7	{L}, {U, D, R}
8	{R}, {U, D, L}
9	{U}, {D}, {L, R}
10	{U}, {L}, {D, R}
11	{U}, {R}, {D, L}
12	{D}, {L}, {U, R}
13	{D}, {R}, {U, L}
14	{L}, {R}, {U, D}

We furthermore wished to include the possibility that neuronal responses depend simultaneously on two or more trial conditions, e.g., both the current-trial gap and past-gap directions. As the number of all possible combinations of all trial-condition categories is exceedingly large, we used a greedy model selection procedure to select a parsimonious set of trial conditions that describes a given neuron's response sufficiently well. Starting from the unspecialized model as the “parent”, we first constructed a set of candidate “child” models, each of which corresponds to a particular choice of categorization for a particular trial condition (e.g., one of these would be {U,D},{L},{R} for saccade direction). We then selected the child model with the highest (cross-validated) likelihood L relative to its parent model, and if this is sufficiently high ($L_{child} > 10 L_{parent}$), we retain this child model as the best model so far. This best model was then used as the parent model for a second repetition of the same procedure. In subsequent repetitions, trial conditions that were already

included in the parent model were no longer considered for the child model (e.g., if the parent model had saccade-direction categories, then its child models will only include additional dependencies on gap, past-gap and past-saccade categories). This procedure terminates when adding categories to the model no longer significantly improves the model likelihood.

Decoding model

We used logistic regression to decode four behavioral quantities from the spiking activity of each neuron as a function of time in the trial (separately for each 200 ms time-bin aligned to different behavioral events). These behavioral quantities are: the C direction, gap direction, the outcome of the trial, i.e., whether the monkey receives a reward at the end of the current trial, and the previous-trial versions of these quantities (past-gap direction and past-outcome). Because the trial outcome (e.g., variable A) and gap directions (variable B) are both behaviorally correlated, neural information about one behavioral quantity (neural activity being correlated with A) can spuriously appear as if it was information about another (neural activity being correlated with B, but only through the *behavioral* correlation between A and B). To control for this issue we used a trial weighting procedure to “re-balance” the dataset so that all pairs of behavioral variables are in effect uncorrelated (Kobak et al., 2016 [DOI](#)). For a given neuron, we first divided the trials in its dataset into categories defined by the three other behavioral quantities of interest (see above). Since the trial outcome can take on two values (rewarded/not rewarded), and gap direction can take on four values (U/D/L/R), this means that there are $2 \times 4 \times 2 \times 4 = 64$ categories of trials. We then computed the weight w_i^c of trial i in category c as $w_i^c = 1/n^c$, where n^c is the total number of trials in category c .

Neuroanatomy

Tracer injections

At the end of the experiments, 18 - 19 days before perfusion, the recording areas of the DN of the animals were injected with 600 nl 0.8% of the tracer Cholera Toxin B (CTb) (Sigma-Aldrich, the Netherlands) dissolved in 0.01 M phosphate buffered saline. Injection position was verified with an electrode that was fixed to the injection line. When performing the tracer injections the tip of the electrode exceeded the tip of the capillary by 1 mm, the positional difference was corrected for by descending this distance deeper. The injectrode was lowered through the tentorium in an extra wide guide tube, allowing simultaneous access of both the electrode and the capillary. After perforation of the tentorium the tissue was left to settle for 20 minutes so as to allow the wider guide tube and injectrode to acquire a stable position. Next, the injectrode was lowered another 5-8 mm over 30 minutes so that the tissue was as stable as possible during the injection. Guided by the earlier neuronal recordings and current activity on the electrode, the tip of the capillary was lowered to a depth that was estimated to be the center of the task-related site. The 600 nl of the CTb solution was injected over a time course of 10 minutes, after which the injectrode was left in place for another 15 minutes. Likewise, the injectrode was retracted into the guide tube over the course of 15-20 minutes.

Perfusion

NHPs were transcardially perfused. The animals were deeply anesthetized with fentanyl and pentobarbital in their home cages, after which they were transported to the perfusion room. To prevent blood coagulation 2000 IE/kg of heparin was injected i.v. with 50 mL saline 15 minutes before the first incision was made. First, the cardiovascular system was flushed with 2-3L of 0.9% NaCl (saline) on a high perfusion speed of ~120 RPM, until the blood ran clear. Subsequently, the body was flushed with 3L fixative of 4% paraformaldehyde in 0.1 M phosphate buffer (pH 7.6) in

saline (PBS) with speed of the perfusion adjusted to 70 RPM, followed by 1.5L of 4% paraformaldehyde with 4% sucrose in 0.1 M PBS (pH 7.6, at 70 RPM). Thereafter, the brain was dissected and post-fixed in 4% paraformaldehyde for up to one week.

Immunocytochemistry

After post-fixation, brains were submerged in 10% sucrose PBS solution at 4°C until they sank. The brains were then embedded in 14% gelatin and 10% sucrose, after which they were fixated in 10% formalin and 30% sucrose overnight, and subsequently in 30% sucrose until they sank. The brains were then cut in 50 µm sections. For light-microscopic staining of the CTb, the sections were first incubated for 20 mins in 3% H₂O₂ (in PBS) to remove endogenous peroxidase activity of blood. Next, the sections were blocked in 10% normal horse serum (NHS) in PBS with 0.5% triton, after which they were incubated at room temperature with anti-CTb (goat, 1:15000, List labs) primary antibody (in PBS with 2% NHS, 0.5% triton) for 48-72 hrs and next with biotinylated rabbit anti-goat (1:200, Vector) secondary antibody (also in PBS with 2% NHS and 0.5% triton) for 1.5 hrs. After washing with PBS, the sections were incubated in Avidin-Biotine complex (1:200 Avidine and Biotine, Vector) in PBS and 0.5% triton for 1.5 hrs at room temperature. After washing with 0.5M PB sections were incubated for several mins with cobalt acetate 0.5% in milliQ or directly washed with milliQ, and subsequently briefly incubated with DAB (33.3 mg/100ml 3,3'-Diaminobenzidine; Sigma) and 17 µL H₂O₂ for visualization of the CTb. After washing with 0.5M PB, all sections were mounted with chrome alum and counterstained with thionin.

Image analysis

Sections were inspected under a Leica (Nussloch, Germany) DM-RB microscope, and relevant sections were identified for further scanning with a Hamamatsu Nanozoomer 2 whole slide imager. The resulting images were then exported as TIFF format for final quantification using NeuroLucida software (MicroBrightField, Inc., Colchester, VT). Sections of the cerebellar cortex (1 in 12), inferior olive (1 in 6) as well as pons (1 in 6) were imaged at 20x for quantification of retrogradely labeled neurons. For identification of the injection site, a 1 in 6 series of the cerebellar nuclei was inspected. For histological delineations we used the Rhesus Monkey Brain in Stereotaxic Coordinates Atlas by (Paxinos et al., 2000 [link](#)) as well as the (Carpenter and Madigan, 1971 [link](#); Gilman, 1972 [link](#)). Sections of the mesencephalon and diencephalon were imaged with the nanozoomer (20x objective) and analyzed with NDP-view (Hamamatsu) software.

Results

Two male rhesus macaques (Mo and Mi) were trained to perform a peripheral Landolt C task (Ignashchenkova et al., 2004 [link](#)), (**Figure 1A** [link](#)), allowing us to assess how DN cells encode visual stimuli and related (preparatory) motor activity. The animals had to maintain central fixation throughout the trial as long as the central fixation dot remained red (see example trial in **Figure 1B** [link](#)). After a delay of ~500 ms after fixation onset, the C-stimulus was randomly presented in one of 4 cardinal directions (Right = 0°, Left = 180°, Up = 90°, Down = 270°) for the duration of 250 ms. The animals had to perceive the direction of the gap of the C (“gap direction”), which was always positioned perpendicular to the stimulus location. Specifically, the gap direction is either Up or Down [U/D] when the C-stimulus was presented in locations L or R [L/R], or the gap direction is either Left or Right [L/R] when the C-stimulus was presented in locations U or D. After the presentation of the C-stimulus, the C was masked for 100 ms to prevent retinal afterimages, and subsequently there was a delay period of 500 - 700 ms in which Mi and Mo had to remember the gap direction. After the delay period, two saccade targets were shown at the same eccentricity; one target was placed in the same direction as the gap direction (i.e., the correct target), and the other one was presented as a distractor. If the animals made a saccade to the correct target within 500 ms after the target onset, they received a juice reward.

Task and behavior

There were significant differences in task performance levels between Mi and Mo that did not depend on the gap direction (**Figure 1C**). Mo consistently outperformed Mi, and this trend was present across different experimental sessions (**Figure 1D**). It should be noted that this analysis uses data from after the monkeys had been fully trained to understand the task structure, so the performance differences do not directly represent the structure of the learning process itself. Still, since for each animal the task sessions were separated by weeks-long breaks, during which the other animal was subjected to electrophysiological recordings while performing the behavioral task, some level of retraining likely took place when task sessions were restarted after a break.

To assess which task variables contributed to the behavioral performance in the individual animals we split the type of tasks in two sets: the upward and downward directions (U/D task) and the leftward and rightward directions (L/R task). This allowed us to apply a logistic regression model with the dependent variable being the fraction of correct responses, and the independent variables being those that specify the task structure. We first explored how the gap direction presented in the current trial was weighted to best predict the animals' performance over the sessions. The weight that Mi assigned to the U/D gap direction increased simultaneously with task performance. In contrast, the weight that was assigned to the L/R stimulus remained constant, suggesting that his overall performance increase was mostly mediated by his improvement in the subset of up-/downward trials in the task (**Figure 1E**, left panel). Mo, on the other hand, exhibited increased weighting of the stimulus in both U/D and L/R trials. Indeed, the overall higher performance of Mo relative to Mi was reflected in the model on Mo's data having higher weights for the gap direction throughout all measured sessions (**Figure 1E**, right panel). We next assessed if any past-trial conditions influenced the behavioral response in the current trial. Therefore, we determined the weight of events gap direction and saccade direction in the previous trial with the same logistic regression approach. According to this model, Mi assigned less weight to the previous saccade over time if the current trial was in the L/R direction ($p = 0.016$; $R^2 = 0.32$, **Supplemental Figure 1**). In contrast, there was no change in event weight to predict the response on the current trial in the past trial variables for Mo.

DN activity

Once the animals reached an above-chance performance level (60%), we identified the site of the DN in both Mi and Mo with the use of structural MRI (**Figure 2A**). Single unit activity was recorded of 304 DN neurons, while the animals performed the Landolt C task. The activity of DN neurons in both Mi and Mo showed correlations with the temporal windows of both the instruction and movement stage (**Figures 2B** and **2D**). For example, during the presentation of the C-stimulus a DN cell could respond with an increase in activity, followed by a pause around the time of the correctly executed saccade (**Figure 2C**). Yet, the level of activity during the burst after the presentation of the C-stimulus could be selective for trials with different directions; for example, it could be higher for trials during which the gap direction was either left or up, compared to those with right or down directions (**Figure 2D**). Differences in spiking for different directions were determined by comparing the amount of spikes in a bin at the moment of largest divergence between directions (see Methods). Changes of neuronal activity in relation to the presentation of the C-stimulus and/or the saccade were common (**Figures 2E** and **2F**). For example, cells could show significant facilitation of their activity in a window associated with the C-stimulus (200 ms moving window, 100 ms steps from 0-800 ms after C-stimulus onset) or the saccade (200 ms moving window, 100 ms steps from -200 ms before to 200 ms after saccade onset) relative to baseline activity from before trial onset (see material and methods for details) (**Figure 2E**). Likewise, cells could also show suppression of their firing following the C-stimulus and the saccade (**Figure 2F**). When we aligned the neurons to the time of maximum change after C-stimulus onset, the population of suppressing neurons ($n = 85$) exhibited the modulation

significantly earlier ($p < 0.0041$; Kolmogorov–Smirnov test) than that of the facilitating units ($n = 90$), (**Figure 2G**). This was not the case ($p = 0.23$) for bursting ($n = 57$) and pausing ($n = 38$) cells during the saccade window.

Encoding of task events

Many DN cells modulated their activity in relation to more than one task events. To fully characterize the neural responses we fitted a generalized linear model (GLM) and analyzed which of the task events were encoded in the spike activity (**Figure 3**). This model predicts the spiking activity time course of individual neurons as a weighted sum of time-dependent basis functions in response to every task event (Park et al., 2014). LASSO regularization was used to select a parsimonious set of task events that should be included in the model (**Figures 3A**, **3B** and **3C**; for details see materials and methods). We found that many cells encoded the gap direction, the saccade or both, while a smaller fraction was sensitive to the events in the previous trial (**Figure 3B** and **3C**). Many units showed firing rate modulation in relation to both particular gap directions and saccades (**Figure 3D** and **3E**). Likewise, a neural response after the saccade in trials where the animal made a mistake was a common feature. Since retinal errors were not expected to be a relevant feature of this task, at least not in the sense of those that may occur in classic saccadic adaptation paradigms (Herzfeld et al., 2018), it is possible that these responses are related to the absence of reward (**Figure 3D** and **3E**), (see also Heffley and Hull, 2019; Kostadinov et al., 2019). Interestingly, these correlations could come with both facilitations and suppressions (**Figure 3D** and **3E**), in line with upbound and downbound spike changes (De Zeeuw, 2021). Despite the level of heterogeneity among the two NHPs in terms of performance level and strategy, the fractions of recorded neurons that were sensitive to particular task parameters, including gap direction, saccade direction or miscellaneous were nearly the same in both animals, indicating that we sampled from the same populations of DN neurons in the two NHPs (**Figure 3C**).

Decoding trial variables

Considering the heterogeneity of the neural activity in relation to the task, we built a decoding model to reveal which parameters of the task could reliably be predicted from the spiking activity of individual cells. For each 200 ms time window in the trial, we fit a logistic regression model to determine at which time points a given trial condition could be decoded from the cell's activity. The condition that was most reliably represented was the trial outcome, i.e., whether the monkey made a correct or incorrect choice (**Figure 4A**). The proportion of cells from which the trial outcome could be decoded increased at about 1 second following the presentation of the C-stimulus to the monkey, and this persisted throughout the delay period and remained present for more than a second after the monkey had made his choice (**Figure 4A**, see different columns). Compared to Mo, Mi had more cells from which the trial outcome could be decoded around the time of the reward. This may be due to the overall poorer performance of Mi, resulting in more error trials which may result in stronger neural responses. Instead, Mo showed many cells in which we could decode information about the C-stimulus, including the gap direction, which is relevant to making a correct choice. Possibly, this was particularly visible in Mo as his DN activity was generally better entrained to the stimulus, leading to a better overall behavioral performance (**Figure 4B**). In line with the findings of the behavioral model, Mi instead had more cells than Mo that maintained information about the choice he had made in the previous trial, well into the time-frame of the ongoing trial. This information was notably present at about 1 to 1.5 seconds after fixation at the center, and gradually disappeared from the onset of the C stimulus and beyond (**Figure 4C**). Lastly, there were almost no cells from which the gap direction of the past trial could be decoded in either NHP (**Figure 4D**).




Figure 3.

Encoding of task events by DN neurons.

A) Schematic of a generalized linear model (GLM) for the spike count of a given neuron as being Poisson-distributed around behavior-dependent mean rates. The neuron is predicted to have time-varying changes in activity triggered by one or more behavioral events, and these changes are flexibly modeled as weighted sums of event-aligned basis functions $b_i^k(t)$. The set of events that the neuron responds to, as well as any trial-type specificity of these responses, was selected so as to obtain the most parsimonious model that produced a good fit. **B**) Distribution across cells (two colors for the two monkeys) of a “modulation index” (modIdx) score for how much trial types influence neural responses (see also examples for individual cells in panels D and E). First, we constructed an “unspecialized” GLM model that predicts the cell’s activity rate in trial j to be $\lambda_j(t)$, independently of trial type. This is compared to a trial-type dependent GLM prediction $\lambda_j^{(c)}(t)$ for that cell, where the significant trial-type categories c for the model were parsimoniously selected as described in the Methods. The modulation index for the cell is then defined using the ratio of trial-type specific to unspecialized model predictions: $\rho_j^{(c)}(t) \equiv \lambda_j^{(c)}(t) / \lambda_j(t) - 1$. To produce a single summary number per cell, we computed modIdx^2 as the time- and trial-type average of $[\rho_j^{(c)}(t)]^2$. The four subplots of this panel show the distribution of modIdx scores for cells with GLM predictions $\lambda_j^{(c)}(t)$ where the model-selected trial category c included gap direction (top left), saccade direction (top right), past-trial gap direction (bottom left), and past-trial saccade direction (bottom right). **C**) Proportions of cells with GLM models that had significant dependencies on various trial types as indicated in the legend of panel A. The miscellaneous 1- to 4-category proportions were pooled over cells with infrequently occurring trial-type dependencies (<2% cells per category); for example, the “misc. 3-cat.” cells included those with simultaneous trial-type dependencies on gap, past-trial gap, and past-trial saccade conditions. **D**) GLM prediction for the firing rate vs. time in a trial of 3 example neurons recorded in Monkey Mi. For each cell (one column), the predicted firing rate is shown for trials of 4 different gap directions (differently colored lines), and separately for correct trials (top row) and error trials (bottom row). The onset of various events in the trial are indicated with vertical dashed lines. **E**) Same as D, but for 3 example cells from monkey Mo.




Figure 4.

Decoding of trial conditions from neural firing rates.

A) Fraction of cells from which the trial outcome can be significantly decoded (see Methods) using data in 200 ms time bins aligned to various events in the trial (columns). * indicate bins in which this fraction is significantly different between the two monkeys ($p < 0.05$; Fisher's exact test, 2-tailed). **B-D)** same as A but for the decoding of gap direction, outcome of the past trial and gap direction of the past trial, respectively.

Identification of recording site in DN and its connectivity

The area of the DN with the most successful recordings was confirmed post mortem with the use of CTb injections (**Figure 5A** [↗](#)). In both cases, the injections were focused on the centrolateral part of the DN positioned in the center of its rostrocaudal axis; given the distribution of the modulating cells according to the stereotactic coordinates with respect to the 2D-grid of the recording chamber and the recording depths, we estimated that the functionally relevant recording area covered a vertical column of approximately 2 mm long, 2 mm wide and 3 mm deep, as all modulated cells were recorded in a square of that size. Despite a similar location for the core of the injection site, the size of the injection was bigger in Mo as compared to Mi. Accordingly, when we quantified the number of retrogradely labeled neurons in 1 out of 4 sections of the inferior olive (IO), so as to assess the relative effective size of the DN injections in both animals, we found more labeled cells in Mo (**Figure 5B** [↗](#) and **5C** [↗](#)). In Mi the dorsolateral and ventrolateral leaf of the principal olivary nucleus (PO) contained 38 and 78 cells, respectively, while the ventrolateral outgrowth of the PO contained only 4 cells. Instead, in Mo we counted 1977, 763 and 312 neurons in the dorsolateral leaf, ventrolateral leaf, and ventrolateral outgrowth of the PO, respectively (**Figure 5C** [↗](#)). The labeling in the PO of the IO was in line with the retrograde labeling in the pons, which was predominantly confined to the medial pontine nuclei in both Mi and Mo, with Mo showing more labeled cells (**Figure 5D** [↗](#)). Likewise, Mo showed more retrogradely labeled Purkinje cells (PCs; $n = 1314$) than Mi ($n = 84$), when we screened 1 out of 24 sections of the cerebellar cortex (**Figure 5E** [↗](#) and **5F** [↗](#)). Most of the labeled PCs appeared to be located in the D1 microzones of the cerebellar cortex (Voogd et al., 2012 [↗](#)).

Interestingly, the CTb injections into the DN also provided anterograde labeling of the climbing fibers, the distribution of which overlapped to a large extent with that of the retrograde labeling of the PCs in both Mi and Mo (lateral panels in **Figure 5F** [↗](#)). In addition, we found anterograde CTb labeling in the higher brainstem; this included for example the parvocellular and magnocellular red nucleus, zona incerta and deep mesencephalic nucleus as well as various thalamic nuclei, such as the ventrolateral thalamic nucleus (VL), pulvinar nucleus (Pul), mediodorsal thalamic nucleus (MD), and centromedial and centrolateral thalamic nuclei (**Figure 5G** [↗](#)). Instead, we did not find substantial labeling in the superior colliculus, ventral tegmental area or hypothalamus. The labeled areas we found are consistent with previous studies illustrating disynaptic pathways from the DN to cerebral cortical areas involved in visual processing such as the frontal eye field area (FEF; Middleton and Strick, 2001 [↗](#)).

Discussion

We investigated to what extent the activity of DN cells in the cerebellum of NHPs encodes different aspects of a complex visual-motor Landolt C task. Task performance was related to individual behavioral strategies of animals that changed over time during and after post-break relearning of the task. We found that DN cells connected with the D1 and D2 zones in the cerebellar cortex have highly diverse activity patterns. Many DN neurons respond in a visual direction- or movement direction-selective manner more than half a second before an upcoming saccade, with relatively short latencies after the animal receives the visual information that instructs saccade direction. The activity of DN cells often bridged different experimental epochs and frequently showed both upbound and downbound modulations (De Zeeuw, 2021 [↗](#)), similar to PCs in the lateral cerebellum that project to the DN (Avila et al., 2022 [↗](#)). In these ways, multi-modality appears to be a common property of neurons in the lateral cerebellum.




Figure 5.

Anatomical characterization of recording area and its input and output connectivity.

A) CTb injections in the dentate nucleus of Mi and Mo displays the site of electrophysiological recordings of its neurons at the lateral edge of the DN. **B)** Pictures of retrogradely labeled cells in the Principal Nucleus of the inferior olive (IO); insets show overview of IO. **C)** Quantification of retrogradely labeled cells in different subnuclei of the IO in Mi (blue) and Mo (pink). **D)** Examples of retrogradely labeled cells in the medial pons of Mi (left) and Mo (right); insets show higher magnification of labeled cells. **E)** Schematic representations of cerebellar cortical areas containing retrogradely labeled Purkinje cells in Mi (left) and Mo (right). **F)** Examples of retrogradely labeled Purkinje cells for both animals (Mi and Mo in panels on the left and right, respectively); insets show areas of the higher magnifications with the labeled cells. **G)** Example of anterograde CTb labeling of cerebellar fibers with dense varicosities (arrowheads) in the thalamus of Mo. Panels i and ii on the right correspond to insets in the left panel; panels iii and iv are examples of anterograde CTb labeling outside of the left panel in the anterior pulvinar and red nucleus, respectively. VL, LD, CL, APuL, VPL, CM and MD indicate ventrolateral, laterodorsal, centrolateral, anterior pulvinar, ventroposterior lateral, centromedial and mediodorsal nucleus of the thalamus, respectively. diPO, viPO: dorsolateral-, ventrolateral Principle Olive, vlo: ventrolateral outgrowth; MAO and DAO: medial and dorsal accessory olive. Scale bars in A, B, D, F and G indicate 1250 μ m, 100 μ m, 1000 μ m, 50 μ m and 1000 μ m, respectively.

Visuomotor DN neurons connect to cerebellar modules in D1 and D2 zone

The distribution of the retrogradely labeled PCs in lateral lobules VII and VIII as well as crus I/II indicates that the region where we recorded in the DN forms part of the cerebellar D1 and D2 microcomplex (De Zeeuw, 2021 [↗](#); De Zeeuw and Ten Brinke, 2015 [↗](#)). This conclusion is supported by the retrograde CTb labeling we found in the part of the medial pons that is known to receive direct input from the prefrontal cortex and to activate the D zones during decision making and motor planning (Buckner et al., 2011 [↗](#); Prevosto et al., 2010 [↗](#); Ramnani, 2012 [↗](#); Salmi et al., 2010 [↗](#); Stoodley and Schmahmann, 2010 [↗](#); Strick et al., 2009 [↗](#); Wang et al., 2022 [↗](#)). Likewise, most of the retrograde labeling in the olive following the injections in the DN was present in the dlPO and vlPO, which are also part of the D1 and D2 module (Voogd et al., 2012 [↗](#); Voogd and Ruigrok, 2012 [↗](#)).

Even though both Mi and Mo showed retrograde CTb labeling patterns consistent with involvement of the D modules, the density of labeling differed substantially, presumably due to differential efficacy of the CTb injections (**Figure 5A** [↗](#)). Accordingly, we observed the same differential pattern when we analyzed the anterograde labeling results. Here too, Mo showed more densely labeled fibers in the various projection regions than Mi. Most anterogradely labeled fibers were found in the pulvinar, mediodorsal thalamus and ventrolateral thalamus, which are known to connect to various parts of the prefrontal cortex, such as FEF and Brodmann area 46 (Kipping et al., 2013 [↗](#); Middleton and Strick, 2001 [↗](#); Romanski et al., 1997 [↗](#)), and/or the parietal eye field (PEF), an established hub in the dorsal attention network (Andersen et al., 1992 [↗](#); Hanks et al., 2015 [↗](#); Roitman and Shadlen, 2002 [↗](#); Shadlen and Newsome, 2001 [↗](#); Svoboda and Li, 2018 [↗](#)). Interestingly, the dorsal attention network has been shown to connect to lobules VII and VIII (Brissenden et al., 2016 [↗](#); Buckner et al., 2011 [↗](#)), precisely the location of the cerebellar cortex where retrograde labeling was detected following the DN injections. Taken together, the anatomical findings suggest that the DN region under study is at least partly integrated in a functional closed loop between cerebellum and cerebral cortex that appears to be involved in decision making, planning and attention.

Visuomotor processing by the DN

In the current study, the animals had to read a peripherally located C, while fixating their eyes on a central fixation point. After a delay, the primates had to move their eyes to a virtual point in space dependent on the cue implied by the peripheral C-stimulus. The spatial and temporal dissociation of cue and action in this task allows us to dissociate neural activity related to the visual and attentional processes from that related to motor coordination (Avila et al., 2022 [↗](#); Goldberg and Segraves, 1987 [↗](#)). Our data reveal that individual DN cells can encode both the direction of the visual cue and the associated direction of the subsequent saccade, suggesting that different task-specific activations in the cerebellar cortex may be relayed downstream to the same target neurons in the cerebellar nuclei. Our data are compatible with and expand upon fMRI studies, which have demonstrated that lateral lobules VIIb/VIIIa upstream of the DN exhibit functional properties characteristic of the dorsal attention network in the cerebral cortex (Brissenden et al., 2018 [↗](#), 2016 [↗](#); van Es et al., 2019 [↗](#)). Indeed, this part of the cerebellar hemispheres shows task-specific activation in that their responses depend on representation of visuospatial location and/or load of working memory (Brissenden et al., 2018 [↗](#), 2016 [↗](#); van Es et al., 2019 [↗](#)).

Our findings are further corroborated by electrophysiological studies that have revealed a role of the cerebellar nuclei, including that of the DN, in motor planning (Chabrol et al., 2019 [↗](#); Deverett et al., 2019 [↗](#), 2018 [↗](#); Gao et al., 2018 [↗](#)). The DN cells recorded in the current study may contribute to the preparation of directed saccades through their projections to the FEF via the thalamus (Lindeman et al., 2021 [↗](#); Middleton and Strick, 2001 [↗](#); Romanski et al., 1997 [↗](#)). Similar

to our DN neurons, FEF neurons encode multiple aspects of saccade planning and processing, including target identification in covert visual searches, as well as decision making among multiple choices (Ding and Gold, 2012 [DOI](#); Gregoriou et al., 2012 [DOI](#); Hanes and Schall, 1996 [DOI](#); Monosov and Thompson, 2009 [DOI](#); Schall and Hanes, 1993 [DOI](#)). In addition, the FEF forms, together with the supplementary eye fields and parietal eye fields, an integral part of the dorsal attention network that is required for our visuomotor task (Brissenden et al., 2018 [DOI](#), 2016 [DOI](#); Buckner et al., 2011 [DOI](#); van Es et al., 2019 [DOI](#)). Intriguingly, we found that the activity of many DN neurons exhibited facilitation or suppression before an upcoming saccade (**Figure 2G** [DOI](#) and **2H** [DOI](#)), similarly to how some FEF neurons show selective ramping activity before the upcoming eye movement (Basu and Murthy, 2020 [DOI](#); Raghavan and Joshua, 2017 [DOI](#)). Such ramping activity is one of the main signatures of motor preparation in general (Chabrol et al., 2019 [DOI](#); Devereux et al., 2019 [DOI](#); Gao et al., 2018 [DOI](#); Hanks et al., 2015 [DOI](#); Svoboda and Li, 2018 [DOI](#)).

The multimodal character of the DN activity

The cerebellum receives inputs from many different external sensory as well as internal cognitive modalities, all of which can be integrated during the preparation and execution of complex tasks (De Zeeuw, 2021 [DOI](#)). In the cerebellar cortex such integration can take place at both the input and output stage. At the input stage, i.e., at the level of the granule cells in the granular layer, activation of different sensory modalities has been shown to enhance spiking output (Giovannucci et al., 2017 [DOI](#); Ishikawa et al., 2015 [DOI](#)). Moreover, given that many individual granule cells can receive inputs from both the cuneate nucleus relaying sensory proprioceptive signals and the pontine nuclei mediating presumptive motor command signals (Guo et al., 2021 [DOI](#); Huang et al., 2013 [DOI](#); Wagner et al., 2019 [DOI](#)), it is likely that sensory, motor and cognitive signals can also be integrated at the level of the cerebellar input layer. At the level of the Purkinje cells in the molecular layer with inputs from thousands of parallel fibers contacting single Purkinje cell dendrites, the opportunities for convergence and integration of different modalities are even greater (Gao et al., 2012 [DOI](#)).

Our results demonstrate that integration of various forms of sensory, motor and/or cognitive signals also occurs at the level of the DN neurons downstream of Purkinje cells in the D1 and D2 zones. Indeed, single DN neurons can encode the direction of a visual stimulus, as well as the preparation and execution of a saccadic eye movement into a particular direction. The encoding of both stimulus and saccade directions is present in the biggest proportion of task-modulated neurons in both monkeys. The decoding analysis showed similar outcomes, albeit with greater variability between Mi and Mo. To what extent the integration of different modalities is taking place at the level of the DN itself, mediated by converging inputs from the Purkinje cells, mossy fibers and/or climbing fiber collaterals, or solely by events upstream in the cerebellar cortex, remains to be elucidated.

Mechanisms underlying covert attention

Our electrophysiological data on DN encoding during the Landolt C task are in line with fMRI studies showing covert attention signals in the cerebellum (Brissenden et al., 2018 [DOI](#), 2016 [DOI](#); van Es et al., 2019 [DOI](#)). These neuro-imaging studies, which have also implemented a consistent distractor stimulus to evaluate the validity of the task (Posner, 1980 [DOI](#)), point towards a coordination of cerebral and cerebellar cortical activation during covert attention. In this respect, both our electrophysiological study and the imaging studies of others elaborate on the premotor theory of covert attention, which has so far mainly highlighted the relevance of processing in the cerebral cortex (Corbetta et al., 1998 [DOI](#); Nobre et al., 2000 [DOI](#); Rizzolatti et al., 1987 [DOI](#)). More specifically, the premotor theory highlights the relevance of circuits in the frontal and parietal lobes that are activated during both attention shifts and subsequent movements, so as to strengthen the association process. Our data show that most of the neurons in the main output nucleus of the cerebellum, the DN, exhibit heterogeneous modulations during putative attention shifts made when the animals read the peripheral C stimulus, as well as during the subsequent

goal-directed saccadic eye movements that depend on this preceding cue in the peripheral visual field. The converging coordination of cerebellum and cerebral cortex of higher order mental processes such as covert attention also occurs during other cognitive processes. For example, not only the mirror neuron system in cerebral cortex but also connected areas in the cerebellum are actively involved in perceiving and interpreting the action of others (Abdelgabar et al., 2019 [DOI](#)). Interestingly, just like during the Landolt C task, lobules VII and VIII in the cerebellar hemispheres are recruited when evaluating the kinematics of goal-directed hand actions of others. Moreover, patients suffering from spinocerebellar ataxia type 6 are severely impaired in performing such action observation tasks (Abdelgabar et al., 2019 [DOI](#)). These data raise the question as to how the cerebellum may contribute to overt and covert attentional tasks. One can hypothesize that cerebellar processing helps the frontal premotor and parietal cortices to map visual input, from high level visual regions, onto the motor machinery involved in performing related goal-directed actions. This suggests that much like action control (Wolpert and Ghahramani, 2000 [DOI](#)), complex mental observation, either overt or covert, relies on a cortico-cerebellar loop that maps sensory input onto motor control structures (inverse models) and motor programs to expected sensory input (forward models). This loop may bring descending information from our visual cortical networks to the cerebellum and ascending information from the cerebellum back to premotor areas in the prefrontal cerebral cortex.

Acknowledgements

Financial support was provided by the Netherlands Organization for Scientific Research (NWO-ALW 824.02.001; CIDZ), the Dutch Organization for Medical Sciences (ZonMW 91120067; CIDZ and Vidi/ZonMw/917.18.380,2018; AB), Medical Neuro-Delta (MD 01092019-31082023; CIDZ), INTENSE LSH-NWO (TTW/00798883; CIDZ), ERC-adv (GA-294775 CIDZ) and ERC-POC (nrs. 737619 and 768914; CIDZ); The NIN Vriendenfonds for Albinism the Dutch NWO Gravitation Program (DBI2; CIDZ) and Erasmus MC Convergence Flagship (Integrative Neuromedicine Convergence Health and Technology 2022; AB). We thank Kor Brandsma and Anneke Ditewig for biotechnical assistance and animal caretaking; Beerend Winkelman for fruitful discussions; Masaki Tanaka and Jun Kunimatsu for sharing the design of the injectrode used for the tracer injections; and Erika Goedknecht for her indispensable contributions to the processes of anatomical tracing and perfusion. Finally, we would like to thank Peter Strick for his advice on the anatomical tracing part of the study.


Contributions

N.A. Flierman, A. Badura, P.R. Roelfsema and C.I. De Zeeuw designed the system's physiological experiments; N.A. Flierman performed the recordings; N.A. Flierman and A. Badura performed the tracer injections; N.A. Flierman, A. Badura, W.S. van Hoogstraten, T.J.H. Ruigrok and C.I. De Zeeuw designed the neural tracing experiments and performed the stainings and analysis of the anatomy; N.A. Flierman analyzed the data; Sue Ann Koay did the encoding and decoding analyses; P.R. Roelfsema was responsible for the housing of the NHPs, N.A. Flierman, A. Badura and C.I. De Zeeuw wrote the manuscript; A. Badura and C.I. De Zeeuw coordinated the project; C.I. De Zeeuw and A. Badura obtained funding from NWO, LSH, NIN and EU to execute the project.

Ethical statement

All experimental and surgical procedures complied with the NIH Guide for the Care and Use of Laboratory Animals (National Institutes of Health, Bethesda, Maryland), and were approved by the institutional animal care and use committee of the Royal Netherlands Academy of Arts and Sciences (AVD8010020184587).

Supplementary Material



Supplementary Figure 1.

Weight to predict response on the current trial from previous trial parameters over time; the trials could be from either a L/R or a U/D task, hence the two conditions per panel. The same logistic regression model as in [Figure 1](#) was applied to determine the influence of previous trial characteristics on current trial behavior (See [Supplementary Table 1](#) for the statistical tests).

Mo				
		Gap L/R n-1	Gap U/D n-1	Sac n- 1
LR	R	0,27	0,03	0,00
	p	0,03	0,47	0,98
UD	R	0,00	0,07	0,00
	p	0,93	0,30	0,97

Mi				
		Gap L/R n-1	Gap U/D n-1	Sac n- 1
LR	R	0,00	0,05	0,32
	p	0,86	0,37	0,016
UD	R	0,02	0,11	0,13
	p	0,59	0,19	0,16

Supplementary Tables 1 and 2.

Regression analyses applied to the data from **Supplementary Figure1** [↗](#). Only two of the regression models do significantly better than a constant model (F-test, highlighted in bold). Both of those do not survive a Bonferroni correction ($\alpha = 0.0083$ for 0.05/6 test per monkey).

References

- Abdelgabar AR, Suttrup J, Broersen R, Bhandari R, Picard S, Keyzers C, De Zeeuw CI, Gazzola V (2019) **Action perception recruits the cerebellum and is impaired in patients with spinocerebellar ataxia** *Brain* **142**:3791–3805
- Andersen RA, Brotchie PR, Mazzoni P (1992) **Evidence for the lateral intraparietal area as the parietal eye field** *Curr Opin Neurobiol* **2**:840–846
- Avila E, Flierman NA, Holland PJ, Roelfsema PR, Frens MA, Badura A, De Zeeuw CI (2022) **Purkinje Cell Activity in the Medial and Lateral Cerebellum During Suppression of Voluntary Eye Movements in Rhesus Macaques** *Front Cell Neurosci* **16**
- Baier B, Dieterich M, Stoeter P, Birklein F, Müller NG (2010) **Anatomical Correlate of Impaired Covert Visual Attentional Processes in Patients with Cerebellar Lesions** *J Neurosci* **30**:770–776
- Basu D, Murthy A (2020) **Parallel programming of saccades in the macaque frontal eye field: are sequential motor plans coactivated?** *J Neurophysiol* **123**:107–119
- Brissenden JA, Levin EJ, Osher DE, Halko MA, Somers DC (2016) **Functional Evidence for a Cerebellar Node of the Dorsal Attention Network** *J Neurosci* **36**:6083–6096
- Brissenden JA, Tobyn SM, Osher DE, Levin EJ, Halko MA, Somers DC (2018) **Topographic Cortico-cerebellar Networks Revealed by Visual Attention and Working Memory** *Curr Biol* **28**:3364–3372
- Buckner RL, Krienen FM, Castellanos A, Diaz JC, Yeo BTT (2011) **The organization of the human cerebellum estimated by intrinsic functional connectivity** *J Neurophysiol* **106**:2322–2345
- Carpenter MB, Madigan JC (1971) **Cerebellum of the rhesus monkey: atlas of lobules, laminae, and folia, in sections**
- Chabrol FP, Blot A, Mrcic-Flogel TD (2019) **Cerebellar Contribution to Preparatory Activity in Motor Neocortex** *Neuron* **103**:506–519
- Chen X, Possel JK, Wacongne C, van Ham AF, Klink PC, Roelfsema PR (2017) **3D printing and modeling of customized implants and surgical guides for non-human primates** *J Neurosci Methods* **286**:38–55
- Corbetta M *et al.* (1998) **A common network of functional areas for attention and eye movements** *Neuron* **21**:761–773
- Courchesne E, Townsend J, Akshoomoff NA, Saitoh O, Yeung-Courchesne R, Lincoln AJ, James HE, Haas RH, Schreibman L, Lau L (1994) **Impairment in shifting attention in autistic and cerebellar patients** *Behav Neurosci* **108**:848–865
- Craighero L, Nascimben M, Fadiga L (2004) **Eye position affects orienting of visuospatial attention** *Curr Biol* **14**:331–333

- Deverett B, Kislin M, Tank DW, Wang SS-H (2019) **Cerebellar disruption impairs working memory during evidence accumulation** *Nat Commun* **10**
- Deverett B, Koay SA, Oostland M, Wang SS-H (2018) **Cerebellar involvement in an evidence-accumulation decision-making task** *Elife* **7**
- De Zeeuw CI (2021) **Bidirectional learning in upbound and downbound microzones of the cerebellum** *Nat Rev Neurosci* **22**:92–110
- De Zeeuw CI, Ten Brinke MM (2015) **Motor Learning and the Cerebellum** *Cold Spring Harb Perspect Biol* **7**
- Ding L, Gold JI (2012) **Neural correlates of perceptual decision making before, during, and after decision commitment in monkey frontal eye field** *Cereb Cortex* **22**:1052–1067
- Flierman NA, Ignashchenkova A, Negrello M, Thier P, De Zeeuw CI, Badura A (2019) **Glissades Are Altered by Lesions to the Oculomotor Vermis but Not by Saccadic Adaptation** *Front Behav Neurosci* **13**
- Gao Z, Davis C, Thomas AM, Economo MN, Abrego AM, Svoboda K, De Zeeuw CI, Li N (2018) **A cortico-cerebellar loop for motor planning** *Nature* **563**:113–116
- Gao Z, van Beugen BJ, De Zeeuw CI (2012) **Distributed synergistic plasticity and cerebellar learning** *Nat Rev Neurosci* **13**:619–635
- Gilman S (1972) **Cerebellum of the Rhesus Monkey: Atlas of Lobules, Laminae, and Folia, in Sections** *Arch Neurol* **26**:192–192
- Giovannucci A *et al.* (2017) **Cerebellar granule cells acquire a widespread predictive feedback signal during motor learning** *Nat Neurosci* **20**:727–734
- Goldberg ME, Segraves MA (1987) **Visuospatial and motor attention in the monkey** *Neuropsychologia* **25**:107–118
- Gregoriou GG, Gotts SJ, Desimone R (2012) **Cell-type-specific synchronization of neural activity in FEF with V4 during attention** *Neuron* **73**:581–594
- Guo J-Z, Sauerbrei BA, Cohen JD, Mischiati M, Graves AR, Pisanello F, Branson KM, Hantman AW (2021) **Disrupting cortico-cerebellar communication impairs dexterity** *Elife* **10** <https://doi.org/10.7554/eLife.65906>
- Hanes DP, Schall JD (1996) **Neural control of voluntary movement initiation** *Science* **274**:427–430
- Hanks TD, Kopec CD, Brunton BW, Duan CA, Erlich JC, Brody CD (2015) **Distinct relationships of parietal and prefrontal cortices to evidence accumulation** *Nature* **520**:220–223
- Hanning NM, Deubel H (2020) **Attention capture outside the oculomotor range** *Curr Biol* **30**:R1353–R1355
- Heffley W, Hull C (2019) **Classical conditioning drives learned reward prediction signals in climbing fibers across the lateral cerebellum** *Elife* **8** <https://doi.org/10.7554/eLife.46764>

- Herzfeld DJ, Kojima Y, Soetedjo R, Shadmehr R (2018) **Encoding of error and learning to correct that error by the Purkinje cells of the cerebellum** *Nat Neurosci* **21**:736–743
- Huang C-C, Sugino K, Shima Y, Guo C, Bai S, Mensh BD, Nelson SB, Hantman AW (2013) **Convergence of pontine and proprioceptive streams onto multimodal cerebellar granule cells** *Elife* **2**
- Huang J, Zhang T (2010) **The benefit of group sparsity** *aos* **38**:1978–x2004
- Ignashchenkova A, Dicke PW, Haarmeier T, Thier P (2004) **Neuron-specific contribution of the superior colliculus to overt and covert shifts of attention** *Nat Neurosci* **7**:56–64
- Inoshita T, Hirano T (2018) **Occurrence of long-term depression in the cerebellar flocculus during adaptation of optokinetic response** *Elife* **7** <https://doi.org/10.7554/eLife.36209>
- Ishikawa T, Shimuta M, Häusser M (2015) **Multimodal sensory integration in single cerebellar granule cells in vivo** *Elife* **4** <https://doi.org/10.7554/eLife.12916>
- Ivry RB, Diener HC (1991) **Impaired velocity perception in patients with lesions of the cerebellum** *J Cogn Neurosci* **3**:355–366
- Kipping JA, Grodd W, Kumar V, Taubert M, Villringer A, Margulies DS (2013) **Overlapping and parallel cerebello-cerebral networks contributing to sensorimotor control: an intrinsic functional connectivity study** *Neuroimage* **83**:837–848
- Kobak D, Brendel W, Constantinidis C, Feierstein CE, Kepecs A, Mainen ZF, Qi X-L, Romo R, Uchida N, Machens CK (2016) **Demixed principal component analysis of neural population data** *Elife* **5** <https://doi.org/10.7554/eLife.10989>
- Kostadinov D, Beau M, Pozo MB, Häusser M (2019) **Predictive and reactive reward signals conveyed by climbing fiber inputs to cerebellar Purkinje cells** *Nat Neurosci* **22**:950–962
- Kunimatsu J, Suzuki TW, Ohmae S, Tanaka M (2018) **Different contributions of preparatory activity in the basal ganglia and cerebellum for self-timing** *Elife* **7** <https://doi.org/10.7554/eLife.35676>
- Kunimatsu J, Suzuki TW, Tanaka M (2016) **Implications of Lateral Cerebellum in Proactive Control of Saccades** *Journal of Neuroscience* **36**:7066–7074
- Lindeman S, Hong S, Kros L, Mejias JF, Romano V, Oostenveld R, Negrello M, Bosman LWJ, De Zeeuw CI (2021) **Cerebellar Purkinje cells can differentially modulate coherence between sensory and motor cortex depending on region and behavior** *Proc Natl Acad Sci U S A* **118** <https://doi.org/10.1073/pnas.2015292118>
- Mairal J, Bach F, Ponce J, Sapiro G, Jenatton R, Obozinski G, Spams: A sparse modeling software, v2. 3 (2014) **Mairal J, Bach F, Ponce J, Sapiro G, Jenatton R, Obozinski G. . Spams: A sparse modeling software, v2. 32014. . URL <http://spams-devel.gforge.inria.fr/downloads.html>.**
- Middleton FA, Strick PL (2001) **Cerebellar projections to the prefrontal cortex of the primate** *J Neurosci* **21**:700–712

- Monosov IE, Thompson KG (2009) **Frontal eye field activity enhances object identification during covert visual search** *J Neurophysiol* **102**:3656–3672
- Nicolson RI, Fawcett AJ, Dean P (2001) **Developmental dyslexia: the cerebellar deficit hypothesis** *Trends Neurosci* **24**:508–511
- Nobre AC, Gitelman DR, Dias EC, Mesulam MM (2000) **Covert Visual Spatial Orienting and Saccades: Overlapping Neural Systems** *Neuroimage* **11**:210–216
- Park IM, Meister MLR, Huk AC, Pillow JW (2014) **Encoding and decoding in parietal cortex during sensorimotor decision-making** *Nat Neurosci* **17**:1395–1403
- Paxinos G, Huang X-F, Toga AW (2000) **The Rhesus Monkey Brain in Stereotaxic Coordinates**
- Posner MI (1980) **Orienting of attention** *Q J Exp Psychol* **32**:3–25
- Prevosto V, Graf W, Ugolini G (2010) **Cerebellar inputs to intraparietal cortex areas LIP and MIP: functional frameworks for adaptive control of eye movements, reaching, and arm/eye/head movement coordination** *Cereb Cortex* **20**:214–228
- Raghavan RT, Joshua M (2017) **Dissecting patterns of preparatory activity in the frontal eye fields during pursuit target selection** *J Neurophysiol* **118**:2216–2231
- Ramnani N (2012) **Frontal lobe and posterior parietal contributions to the cortico-cerebellar system** *Cerebellum* **11**:366–383
- Rizzolatti G, Riggio L, Dascola I, Umiltà C (1987) **Reorienting attention across the horizontal and vertical meridians: evidence in favor of a premotor theory of attention** *Neuropsychologia* **25**:31–40
- Roitman JD, Shadlen MN (2002) **Response of neurons in the lateral intraparietal area during a combined visual discrimination reaction time task** *J Neurosci* **22**:9475–9489
- Romanski LM, Giguere M, Bates JF, Goldman-Rakic PS (1997) **Topographic organization of medial pulvinar connections with the prefrontal cortex in the rhesus monkey** *J Comp Neurol* **379**:313–332
- Salmi J, Pallesen KJ, Neuvonen T, Brattico E, Korvenoja A, Salonen O, Carlson S (2010) **Cognitive and motor loops of the human cerebro-cerebellar system** *J Cogn Neurosci* **22**:2663–2676
- Scerra VE, Costello MG, Salinas E, Stanford TR (2019) **All-or-None Context Dependence Delineates Limits of FEF Visual Target Selection** *Curr Biol* **29**:294–305
- Schall JD, Hanes DP (1993) **Neural basis of saccade target selection in frontal eye field during visual search** *Nature*
- Shadlen MN, Newsome WT (2001) **Neural basis of a perceptual decision in the parietal cortex (area LIP) of the rhesus monkey** *J Neurophysiol* **86**:1916–1936
- Stoodley CJ, Schmahmann JD (2010) **Evidence for topographic organization in the cerebellum of motor control versus cognitive and affective processing** *Cortex* **46**:831–844
- Strick PL, Dum RP, Fiez JA (2009) **Cerebellum and Nonmotor Function** *Annu Rev Neurosci* **32**:413–434

- Svoboda K, Li N (2018) **Neural mechanisms of movement planning: motor cortex and beyond** *Curr Opin Neurobiol* **49**:33–41
- Tanaka M (2006) **Inactivation of the central thalamus delays self-timed saccades** *Nat Neurosci* **9**:20–22
- van Es DM, van der Zwaag W, Knapen T. (2019) **Topographic Maps of Visual Space in the Human Cerebellum** *Curr Biol* **29**:1689–1694
- Voogd J, Ruigrok TJH, Cerebellum and Precerebellar Nuclei, Mai JK, Paxinos G (2012) **The Human Nervous System**
- Voogd J, Schraa-Tam CKL, van der Geest JN, De Zeeuw CI (2012) **Visuomotor cerebellum in human and nonhuman primates** *Cerebellum* **11**:392–410
- Wagner MJ, Kim TH, Kadmon J, Nguyen ND, Ganguli S, Schnitzer MJ, Luo L (2019) **Shared Cortex-Cerebellum Dynamics in the Execution and Learning of a Motor Task** *Cell* **177**:669–682
- Wang X, Novello M, Gao Z, Ruigrok TJH, De Zeeuw CI (2022) **Input and output organization of the mesodiencephalic junction for cerebro-cerebellar communication** *J Neurosci Res* **100**:620–637
- Wardak C, Olivier E, Duhamel J-R (2011) **The relationship between spatial attention and saccades in the frontoparietal network of the monkey** *Eur J Neurosci* **33**:1973–1981
- Wolpert DM, Ghahramani Z (2000) **Computational principles of movement neuroscience** *Nat Neurosci* **3**:1212–1217

Editors

Reviewing Editor

Masaki Tanaka

Hokkaido University School of Medicine, Sapporo, Japan

Senior Editor

Tirin Moore

Howard Hughes Medical Institute, Stanford University, Stanford, United States of America

Reviewer #1 (Public Review):

Summary:

Recordings were made from the dentate nucleus of two monkeys during a decision-making task. Correlates of stimulus position and stimulus information were found to varying degrees in the neuronal activities.

Strengths:

A difficult decision-making task was examined in two monkeys.

Weaknesses:

One of the monkeys did not fully learn the task. The manuscript lacked a coherent hypothesis to be tested, and no attempt was made to consider the possibility that this part of the brain may have little to do with the task that was being studied.

<https://doi.org/10.7554/eLife.99696.1.sa2>

Reviewer #2 (Public Review):

The authors trained monkeys to discriminate peripheral visual cues and associate them with planning future saccades of an indicated direction. At the same time, the authors recorded single-unit neural activity in the cerebellar dentate nucleus. They demonstrated that substantial fractions of DN cells exhibited sustained modulation of spike rates spanning task epochs and carrying information about stimulus, response, and trial outcome. Finally, tracer injections demonstrated this region of the DN projects to a large number of targets including several known to interconnect the visual attention network. The data compellingly demonstrate the authors' central claims, and the analyses are well-suited to support the conclusions. Importantly, the study demonstrates that DN cells convey many motor and nonmotor variables related to task execution, event sequencing, visual attention, and arguably decision-making/working memory.

<https://doi.org/10.7554/eLife.99696.1.sa1>

Author response:

General comment:

"This important study examined neuronal activity in the dentate nucleus of the cerebellum when monkeys performed a difficult perceptual decision-making task. The authors provide convincing evidence that the cerebellum represents sensory, motor, and behavioral outcome signals that are sent to the attentional system, but further analysis focusing on the disparity of performance between animals would improve the quality of the paper. This paper is of great general interest in that it shows the involvement of the cerebellum in cognitive processes at the neuronal level."

We thank you for these general comments, and we agree with all of them.

Public Reviews (Reviewer #1):

Summary:

Recordings were made from the dentate nucleus of two monkeys during a decision-making task. Correlates of stimulus position and stimulus information were found to varying degrees in the neuronal activities.

We agree with this summary.

Strengths:

A difficult decision-making task was examined in two monkeys.

We agree with this statement.

Weaknesses:

One of the monkeys did not fully learn the task. The manuscript lacked a coherent hypothesis to be tested, and no attempt was made to consider the possibility that this part of the brain may have little to do with the task that was being studied.

We understand these comments. It is correct that one of the monkeys did not fully learn the task, but it should be noted that both monkeys learned significantly above chance level, and we therefore find the recordings of both monkeys useful. We tested the hypothesis that neurons of the nucleus dentate can dynamically modulate their activity during a visual attention task, comprising not only sensorimotor but also cognitive attentional components. We agree that this hypothesis should be spelled out more explicitly in the introduction, which we will do in the revised version. We also appreciate the comment of this Reviewer that in our original submission we did not show our attempt to consider the possibility that this part of the brain may have little to do with the task that was being studied. We in fact did consider this possibility in that we applied muscimol to the dentate nucleus in one of the monkeys. The data of this one successful experiment show that the behaviour was reversibly affected in line with our hypothesis. Given that this only concerned one of the monkeys, we preferred not to present these data in the article. However, as the Reviewer correctly points out that this question remains hanging in the air, we will show them in our formal rebuttal letter. Please note that we decided to focus at the end of our research project on the tracing experiments, showing in both monkeys the connections of the dentate nucleus with the regions that are involved in attention. As a result, both monkeys have been sacrificed and we cannot expand upon our muscimol experiments anymore (which would have been useful indeed).

Last but not least, given the comments of the Reviewers, we will also add a Supplementary figure to Figure 2, in which we will present the data for both monkeys separately and provide our interpretation. This may help to strengthen our conclusions.

Public Reviews (Reviewer #2):

The authors trained monkeys to discriminate peripheral visual cues and associate them with planning future saccades of an indicated direction. At the same time, the authors recorded single-unit neural activity in the cerebellar dentate nucleus. They demonstrated that substantial fractions of DN cells exhibited sustained modulation of spike rates spanning task epochs and carrying information about stimulus, response, and trial outcome. Finally, tracer injections demonstrated this region of the DN projects to a large number of targets including several known to interconnect the visual attention network. The data compellingly demonstrate the authors' central claims, and the analyses are well-suited to support the conclusions. Importantly, the study demonstrates that DN cells convey many motor and nonmotor variables related to task execution, event sequencing, visual attention, and arguably decision-making/working memory.

We thank the Reviewer for this positive and constructive feedback.

<https://doi.org/10.7554/eLife.99696.1.sa0>